

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

Physics Letters B

www.elsevier.com/locate/physletb

Spontaneous spacetime reduction and unitary weak boson scattering at the LHC

Hong-Jian He^{a,b,c,*}, Zhong-Zhi Xianyu^a^a Institute of Modern Physics and Center for High Energy Physics, Tsinghua University, Beijing 100084, China^b Center for High Energy Physics, Peking University, Beijing 100871, China^c Kavli Institute for Theoretical Physics China, CAS, Beijing 100190, China

ARTICLE INFO

Article history:

Received 31 October 2012

Received in revised form 9 December 2012

Accepted 17 January 2013

Available online 4 February 2013

Editor: B. Grinstein

ABSTRACT

Theories of quantum gravity predict spacetime dimensions to become reduced at high energies, a striking phenomenon known as spontaneous dimensional reduction (SDR). It has found nice applications to astrophysics and collider phenomenologies. We construct an effective electroweak theory based on the standard model (SM) and incorporate the TeV-scale SDR, which exhibits good high energy behavior and ensures the unitarity of weak gauge boson scattering. This also provides a natural solution to the hierarchy problem in the presence of a SM-like Higgs boson. We demonstrate that this model predicts the unitary longitudinal weak boson scattering, and can be discriminated from the conventional 4d SM by the WW scattering experiments at the CERN LHC.

© 2013 Elsevier B.V. Open access under [CC BY](http://creativecommons.org/licenses/by/3.0/) license.

1. Introduction

Spontaneous dimensional reduction (SDR) [1] is a striking phenomenon, showing that the spacetime dimensions effectively equal $3 + 1$ at low energies, but get reduced toward $1 + 1$ at high energies. This is predicted by a number of quantum gravity approaches [1], including the causal dynamical triangulation, the exact renormalization group method, the loop quantum gravity, the high-temperature string theory, and Hořava–Lifshitz gravity, etc. The SDR is expected to greatly improve the ultraviolet (UV) behavior of the standard model (SM) of particle physics. This property of the quantum gravity has called phenomenological interests recently, and has found applications to astrophysics and collider phenomenologies in a different context [2]. In addition, some hints of a TeV-scale SDR have been noted [2] from the observations of alignment of high energy cosmic rays [3].

The $SU(2)_L \otimes U(1)_Y$ gauge structure of the SM is well established for electroweak forces. In the conventional SM, it is linearly realized and spontaneously broken by the Higgs mechanism [4], leading to massive weak gauge bosons (W^\pm, Z^0). A Higgs boson is predicted and is crucial for the renormalizability and unitarity of the SM. Recently, ATLAS and CMS Collaborations have found signals for a new particle with mass around 125 GeV at the LHC, which are somewhat different from the SM Higgs boson (especially in the

* Corresponding author at: Institute of Modern Physics and Center for High Energy Physics, Tsinghua University, Beijing 100084, China.

E-mail addresses: hjhe@tsinghua.edu.cn (H.-J. He), xianyuzhongzhi@gmail.com (Z.-Z. Xianyu).

diphoton mode), although still consistent with the SM expectations within about 2σ statistics [5]. Hence, the true mechanism of electroweak symmetry breaking is awaiting further explorations at the LHC, and the possible new physics beyond the SM Higgs boson is highly anticipated. Given the fact that no other new particles have been detected so far, it is tantalizing to explore alternative new physics sources at the TeV scale, beyond the conventional proposals such as extra dimensions, supersymmetry and strong dynamics at the TeV scale.

In this Letter, we will explore the quantum gravity effect of SDR at the TeV scale, and study its applications to the electroweak sector of the SM. We conjecture that the TeV-scale SDR can play a key role to unitarize weak gauge boson scattering in the theory without or with a Higgs boson. As the first example, we will show that the perturbative unitarity is maintained by the TeV-scale SDR in scenarios without a Higgs boson, where the recently observed 125 GeV boson [5] can be something else, such as a dilaton-like particle [6]. Without a Higgs boson, the SM electroweak gauge symmetry $SU(2)_L \otimes U(1)_Y$ becomes nonlinearly realized [7] and the three Goldstone bosons are converted to the longitudinal polarizations of (W^\pm, Z^0) after spontaneous symmetry breaking. Such a minimal Higgsless SM loses traditional renormalizability [7] and violates unitarity of WW scattering at TeV scale [8], so it is incomplete. We show that the TeV-scale SDR can provide a new way to unitarize the WW scattering, and will be discriminated from the SM at the LHC.

Then, we study the SM with a non-standard Higgs boson of mass around 125 GeV under the TeV-scale SDR (called the Higgsful SM). We will show that the corresponding WW scattering cross sections become unitary at TeV scale under the SDR, but exhibit

different behaviors from the conventional 4d SM. We note that different ways of unitarizing the longitudinal WW scattering around TeV scale reflect the underlying mechanisms of electroweak symmetry breaking (EWSB), and will be discriminated by the WW scattering experiments as a key task of the LHC [9].

2. The TeV scale SDR

Despite lacking a full theory of quantum gravity that could precisely describe the SDR, we are modest and approach this problem by using the *effective theory formulation* [10]. In particular, to mimic the result from the causal dynamical triangulation [11], we parameterize the spacetime dimension $n = n(\mu)$ as a smooth function of the energy scale μ (which we call the dimensional flow by following Calcagni [12]), such that $n(\mu) \rightarrow 4$ under $\mu \rightarrow 0$ in the infrared region as supported by all low energy experiments, and $n(\mu) \rightarrow 2$ at a certain UV scale Λ_{UV} . We can make a simple choice for the dimensional flow,

$$n(\mu) = 4 - 2 \left(\frac{\mu}{\Lambda_{UV}} \right)^\gamma, \quad (\mu \leq \Lambda_{UV}), \quad (1)$$

where the index $\gamma > 1$ is a model-dependent parameter, determined by the nonperturbative dynamics of quantum gravity. As simple realizations we may set $\gamma = 2$ or 1.5. Before finding a unique full theory of quantum gravity, other variations of (1) are possible [11,12], but this will not affect the main physics features of the present analysis. An easy choice for Λ_{UV} would be the Planck scale. But it is a very interesting and intriguing possibility that the nonperturbative dynamics of quantum gravity drives Λ_{UV} down to \mathcal{O} (TeV) [2]. If this happens, a number of difficulties associated with the EWSB and W/Z mass-generations in the SM can be resolved without introducing additional *ad hoc* hypothetical dynamics.

When the quantum gravity effects show up at the TeV scale, they will induce effective operators causing sizable anomalous Higgs couplings to WW (ZZ) and fermions in the low energy effective theory. This will violate perturbative unitarity at TeV energy scale in the conventional 4d setup [13,14]. However, we show that under the TeV-scale SDR, the weak boson scattering amplitudes will still be unitarized through the reduction of spacetime dimensions. Furthermore, the presence of the TeV-scale SDR will also provide a natural solution to the hierarchy problem since the quadratically divergent radiative corrections to Higgs mass is rendered to be logarithmic in $n = 2$ spacetime and thus harmless.

3. The standard model with SDR

As an effective theory description of the SDR, we encode the information of dimensional flow $n = n(\mu)$ into the measure of spacetime integral $d\rho$, and replace all integral measure d^4x in the action functional by $d\rho$. A rigorous mathematical construction of $d\rho$ is given by Ref. [12], but the detail is not needed here. All we need to know is that the mass-dimension of this measure is $[d\rho] = -n$, where $n = n(\mu)$ is the dimensional flow in (1). It is enough to define the measure $d\rho$ formally by $d^n x$, with n a scale-dependent quantity. Thus, we can write the action of the theory, $S = \int d^n x \mathcal{L} = \int d^n x (\mathcal{L}_G + \mathcal{L}_F)$, where \mathcal{L}_G and \mathcal{L}_F are the gauge and fermion parts of the SM Lagrangian. We will focus on the gauge sector for the current study. We first consider the gauge Lagrangian with Higgs boson removed,

$$\begin{aligned} \mathcal{L}_G = & -\frac{1}{4} W_{\mu\nu}^a W^{\mu\nu a} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} + M_W^2 W_\mu^+ W^{-\mu} \\ & + \frac{1}{2 \cos^2 \theta_w} M_W^2 Z_\mu Z^\mu, \end{aligned} \quad (2)$$

where $W_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g \epsilon^{abc} W_\mu^b W_\nu^c$ and $B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$. In the above, $\theta_w = \arctan(g'/g)$ represents the weak mixing angle and connects the gauge-eigenbasis (W_μ^3, B_μ) to the mass-eigenbasis (Z_μ^0, A_μ) . Eq. (2) contains W/Z mass-terms in unitary gauge and can be made gauge-invariant in the nonlinear realization of $SU_L(2) \otimes U(1)_Y$ gauge symmetry,

$$\mathcal{L}_\Sigma = \frac{1}{4} v^2 \text{tr}[(D^\mu \Sigma)^\dagger (D_\mu \Sigma)], \quad (3)$$

where $D_\mu \Sigma = \partial_\mu \Sigma + \frac{1}{2} g W_\mu^a \tau^a \Sigma - \frac{1}{2} g' B_\mu \Sigma \tau^3$, $\Sigma = \exp[i\tau^a \pi^a / v]$ with $\{\pi^a\}$ the Goldstone bosons. Eq. (3) gives $M_W = \frac{1}{2} g v$, where the parameter v will be fixed by the low energy Fermi constant $G_F = (\sqrt{2} v^2)^{-1}$. The Lagrangian (2) derives directly from the SM in unitary gauge after removing the Higgs boson; while Eq. (3) is just the lowest order electroweak chiral Lagrangian of the SM [7].

We can further embed the Higgs boson as a singlet scalar h^0 in this formulation by extending the Lagrangian (3) as in [13,14],

$$\begin{aligned} \mathcal{L}_H = & \frac{1}{4} (v^2 + 2\kappa v h + \kappa' h^2) \text{tr}[(D^\mu \Sigma)^\dagger (D_\mu \Sigma)] \\ & + \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{1}{2} M_h^2 h^2 - \frac{\lambda_3}{3!} v h^3 + \frac{\lambda_4}{4!} h^4, \end{aligned} \quad (4)$$

where $\Delta\kappa \equiv \kappa - 1$ and $\Delta\kappa' \equiv \kappa' - 1$ denote the anomalous gauge couplings of the Higgs boson h^0 with WW and ZZ . The conventional 4d SM is just a special case, $\Delta\kappa = \Delta\kappa' = 0$, and $\lambda_3 = \lambda_4 = \lambda_0 = 3M_h^2/v^2$, in the general effective Lagrangian (4). For nonzero anomalous couplings $\Delta\kappa, \Delta\kappa' \neq 0$ and/or $\lambda_3, \lambda_4 \neq \lambda_0$, as induced by the effects of TeV-scale SDR, the scalar field h^0 becomes a non-standard Higgs boson. We will study how to discriminate such as non-SM Higgs particle from the conventional 4d SM in Section 5.

We also note that our Lagrangian \mathcal{L} is manifestly Lorentz invariant and thus all particles' dispersion relations remain unchanged, as in [12]. This is because our formulation is based on the framework of [12], where it is shown that the action can be constructed in such a way that the Lagrangian density \mathcal{L} lies in 3+1 dimensional spacetime and respects the (3+1)d Poincaré symmetry, while the effect of SDR is fully governed by a properly defined integral measure $d\rho$. In such a scenario, scalar, spinor and vector fields are linear representations of (3+1)d Lorentz group $SO(3,1)$ (up to a gauge transformation for gauge fields). Practically, this is similar to the conventional dimensional reduction regularization method [15], which maintains the 4d Lorentz symmetry and continues physics to $n < 4$. Thus, our model is free from Lorentz-violation constraints in the cosmic ray observations and collider experiments at the tree level. In addition, our present study focuses on longitudinal weak boson scattering, and their amplitudes are equivalent to that of the corresponding Goldstone bosons at high energies according to the equivalence theorem [16]. The amplitudes of scalar Goldstone bosons are cleanly defined in general dimension n . Manipulations of scalar fields do not involve contracting or counting Lorentz indices, and thus do not rely on details of realizing the SDR.

In general n -dimensional spacetime, the action functional should remain dimensionless. Hence the Lagrangian has the mass-dimension $[\mathcal{L}] = n$, and the gauge coupling g has mass-dimension $[g] = (4-n)/2$. We can always define a new dimensionless coupling \tilde{g} and transfer the mass-dimension of g to another mass-parameter. Since the gauge coupling g in (2) becomes super-renormalizable for $n < 4$ and thus insensitive to the UV, it is natural to scale g by the W mass M_W [which is the only dimensional parameter of the Lagrangian (2) in 4d],

$$g = \tilde{g} M_W^{(4-n)/2}, \quad (5)$$

with \tilde{g} being dimensionless. The value of \tilde{g} is given by that of g at $n = 4$. We will concentrate on the tree-level analysis in this study, so \tilde{g} is a scale-independent constant.

In fact, the scaling (5) is well justified for more reasons. We may easily wonder why we could not use the UV cutoff Λ_{UV} in the scaling of g as a replacement of the infrared mass-parameter M_W of the theory. This is because in spacetime dimension $n < 4$, the gauge coupling g is super-renormalizable with positive mass-dimension $[g] = (4 - n)/2 > 0$. Such a super-renormalizable coupling must be insensitive to the UV cutoff of the theory, contrary to a non-renormalizable coupling with negative mass-dimension and thus naturally suppressed by negative powers of the UV cutoff Λ_{UV} (e.g., in $n > 4$ or in association with certain higher-dimensional operators). It is easy to imagine that for a super-renormalizable theory in dimension $n < 4$, if its coupling g were scaled as $g = \tilde{g}\Lambda_{UV}^{(4-n)/2}$, it would even make tree-level amplitude UV divergent and blow up as $\Lambda_{UV} \rightarrow \infty$; this is clearly not true. On the other hand, it is well known that a non-renormalizable coupling g with negative mass-dimension $[g] \equiv -p < 0$ should be scaled as $g = \tilde{g}/\Lambda_{UV}^p$, and thus the tree-level amplitude naturally approaches zero when $\Lambda_{UV} \rightarrow \infty$, as expected.

Since spacetime dimension flows to $n = 2$ in the UV limit, we observe that the theory (2) is well-behaved at high energies. This is because all gauge couplings of the Lagrangian (2) in $n < 4$ dimensions become super-renormalizable, and the gauge boson propagators scale as $1/p^2$ in high momentum limit under the R_ξ gauge-fixing. So we only concern about gauge boson mass-terms in (2) or (3), which is the origin of nonrenormalizability and unitarity violation in 4d. But, in our construction the spacetime dimension flows to $n = 2$ in high energy limit where (3) just describes a 2d gauged nonlinear sigma model and is renormalizable, as is well known.

We also note that in $n < 4$ dimensions gauge bosons can acquire masses via new mechanisms other than the Higgs mechanism. For instance, in the 2d Schwinger model [17], radiative corrections to the vacuum polarization from a massless-fermion loop generate a nonzero photon mass, $m_\gamma = \frac{e}{\sqrt{\pi}}$. Also, the 3d Chern-Simons term induces a topological mass for the corresponding gauge field [18], $m_{CS} = \kappa e^2$. Hence, it is natural to have an explicit mass-term of vector boson in a lower dimensional field theory. We will further demonstrate below that such a mass-term is indeed harmless in a Higgsless SM with the TeV-scale SDR, and the unitarity of high energy longitudinal WW scattering is ensured.

4. Longitudinal weak boson scattering under SDR

As a simple illustration of the unitarization mechanism of longitudinal WW scattering under the SDR, we first present the analysis in the SM without Higgs boson (called the Higgsless SM and denoted by HLSM-SDR). It is noted that such a scenario is still consistent with the current LHC data since the 125 GeV new boson can be something else, such a dilaton-like particle [6]. The effect of SDR is most clearly seen in this case. After this, we will further extend this mechanism to the Higgsful SM under the SDR (including a Higgs boson and called the HFSM-SDR), in the next section.

The minimal 4d Higgsless SM violates unitarity at TeV scale, because the SM Higgs boson plays the key role to unitarize the bad high energy behaviors of the longitudinal WW scattering. For instance, without Higgs boson, the amplitude of $W_L^+ W_L^- \rightarrow Z_L^0 Z_L^0$ has non-canceled E^2 term,

$$\mathcal{T}_{HL} = g^2 E_{cm}^2 / (4M_W^2) + \mathcal{O}(E_{cm}^0), \quad (6)$$

where E_{cm} is the c.m. energy. This bad E^2 behavior leads to unitarity violation at TeV scale. In contrast, for the conventional 4d SM, this E^2 term is exactly canceled by the contribution of the

s -channel Higgs-exchange, which is the key to ensure the SM unitarity.

In lower dimensions, the longitudinal amplitudes remain the same as in 4d. But, we observe that the form of partial wave expansion changes, due to the phase-space reduction for final state. Hence, the E^2 -cancellation described above is no longer essential for ensuring the unitarity. This is an essential feature of the unitarization mechanism through SDR, i.e., the WW scattering amplitudes remain unitary at high energies under SDR, even without a Higgs boson.

To be explicit, we recall that unitarity condition for S -matrix arises from probability conservation, $SS^\dagger = S^\dagger S = 1$. This leads to $\mathcal{T}^\dagger \mathcal{T} = 2\Im \mathcal{T}$, where \mathcal{T} is defined via $S = 1 + i\mathcal{T}$, and is related to the amplitude \mathcal{T} via $\mathcal{T} = (2\pi)^n \delta^n(p_f - p_i) \mathcal{T}$ with p_i (p_f) the total momentum of the initial (final) state. For $2 \rightarrow 2$ scattering, \mathcal{T} depends only on the c.m. energy E_{cm} and scattering angle θ . Thus, in this case we can always expand $\mathcal{T}(E_{cm}, \theta)$ in terms of partial waves $a_\ell(E_{cm})$ for $n > 3$ dimensions,

$$\begin{aligned} \mathcal{T} &= \lambda_n E_{cm}^{4-n} \sum_\ell \frac{1}{N_\ell^\nu} C_\ell^\nu(1) C_\ell^\nu(\cos\theta) a_\ell, \\ a_\ell &= \frac{E_{cm}^{n-4}}{\lambda_n C_\ell^\nu(1)} \int_0^\pi d\theta \sin^{n-3} \theta C_\ell^\nu(\cos\theta) \mathcal{T}, \end{aligned} \quad (7)$$

with

$$\begin{aligned} \lambda_n &= 2(16\pi)^{n/2-1} \Gamma\left(\frac{n}{2} - 1\right), \quad \nu = \frac{1}{2}(n - 3), \\ N_\ell^\nu &= \frac{\pi \Gamma(\ell + 2\nu)}{2^{2\nu-1} \ell! (\ell + \nu) \Gamma^2(\nu)}, \end{aligned}$$

and $C_\ell^\nu(x)$ is the Gegenbauer polynomial of order ν and degree ℓ . This partial wave expansion holds for $n > 3$ because the eigenfunctions of rotation generators (namely the Gegenbauer function) are not well defined below $n = 3$. The appearance of factor E_{cm}^{4-n} in the expansion of \mathcal{T} is expected, since the S -matrix of $2 \rightarrow 2$ scattering has a mass-dimension $4 - n$ in n dimensions, and the partial wave amplitude a_ℓ is dimensionless by definition. Then, we can derive unitarity conditions for the elastic and inelastic partial waves, $|\Re a_\ell^{\text{el}}| \leq \frac{\rho_e}{2}$, $|a_\ell^{\text{el}}| \leq \rho_e$, and $|a_\ell^{\text{inel}}| \leq \sqrt{\rho_i \rho_e}/2$, where ρ_e (ρ_i) is a symmetry factor of final state in $2 \rightarrow 2$ elastic (inelastic) scattering, and equals $1!$ ($2!$) for the final state particles being nonidentical (identical) [19].

With these, we perform the coupled channel analysis for electrically neutral channels. There are two relevant initial/final states, $|W_L^+ W_L^- \rangle$ and $\frac{1}{\sqrt{2}} |Z_L^0 Z_L^0 \rangle$, and the corresponding amplitudes form a 2×2 matrix,

$$\mathcal{T}_{\text{coup}} = \frac{g^2 E_{cm}^2}{8M_W^2} \begin{pmatrix} 1 + \cos\theta & \sqrt{2} \\ \sqrt{2} & 0 \end{pmatrix}. \quad (8)$$

Thus, we derive the s -wave amplitude from (8) in n -dimensions and extract the maximal eigenvalue,

$$|a_0^{\text{max}}| = \frac{\tilde{g}^2}{2^{n+1} \pi^{(n-3)/2} \Gamma(\frac{n-1}{2})} \left(\frac{E_{cm}}{2M_W} \right)^{n-2}. \quad (9)$$

Although the partial wave expansion (7) holds for $n > 3$, we can make analytical continuation of (9) as a function of dimension n to the full range $2 \leq n \leq 4$. Here, we perform the analytic continuation on the complex plane of spacetime dimension n , while the one-to-one mapping between n and μ is only defined within the real interval $2 \leq n \leq 4$.

In Fig. 1, we present the unitarity constraint for the standard model without a Higgs boson, under Eq. (1) with $\gamma = 1.5$, where




Fig. 1. Partial wave amplitude of coupled channel scattering versus c.m. energy E_{cm} . Predictions of the HLSM–SDR are shown by (red, purple, blue) curves from bottom to top, for $\Lambda_{\text{UV}} = (4, 5, 6)$ TeV. For comparison, the amplitudes for the 4d Higgsless SM and for the conventional 4d SM (with a 600 GeV Higgs boson) are depicted by the black curves. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this Letter.)

we have varied the transition scale $\Lambda_{\text{UV}} = (4, 5, 6)$ TeV. The shaded yellow region is excluded by the unitarity bound $|a_0^{\text{max}}| \leq 1$. The s -waves of HLSM–SDR always have a rather broad “lump” around 1.5–5 TeV and then fall off quickly, exhibiting desired unitary high energy behaviors. For comparison, we also show the results of the 4d SM with a 600 GeV Higgs boson, and the naive 4d Higgsless SM which breaks unitarity at $E_{\text{cm}} \simeq 1.74$ TeV.

Next, we compute the cross section for $W_L^+ W_L^- \rightarrow Z_L^0 Z_L^0$ and $W_L^+ W_L^+ \rightarrow W_L^+ W_L^+$, as shown in Fig. 2. The 4d unitarity condition for inelastic cross sections is, $\sigma_{\text{inel}} \leq 4\pi \rho_e E_{\text{cm}}^{-2}$. We derive the generalized form in n -dimensions,

$$\sigma_{\text{inel}} \leq \frac{\lambda_n \rho_e}{4N_0^v E_{\text{cm}}^{n-2}}, \quad (10)$$

where λ_n and N_0^v are defined below (7). Fig. 2 shows how the SDR works as a new mechanism to successfully unitarize the high energy behaviors of cross sections without invoking extra hypothesized particle (such as the SM Higgs boson). Furthermore, the new

predictions of the HLSM–SDR are universal and show up in all WW scattering channels. This is an essential feature of our model and will be crucial for discriminating the HLSM–SDR from all other models of the EWSB at the LHC.

Here we note that in n -dimensions the cross section σ has its mass-dimensions equal $[\sigma] = 2 - n$, while the experimentally measured cross section σ_{exp} always has mass-dimension -2 , as the detectors record events in 4d. So we need to convert the theory cross section σ under the SDR to σ_{exp} , where the extra mass-dimensions of σ should be scaled by the involved energy scale E_{cm} of the reaction, $\sigma_{\text{exp}} = \sigma E_{\text{cm}}^{n-4}$.

As a final remark, it was found [19] in 4d that varying the phase space may strongly alter the unitarity limit. Ref. [19] observed that the *enlarged phase space* of $2 \rightarrow \mathcal{N}$ scattering (due to properly increasing the number \mathcal{N} of gauge bosons in the final state) will enhance the cross section and result in a new class of much stronger unitarity bounds for all light SM fermions. Interestingly, the current study just shows the other way around: for $2 \rightarrow 2$ scattering, *reduction of the phase space* of final states from decreasing the spacetime dimension n can significantly reduce the partial wave amplitudes and cross sections, leading to the unitarity restoration.

5. Weak boson scattering in Higgsful SM with SDR

In this section, we extend our mechanism in Section 4 to the Higgsful SM with SDR (HFSM–SDR). In this case, the quantum-gravity-induced SDR at TeV scales provides a natural solution to the hierarchy problem that plagues the Higgs boson in the conventional 4d SM. For TeV-scale SDR, it is expected that new physics effects induced by the quantum gravity will show up in the low energy effective theory. So, the Higgs boson can have anomalous couplings with WW and ZZ gauge bosons, and thus behaves as non-SM-like.

In unitary gauge, we can write down the leading anomalous gauge interactions of the Higgs boson [13,14] from the effective Lagrangian (4),

$$\left(\Delta\kappa v h + \frac{1}{2} \Delta\kappa' h^2 \right) \left[\frac{2M_W^2}{v^2} W_\mu^+ W^{-\mu} + \frac{M_Z^2}{v^2} Z_\mu Z^\mu \right], \quad (11)$$




Fig. 2. Cross sections versus center-of-mass energy E_{cm} for processes (a) $W_L^+ W_L^- \rightarrow Z_L^0 Z_L^0$ and (b) $W_L^+ W_L^+ \rightarrow W_L^+ W_L^+$. In each plot, predictions of the HLSM–SDR are shown by (red, purple, blue) curves, for $\Lambda_{\text{UV}} = (4, 5, 6)$ TeV. As comparison, results of the conventional 4d SM with a light (heavy) Higgs boson of mass $M_h = 125$ GeV (600 GeV) are depicted by black dashed-curves; the result of the 4d Higgsless SM is given by black solid-curve. Shaded regions in yellow and light-blue represent unitarity violation in 4d and in the HLSM–SDR, respectively. The three blue dashed-lines, from bottom to top, show the unitarity bounds for $\Lambda_{\text{UV}} = (4, 5, 6)$ TeV. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this Letter.)




Fig. 3. Cross sections of $W_L^+ W_L^- \rightarrow Z_L^0 Z_L^0$ versus center-of-mass energy E_{cm} . In each plot, predictions of the HFSM-SDR with Higgs mass $M_h = 125$ GeV [5] and anomalous couplings $\Delta\kappa = +0.3$ [plot (a)] and $\Delta\kappa = -0.3$ [plot (b)] are shown by (red, purple, blue) curves, for $\Lambda_{\text{UV}} = (4, 5, 6)$ TeV. As comparison, results of the conventional 4d-SM with $M_h = 125$ GeV (labeled by “SM”) and the 4d-SM with the same anomalous coupling (labeled by “SM + AC”) are depicted by black curves. Shaded regions are the same as in Fig. 2. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this Letter.)

where the anomalous couplings $\Delta\kappa, \Delta\kappa' \neq 0$ represent new physics. Besides the hierarchy problem, the conventional 4d SM also suffers constraints from the Higgs vacuum instability and the triviality of Higgs self-coupling. If such a 4d SM would be valid up to Planck scale, then the SM Higgs boson mass is bounded within the range [20], $133\text{GeV} \lesssim M_h \lesssim 180$ GeV. Hence, a Higgs mass outside this window will indicate a non-standard Higgs boson in association with *new physics*. The Higgs boson in our present model under the TeV-scale SDR has anomalous couplings induced from quantum gravity and thus behaves as non-SM-like. With the recent LHC data [5], model-independent fits already put some interesting constraints on the Higgs anomalous couplings. Using the fitting result of [21], we find that for $M_h = 125$ GeV, the $\Delta\kappa$ in (11) is bounded within the range, $\Delta\kappa = 0.2_{-0.5}^{+0.4}$.

In the conventional 4d SM with (11), it was found [14] that the WW scattering has non-canceled large E^2 behavior and will eventually violate unitarity at TeV scale. But in our new model, the TeV-scale SDR can always unitarize WW scattering and predicts different behaviors for cross sections, as shown in Fig. 3 for $\gamma = 1.5$. In Fig. 3, we study WW scattering process for probing a non-SM Higgs boson with mass $M_h = 125$ GeV [5] and anomalous coupling $\Delta\kappa = \pm 0.3$. We find that the cross sections under SDR-unitarization (middle colored curves) have sizable excesses above the 4d SM with a 125 GeV Higgs boson ($\kappa = 0$, flat black curve). Then they fall off in the 2–4 TeV region, consistent with the corresponding unitarity limits. Fig. 3 also shows that for the usual non-unitarized 4d SM with nonzero anomalous coupling $\kappa \neq 0$, the cross section (upper black curve) rapidly increases and eventually violate unitarity around $E_{\text{cm}} = 2$ TeV for this scattering channel.

Before concluding this section, we also clarify the validity range of our effective theory of the SDR. This validity range lies between the WW (ZZ) threshold (around 160–180 GeV) and the UV-cutoff $\Lambda_{\text{UV}} = \mathcal{O}(5)$ TeV. It is clearly shown in our Fig. 2 and Fig. 3, where the relevant scattering energy E_{cm} (for our model to be discriminated from the 4d-SM and 4d-HLSM at the LHC) is always within 0.2–3 TeV, which is *significantly below* $4 \text{ TeV} \leq \Lambda_{\text{UV}}$. Moreover, within this energy region 0.2–3 TeV (relevant to the LHC test), we can explicitly derive the dimensional flow from Eq. (1), $n \simeq 3.98 - 3.07$ (under the typical input of $\Lambda_{\text{UV}} = 5$ TeV and $\gamma = 1.5$), which is *significantly above*

$n = 2$.¹ This clearly shows that for our effective theory study we do not need to invoke any detailed UV dynamics at or above Λ_{UV} .

6. Conclusions

We have studied the exciting possibility for the onset of the SDR at TeV scale. We demonstrated that the TeV-scale SDR can play a key role to unitarize longitudinal weak boson scattering. We have constructed an effective theory of the SM under the SDR, either without a Higgs boson or with a light non-standard Higgs boson.

In the first construction, it nonlinearly realizes the electroweak gauge symmetry and its spontaneous breaking. The model becomes manifestly renormalizable at high energies by power counting. We found that the non-canceled E^2 contributions to the WW scattering are unitarized by the SDR at TeV scales (Fig. 1), and the scattering cross sections exhibit different behaviors (Fig. 2). This will be probed at the LHC. Here the recent observation of a 125 GeV boson at the LHC (8 TeV) could be something else, such as a dilaton-like particle [6].

For the second construction of the Higgsful SM with SDR, we studied the WW scattering with a light non-standard Higgs boson of mass 125 GeV. It has effective anomalous couplings with gauge bosons as induced from the TeV-scale quantum gravity effects [cf. Eq. (11)]. Fig. 3 showed that the $W_L^+ W_L^- \rightarrow Z_L^0 Z_L^0$ cross section with anomalous Higgs couplings has distinctive invariant-mass distributions from the naive SM Higgs boson over the 0.2–3 TeV energy regions. This will be definitively probed by the next LHC runs at 14 TeV collision energy with higher luminosity.

In passing, we note that the unitarity of WW scattering in generic 4d Technicolor theories was recently studied in Ref. [22]. For future works, it is useful to further develop a method for quantizing field theories with SDR and compute the sub-leading effect of loop corrections in fractional spacetime [12], which is expected to have better UV behavior than the usual 4d SM and

¹ Note that only at $n = 2$ and its vicinity, we have $E_{\text{cm}} \rightarrow \Lambda_{\text{UV}}$ and thus our effective theory should be replaced by a full theory of quantum gravity.

thus agree better with the precision data. This is fully beyond the current scope and will be further explored in future works. A systematical expansion of our study in the present Letter is given elsewhere [23].

Note added

As the final remark, our effective theory construction is also partly motivated by the asymptotic safety (AS) scenario of quantum general relativity (QGR) à la Weinberg [24,25]. In the AS scenario, the theory is originally defined in $(3+1)d$, while solving the exact renormalization group equation of QGR points to nontrivial UV fixed point, under which the graviton two-point function exhibits effective two-dimensional UV behavior [25]. Here, the SDR is reflected in anomalous scalings of the fields, as well as physical variables like the spacetime curvature. Such anomalous scalings share the similarity with our effective theory construction, while the field contents are still defined in $(3+1)d$ and respect the $(3+1)d$ Lorentz symmetry. Our effective theory is a simplified formulation at low energy, so it does not rely on any detailed UV dynamics of the AS scenario. It is interesting to further study the quantitative connection between the SDR and the AS scenario. We also note that the Horava–Lifshitz model [26] of quantum gravity can provide a concrete field-theoretical realization of SDR with UV-completion, which has relatively tractable Lagrangian. Thus, the various scaling properties in our effective theory are expected to arise from the formulation of the Horava–Lifshitz model. We will consider these two interesting scenarios for future works.

Acknowledgements

We are grateful to Gianluca Calcagni, Steven Carlip and Dejan Stojkovic for discussing the spontaneous dimensional reduction, to Daniel Litim for discussing the asymptotic safety, and to Petr Horava for discussing the Horava–Lifshitz gravity. We thank Francesco Sannino and Chris Quigg for discussions during their visits to Tsinghua HEP Center. This work was supported by National NSF of China (under grants 11275101, 11135003, 10625522, 10635030) and National Basic Research Program (under grant 2010CB833000).

References

- [1] For reviews, S. Carlip, arXiv:1009.1136 [gr-qc]; S. Carlip, arXiv:0909.3329 [gr-qc], and references therein.
- [2] Applications of TeV scale vanishing dimensions to certain astrophysics and collider phenomenology were recently studied in a different context, J. Mureika, D. Stojkovic, Phys. Rev. Lett. 106 (2011) 101101, arXiv:1102.3434; L. Anchordoqui, D.C. Dai, M. Fairbairn, G. Landsberg, D. Stojkovic, Mod. Phys. Lett. A 27 (2012) 1250021, arXiv:1003.5914 [hep-ph]; L. Anchordoqui, D.C. Dai, H. Goldberg, G. Landsberg, G. Shaughnessy, D. Stojkovic, T.J. Weiler, Phys. Rev. D 83 (2011) 114046, arXiv:1012.1870 [hep-ph].
- [3] T. Antoni, et al., KASCADE Collaboration, Phys. Rev. D 71 (2005) 072002, arXiv:hep-ph/0503218; R.A. Mukhamedshin, JHEP 0505 (2005) 049; L.T. Baradzei, et al., Pamir Collaboration, Izv. Akad. Nauk SSSR, Ser. Fiz. 50 (1986) 2125, Bull. Acad. Sci. USSR, Phys. Ser. 50 (11) (1986) 46.
- [4] F. Englert, R. Brout, Phys. Rev. Lett. 13 (1964) 321; P.W. Higgs, Phys. Lett. 12 (1964) 132; P.W. Higgs, Phys. Rev. Lett. 13 (1964) 508; P.W. Higgs, Phys. Rev. 145 (1966) 1156; G.S. Guralnik, C.R. Hagen, T.W. Kibble, Phys. Rev. Lett. 13 (1964) 585.
- [5] G. Aad, et al., ATLAS Collaboration, Phys. Lett. B 716 (2012) 1, arXiv:1207.7214 [hep-ex]; S. Chatrchyan, et al., CMS Collaboration, Phys. Lett. B 716 (2012) 30, arXiv:1207.7235 [hep-ex].
- [6] For recent studies, S. Matsuzaki, K. Yamawaki, Phys. Rev. D 86 (2012) 035025, arXiv:1206.6703; S. Matsuzaki, K. Yamawaki, Phys. Rev. D 86 (2012) 115004, arXiv:1209.2017; arXiv:1207.5911; B. Bellazzini, C. Csaki, J. Hubisz, J. Serra, J. Terning, arXiv:1209.3299, and references therein.
- [7] T. Appelquist, C. Bernard, Phys. Rev. D 22 (1980) 200.
- [8] J.M. Cornwall, D.N. Levin, G. Tiktopoulos, Phys. Rev. Lett. 30 (1973) 1268; B.W. Lee, C. Quigg, H.B. Thacker, Phys. Rev. Lett. 38 (1977) 883; B.W. Lee, C. Quigg, H.B. Thacker, Phys. Rev. D 16 (1977) 1519.
- [9] For reviews, M.S. Chanowitz, arXiv:hep-ph/9812215; M.S. Chanowitz, Czech. J. Phys. 55 (2005) B45, arXiv:hep-ph/0412203, and references therein.
- [10] For a recent review of the concept of effective field theory, S. Weinberg, PoS CD 09 (2009) 001, arXiv:0908.1964 [hep-th].
- [11] J. Ambjørn, J. Jurkiewicz, R. Loll, Phys. Rev. Lett. 95 (2005) 171301, arXiv:hep-th/0505113; J. Ambjørn, J. Jurkiewicz, R. Loll, Phys. Rev. D 72 (2005) 064014, arXiv:hep-th/0505154.
- [12] G. Calcagni, Phys. Rev. Lett. 104 (2010) 251301, arXiv:0912.3142; G. Calcagni, JHEP 1003 (2010) 120, arXiv:1001.0571; G. Calcagni, JHEP 1201 (2012) 065, arXiv:1107.5041; G. Calcagni, AIP Conf. Proc. 1483 (2012) 31, arXiv:1209.1110 [hep-th], and references therein.
- [13] R.S. Chivukula, V. Koulovassilopoulos, Phys. Lett. B 309 (1993) 371, arXiv:hep-ph/9304293.
- [14] H.J. He, Y.P. Kuang, C.P. Yuan, B. Zhang, Phys. Lett. B 554 (2003) 64, arXiv:hep-ph/0211229; H.J. He, Y.P. Kuang, C.P. Yuan, B. Zhang, in: The Proceedings of the “Workshop on Physics at TeV Colliders, May 26–June 6, 2003, Les Houches, France, arXiv:hep-ph/0401209.
- [15] W. Siegel, Phys. Lett. B 84 (1979) 193.
- [16] For a comprehensive review on this subject, H.J. He, Y.P. Kuang, C.P. Yuan, DESY-97-056, arXiv:hep-ph/9704276, and references therein.
- [17] J. Schwinger, Phys. Rev. 128 (1962) 2425.
- [18] S. Deser, R. Jackiw, S. Templeton, Ann. Phys. 140 (1982) 372.
- [19] D.A. Dicus, H.J. He, Phys. Rev. D 71 (2005) 093009, arXiv:hep-ph/0409131; D.A. Dicus, H.J. He, Phys. Rev. Lett. 94 (2005) 221802, arXiv:hep-ph/0502178.
- [20] For a review, C. Quigg, arXiv:0905.3187 [hep-ph], and references therein.
- [21] E.g., D. Carmi, A. Falkowski, E. Kuflik, T. Volansky, J. Zupan, JHEP 1210 (2012) 196, arXiv:1207.1718 [hep-ph], and references therein.
- [22] R. Foadi, F. Sannino, Phys. Rev. D 78 (2008) 037701, arXiv:0801.0663 [hep-ph]; R. Foadi, M. Jarvinen, F. Sannino, Phys. Rev. D 79 (2009) 035010, arXiv:0811.3719 [hep-ph].
- [23] Hong-jian He, Zhong-Zhi Xianyu, arXiv:1112.1028 [hep-ph].
- [24] S. Weinberg, in: S.W. Hawking, W. Israel (Eds.), General Relativity: An Einstein Centenary Survey, Cambridge University Press, 1979, p. 790.
- [25] For a recent review, D.F. Litim, Phil. Trans. Roy. Soc. Lond. A 369 (2011) 2759, arXiv:1102.4624 [hep-th], and references therein.
- [26] P. Horava, Phys. Rev. Lett. 102 (2009) 161301; P. Horava, Phys. Rev. D 79 (2009) 084008.