

RESEARCH

Open Access

Existence results for fractional q -difference equations with nonlocal q -integral boundary conditions

Yulin Zhao^{1*}, Haibo Chen² and Qiming Zhang¹

*Correspondence:

zhaoylch@sina.com

¹School of Science, Hunan University of Technology, Zhuzhou, Hunan 412007, China
Full list of author information is available at the end of the article**Abstract**

In this paper, we discuss the existence of positive solutions for nonlocal q -integral boundary value problems of fractional q -difference equations. By applying the generalized Banach contraction principle, the monotone iterative method, and Krasnoselskii's fixed point theorem, some existence results of positive solutions are obtained. In addition, some examples to illustrate our results are given.

MSC: 39A13; 34B18; 34A08**Keywords:** fractional q -difference equation; nonlocal boundary value problems; existence; positive solutions

1 Introduction

Studies on q -difference equations appeared already at the beginning of the twentieth century in intensive works especially by Jackson [1], Carmichael [2] and other authors such as Poincare, Picard, Ramanujan. Up to date, q -difference equations have evolved into a multidisciplinary subject; for example, see [3–6] and the references therein. For some recent work on q -difference equations, we refer the reader to the papers [7–20], and basic definitions and properties of q -difference calculus can be found in the book [21]. On the other hand, fractional differential equations have gained importance due to their numerous applications in many fields of science and engineering including fluid flow, rheology, diffusive transport akin to diffusion, electrical networks, probability, *etc.* For details, see [22, 23]. Many researchers studied the existence of solutions to fractional boundary value problems; see, for example, [24–33] and the references therein.

The fractional q -difference calculus had its origin in the works by Al-Salam [34] and Agarwal [35]. More recently, perhaps due to the explosion in research within the fractional differential calculus setting, new developments in this theory of fractional q -difference calculus were made, specifically, q -analogues of the integral and differential fractional operators properties such as the Mittag-Leffler function, the q -Laplace transform, and q -Taylor's formula [12, 21, 36, 37], just to mention some.

However, the theory of boundary value problems for nonlinear q -difference equations is still in the initial stage and many aspects of this theory need to be explored. Recently, there have been some paper considering the existence of solutions to boundary value problems of fractional q -difference equations, for example, [10, 16–20, 38] and the references therein.

In [16], Ferreira considered the Dirichlet type nonlinear q -difference boundary value problem

$$\begin{cases} D_q^\alpha u(t) + f(u(t)) = 0, & 0 < t < 1, 1 < \alpha \leq 2, \\ u(0) = u(1) = 0. \end{cases}$$

By applying a fixed point theorem in cones, sufficient conditions for the existence of non-trivial solutions were enunciated.

In [20], Graef and Kong investigated the boundary value problem with fractional q -derivatives

$$\begin{cases} (D_q^\alpha u)(t) + f(t, u(t)) = 0, & 0 < t < 1, n - 1 < \alpha \leq n, n \in \mathbf{N}, \\ (D_q^i u)(0) = 0, & i = 0, \dots, n - 2, \quad bD_q u(1) = \sum_{j=1}^m a_j D_q u(t_j) + \lambda, \end{cases}$$

where $\lambda \geq 0$ is a parameter, and the uniqueness, existence, and nonexistence of positive solutions are considered in terms of different ranges of λ .

Furthermore, Ahmad, Ntouyas, and Purnaras [10] studied the following nonlinear fractional q -difference equation with nonlocal boundary conditions:

$$\begin{cases} ({}_C D_q^\alpha u)(t) = f(t, u(t)), & 0 \leq t \leq 1, 1 < \alpha \leq 2, \\ a_1 u(0) - b_1 D_q u(0) = c_1 u(\eta_1), & a_2 u(1) + b_2 D_q u(1) = c_2 u(\eta_2), \end{cases}$$

where ${}_C D_q^\alpha$ is the fractional q -derivative of the Caputo type and $a_i, b_i, c_i, \eta_i \in \mathbf{R}$. The existence of solutions for the problem is shown by applying some well-known tools of fixed point theory such as Banach's contraction principle, Krasnoselskii's fixed point theorem, and the Leray-Schauder nonlinear alternative.

In this paper, we deal with the following nonlocal q -integral boundary value problem of nonlinear fractional q -derivatives equation:

$$\begin{cases} (D_q^\alpha u)(t) + f(t, u(t)) = 0, & t \in (0, 1), \\ u(0) = 0, \quad u(1) = \mu I_q^\beta u(\eta) = \mu \int_0^\eta \frac{(\eta - qs)^{(\beta-1)}}{\Gamma_q(\beta)} u(s) d_qs, \end{cases} \tag{1.1}$$

where $q \in (0, 1)$, $1 < \alpha \leq 2$, $0 < \beta \leq 2$, $0 < \eta < 1$, and $\mu > 0$ is a parameter, D_q^α is the q -derivative of Riemann-Liouville type of order α , $f : [0, 1] \times \mathbf{R}^+ \rightarrow \mathbf{R}^+$ is continuous, in which $\mathbf{R}^+ = [0, +\infty)$. To the authors' knowledge, no one has studied the existence of positive solutions for the fractional q -difference boundary value problem (1.1). In the present work, we gave the corresponding Green's function of the boundary value problem (1.1) and its properties. By using the generalized Banach contraction principle, the monotone iterative method, and Krasnoselskii's fixed point theorem, some existence results of positive solutions to the above boundary value problems are enunciated.

2 Preliminaries on q -calculus and lemmas

For the convenience of the reader, below we recall some known facts on fractional q -calculus. The presentation here can be found in, for example, [1, 21, 36, 37].

Let $q \in (0, 1)$ and define

$$[a]_q = \frac{1 - q^a}{1 - q}, \quad a \in \mathbf{R}.$$

The q -analogue of the power function $(a - b)^n$ with $n \in \mathbf{N}_0 := \{0, 1, 2, \dots\}$ is

$$(a - b)^{(0)} = 1, \quad (a - b)^{(n)} = \prod_{k=0}^{n-1} (a - bq^k), \quad n \in \mathbf{N}, a, b \in \mathbf{R}.$$

More generally, if $\gamma \in \mathbf{R}$, then

$$(a - b)^{(\gamma)} = a^\gamma \prod_{k=0}^{\infty} \frac{a - bq^k}{a - bq^{\gamma+k}}, \quad a \neq 0. \tag{2.1}$$

Clearly, if $b = 0$, then $a^{(\gamma)} = a^\gamma$. The q -gamma function is defined by

$$\Gamma_q(x) = \frac{(1 - q)^{(x-1)}}{(1 - q)^{x-1}}, \quad x \in \mathbf{R} \setminus \{0, -1, -2, \dots\},$$

and satisfies $\Gamma_q(x + 1) = [x]_q \Gamma_q(x)$.

The q -derivative of a function f is defined by

$$(D_q f)(x) = \frac{f(qx) - f(x)}{(q - 1)x}, \quad (D_q f)(0) = \lim_{x \rightarrow 0} (D_q f)(x),$$

and the q -derivatives of higher order by

$$(D_q^0 f)(x) = f(x), \quad (D_q^n f)(x) = D_q(D_q^{n-1} f)(x), \quad n \in \mathbf{N}.$$

The q -integral of a function f defined in the interval $[0, b]$ is given by

$$(I_q f)(x) = \int_0^x f(s) d_q s = x(1 - q) \sum_{k=0}^{\infty} f(xq^k) q^k, \quad x \in [0, b],$$

provided the sum converges absolutely.

If $a \in [0, b]$ and f is defined in the interval $[0, b]$, then its integral from a to b is defined by

$$\int_a^b f(s) d_q s = \int_0^b f(s) d_q s - \int_0^a f(s) d_q s.$$

Obviously, if $f(x) \geq g(x)$ on $[a, b]$, then $\int_a^b f(x) d_q x \geq \int_a^b g(x) d_q x$.

Similar as done for derivatives, an operator I_q^n is given by

$$(I_q^0 f)(x) = f(x), \quad (I_q^n f)(x) = I_q(I_q^{n-1} f)(x), \quad n \in \mathbf{N}.$$

The fundamental theorem of calculus applies to these operators I_q and D_q , *i.e.*,

$$(D_q I_q f)(x) = f(x),$$

and if f is continuous at $x = 0$, then

$$(I_q D_q f)(x) = f(x) - f(0). \tag{2.2}$$

The following formulas will be used later, namely, the integration by parts formula

$$\int_0^x f(s)(D_q g)(s) d_q s = [f(s)g(s)]_{s=0}^{s=x} - \int_0^x (D_q f)(s)g(qs) d_q s,$$

and

$$[a(t-s)]^{(\gamma)} = a^\gamma (t-s)^{(\gamma)}, \tag{2.3}$$

$${}_t D_q (t-s)^{(\gamma)} = [\gamma]_q (t-s)^{(\gamma-1)}, \tag{2.4}$$

$${}_s D_q (t-s)^{(\gamma)} = -[\gamma]_q (t-qs)^{(\gamma-1)}, \tag{2.5}$$

$$\left({}_x D_q \int_0^x f(x,s) d_q s \right) (x) = \int_0^x {}_x D_q f(x,s) d_q s + f(qx, x), \tag{2.6}$$

where ${}_t D_q$ denotes the derivative with respect to the variable t .

Definition 2.1 Let $\alpha \geq 0$ and f be a function defined on $[0, 1]$. The fractional q -integral of Riemann-Liouville type is $(I_q^\alpha f)(x) = f(x)$ and

$$(I_q^\alpha f)(x) = \frac{1}{\Gamma_q(\alpha)} \int_0^x (x-qs)^{(\alpha-1)} f(s) d_q s, \quad \alpha > 0, x \in [0, 1].$$

Definition 2.2 The fractional q -derivative of the Riemann-Liouville type of order $\alpha \geq 0$ is defined by $(D_q^\alpha f)(x) = f(x)$ and

$$(D_q^\alpha f)(x) = (D_q^{[\alpha]} I_q^{[\alpha]-\alpha} f)(x), \quad \alpha > 0,$$

where $[\alpha]$ is the smallest integer greater than or equal to α .

Lemma 2.3 Assume that $\gamma \geq 0$ and $a \leq b \leq t$, then $(t-a)^{(\gamma)} \geq (t-b)^{(\gamma)}$.

Lemma 2.4 Let $\alpha, \beta \geq 0$ and f be a function defined on $[0, 1]$. Then the following formulas hold:

- (1) $(I_q^\beta I_q^\alpha f)(x) = (I_q^{\alpha+\beta} f)(x),$
- (2) $(D_q^\alpha I_q^\alpha f)(x) = f(x).$

Lemma 2.5 ([16]) Let $\alpha > 0$ and n be a positive integer. Then the following equality holds:

$$(I_q^\alpha D_q^n f)(x) = (D_q^n I_q^\alpha f)(x) - \sum_{k=0}^{n-1} \frac{x^{\alpha-n+k}}{\Gamma_q(\alpha+k-n+1)} (D_q^k f)(0).$$

Lemma 2.6 ([36]) Let $\alpha \in \mathbf{R}^+, \lambda \in (-1, +\infty)$, the following is valid:

$$I_q^\alpha ((t-a)^{(\lambda)}) = \frac{\Gamma_q(\lambda+1)}{\Gamma_q(\alpha+\lambda+1)} (t-a)^{(\alpha+\lambda)}, \quad 0 < a < t < b.$$

Particularly, for $\lambda = 0, a = 0$, using q -integration by parts, we have

$$\begin{aligned} (I_q^\alpha 1)(t) &= \frac{1}{\Gamma_q(\alpha)} \int_0^t (t - qs)^{(\alpha-1)} d_qs = \frac{1}{\Gamma_q(\alpha)} \int_0^t \frac{{}_sD_q((t-s)^{(\alpha)})}{-[\alpha]_q} d_qs \\ &= -\frac{1}{\Gamma_q(\alpha+1)} \int_0^t {}_sD_q((t-s)^{(\alpha)}) d_qs = \frac{1}{\Gamma_q(\alpha+1)} t^{(\alpha)}. \end{aligned}$$

Obviously, we have $\int_0^t (t - qs)^{(\alpha-1)} d_qs = \frac{1}{[\alpha]_q} t^{(\alpha)}$.

In order to define the solution for the problem (1.1), we need the following lemmas.

Lemma 2.7 *Let $M := \Gamma_q(\alpha + \beta) - \mu \Gamma_q(\alpha) \eta^{\alpha+\beta-1} > 0$. Then, for a given $y \in C[0, 1]$, the unique solution of the boundary value problem*

$$(D_q^\alpha u)(t) + y(t) = 0, \quad t \in (0, 1), 1 < \alpha \leq 2, \tag{2.7}$$

subject to the boundary condition

$$\begin{aligned} u(0) &= 0, \\ u(1) &= \mu I_q^\beta u(\eta) = \mu \int_0^\eta \frac{(\eta - qs)^{(\beta-1)}}{\Gamma_q(\beta)} u(s) d_qs, \quad 0 < \beta \leq 2, 0 < \eta < 1, \end{aligned} \tag{2.8}$$

is given by

$$u(t) = \int_0^1 G(t, qs) y(s) d_qs, \tag{2.9}$$

where

$$G(t, qs) = g(t, qs) + \frac{\mu t^{\alpha-1}}{M} H(\eta, qs), \tag{2.10}$$

$$g(t, s) = \begin{cases} \frac{t^{\alpha-1}(1-s)^{(\alpha-1)} - (t-s)^{(\alpha-1)}}{\Gamma_q(\alpha)}, & 0 \leq s \leq t \leq 1, \\ \frac{t^{\alpha-1}(1-s)^{(\alpha-1)}}{\Gamma_q(\alpha)}, & 0 \leq t \leq s \leq 1, \end{cases} \tag{2.11}$$

and

$$H(\eta, s) = \begin{cases} \eta^{\alpha+\beta-1}(1-s)^{(\alpha-1)} - (\eta-s)^{(\alpha+\beta-1)}, & 0 \leq s \leq \eta < 1, \\ \eta^{\alpha+\beta-1}(1-s)^{(\alpha-1)}, & 0 < \eta \leq s \leq 1. \end{cases} \tag{2.12}$$

Proof Since $1 < \alpha \leq 2$, we take $n = 2$. In view of Definition 2.1 and Lemma 2.4, we have

$$(D_q^\alpha u)(t) = -y(t) \Leftrightarrow (I_q^\alpha D_q^2 I_q^{2-\alpha} u)(t) = -(I_q^\alpha y)(t).$$

Then it follows from Lemma 2.5 that the solution $u(t)$ of (2.7) and (2.8) is given by

$$u(t) = c_1 t^{\alpha-1} + c_2 t^{\alpha-2} - \int_0^t \frac{(t - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} y(s) d_qs \tag{2.13}$$

for some constants $c_1, c_2 \in \mathbf{R}$. Since $u(0) = 0$, we have $c_2 = 0$.

Using the Riemann-Liouville integral of order β for (2.13), we have

$$\begin{aligned} (I_q^\beta u)(t) &= \int_0^t \frac{(t-qs)^{(\beta-1)}}{\Gamma_q(\beta)} \left(c_1 s^{\alpha-1} - \int_0^s \frac{(s-qx)^{(\alpha-1)}}{\Gamma_q(\alpha)} y(x) d_q x \right) d_q s \\ &= - \int_0^t \frac{(t-qs)^{(\alpha+\beta-1)}}{\Gamma_q(\alpha+\beta)} y(s) d_q s + c_1 \int_0^t \frac{(t-qs)^{(\beta-1)}}{\Gamma_q(\beta)} s^{\alpha-1} d_q s \\ &= - \int_0^t \frac{(t-qs)^{(\alpha+\beta-1)}}{\Gamma_q(\alpha+\beta)} y(s) d_q s + c_1 \frac{\Gamma_q(\alpha)}{\Gamma_q(\alpha+\beta)} t^{\alpha+\beta-1}, \end{aligned}$$

where we have used Lemma 2.4 and Lemma 2.6. Using the boundary condition $u(1) = \mu I_q^\beta u(\eta)$, we get

$$c_1 = \frac{\Gamma_q(\alpha+\beta)}{M} \left(\int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} y(s) d_q s - \mu \int_0^\eta \frac{(\eta-qs)^{(\alpha+\beta-1)}}{\Gamma_q(\alpha+\beta)} y(s) d_q s \right).$$

Hence, we have

$$\begin{aligned} u(t) &= \frac{\Gamma_q(\alpha+\beta)t^{\alpha-1}}{M} \left(\int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} y(s) d_q s - \mu \int_0^\eta \frac{(\eta-qs)^{(\alpha+\beta-1)}}{\Gamma_q(\alpha+\beta)} y(s) d_q s \right) \\ &\quad - \int_0^t \frac{(t-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} y(s) d_q s \\ &= \int_0^1 \frac{t^{\alpha-1}(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} y(s) d_q s - \int_0^t \frac{(t-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} y(s) d_q s \\ &\quad + \frac{\mu t^{\alpha-1}}{M} \left(\int_0^1 \eta^{\alpha+\beta-1} (1-qs)^{(\alpha-1)} y(s) d_q s - \int_0^\eta (\eta-qs)^{(\alpha+\beta-1)} y(s) d_q s \right) \\ &= \int_0^1 g(t,qs) y(s) d_q s + \frac{\mu t^{\alpha-1}}{M} \int_0^1 H(\eta,qs) d_q s \\ &= \int_0^1 G(t,qs) y(s) d_q s. \end{aligned}$$

This completes the proof of the lemma. □

Remark 2.8 For the special case where $\mu = 0$, Lemma 2.7 has been obtained by Ferreira [16].

Lemma 2.9 ([16]) *The function $g(t, s)$ defined by (2.11) satisfies the following properties:*

$$g(t,qs) \geq 0 \quad \text{and} \quad g(t,qs) \leq g(qs,qs) \quad \text{for all } 0 \leq t, s \leq 1.$$

Lemma 2.10 *The function $G(t, qs)$ defined by (2.10) satisfies the following properties:*

- (i) G is a continuous function and $G(t, qs) \geq 0$ for $(t, s) \in [0, 1] \times [0, 1]$.
- (ii) There exists a positive function $\rho \in C((0, 1), (0, +\infty))$ such that

$$\max_{0 \leq t \leq 1} G(t,qs) \leq \rho(s),$$

where

$$\rho(s) = g(qs, qs) + \frac{\mu}{M}H(\eta, qs), \quad s \in (0, 1).$$

Proof It is easy to prove that the statement (i) holds. On the other hand, we note that $g(t, s)$ defined by (2.11) is decreasing with respect to t for $s \leq t$ and increasing with respect to t for $t \leq s$. Hence, we have

$$\begin{aligned} \max_{0 \leq t \leq 1} G(t, qs) &= \max_{0 \leq t \leq 1} \left(g(t, qs) + \frac{\mu t^{\alpha-1}}{M}H(\eta, qs) \right) \\ &\leq g(qs, qs) + \frac{\mu}{M}H(\eta, qs) := \rho(s), \quad s \in (0, 1). \end{aligned}$$

The proof is completed. □

3 The main results

Let $X = C([0, 1])$ be a Banach space endowed with the norm $\|u\|_X = \max_{0 \leq t \leq 1} |u(t)|$. Define the cone $P \subset X$ by $P = \{u \in X : u(t) \geq 0, 0 \leq t \leq 1\}$.

Define the operator $T : P \rightarrow X$ as follows:

$$(Tu)(t) = \int_0^1 G(t, qs)f(s, u(s)) d_qs. \tag{3.1}$$

It follows from the nonnegativeness and continuity of G and f that the operator $T : P \rightarrow X$ satisfies $T(P) \subset P$ and is completely continuous.

Theorem 3.1 *Suppose that $f : [0, 1] \times \mathbf{R}^+ \rightarrow \mathbf{R}^+$ is continuous and there exists a function $h : [0, 1] \rightarrow \mathbf{R}^+$ such that*

$$|f(t, u) - f(t, v)| \leq h(t)|u - v|, \quad t \in [0, 1], u, v \in \mathbf{R}^+. \tag{3.2}$$

Then the BVP (1.1) has a unique positive solution provided

$$\int_0^1 s^{\alpha-1}(1 - qs)^{(\alpha-1)}h(s) d_qs < \frac{\Gamma_q(\alpha)M}{2\Gamma_q(\alpha + \beta)}. \tag{3.3}$$

Proof We will prove that under the assumptions (3.2) and (3.3), T^m is a contraction operator for m sufficiently large.

By (2.10), (2.11), and (2.12), for $u, v \in P$, we obtain the estimate

$$\begin{aligned} |(Tu)(t) - (Tv)(t)| &\leq \int_0^1 G(t, qs)|f(s, u(s)) - f(s, v(s))| d_qs \\ &= \int_0^1 \left[g(t, qs) + \frac{\mu t^{\alpha-1}}{M}H(\eta, qs) \right] |f(s, u(s)) - f(s, v(s))| d_qs \\ &\leq \int_0^1 \frac{t^{\alpha-1}(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} h(s) d_qs \cdot \|u - v\|_X \\ &\quad + \mu\eta^{\alpha+\beta-1} \int_0^1 \frac{t^{\alpha-1}(1 - qs)^{(\alpha-1)}}{M} h(s) d_qs \cdot \|u - v\|_X \end{aligned}$$

$$\begin{aligned}
 &= \frac{\Gamma_q(\alpha + \beta)t^{\alpha-1}\|u - v\|_X}{\Gamma_q(\alpha)M} \int_0^1 (1 - qs)^{(\alpha-1)}h(s) d_qs \\
 &= \frac{\Lambda_1\Gamma_q(\alpha + \beta)t^{\alpha-1}}{\Gamma_q(\alpha)M} \|u - v\|_X,
 \end{aligned}$$

where $\Lambda_1 = \int_0^1 (1 - qs)^{(\alpha-1)}h(s) d_qs$.

Consequently,

$$\begin{aligned}
 |(T^2u)(t) - (T^2v)(t)| &\leq \int_0^1 G(t, qs)|f(s, (Tu)(s)) - f(s, (Tv)(s))| d_qs \\
 &\leq \frac{\Lambda_1\Gamma_q(\alpha + \beta)\|u - v\|_X}{\Gamma_q(\alpha)M} \int_0^1 G(t, qs)s^{\alpha-1}h(s) d_qs \\
 &\leq \frac{\Lambda_1[\Gamma_q(\alpha + \beta)]^2t^{\alpha-1}\|u - v\|_X}{[\Gamma_q(\alpha)M]^2} \int_0^1 s^{\alpha-1}(1 - qs)^{(\alpha-1)}h(s) d_qs \\
 &= \frac{\Lambda_1\Lambda_2[\Gamma_q(\alpha + \beta)]^2t^{\alpha-1}}{[\Gamma_q(\alpha)M]^2} \|u - v\|_X,
 \end{aligned}$$

where $\Lambda_2 = \int_0^1 s^{\alpha-1}(1 - qs)^{(\alpha-1)}h(s) d_qs$.

By introduction, we have

$$|(T^m u)(t) - (T^m v)(t)| \leq \frac{\Lambda_1\Lambda_2^{m-1}[\Gamma_q(\alpha + \beta)]^m t^{\alpha-1}}{[\Gamma_q(\alpha)M]^m} \|u - v\|_X.$$

According to (3.3), we can choose m sufficiently large such that

$$\frac{\Lambda_1\Lambda_2^{m-1}[\Gamma_q(\alpha + \beta)]^m}{[\Gamma_q(\alpha)M]^m} = \frac{\Lambda_1}{\Lambda_2} \left[\frac{\Lambda_2\Gamma_q(\alpha + \beta)}{\Gamma_q(\alpha)M} \right]^m < \left(\frac{1}{2}\right)^m \frac{\Lambda_1}{\Lambda_2} < \frac{1}{2},$$

which implies

$$\|(T^m u)(t) - (T^m v)(t)\|_X < \frac{1}{2} \|u - v\|_X.$$

Hence, it follows from the generalized Banach contraction principle that the BVP (1.1) has a unique positive solution. □

Remark 3.2 When $h(t) \equiv h_0$ is a constant, the condition (3.2) reduces to a Lipschitz condition.

For the sake of convenience, we set

$$\begin{aligned}
 l &= \left(\frac{\mu\tau_1^{\alpha-1}}{M} \int_{\tau_1}^{\tau_2} H(\eta, qs) d_qs \right)^{-1}, \\
 L &= \left(\int_0^1 \left(\frac{q^{\alpha-1}(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} + \frac{\mu}{M} H(\eta, qs) \right) d_qs \right)^{-1},
 \end{aligned}$$

where $\tau_1 = q^{m_1}$, $\tau_2 = q^{m_2}$ with $m_1, m_2 \in \mathbf{N}_0$, $m_1 > m_2$.

Theorem 3.3 *Suppose that there exists $0 < \xi_1 < \xi_2$ such that*

$(H_f) f(t, u) : [0, 1] \times [0, \xi_2] \rightarrow \mathbf{R}^+$ is continuous and nondecreasing relative to u , and

$$\max_{0 \leq t \leq 1} f(t, \xi_2) \leq L\xi_2, \tag{3.4}$$

$$\min_{\tau_1 \leq t \leq \tau_2} f(t, \xi_1) \geq l\xi_1. \tag{3.5}$$

Then the BVP (1.1) has one positive solution u^ satisfying*

$$\xi_1 \leq \|u^*\|_X \leq \xi_2 \quad \text{and} \quad \lim_{m \rightarrow \infty} T^m u_0 = u^*, \quad u_0(t) = \xi_2.$$

Proof We denote $\Omega[\xi_1, \xi_2] = \{u \in P : \xi_1 \leq u \leq \xi_2\}$. In what follows, we first show that $T : \Omega[\xi_1, \xi_2] \rightarrow \Omega[\xi_1, \xi_2]$.

Let $u \in \Omega[\xi_1, \xi_2]$; then $0 < \xi_1 \leq u(t) \leq \|u\|_X \leq \xi_2$. By assumption (H_f) , we have

$$0 \leq f(t, u(t)) \leq f(t, \xi_2) \leq \max_{0 \leq t \leq 1} f(t, \xi_2) \leq L\xi_2, \quad 0 \leq t \leq 1,$$

$$f(t, u(t)) \geq f(t, \xi_1) \geq \min_{\tau_1 \leq t \leq \tau_2} f(t, \xi_1) \geq l\xi_1.$$

Hence, for any $u \in \Omega[\xi_1, \xi_2]$,

$$\begin{aligned} \|Tu\|_X &= \max_{0 \leq t \leq 1} \int_0^1 |G(t, qs)f(s, u(s))| d_qs \\ &\leq \int_0^1 \rho(s)f(s, u(s)) d_qs \\ &\leq L\xi_2 \int_0^1 \rho(s) d_qs \\ &\leq L\xi_2 \int_0^1 \left(\frac{q^{\alpha-1}(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} + \frac{\mu}{M} H(\eta, qs) \right) d_qs = \xi_2, \end{aligned}$$

and

$$\begin{aligned} \|Tu\|_X &\geq \min_{\tau_1 \leq t \leq \tau_2} \left(\int_0^1 |g(t, qs)f(s, u(s))| d_qs + \frac{\mu t^{\alpha-1}}{M} \int_0^1 |H(\eta, qs)f(s, u(s))| d_qs \right) \\ &\geq \frac{\mu \tau_1^{\alpha-1}}{M} \left(\int_{\tau_1}^{\tau_2} H(\eta, qs) d_qs \right) \cdot l\xi_1 = \xi_1. \end{aligned}$$

Thus, we get $T : \Omega[\xi_1, \xi_2] \rightarrow \Omega[\xi_1, \xi_2]$.

Let $u_0(t) = \xi_2$, $0 \leq t \leq 1$; then $u_0 \in \Omega[\xi_1, \xi_2]$. Let $u_1(t) = Tu_0(t)$; then $u_1 \in \Omega[\xi_1, \xi_2]$. We denote $u_{m+1} = Tu_m = T^{m+1}u_0$, $m = 0, 1, 2, \dots$. According to $T : \Omega[\xi_1, \xi_2] \rightarrow \Omega[\xi_1, \xi_2]$, we have $u_n \in \Omega[\xi_1, \xi_2]$, $n = 0, 1, 2, \dots$. Since T is completely continuous, we assert that $\{u_m\}_{m=1}^\infty$ has a convergent subsequence $\{u_{m_k}\}_{k=1}^\infty$ and there exists $u^* \in \Omega[\xi_1, \xi_2]$ such that $\lim_{k \rightarrow +\infty} u_{m_k} = u^*$. Since $u_1 \in \Omega[\xi_1, \xi_2]$, then $\xi_1 \leq u_1 \leq \|u_1\|_X \leq \xi_2 = u_0$. According to the definition of T and (H_f) , we have

$$\begin{aligned} u_1(t) &= (Tu_0)(t) \\ &= \int_0^1 G(t, qs)f(s, u_0(s)) d_qs \end{aligned}$$

$$\begin{aligned} &\leq \int_0^1 \rho(s)f(s, \xi_2) d_q s \\ &\leq L\xi_2 \int_0^1 \left(\frac{q^{\alpha-1}(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} + \frac{\mu}{M}H(\eta, qs) \right) d_q s = \xi_2 = u_0(t), \end{aligned}$$

which implies

$$u_2(t) = Tu_1(t) \leq Tu_0(t) = u_1(t), \quad 0 \leq t \leq 1.$$

By introduction, we have $u_{m+1}(t) \leq u_m(t)$ for $0 \leq t \leq 1$, $m = 0, 1, 2, \dots$. Thus, there exists $u^* \in \Omega[\xi_1, \xi_2]$ such that $\lim_{m \rightarrow +\infty} u_m = u^*$. From the continuity of T and $u_{m+1} = Tu_m$, we have $u^* = Tu^*$. The proof is completed. \square

Our next existence result is based on Krasnoselskii's fixed point theorem [39].

Lemma 3.4 (Krasnoselskii's) *Let E be a Banach space, and let $P \subset E$ be a cone. Assume Ω_1, Ω_2 are open subsets of E with $\theta \in \Omega_1 \subset \bar{\Omega}_1 \subset \Omega_2$, and let $T : P \cap (\bar{\Omega}_2 \setminus \Omega_1) \rightarrow P$ be a completely continuous operator such that*

$$\|Tu\| \geq \|u\|, \quad u \in P \cap \partial\Omega_1 \quad \text{and} \quad \|Tu\| \leq \|u\|, \quad u \in P \cap \partial\Omega_2.$$

Then T has at least one fixed point in $P \cap (\bar{\Omega}_2 \setminus \Omega_1)$.

Theorem 3.5 *Let $f(t, u)$ be a nonnegative continuous function on $[0, 1] \times \mathbf{R}^+$. In addition, we assume that*

(H1) *There exists a positive constant r_1 such that*

$$f(t, u) \geq \kappa r_1, \quad \text{for } (t, u) \in [\tau_3, \tau_4] \times [0, r_1],$$

where $\tau_3 = q^{m_3}, \tau_4 = q^{m_4}$ with $m_3, m_4 \in \mathbf{N}_0, m_3 > m_4$, and

$$\kappa \geq \left(\int_{\tau_3}^{\tau_4} \left(g(qs, qs) + \frac{\mu}{M}H(\eta, qs) \right) d_q s \right)^{-1}.$$

(H2) *There exists a positive constant r_2 with $r_2 > r_1$ such that*

$$f(t, u) \leq Lr_2, \quad \text{for } (t, u) \in [0, 1] \times [0, r_2].$$

Then the BVP (1.1) has at least one positive solution u_0 satisfying $0 < r_1 \leq \|u_0\|_X \leq r_2$.

Proof By Lemma 2.9, we obtain that $\max_{0 \leq t \leq 1} g(t, qs) = g(qs, qs)$. Let $\Omega_1 = \{u \in X : \|u\|_X < r_1\}$. For any $u \in X \cap \partial\Omega_1$, according to (H1) and the definitions of τ_3 and τ_4 , we obtain

$$\begin{aligned} \|Tu\|_X &= \max_{0 \leq t \leq 1} \int_0^1 g(t, qs)f(s, u(s)) d_q s + \max_{0 \leq t \leq 1} \int_0^1 \frac{\mu t^{\alpha-1}}{M}H(\eta, qs)f(s, u(s)) d_q s \\ &= \int_0^1 \left[g(qs, qs) + \frac{\mu}{M}H(\eta, qs) \right] f(s, u(s)) d_q s \end{aligned}$$

$$\begin{aligned} &\geq \int_{\tau_3}^{\tau_4} \left(g(qs, qs) + \frac{\mu}{M} H(\eta, qs) \right) f(s, u(s)) d_qs \\ &\geq \kappa r_1 \int_{\tau_3}^{\tau_4} \left(g(qs, qs) + \frac{\mu}{M} H(\eta, qs) \right) d_qs \\ &\geq \|u\|_X = r_1. \end{aligned}$$

Let $\Omega_2 = \{u \in X : \|u\|_X < r_2\}$. For any $u \in X \cap \partial\Omega_2$, by (H2) and Lemma 2.10, we have

$$\begin{aligned} \|Tu\|_X &= \max_{0 \leq t \leq 1} \int_0^1 G(t, qs) f(s, u(s)) d_qs \\ &\leq Lr_2 \int_0^1 \rho(s) d_qs \\ &\leq Lr_2 \int_0^1 \left(\frac{q^{\alpha-1}(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} + \frac{\mu}{M} H(\eta, qs) \right) d_qs = \|u\|_X = r_2. \end{aligned}$$

Now, an application of Lemma 3.4 concludes the proof. □

Theorem 3.6 *Assume that there exist $m + 1$ positive numbers $0 < r_1 < r_2 < \dots < r_{m+1}$ such that*

(H3) $f(t, u) > \kappa r_{2j-1}$ for $(t, u) \in [\tau_3, \tau_4] \times [0, r_{2j-1}]$ and $f(t, u) < Lr_{2j}$ for $(t, u) \in [0, 1] \times [0, r_{2j}]$, $j = 1, 2, \dots, [\frac{m+1}{2}]$; or

(H4) $f(t, u) > \kappa r_{2j}$ for $(t, u) \in [\tau_3, \tau_4] \times [0, r_{2j}]$ and $f(t, u) < Lr_{2j-1}$ for $(t, u) \in [0, 1] \times [0, r_{2j-1}]$, $j = 1, 2, \dots, [\frac{m+1}{2}]$,

where τ_3, τ_4, κ are given in (H1).

Then the BVP (1.1) has at least n positive solutions u_1, u_2, \dots, u_m with $r_j < \|u_j\|_X < r_{j+1}$, $j = 1, \dots, m$.

Proof Suppose that the condition (H3) holds. According to the continuity of f , for every pair (r_j, r_{j+1}) , there exists (a_j, b_j) with $r_j < a_j < b_j < r_{j+1}$ such that

$$\begin{aligned} f(t, u) &\geq \kappa b_{2j-1}, & f(t, u) &\leq La_{2j-1} \quad \text{and} \\ f(t, u) &\geq \kappa a_{2j}, & f(t, u) &\leq Lb_{2j}, \quad j = 1, 2, \dots, \left[\frac{m+1}{2} \right]. \end{aligned}$$

It follows from Theorem 3.5 that every pair (a_j, b_j) presents a positive solution of the BVP (1.1) such that $a_j \leq \|u_j\|_X < b_j$, $j = 1, 2, \dots, m$.

When the condition (H4) holds, the proofs are similar to those in the case (H3). The proof is completed. □

4 Examples

Example 4.1 The fractional boundary value problem

$$\begin{cases} D_{0.5}^{1.5} u(t) + \frac{e^t(1+u)}{2(1+e^t)} + t \sin^2 t + 4 = 0, & 0 < t < 1, \\ u(0) = 0, & u(1) = 0.25 I_{0.5}^{0.5} u(0.75) \end{cases} \quad (4.1)$$

has a unique positive solution.

Proof In this case, $\alpha = 1.5$, $\beta = q = 0.5$, $\mu = 0.25$, $\eta = 0.75$. Let

$$f(t, u) = \frac{e^t(1 + u)}{2(1 + e^t)} + t \sin^2 t + 4, \quad (t, u) \in [0, 1] \times [0, +\infty),$$

and $h(t) = \frac{e^t}{2(1+e^t)}$. It is easy to prove that

$$|f(t, u) - f(t, v)| \leq h(t)|u - v|, \quad \text{for } (t, u), (t, v) \in [0, 1] \times [0, +\infty).$$

By simple calculation, we get

$$\begin{aligned} M &= \Gamma_q(\alpha + \beta) - \mu \Gamma_q(\alpha) \eta^{\alpha+\beta-1} \approx 0.8072, \\ \int_0^1 s^{\alpha-1} (1 - qs)^{(\alpha-1)} h(s) d_qs &= \int_0^1 s^{\alpha-1} (1 - qs)^{(\alpha-1)} \cdot \frac{e^s}{2(1 + e^s)} d_qs \\ &\leq \frac{1}{2} \int_0^1 s^{\alpha-1} (1 - qs)^{(\alpha-1)} d_qs \\ &\leq \frac{1}{2} \int_0^1 (1 - qs)^{(\alpha-1)} d_qs \approx 0.3867, \end{aligned}$$

and

$$\begin{aligned} \int_0^1 (1 - qs)^{(\alpha-1)} h(s) d_qs &= \int_0^1 (1 - qs)^{(\alpha-1)} \cdot \frac{e^s}{2(1 + e^s)} d_qs \\ &\leq \frac{1}{2} \int_0^1 (1 - qs)^{(\alpha-1)} d_qs \approx 0.3867, \end{aligned}$$

which implies that

$$\int_0^1 s^{\alpha-1} (1 - qs)^{(\alpha-1)} h(s) d_qs < \frac{\Gamma_q(\alpha)M}{2\Gamma_q(\alpha + \beta)} \approx 0.4512.$$

Obviously, for any $m \geq 3$, we have

$$\frac{\Lambda_1 \Lambda_2^{m-1} [\Gamma_q(\alpha + \beta)]^m}{[\Gamma_q(\alpha)M]^m} \leq \frac{0.3867 \times \Gamma_q(\alpha + \beta)}{2^{m-1} \Gamma_q(\alpha)M} < 0.2329 < \frac{1}{2}.$$

Thus Theorem 3.1 implies that the boundary value problem (4.1) has a unique positive solution. \square

Example 4.2 Consider the following fractional boundary value problem:

$$\begin{cases} D_{0.5}^{1.5} u(t) + 21\sqrt{u} - \frac{t^2}{10} + \frac{t}{10} = 0, & 0 < t < 1, \\ u(0) = 0, & u(1) = 1.25 I_{0.5}^{0.5} u(0.25), \end{cases} \quad (4.2)$$

where $\alpha = 1.5$, $\beta = q = 0.5$, $\mu = 1.25$, $\eta = 0.25$. Choosing $m_1 = 2$, $m_2 = 1$, then $\tau_1 = 0.25$, $\tau_2 = 0.5$.

A simple computation showed $M \approx 0.6786$. By Lemma 2.6 and with the aid of a computer, we obtain that

$$l = \left(\frac{\mu \tau_1^{\alpha-1}}{M} \int_{\tau_1}^{\tau_2} H(\eta, qs) d_qs \right)^{-1} \approx 18.9553,$$

and

$$L = \left(\int_0^1 \left(\frac{q^{\alpha-1}(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} + \frac{\mu}{M} H(\eta, qs) \right) d_qs \right)^{-1} \approx 1.1770.$$

Let $f(t, u) = -\frac{t^2}{10} + \frac{t}{10} + 21\sqrt{u}$. Take $\xi_1 = 1, \xi_2 = 340$, then $f(t, u)$ satisfies

- (i) $f(t, u) : [0, 1] \times [0, 340] \rightarrow \mathbf{R}^+$ is continuous and nondecreasing relative to u ;
- (ii) $\max_{0 \leq t \leq 1} f(t, \xi_2) = f(\frac{1}{2}, 340) \approx 387.2459 < L\xi_2 \approx 400.1800$;
- (iii) $\min_{\frac{1}{4} \leq t \leq \frac{1}{2}} f(t, \xi_1) = f(\frac{1}{4}, 1) \approx 21.0188 > l\xi_1 \approx 18.9553$.

So, by Theorem 3.3, the problem (4.2) has one positive solution u^* satisfying

$$1 \leq \|u^*\|_X \leq 340 \quad \text{and} \quad \lim_{m \rightarrow \infty} T^m u_0 = u^*, \quad u_0(t) = 340.$$

Example 4.3 Consider the following fractional boundary value problem:

$$\begin{cases} D_{0.5}^{1.5} u(t) + \frac{u^2}{100} + \frac{\sin^2 t}{20} + \frac{t}{10} + 1 = 0, & 0 < t < 1, \\ u(0) = 0, \quad u(1) = 0.75 I_{0.5}^{1.5} u(0.75), \end{cases} \quad (4.3)$$

where $\alpha = \beta = 1.5, q = 0.5, \mu = \eta = 0.75$. Choosing $m_3 = 3, m_4 = 1$, then $\tau_3 = 0.125, \tau_4 = 0.5$.

By calculation, we get $M \approx 1.0661$. By Lemma 2.6 and with the aid of a computer, we obtain that

$$\begin{aligned} & \left(\int_{\tau_3}^{\tau_4} \left(g(qs, qs) + \frac{\mu}{M} H(\eta, qs) \right) d_qs \right)^{-1} \\ & \leq \left(\int_{\tau_3}^{\tau_4} \left(\frac{(q\tau_3)^{\alpha-1}(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} d_qs + \frac{\mu}{M} H(\eta, qs) \right) d_qs \right)^{-1} \approx 3.4698, \end{aligned}$$

and

$$L = \left(\int_0^1 \left(\frac{q^{\alpha-1}(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} + \frac{\mu}{M} H(\eta, qs) \right) d_qs \right)^{-1} \approx 1.4966.$$

Let $f(t, u) = \frac{u^2}{100} + \frac{\sin^2 t}{20} + \frac{t}{10} + 1$. Take $\kappa = 7$ and $r_1 = \frac{1}{7}, r_2 = \frac{9}{10}$. Then $f(t, u)$ satisfies

- (i) $f(t, u) : [0, 1] \times \mathbf{R}^+ \rightarrow \mathbf{R}^+$ is continuous;
- (ii) $f(t, u) \leq \frac{u^2}{100} + \frac{1}{20} + \frac{1}{10} + 1 \leq 1.1581 < Lr_2 \approx 1.3469, (t, u) \in [0, 1] \times [0, \frac{9}{10}]$;
- (iii) $f(t, u) = \frac{u^2}{100} + \frac{\sin^2 t}{20} + \frac{t}{10} + 1 \geq 1 + \frac{\tau_1}{10} = 1.0125 > \kappa r_1 = 1, (t, u) \in [0.125, 0.5] \times [0, \frac{1}{7}]$.

So, by Theorem 3.5, the problem (4.3) has at least one positive solution u_0 with $\frac{1}{7} \leq \|u_0\|_X \leq \frac{9}{10}$.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Each of the authors, YZ, HC, and QZ, contributed to each part of this study equally and read and approved the final version of the manuscript.

Author details

¹School of Science, Hunan University of Technology, Zhuzhou, Hunan 412007, China. ²Department of Mathematics, Central South University, Changsha, Hunan 410075, China.

Acknowledgements

Dedicated to Professor Hari M Srivastava.

The authors are highly grateful for the referee's careful reading and comments on this paper. The research is supported by the National Natural Science Foundation of China (11271372, 11201138); it is also supported by the Hunan Provincial Natural Science Foundation of China (12JJ2004, 13JJ3106), and the Scientific Research Fund of Hunan Provincial Education Department (12B034).

Received: 1 December 2012 Accepted: 17 February 2013 Published: 4 March 2013

References

1. Jackson, FH: q -difference equations. *Am. J. Math.* **32**, 305-314 (1910)
2. Carmichael, RD: The general theory of linear q -difference equations. *Am. J. Math.* **34**, 147-168 (1912)
3. Finkelstein, R, Marcus, E: Transformation theory of the q -oscillator. *J. Math. Phys.* **36**, 2652-2672 (1995)
4. Freund, PGO, Zabrodin, AV: The spectral problem for the q -Knizhnik-Zamolodchikov equation and continuous q -Jacobi polynomials. *Commun. Math. Phys.* **173**, 17-42 (1995)
5. Han, G, Zeng, J: On a q -sequence that generalizes the median Genocchi numbers. *Ann. Sci. Math. Qué.* **23**, 63-72 (1999)
6. Floreanini, R, Vinet, L: Quantum symmetries of q -difference equations. *J. Math. Phys.* **36**, 3134-3156 (1995)
7. Bangerezako, G: Variational q -calculus. *J. Math. Anal. Appl.* **289**, 650-665 (2004)
8. Dobrogowska, A, Odziejewicz, A: Second order q -difference equations solvable by factorization method. *J. Comput. Appl. Math.* **193**, 319-346 (2006)
9. Ismail, MEH, Simeonov, P: q -difference operators for orthogonal polynomials. *J. Comput. Appl. Math.* **233**, 749-761 (2009)
10. Ahmad, B, Ntouyas, SK, Purnaras, I: Existence results for nonlocal boundary value problems of nonlinear fractional q -difference equations. *Adv. Differ. Equ.* **2012**, 140 (2012)
11. El-Shahed, M, Hassan, HA: Positive solutions of q -difference equation. *Proc. Am. Math. Soc.* **138**, 1733-1738 (2010)
12. Annaby, MH, Mansour, ZS: q -Taylor and interpolation series for Jackson q -difference operators. *J. Math. Anal. Appl.* **344**, 472-483 (2008)
13. Ahmad, B, Ntouyas, SK: Boundary value problems for q -difference inclusions. *Abstr. Appl. Anal.* **2011**, Article ID 292860 (2011)
14. Gasper, G, Rahman, M: Some systems of multivariable orthogonal q -Racah polynomials. *Ramanujan J.* **13**, 389-405 (2007)
15. Gauchman, H: Integral inequalities in q -calculus. *Comput. Math. Appl.* **47**, 281-300 (2004)
16. Ferreira, RAC: Nontrivial solutions for fractional q -difference boundary value problems. *Electron. J. Qual. Theory Differ. Equ.* **2010**, 70 (2010)
17. Ferreira, RAC: Positive solutions for a class of boundary value problems with fractional q -differences. *Comput. Math. Appl.* **61**, 367-373 (2011)
18. El-Shahed, M, Al-Askar, F: Positive solutions for boundary value problem of nonlinear fractional q -difference equation. *ISRN Math. Anal.* **2011**, Article ID 385459 (2011)
19. Liang, S, Zhang, J: Existence and uniqueness of positive solutions for three-point boundary value problem with fractional q -differences. *J. Appl. Math. Comput.* **40**, 277-288 (2012)
20. Graef, JR, Kong, L: Positive solutions for a class of higher order boundary value problems with fractional q -derivatives. *Appl. Math. Comput.* **218**, 9682-9689 (2012)
21. Kac, V, Cheung, P: *Quantum Calculus*. Springer, New York (2002)
22. Podlubny, I: *Fractional Differential Equations*. Academic Press, San Diego (1999)
23. Kibas, AA, Srivastava, HM, Trujillo, JJ: *Theory and Applications of Fractional Differential Equations*. Elsevier, Amsterdam (2006)
24. Li, C, Luo, X, Zhou, Y: Existence of positive solutions of the boundary value problem for nonlinear fractional differential equations. *Comput. Math. Appl.* **59**, 1363-1375 (2010)
25. Wang, G, Ahmad, B, Zhang, L: Some existence results for impulsive nonlinear fractional differential equations with mixed boundary conditions. *Comput. Math. Appl.* **62**, 1389-1397 (2011)
26. Ahmad, B, Sivasundaram, S: On four-point nonlocal boundary value problems of nonlinear integro-differential equations of fractional order. *Appl. Math. Comput.* **217**, 480-487 (2010)
27. Staněk, S: The existence of positive solutions of singular fractional boundary value problems. *Comput. Math. Appl.* **62**, 1379-1388 (2011)
28. Zhao, Y, Huang, L, Wang, X, Zhu, X: Existence of solutions for fractional integro-differential equation with multipoint boundary value problem in Banach spaces. *Abstr. Appl. Anal.* **2012**, Article ID 172963 (2012)
29. Zhao, Y, Chen, H, Huang, L: Existence of positive solutions for nonlinear fractional functional differential equation. *Comput. Math. Appl.* **64**, 3456-3467 (2012)
30. Bai, Z, Lü, H: Positive solutions for boundary value problem of nonlinear fractional differential equation. *J. Math. Anal. Appl.* **311**, 495-505 (2005)
31. Goodrich, CS: On discrete sequential fractional boundary value problems. *J. Math. Anal. Appl.* **385**, 111-124 (2012)
32. Zhang, S: Positive solutions to singular boundary value problem for nonlinear fractional differential equation. *Comput. Math. Appl.* **59**, 1300-1309 (2010)

33. Zhang, X, Liu, L, Wu, Y: Multiple positive solutions of a singular fractional differential equation with negatively perturbed term. *Math. Comput. Model.* **55**, 1263-1274 (2012)
34. Al-Salam, WA: Some fractional q -integrals and q -derivatives. *Proc. Edinb. Math. Soc.* **15**, 135-140 (1966/1967)
35. Agarwal, RP: Certain fractional q -integrals and q -derivatives. *Proc. Camb. Philos. Soc.* **66**, 365-370 (1969)
36. Rajković, PM, Marinković, SD, Stanković, MS: Fractional integrals and derivatives in q -calculus. *Appl. Anal. Discrete Math.* **1**, 311-323 (2007)
37. Annaby, MH, Mansour, IS: *q -Fractional Calculus and Equations*. Springer, Berlin (2012)
38. Zhao, Y, Ye, G, Chen, H: Multiple positive solutions of a singular semipositone integral boundary value problem for fractional q -derivatives equation. *Abstr. Appl. Anal.* **2013**, Article ID 643571 (2013)
39. Guo, DJ, Lakshmikantham, V: *Nonlinear Problems in Abstract Cones*. Academic Press, San Diego (1988)

doi:10.1186/1687-1847-2013-48

Cite this article as: Zhao et al.: Existence results for fractional q -difference equations with nonlocal q -integral boundary conditions. *Advances in Difference Equations* 2013 2013:48.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
