




Contents lists available at ScienceDirect

Topology and its Applications

www.elsevier.com/locate/topol


$\mathcal{P}(\mathbb{R})$, ordered by homeomorphic embeddability, does not represent all posets of cardinality 2^c

 Robin W. Knight^a, Aisling E. McCluskey^{b,*}
^a *Mathematical Institute, Oxford, UK*^b *Department of Mathematics, National University of Ireland, Galway, Ireland*

ARTICLE INFO

Article history:

Received 24 July 2006

Received in revised form 3 August 2007

MSC:

06A06

54H10

Keywords:

Partial order

Ordering by homeomorphic embeddability

Forcing

ABSTRACT

We prove it to be consistent that there is a poset of cardinality 2^c which is not realizable in $\mathcal{P}(\mathbb{R})$, ordered by homeomorphic embeddability. This addresses and answers resolutely (and in the negative) the open question of whether there is a ZFC theorem that all posets of cardinality 2^c can be represented by subspaces of the real line ordered by homeomorphic embeddability. This question arises from the pioneering work of Banach, Kuratowski and Sierpiński in the area and this result complements the recent work of [A.E. McCluskey, D. Shakhmatov, It is consistent that all posets of cardinality 2^c can be realized within $\mathcal{P}(\mathbb{R})$, preprint], thereby providing a proof of independence.

© 2009 Elsevier B.V. All rights reserved.

1. Introduction

The ordering by embeddability of topological spaces, although a fundamental notion in topology, has been remarkably little understood for some years. This ordering is that introduced into a family of topological spaces by writing $X \hookrightarrow Y$ whenever X is homeomorphic to a subspace of Y . Its subtlety and relative intractability are well illustrated by the problem of recognizing which order-types are those of collections of subspaces of the real line \mathbb{R} (see [4–8]). A partially-ordered set (poset) P is *realized* (or *realizable*) within a family \mathcal{F} of topological spaces whenever there is an injection $\theta : P \rightarrow \mathcal{F}$ for which $p \leq q$ if and only if $\theta(p) \hookrightarrow \theta(q)$. Discussion of realizability in the powerset $\mathcal{P}(\mathbb{R})$ can be traced back to Banach, Kuratowski and Sierpiński [2,3,11], whose work on the extensibility of continuous maps over G_δ subsets (in the context of Polish spaces) revealed *inter alia* that it is possible to realize, within $\mathcal{P}(\mathbb{R})$, (i) the antichain of cardinality 2^c [2, p. 205] and (ii) the ordinal c^+ [3, p. 199]. Renewed interest in the problem was initiated in [4] in which it was shown that every poset of cardinality c or less can be realized within $\mathcal{P}(\mathbb{R})$. Till now, the question of precisely which posets of cardinalities exceeding c can be so realized had been unresolved. Indeed, the question ultimately revealed itself to be set-theoretic in nature. Accordingly, this paper seeks to answer this question by forcing to construct a poset of cardinality 2^c which cannot be realized within $\mathcal{P}(\mathbb{R})$. This, together with [9], then provides a clearer picture of the order-theoretic structure of $\mathcal{P}(\mathbb{R})$, namely that the statement that all posets of cardinality 2^c can be realized within $\mathcal{P}(\mathbb{R})$ is independent of ZFC.

2. Forcing argument

All forcing-related notation in this section is taken from [1].

* Corresponding author. Tel.: +353 91 493162.

E-mail addresses: knight@maths.ox.ac.uk (R.W. Knight), aisling.mccluskey@nuigalway.ie (A.E. McCluskey).

Theorem 2.1. *It is consistent with ZFC that there exists a partially ordered set of cardinality 2^c that is not realizable in $\mathcal{P}(\mathbb{R})$.*

Proof. We assume that in the ground model $2^{\aleph_0} = \aleph_1$, $2^{\aleph_1} = \aleph_3$, and $2^{\aleph_2} = \aleph_4$. We will construct a bipartite graph, and construct a partially-ordered set from it that cannot be realized in $\mathcal{P}(\mathbb{R})$. Our graph will have a set of vertices

$$V = \omega_2 \times \{0\} \cup \omega_3 \times \{1\},$$

and the edge relation E will be a subset of $(\omega_2 \times \{0\}) \times (\omega_3 \times \{1\})$. We will code E by a relation $F \subseteq \omega_2 \times \omega_3$. The partially-ordered set will then be V , ordered so that $\langle \alpha, i \rangle \leq \langle \beta, j \rangle$ if and only if either $\langle \alpha, i \rangle = \langle \beta, j \rangle$, or $i = 0, j = 1$, and $\langle \langle \alpha, i \rangle, \langle \beta, j \rangle \rangle \in E$, that is to say, iff $\langle \alpha, \beta \rangle \in F$.

We create F by forcing. The forcing poset will be

$$\mathbb{P} = \text{Fn}(\omega_2 \times \omega_3, 2, \aleph_3).$$

Since \mathbb{P} is c^+ -closed, forcing with \mathbb{P} creates no new reals and no new sets of reals. Let \dot{F} be a name for the generic subset of $\omega_2 \times \omega_3$ that we have created by forcing over \mathbb{P} . We show that, in a generic extension, the associated partial order, as defined above, is not realizable, by considering possible maps $\phi : V \rightarrow \mathcal{P}(\mathbb{R})$; we aim to show that it is impossible that $\phi(\alpha, 0)$ is (homeomorphically) embeddable in $\phi(\beta, 1)$ iff $\langle \alpha, \beta \rangle \in F$.

In other words, we show that for every name $\dot{\phi}$ for such a map, for all $p \in \mathbb{P}$, there exists $q \leq p$ such that, for some α, β , either

$$q \Vdash \langle \alpha, \beta \rangle \in \dot{F} \quad \wedge \quad \dot{\phi}(\check{\alpha}, \check{0}) \not\hookrightarrow \dot{\phi}(\check{\beta}, \check{1}),$$

or

$$q \Vdash \langle \alpha, \beta \rangle \notin \dot{F} \quad \wedge \quad \dot{\phi}(\check{\alpha}, \check{0}) \hookrightarrow \dot{\phi}(\check{\beta}, \check{1}).$$

So, given $\dot{\phi}$ and p , we first exploit \aleph_3 -closure of \mathbb{P} to extend p to a condition $r \leq p$ such that for all $\alpha < \omega_2$, there exists $A_\alpha \in \mathcal{P}(\mathbb{R})$ such that

$$r \Vdash \dot{\phi}(\check{\alpha}, \check{0}) = \check{A}_\alpha.$$

For each subset C of \mathbb{R} , let

$$D_C = \{\alpha : A_\alpha \hookrightarrow C\}.$$

Now $\mathcal{P}(\mathbb{R})$ has only \aleph_3 elements, while $2^{\aleph_2} = \aleph_4$. Therefore there exists a subset Δ of ω_2 such that for all $C, \Delta \neq D_C$.

Now pick $\beta \in \omega_3$ such that for all $\alpha \in \omega_2, \langle \alpha, \beta \rangle \notin \text{dom } r$, which we can do since r has cardinality at most \aleph_2 .

Define $s \leq r$ so that if $\alpha \in \Delta, s(\alpha, \beta) = 1$, and if $\alpha \notin \Delta$, then $s(\alpha, \beta) = 0$.

Now find $q \leq s$ such that for some $C \subseteq \mathbb{R}$,

$$q \Vdash \dot{\phi}(\check{\beta}, \check{1}) = \check{C}.$$

Now, recall that $\Delta \neq D_C$. Therefore, either there exists $\alpha \in \Delta$ such that $\alpha \notin D_C$, or there exists $\alpha \notin \Delta$ such that $\alpha \in D_C$.

In the first case,

$$q \Vdash \langle \alpha, \beta \rangle \in \dot{F} \quad \wedge \quad \dot{\phi}(\check{\alpha}, \check{0}) \not\hookrightarrow \dot{\phi}(\check{\beta}, \check{1}),$$

and in the second case,

$$q \Vdash \langle \alpha, \beta \rangle \notin \dot{F} \quad \wedge \quad \dot{\phi}(\check{\alpha}, \check{0}) \hookrightarrow \dot{\phi}(\check{\beta}, \check{1}),$$

and the proof is complete. \square

3. Future hopes

In its most general form, the question of determining, given a poset P and a topological space X , if there are subspaces of X whose embeddability ordering precisely matches that of the ordering within P , is relatively unexplored and deserving of investigation. We now know that for a space as familiar as \mathbb{R} , the order-theoretic structure of its powerset $\mathcal{P}(\mathbb{R})$ is inherently set-theoretic and ultimately cannot be determined in ZFC. A surprising observation is that the topological nature of \mathbb{R} appears to be irrelevant. In fact, the argument presented applies to any space of cardinality c in the ground model. What is true then for spaces of higher cardinality? Further, for such poset representations within $\mathcal{P}(\mathbb{R})$ as are possible, is it possible to restrict the representative subspaces to some ‘nice’ family of subsets of \mathbb{R} ? In [10], further work along these lines is presented.

More generally, the relation of embeddability is of interest in that in studying it, we are studying the structure of the category of topological spaces, or rather, of those spaces which are homeomorphic to subsets of \mathbb{R} . Since all subspaces of \mathbb{R} which include an interval are equivalent under the relation of embeddability, what we are essentially studying is the structure of the category of all zero-dimensional separable metric spaces. It would be natural, therefore, to broaden our study to include other categories of topological spaces, such as zero-dimensional spaces of weight κ , or spaces of cardinality κ , and ask what can be said about these.

Acknowledgement

It is a pleasure to record our indebtedness to Stephen Watson, York University, Toronto for valuable conversations and insights in pursuit of this result.

References

- [1] K. Kunen, *Set Theory: An Introduction to Independence Proofs*, North-Holland, 1980.
- [2] C. Kuratowski, Sur la puissance de l'ensemble des "nombres de dimension" au sens de M. Fréchet, *Fund. Math.* 8 (1926) 201–208.
- [3] C. Kuratowski, W. Sierpiński, Sur un problème de M. Fréchet concernant les dimensions des ensembles linéaires, *Fund. Math.* 8 (1926) 193–200.
- [4] P.T. Matthews, T.B.M. McMaster, Families of spaces having prescribed embeddability order-type, *Rend. Istit. Mat. Univ. Trieste* 25 (1993) 345–352.
- [5] P.T. Matthews, T.B.M. McMaster, On chains and posets within the power set of a continuum, *Int. J. Math. Sci.* 18 (1995) 305–309.
- [6] A.E. McCluskey, T.B.M. McMaster, Representing quasi-orders by embeddability ordering of families of topological spaces, *Proc. Amer. Math. Soc.* 127 (5) (1999) 1275–1279.
- [7] A.E. McCluskey, T.B.M. McMaster, Realising quasiordered sets by subspaces of 'continuum-like' spaces, *Order* 15 (2) (1999) 143–149.
- [8] A.E. McCluskey, T.B.M. McMaster, W.S. Watson, Representing set-inclusion by embeddability (among the subspaces of the real line), *Topology Appl.* 96 (1) (1999) 89–92.
- [9] A.E. McCluskey, D. Shakhmatov, It is consistent that all posets of cardinality 2^c can be realized within $\mathcal{P}(\mathbb{R})$, preprint.
- [10] A.E. McCluskey, D. Shakhmatov, More on embeddability of posets, preprint.
- [11] W. Sierpiński, Sur un problème concernant les types de dimensions, *Fund. Math.* 19 (1932) 65–71.