

Erratum

In the November 2000 article by Suzuki et al., “Gut Cryptopatches: Direct Evidence of Extrathymic Anatomical Sites for Intestinal Lymphopoiesis,” there were errors in Table 1, which was published in the Results section of this paper. We would therefore like to publish a correction. The correct table appears on the following page.

**Kenji Suzuki,^{1,2} Takatoku Oida,^{1,3,8} Hiromasa Hamada,^{1,4} Osamu Hitotsumatsu,⁵
Mamoru Watanabe,⁶ Toshifumi Hibi,⁵ Hiroshi Yamamoto,⁴ Eiro Kubota,²
Shuichi Kaminogawa,³ and Hiromichi Ishikawa^{1,7}**

¹Department of Microbiology
Keio University School of Medicine
Tokyo 160-8582

²Second Department of Oral and Maxillofacial Surgery
Kanagawa Dental School
Kanagawa 238-8580

³Department of Applied Biological Chemistry
The University of Tokyo
Tokyo 113-8657

⁴Department of Immunology
Graduate School of Pharmaceutical Science
Osaka University
Osaka 565-0871

⁵Department of Internal Medicine
Keio University School of Medicine
Tokyo 160-8582

⁶Department of Gastroenterology and Hepatology
Graduate School
Tokyo Medical and Dental University
Tokyo 113-8519
Japan

⁷To whom correspondence should be addressed (e-mail: ishikawa@sun.microb.med.keio.ac.jp).

⁸Present address: Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts 02115.

Table 1. Characterization of IEL from BM Chimeric Mice^a

Host	Host No.	Irradiation (6 Gy)	Days After Reconstitution	BM-Derived Ly5.1 ⁺ IEL %	No. of Cells ($\times 10^6$)	TCR- $\alpha\beta$	TCR- $\gamma\delta$	CD4	CD8 $\alpha\alpha$	CD8 $\alpha\beta$
	1	+	22	95	99.0	0.59	14.9	2.48	52.5	1.29
	2	+	22	97	107	1.28	25.7	2.03	64.2	0.96
	3	+	22	99	119	2.62	34.5	2.62	84.5	1.07
	4	+	35	98	157	7.22	58.1	1.57	92.6	1.26
	5	+	35	98	98.0	1.67	40.2	2.94	55.9	1.96
	6	+	60	99	139	16.7	54.2	11.8	80.6	2.92
	7	+	60	98	172	9.98	96.3	6.19	119	2.41
	8	+	83	99	168	20.2	104	12.9	116	7.56
	9	+	83	99	221	46.4	135	15.9	183	6.85
	10	-	25	62	62.0	4.15	14.3	1.36	31.6	0.50
	11	-	25	53	74.2	8.16	11.1	2.23	28.2	0.59
	12	-	35	73	80.3	12.8	45.0	2.57	62.6	0.64
	13	-	46	86	73.1	17.5	30.0	3.14	54.8	1.46
	14	-	46	87	120	36.0	56.4	3.00	92.4	1.44
	15	-	60	90	126	17.6	84.4	8.06	111	0.88
	16	-	60	97	233	17.5	163	5.83	186	5.13
	17	-	77	89	134	20.1	83.1	6.97	96.5	1.47
	18	-	83	95	200	34.0	126	8.40	164	4.80
	1	+	35	99	218	63.2	78.5	6.54	129	26.2
	2	+	60	99	351	91.3	214	23.2	228	45.6
	3	+	60	98	258	41.3	165	13.7	181	28.4
	4	+	83	98	245	71.1	169	19.6	176	36.8
	5	+	83	99	396	111	261	19.8	293	47.5
									125 \pm 42.6	4.94 \pm 2.65
									11.7 \pm 4.06	139 \pm 42.5
									7.32 \pm 1.16	3.07 \pm 2.21
									19.1 \pm 3.95	220 \pm 54.3
									19.1 \pm 3.95	39.6 \pm 8.79

^a Two-color flow cytometric analyses were performed on IEL isolated from various BM chimeric mice. IEL were incubated first with biotinylated anti-Ly5.1 mAb and then with streptavidin-PE and FITC-conjugated anti-TCR- $\alpha\beta$, anti-TCR- $\gamma\delta$, anti-CD4, anti-CD8 α or anti-CD8 β mAb. Absolute numbers of Ly5.1⁺ IEL were calculated on the basis of total numbers of IEL recovered, and those of BM-derived $\alpha\beta$, $\gamma\delta$, CD4 and CD8 $\alpha\beta$ IEL subsets were calculated on the basis of total numbers of Ly5.1⁺ IEL. Absolute numbers of CD8 $\alpha\alpha$ IEL were calculated from (numbers of Ly5.1⁺ CD8 $\alpha\alpha$ IEL - numbers of Ly5.1⁺ CD8 $\alpha\beta$ IEL). Dead cells were excluded by PI gating.