

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Forensic Science International: Genetics

journal homepage: www.elsevier.com/locate/fsig

Verifying likelihoods for low template DNA profiles using multiple replicates

Christopher D. Steele^{a,*}, Matthew Greenhalgh^b, David J. Balding^a^a UCL Genetics Institute, Darwin Building, Gower Street, London WC1E 6BT, UK^b Orchid Cellmark Ltd., Abingdon Business Park, Blacklands Way, Abingdon OX14 1YX, UK

ARTICLE INFO

Article history:

Received 19 February 2014

Received in revised form 9 June 2014

Accepted 30 June 2014

Keywords:

Low-template DNA

DNA mixtures

Likelihood ratio

Replicates

Forensic

likeLTD

ABSTRACT

To date there is no generally accepted method to test the validity of algorithms used to compute likelihood ratios (LR) evaluating forensic DNA profiles from low-template and/or degraded samples. An upper bound on the LR is provided by the inverse of the match probability, which is the usual measure of weight of evidence for standard DNA profiles not subject to the stochastic effects that are the hallmark of low-template profiles. However, even for low-template profiles the LR in favour of a true prosecution hypothesis should approach this bound as the number of profiling replicates increases, provided that the queried contributor is the major contributor. Moreover, for sufficiently many replicates the standard LR for mixtures is often surpassed by the low-template LR. It follows that multiple LTDNA replicates can provide stronger evidence for a contributor to a mixture than a standard analysis of a good-quality profile. Here, we examine the performance of the `likeLTD` software for up to eight replicate profiling runs. We consider simulated and laboratory-generated replicates as well as resampling replicates from a real crime case. We show that LRs generated by `likeLTD` usually do exceed the mixture LR given sufficient replicates, are bounded above by the inverse match probability and do approach this bound closely when this is expected. We also show good performance of `likeLTD` even when a large majority of alleles are designated as uncertain, and suggest that there can be advantages to using different profiling sensitivities for different replicates. Overall, our results support both the validity of the underlying mathematical model and its correct implementation in the `likeLTD` software.

© 2014 The Authors. Published by Elsevier Ireland Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

1. Introduction

In forensic DNA profiling, a likelihood ratio (LR) is calculated to measure the support provided by DNA evidence (E) for a proposition H_p favouring the prosecution case, relative to its support for H_d representing the defence case. The LR can be written as

$$LR = \frac{\Pr(E|H_p)}{\Pr(E|H_d)}. \quad (1)$$

Each of H_p and H_d specifies a number of unprofiled contributors and a list of contributors whose DNA profiles are known (included in E). Typically H_p includes a profiled, queried contributor that we designate Q , who is replaced under H_d by an unprofiled individual X . Q may be an alleged offender, or a victim, while X is an

alternative, usually unknown, possible source of the DNA. It usually suffices to limit attention to H_p and H_d that differ only in replacing Q with X , otherwise the LR is difficult to interpret as a measure of the weight of evidence for Q to be a contributor of DNA.

In addition to reference profile(s), of Q and possibly other known contributors, the DNA evidence consists of one or more profiling runs performed on a DNA sample recovered from a crime scene, or from an item thought to have been present when the crime occurred. Each profiling run generates graphical results in an electropherogram (epg), which we assume has been interpreted by a forensic scientist who decides a list of alleles observed at each locus, and also a list of potential alleles about which there is substantial uncertainty, perhaps due to possible stutter. Alleles not on either list are regarded as unobserved in that run.

In low-template DNA (or LTDNA) profiling, each epg can be affected by stochastic effects such as dropout, dropout and stutter [1]. To help assess stochastic effects, it is common to perform multiple profiling runs, possibly varying the laboratory conditions but these are nevertheless referred to as replicates.

* Corresponding author.

E-mail addresses: c.steele.11@ucl.ac.uk (C.D. Steele), mgreenhalgh@cellmark.co.uk (M. Greenhalgh), d.balding@ucl.ac.uk (D.J. Balding).

Joint likelihoods for multiple replicates are obtained by assuming that the replicates are independent conditional on the genotypes of all contributors and parameters ϕ such as the amounts and degradation levels of DNA from each contributor [2]. We can write

$$\Pr(E|H) = \sum_j \Pr(\mathcal{G}_j) \prod_i \Pr(R_i|\mathcal{G}_j, \phi), \quad (2)$$

where R_i is the set of allele designations in the i th replicate run of the crime scene profile (CSP), \mathcal{G}_j denotes the j th set of contributor genotypes, and the summation is over all possible sets of contributor genotypes under H . $\Pr(\mathcal{G}_j)$ is computed under a standard population genetics model [1]. The unknown parameters ϕ can be replaced with estimates, or eliminated by maximisation or integration with respect to a prior distribution.

Currently, there are only limited possibilities to check the validity of an algorithm for evaluating an LTDNA LR (henceforth ltLR). One approach is to evaluate the ltLR when Q is repeatedly replaced by a random profile [3]. In that case H_p is false and we expect the majority of computed ltLRs to be small. Here, we propose to investigate a performance indicator for ltLR algorithms when H_p is true. Under H_d , it may occur that $\mathcal{G}_X = \mathcal{G}_Q$, where \mathcal{G}_X and \mathcal{G}_Q denote the genotypes of X and Q . This occurs with probability π_Q , the match probability for Q . Since $\Pr(E|H_d, \mathcal{G}_X = \mathcal{G}_Q) = \Pr(E|H_p)$, it follows that [4]

$$\begin{aligned} \text{ltLR} &= \frac{\Pr(E|H_p)}{\Pr(E|H_d, \mathcal{G}_X = \mathcal{G}_Q)\pi_Q + \Pr(E|H_d, \mathcal{G}_X \neq \mathcal{G}_Q)(1 - \pi_Q)} \\ &\leq \frac{1}{\pi_Q}. \end{aligned} \quad (3)$$

We will refer to $1/\pi_Q$ as the inverse match probability (IMP).

Consider first that Q is the major contributor to an LTDNA profile. Intuitively, if E implies that $\mathcal{G}_X = \mathcal{G}_Q$ then equality should be achieved in Eq. (3). The key idea of this paper is that if H_p is true then increasing numbers of LTDNA replicates should provide increasing evidence that $\mathcal{G}_X = \mathcal{G}_Q$, and so the ltLR should converge to the IMP. This holds even for mixtures if Q is the major contributor, since differential dropout rates should allow the alleles of Q to be identified from multiple replicates. However, any inadequacies in the underlying mathematical model or numerical approximations may become more pronounced with increasing numbers of replicates, preventing the ltLR from approaching the IMP. Therefore we propose to consider convergence of the ltLR towards the IMP as the number of replicates increases as an indicator of the validity of an algorithm to compute the ltLR when Q is the major contributor.

If Q is not the major contributor, even for many replicates there may remain ambiguity about the alleles of Q so that there remains a gap between the ltLR and IMP. However, the bound (3) still holds, and there is a useful guide to the appropriate value of the ltLR provided by the mixture LR for good-quality CSPs computed using only presence/absence of alleles [5]. If under H_p the contributors are Q and U , where U denotes an unknown, unprofiled individual, and H_d corresponds to two unknown contributors X and U , an example of a mixture LR is

$$\begin{aligned} \text{mixLR} &= \frac{\Pr(\text{CSP} = \text{ABC}, \mathcal{G}_Q = \text{AB}|\text{Q}, \text{U})}{\Pr(\text{CSP} = \text{ABC}, \mathcal{G}_Q = \text{AB}|\text{X}, \text{U})} \\ &= \frac{\Pr(\mathcal{G}_U \text{ is one of AC, BC, CC})}{\Pr((\mathcal{G}_X, \mathcal{G}_U) \text{ is one of (AA, BC), (AC, BB), (AB, CC), (AB, AC), (AB, BC), (AC, BC)})}, \end{aligned} \quad (4)$$

where within-pair ordering is ignored in the denominator. Under the standard population genetics model [6,7] and setting $F_{ST} = 0$,

the mixLR for this example is

$$\frac{\Pr(\text{CSP} = \text{ABC}, \mathcal{G}_Q = \text{AB}|\text{Q}, \text{U})}{\Pr(\text{CSP} = \text{ABC}, \mathcal{G}_Q = \text{AB}|\text{X}, \text{U})} = \frac{2p_A + 2p_B + p_C}{6p_A p_B (p_A + p_B + p_C)}, \quad (5)$$

where the p are population allele probabilities. As expected, $\text{mixLR} < \text{IMP} = 1/2p_A p_B$. See Ref. [8] for further details and examples. Note that the mixLR does not use peak height information.

Multiple LTDNA replicates should allow identification of all alleles present in any contributor, and hence the ltLR should reach the mixLR. In fact, ltLR will typically exceed mixLR because the alleles of different contributors may be distinguished over the multiple replicates through differential dropout rates. Indeed, Ref. [9] propose subsampling to generate different mixture ratios in low-template replicates as a strategy to assist mixture deconvolution. We cast light on this possibility below by considering a real CSP that has been profiled using multiple replicates at two different levels of sensitivity. More generally, we examine the behaviour of ltLR in relation to mixLR and IMP, and the utility of each of these for verifying the validity of ltLR computations.

`likeLTD` is an open-source R package that computes likelihoods for low-template DNA profiles [10]. `likeLTD` allows for the designation of epg peaks as uncertain in addition to the usual allelic/non-allelic classification, but does not directly use epg peak heights. Uncertain alleles are treated as if they were masked in calculation of the likelihood: the presence/absence of the allele is regarded as unknown. The effect of an uncertain call on calculation of the likelihood is illustrated in Table 1. When B is called as uncertain rather than absent and the hypothesised contributor has a B allele, a dropout term D is removed from the likelihood because the dropout status of B is unknown. We use `likeLTD` here both to confirm its good performance in computing ltLRs, and to illustrate the value of the IMP as a strict upper bound and the mixLR as an approximate lower bound. We apply `likeLTD` to lab-based profiling replicates, simulated replicates, and replicates obtained by re-sampling the five actual replicates of a real CSP.

Throughout this paper, ltLR, mixLR and IMP will be reported in units of bans, which is a base 10 logarithmic scale introduced as a measure of weight of evidence by Alan Turing during his wartime code breaking work [11]. Thus 6 bans corresponds to an LR of 1 million on the natural scale.

2. Materials and methods

2.1. Laboratory replicates

Cheek swab samples were obtained from five volunteers, and DNA was extracted using a PrepFiler Express BTA™ Forensic DNA Extraction Kit and the Life Technologies Automate Express™ Instrument as per the manufacturer's recommendations. The samples were then quantified using the Life Technologies Quantifiler® Human DNA Quantification kit as per the manufacturer's recommendations.

Table 1

Likelihood calculations for a CSP when the queried contributor Q has genotype AB and U indicates an allele designated as uncertain. L_p is the likelihood under the prosecution hypothesis, and D is the dropout probability. Under H_d are possible genotypes for the alternative contributor X , where Z is any other allele. L_d is the corresponding contribution to the likelihood under the defence hypothesis, where p_x is the probability of allele x , and D_2 is the homozygote dropout probability.

CSP	L_p	H_d	L_d
A	$D(1 - D)$	AA	$p_A^2(1 - D_2)$
		AZ	$2p_A(1 - p_A)D(1 - D)$
A[B]	$1 - D$	AA	$p_A^2(1 - D_2)$
		AB	$2p_A p_B(1 - D)$
		AZ	$2p_A(1 - p_A - p_B)D(1 - D)$

Table 2

Sample preparation and genotyping protocol for all conditions examined in the lab-based experiments (described in Table 3). Each condition was replicated eight times. The initial DNA concentration (column 3), dilution (column 4) and volume (column 5) generate approximately the DNA mass indicated in column 6. Columns 7 and 8 show the number of PCR cycles and the volume of PCR product added to each well for the genotyping. Columns 9 and 10 show the ratio of Hi-Di™ formamide to GeneSan™ 400HD ROX™ and the volume of the mixture added to each well. Apmr stands for as per manufacturers recommendations.

Condition	Contributor	Init. conc. (ng μL^{-1})	Dilution (%)	Volume (μL)	Mass (pg)	Cycles	Product (μL)	Formamide: ROX	F/ROX mixture (μL)
(i)	B	31.0	1	1.6	500	28	apmr	apmr	apmr
(ii)	B	31.0	0.1	2.0	60				
(iii)	B	31.0	0.01	5.0	15				
	A	23.0	1	17.6	500	28	apmr	apmr	apmr
(iv)	C	18.1	0.1	16	30				
	A	23.0	0.1	22.4	60	28	apmr	apmr	apmr
(v)	C	18.1	1	22.0	500				
	A	23.0	0.1	2.7	60	28	apmr	apmr	apmr
(vi)	B	31.0	0.1	2.0	60				
	C	18.1	0.1	3.5	60				
	A	23.0	0.1	2.7	60	28	1	600:1	9
(vii)	B	31.0	0.1	2.0	60				
	C	18.1	0.1	3.5	60				
	A	23.0	0.1	2.7	60	28	9	366:1	11
(viii)	B	31.0	0.1	2.0	60				
	C	18.1	0.1	3.5	60				
	A	23.0	0.1	2.7	60	30	apmr	apmr	apmr
(ix)	B	31.0	0.1	2.0	60				
	C	18.1	0.1	3.5	60				

Each sample was serially diluted on a log₁₀ scale, and then amplified using the AmpF Φ STR[®] SGM Plus[®] PCR kit as per the manufacturer's recommendations on a Veriti[®] 96-Well Fast Thermal Cycler.

An ABI 3130 Sequencer was used to analyse 1 μL of the PCR products, with 10 second injections at 3 kV; these settings were

used for all subsequent analyses. The results returned from the 3130 sequencer were analysed using GeneMapper[®] ID v3.2 to determine which samples were suitable for further use.

For the one-contributor investigation eight replicates of each of three conditions were created (Table 2). The conditions were created to investigate increasing dropout rate. For the 500 pg and

Table 3

Experimental conditions and hypotheses compared. pg denotes picograms and measures DNA mass; Pr(D) denotes the probability of dropout for a heterozygote allele, while Pr(C) denotes the probability of dropin. Pr(unc) indicates the probability of designating a CSP allele as uncertain. ν indicates the number of uncertain dropins per locus per replicate; see text for further details of "Condition". Q denotes the queried contributor, who is one of A, B or C as indicated in parentheses. X is an unknown alternative to Q under H_d , while U1 and U2 are unknown contributors under both H_p and H_d .

Study	# Contributors	Condition	H_p	H_d	
Lab-based	1	500 pg (i)	Q (B)	X	
		60 pg (ii)	Q (B)	X	
		15 pg (iii)	Q (B) + dropin	X + dropin	
	2	A=500 pg; C=30 pg (iv)	Q (A) + dropin	X + dropin	
			Q (A) + U1	X + U1	
			Q (C) + U1	X + U1	
		A=60 pg; C=500 pg (v)	Q (C) + dropin	X + dropin	
			Q (C) + U1	X + U1	
			Q (A) + U1	X + U1	
	3	28 cycles (vi)	Q (A) + U1 + U2	X + U1 + U2	
			Phase 1 (vii)	Q (A) + U1 + U2	X + U1 + U2
			Phase 2 (viii)	Q (A) + U1 + U2	X + U1 + U2
30 cycles (ix)			Q (A) + U1 + U2	X + U1 + U2	
Simulation	1	Pr _B (D) = 0; Pr(C) = 0	Q (B)	X	
		Pr _B (D) = 0.4; Pr(C) = 0.05	Q (B) + dropin	X + dropin	
		Pr _B (D) = 0.8; Pr(C) = 0.05	Q (B) + dropin	X + dropin	
		Pr(unc) = 0.8; $\nu \sim \text{Pois}(\lambda = 1)$	Q (B)	X	
		Pr(unc) = 0.4; $\nu \sim \text{Pois}(\lambda = 1)$	Q (B)	X	
	2	Pr _{A,C} (D) = {0.2, 0.8}; Pr(C) = 0	Q (A) + dropin	X + dropin	
			Q (A) + U1	X + U1	
			Q (C) + U1	X + U1	
		Pr _{A,C} (D) = {0.2, 0.6}; Pr(C) = 0	Q (A) + dropin	X + dropin	
			Q (A) + U1	X + U1	
			Q (C) + U1	X + U1	
	3	Pr _{A,B,C} (D) = {0.8, 0.5, 0.2}; Pr(C) = 0	Q (A) + U1 + U2	X + U1 + U2	
		Pr _{A,B,C} (D) = {0.5, 0.5, 0.5}; Pr(C) = 0	Q (A) + U1 + U2	X + U1 + U2	
		Pr _{A,B,C} (D) = {0.2, 0.5, 0.8}; Pr(C) = 0	Q (A) + U1 + U2	X + U1 + U2	
	Real-world	≥ 3	Standard and sensitive	Q + U1 + U2	X + U1 + U2
Standard only			Q + U1 + U2	X + U1 + U2	
Sensitive only			Q + U1 + U2	X + U1 + U2	

Table 4

Five replicates of a crime scene profile, three from a sensitive LTDNA profiling technique and two from standard DNA profiling. Alleles shown in [] were called as uncertain.

Locus	Sensitive profiling			Standard profiling	
	Run 1	Run 2	Run 3	Run 4	Run 5
D3	16, [15]	16, [15]	16, 18, [15]	16	16
vWA	15, 16, [17]	15, [14]	15, 18, [14]	15	15
D16	9	9	9, 11, [10]	9	9
D2	17, 19, 24	16, 17, 24, [23]	17, [16]	24	24
D8	8, 13, 15, 16	8, 12, 13, 16, [15]	8, 13, 14, 16, [15]	[8]	
D21	30, 32, 33.2	32, 32.2, 33.2	32, 32.2, 33.2, 34, [31]	[32], [32.2]	[33.2]
D18	12, 17	12, 17, 19	12, 17, [11], [16]	[17]	17
D19	14, 21, [13]	11, 14, [13]	14, [13]	14	14
TH01	6, 9.3	6, 9.3	6, 8, 9.3	[6], [9.3]	[6]
FGA	21	21, [20]	21, 20	21	

60 pg conditions, one-contributor hypotheses were compared, B under H_p and X under H_d , while for the 15 pg condition dropout was also modelled under both hypotheses (Table 3).

For the two-contributor investigation eight replicates of each of two conditions were created (Table 2). The major and minor contributors were reversed between conditions, with an increased DNA contribution from the minor. These samples were amplified and analysed as described previously. Two-contributor hypotheses were compared, with each of A and C in turn playing the role of Q, while the other contributor was treated as unknown. Additionally one-contributor-plus-dropin hypotheses were compared, with only the major contributor playing the role of Q (Table 3).

For the three-contributor investigation eight replicates of each of four conditions were created (Table 2). The conditions were created to investigate different profiling protocols. The Phase 1 and Phase 2 conditions are post-PCR purification protocols designed to enhance the sensitivity of detection of the standard protocol [12], and both involve concentrating the post-PCR product using an Amicon[®] PCR microcon unit according to the manufacturer's recommendations. Phase 1 enhancement increases the amount of formamide in the mixture compared to the manufacturer's recommendations, while Phase 2 enhancement increases the amount of DNA, formamide and ROX compared to Phase 1. For all four conditions (30 cycles, 28 cycles, Phase 1, and Phase 2), three-contributor hypotheses were compared, with A playing the role of Q and the other contributors treated as unknown (Table 3). Dropout was not modelled under either hypothesis, although dropout was included in the simulations. This reflects a realistic challenge for

few replicates with multiple contributors, whereby any dropout alleles may be wrongly attributed to one of the contributors. However the incorrect model will lead to deterioration of inferences for larger numbers of replicates.

2.2. Simulated replicates

All of the conditions that we now describe were simulated in eight replicates, with the whole simulation being performed five times. Initially a number of single-contributor CSPs were simulated using the profile of individual B. The first condition investigated was a "perfect match", in which all eight replicates generated exactly the profile of B. Next, we introduced mild dropout ($\Pr(D) = 0.4$) and severe dropout ($\Pr(D) = 0.8$) of the alleles of B, in each case with dropins included at rate $\Pr(C) = 0.05$ (at most one dropin per locus per replicate). The homozygous dropout probability was set equal to $\Pr(D)^2/2$, as suggested by [13]. We then examined the effect of uncertain allele designations by randomly designating some alleles of B as uncertain, first with $\Pr(\text{unc}) = 0.4$ and then $\Pr(\text{unc}) = 0.8$. In both conditions, at each locus and in each replicate a Poisson mean one number of alleles not in the profile of B was also designated as uncertain, with types randomly selected according to frequencies in the UK Caucasian database. For all these simulated profiles, one-contributor hypotheses were compared, B under H_p and X under H_d .

Next two-contributor CSPs were simulated, based on the profiles of A and C. Two conditions were simulated, both used $\Pr_A(D) = 0.2$, while $\Pr_C(D)$ was initially 0.8 and then 0.6. Dropout was not




Fig. 1. The L_r shown on a logarithmic scale (in bans) from one-contributor CSPs evaluated using from one up to eight replicates. Left: lab-based replicates, with DNA template (in pg) as shown in the legend box. Middle: simulated replicates with dropout (probability $\Pr(D)$) and dropin (probability $\Pr(C)$); the plotted points represent the median from five repetitions of the simulation, and the vertical bars show the range. Right: simulated replicates with uncertain allele calls (probability $\Pr(\text{unc})$) for a true allele to be uncertain, and a Poisson (rate λ) number of non-alleles labelled as uncertain at each locus.

simulated. For shared alleles the dropout probability was the product of the dropout probabilities for each contributor having that allele. Two-contributor hypotheses were compared, with each of A and C in turn taking the role of Q, while the other was treated as unknown in the analysis. Additionally one-contributor-plus-dropin hypotheses were compared, only for A playing the role of Q (Table 3).

Three-contributor CSPs were then simulated under three conditions, with dropout probabilities for Donors A, B and C as shown in Table 3. Dropin was included as for the one-contributor simulations. Three-contributor hypotheses were compared, with A playing the role of Q and the other two contributors being treated as unknown.

2.3. Crime case replicates

We used a CSP from an actual crime investigation, consisting of five replicates: two using standard SGM+ profiling and three generated using an LCN protocol with 34 PCR cycles (Table 4). This example was submitted to us for *likeLTD* analysis, and as is typical only limited information about the profiling protocol was provided by the profiling lab. These details are not required by *likeLTD* because it estimates the unknown parameters from the CSP allele designations. We re-sampled the five actual replicates to generate simulated profiles with up to eight replicates, consisting of standard replicates only, sensitive replicates only, or both. Six distinct alleles were observed at locus D8, but no more than three




Fig. 2. The low-template likelihood ratio (ItLR) from two-contributor CSPs profiled at up to eight replicates. Left: lab-based replicates, with the DNA template from the minor contributor greater in the lower panel (see legend boxes). Right: simulation-based replicates, with the minor contributor having reduced dropout in the lower panel. The simulated CSPs were generated from the profiles of Donors A and C, and the line colours on the graph indicate whether the queried individual (Q) is A (blue) or C (red). Solid lines indicate a two-contributor analysis, with the non-Q individual regarded as unknown (U1). Dashed lines indicate a one-contributor analysis that also allows for dropin (only for Q the major contributor). The inverse match probability is shown with dot-dash lines, coloured according to Q. The mixLR is shown with dotted lines, coloured according to Q. In the legend boxes, H indicates the hypotheses with X an unknown alternative to Q, and Pr(D) indicates the probability of dropout.

replicated alleles were observed at any locus. Three-contributor hypotheses were compared, with all contributors unknown under H_d , and no dropout (Table 3).

3. Results

3.1. One contributor

3.1.1. Lab-based

For the good-template experiments (500 pg), Fig. 1 (left) shows that the ItLR equals the IMP for all numbers of replicates (one through eight). This is the expected result, and the exercise shows that in this simple setting there is no deterioration in the quality of the computed LR for large numbers of replicates. Low DNA template (60 pg) generates an ItLR about 1.6 bans below the IMP for one replicate, but the gap is very small for two replicates and is negligible for larger numbers of replicates. For very low DNA template (15 pg) the ItLR is just under 6 bans for a single replicate, about 6 bans below the IMP. Replicate profiling substantially narrows the gap, but does not completely close it, with a difference of about 3 decibans remaining at eight replicates.

3.1.2. Simulation

The corresponding simulation studies show broadly similar trends to the lab-based data. For both the perfect match ($\text{Pr}(D) = 0$) and mild dropout ($\text{Pr}(D) = 0.4$) conditions, the median ItLR rapidly reaches the IMP but does not exceed it, while under severe dropout ($\text{Pr}(D) = 0.8$) the median ItLR rises towards the IMP but does not reach it (Fig. 1, middle). For the low and high rates of uncertain calls, the IMP is approximately reached at a five and eight replicates, respectively (Fig. 1, right).

3.2. Two contributors

3.2.1. Lab-based

When the minor contributor provides only 30 pg of DNA (Fig. 2, top left panel), then if Q is the major contributor the ItLR is very close to the IMP for all numbers of replicates, whereas if Q is the minor contributor then there remains a substantial gap between ItLR and IMP even at eight replicates. However, even with this very low template, the ItLR exceeds the mixLR beyond five replicates. When the major and minor contributors are reversed, and the amount of DNA from the minor is doubled (Fig. 2, bottom left), then if Q is the minor contributor the ItLR substantially exceeds mixLR from six replicates and rises to within two bans of the IMP at eight replicates. Under both conditions, the two-contributor

analysis gives a very similar result to the one-contributor-with-dropin analysis.

3.2.2. Simulation

When the minor contributor is subject to high dropout (Fig. 2, top right), then if Q is the major contributor the ItLR exceeds the mixLR after one replicate, and rises rapidly to within about 2 bans of the IMP, but the gap narrows only slowly thereafter. The one-contributor-plus-dropin analysis gives an ItLR that is broadly similar to the two contributor analysis, but with a wider range indicating greater variability. If Q is the minor contributor, the median ItLR increases rapidly from a low base, and appears to stabilise after about five replicates, about four bans below the IMP but exceeding the mixLR. The range increases after three replicates, and remains high up to eight replicates.

With reduced dropout for the minor contributor (Fig. 2, bottom right), inferring the presence of a major contributor Q is harder because of additional masking by the minor contributor. The median ItLR in both the two contributor and one-contributor-plus-dropin analyses eventually reaches within 2 bans of the IMP, with the latter showing a greater range. Conversely, the lower dropout rate leads to improved inference for a minor contributor Q, with the median ItLR rising to about three bans below the IMP at eight replicates, and exceeding the mixLR from four replicates. Interestingly, after six replicates the range of the minor contributor ItLR overlaps the range for the major contributor.

3.3. Three contributors

3.3.1. Lab-based

The 30 PCR cycles condition gives the highest ItLR at one replicate but little improvement with additional replicates (Fig. 3, left). The other amplification methods do show an increasing ItLR trend with additional replicates, but in no case did the ItLR reach within four bans of the IMP. As expected, the ItLR for both phase 1 and phase 2 enhancement exceeds that for standard 28 PCR cycles at all numbers of replicates, and phase 2 enhancement ItLR typically gives a small improvement over phase 1 enhancement. For 30 PCR cycles, the ItLR exceeds the mixLR for a single replicate but dips slightly below it at six replicates. For the other conditions, the mixLR is always exceeded from four replicates.

3.3.2. Simulation

All three curves in Fig. 3 (middle) show an increasing trend with number of replicates, with the median ItLR being in the expected order throughout (decreasing ItLR with increasing




Fig. 3. The low-template likelihood ratio (ItLR) for three-contributor crime stains profiled with one to eight replicates. Left: laboratory replicates using four lab techniques indicated in the legend box and described further in Section 2. Middle: simulated replicates with dropout rates for the three contributors as shown in the legend box against $\text{Pr}(D)$, the first value being for the queried contributor. $\text{Pr}(C)$ is the dropout probability. Right: re-sampled actual crime-stain replicates; the original data are two standard profiling replicates, and three replicates using enhanced sensitivity. The ItLR returned from a perfect replicate of the contributors (consisting of every allele from each contributor) is shown with dotted lines; this is not possible for the real-world case, as the true contributors are unknown.

dropout for Q). The median ItLR exceeds the mixLR after one replicate (low dropout), after two replicates (medium dropout) and after four replicates (high dropout). The range is often wide, reflecting a strong dependence of the ItLR on the details of the simulation (in particular the number of alleles shared across contributors).

3.3.3. Real-world case

The ItLR returned when only standard or only sensitive replicates are used shows a similar trend, but nearly five bans lower for the standard replicates (Fig. 3, right). For three or more replicates, using mixed types of replicates is superior even to only using sensitive replicates, coming to within two bans of the IMP. This partly reflects the limited pool of replicates used in the actual crime case, but suggests that using different sensitivities in the profiling replicates may convey an advantage due to different contributors being better distinguished.

4. Discussion

We have shown that ItLR computed by `likeLTD` is bounded above by the IMP in every condition considered, as predicted by theory (Eq. (3)). That the bound is often tight when Q is the major contributor (Figs. 1 and 2 (top)) supports the validity of the underlying mathematical model, and its correct implementation in the `likeLTD` software. Our results should help counter any misconception that combining multiple noisy profiling replicates only compounds the noise: in fact, multiple noisy replicates can fully recover the genotype of a contributor [14].

A novel feature of `likeLTD`, is that it can accommodate uncertain allele designations, which diminishes the problem of an all-or-nothing allele call, therefore mitigating the problem highlighted by [15] of choosing a detection threshold. We have shown (Fig. 1 (right)) that introducing many uncertain allele calls leads to ItLRs that satisfy the bound, which is reasonably tight with as few as three replicates even when 80% of true alleles are designated as uncertain and there are also multiple uncertain non-alleles.

We have further shown that mixLR, the LR computed from knowing every allele that is represented in the profile of at least one contributor to the CSP, is often surpassed after only a handful of replicates. Then, multiple LTDNA replicates provide stronger evidence than a single good quality profile correctly representing the alleles of all contributors, which occurs because the alleles of different contributors can to some extent be distinguished through differential dropout rates in multiple replicates. These results lend support in principle to the proposal of [9].

Fig. 2 shows that, for two-person mixtures, the analysis assuming one-contributor-plus-dropin gave a very good approximation for the lab-based replicates (left panels), and a reasonably good approximation for the simulation replicates, but with more variable ItLR values, as indicated by the wider range.

4.1. Choice of profiling technique

We generated three-contributor CSPs in order to compare different LTDNA profiling techniques. We chose the most challenging condition in which all three contribute the same DNA template, making it impossible to deconvolve the mixture into the genotypes of individual contributors. We found that PCR performed with 28 cycles (regardless of enhancement) is preferable to 30 cycle PCR beyond one replicate (Fig. 3). More PCR cycles introduces more stochasticity in the results, as stated in the AmpF ℓ STR[®] SGM Plus[®] PCR Amplification Kit user guide. We found that enhancement of the post-PCR sample is advantageous, with Phase 2 enhancement providing a small further improvement

over Phase 1 (Fig. 3). These results support those of Forster et al. [16], who demonstrated that increasing PCR cycles increases the size of stutter peaks and the incidence of dropin; we observed no improvement in the WoE for 30 PCR cycles, possibly due to these stochastic effects.

The results from the real crime case (Fig. 3, right) suggest that if possible, a mixture of LTDNA replicates with differing sensitivities should be employed, as this allows better discrimination between the alleles of different contributors and hence a higher ItLR than the same number of replicates all using the same sensitivity.

4.2. Use of replicates

Splitting the sample reduces the quality of results expected in each replicate compared with that which would be obtained from a single profiling run using all available DNA. Grisedale and van Daal [17] favour use of a single run, but their comparison was with a consensus sequence obtained from multiple replicates, rather than the more efficient statistical analysis available through analysing individual replicates. Our results show increasing information obtained from additional replicates, which may tilt the argument towards use of multiple replicates but we have not done a comparison directly addressing this question. To fully test the performance of `likeLTD` in relation to mixLR and IMP we have used up to eight replicates. Taberlet et al. [18] suggest seven replicates to generate a quality profile when the amount of DNA is low, but this many replicates is rarely available for low-template crime samples [15].

Acknowledgements

CDS is funded by a PhD studentship from the UK Biotechnology and Biological Sciences Research Council and Cellmark Forensic Services.

References

- [1] C.D. Steele, D.J. Balding, Statistical evaluation of forensic DNA profile evidence, *Annu. Rev. Stat. Appl.* 1 (2014) 20–21. , <http://dx.doi.org/10.1146/annurev-statistics-022513-115602>.
- [2] J.M. Curran, P. Gill, M.R. Bill, Interpretation of repeat measurement DNA evidence allowing for multiple contributors and population substructure, *Forensic Sci. Int.* 148 (1) (2005) 47.
- [3] P. Gill, H. Haned, A new methodological framework to interpret complex DNA profiles using likelihood ratios, *Forensic Sci. Int. Genet.* (2012), <http://dx.doi.org/10.1016/j.fsigen.2012.11.002>.
- [4] R.G. Cowell, T. Graverson, S. Lauritzen, J. Mortera, Analysis of DNA mixtures with artefacts, in: ArXiv e-Prints, 2013, February.
- [5] B.S. Weir, C.M. Triggs, L. Starling, K.A.J. Stowell, J. Buckleton, Interpreting DNA mixtures, *J. Forensic Sci.* 42 (1997) 213–222.
- [6] P. Gill, C.H. Brenner, J.S. Buckleton, A. Carracedo, M. Krawczak, W.R. Mayr, N. Morling, M. Prinz, P.M. Schneider, B.S. Weir, DNA commission of the International Society of Forensic Genetics: recommendations on the interpretation of mixtures, *Forensic Sci. Int.* 160 (2) (2006) 90–101.
- [7] P. Gill, L. Gusm ao, H. Haned, W.R. Mayr, N. Morling, W. Parson, L. Prieto, M. Prinz, H. Schneider, P.M. Schneider, B.S. Weir, DNA commission of the International Society of Forensic Genetics: recommendations on the evaluation of STR typing results that may include drop-out and/or drop-in using probabilistic methods, *Forensic Sci. Int. Genet.* (2012), <http://dx.doi.org/10.1016/j.fsigen.2012.06.002>.
- [8] D.J. Balding, *Weight-of-Evidence for Forensic DNA Profiles*, Wiley, 2005.
- [9] J. Ballantyne, E.K. Hanson, M.W. Perlin, DNA mixture genotyping by probabilistic computer interpretation of binomially-sampled laser captured cell populations: combining quantitative data for greater identification information, *Sci. Justice* 53 (2) (2013) 103–114. , <http://dx.doi.org/10.1016/j.scijus.2012.04.004>, ISSN 1355-0306.
- [10] D.J. Balding, Evaluation of mixed-source, low-template DNA profiles in forensic science, *Proc. Natl. Acad. Sci. U. S. A.* (2013), <http://dx.doi.org/10.1073/pnas.1219739110>.
- [11] J. Good Irving, Studies in the history of probability and statistics. xxxvii am Turing's statistical work in world war ii, *Biometrika* 66 (2) (1979) 393–396.
- [12] A.D. Roeder, P. Elsmore, M. Greenhalgh, A. McDonald, Maximizing DNA profiling success from sub-optimal quantities of DNA: a staged approach, *Forensic Sci. Int. Genet.* 3 (2) (2009) 128–137. , <http://dx.doi.org/10.1016/j.fsigen.2008.12.004>.
- [13] D.J. Balding, J. Buckleton, Interpreting low template DNA profiles, *Forensic Sci. Int. Genet.* 4 (1) (2009) 1–10. , <http://dx.doi.org/10.1016/j.fsigen.2009.03.003>.

- [14] L. Schneps, C. Colmez, *Math on trial: how numbers get used and abused in the courtroom*, Wiley Online Library, 2013.
- [15] B. Budowle, A.J. Eisenberg, A. van Daal, Validity of low copy number typing and applications to forensic science, *Croat. Med. J.* 50 (3) (2009) 207–217. , <http://dx.doi.org/10.3325/cmj.2009.50.207>.
- [16] L. Forster, J. Thomson, S. Kutranov, Direct comparison of post-28-cycle PCR purification and modified capillary electrophoresis methods with the 34-cycle “low copy number”(LCN) method for analysis of trace forensic DNA samples, *Forensic Sci. Int. Genet.* 2 (4) (2008) 318–328. . <http://dx.doi.org/10.1016/j.fsi-gen.2008.04.005>.
- [17] K.S. Grisedale, A. van Daal, Comparison of STR profiling from low template DNA extracts with and without the consensus profiling method, *Invest. Genet.* 3 (2012) 14, <http://dx.doi.org/10.1186/2041-2223-3-14>.
- [18] P. Taberlet, S. Griffin, B. Goossens, S. Questiau, V. Manceau, N. Escaravage, L.P. Waits, J. Bouvet, Reliable genotyping of samples with very low DNA quantities using PCR, *Nucleic Acids Res.* 24 (16) (1996) 3189–3194.