



Convergent series for lattice models with polynomial interactions

Aleksandr S. Ivanov^{a,b}, Vasily K. Sazonov^{c,*}

^a *M. V. Lomonosov Moscow State University, Faculty of Physics, Leninskie Gory, 119991, Moscow, Russia*

^b *Institute for Nuclear Research RAS, 60-letiya Oktyabrya prospekt 7a, 117312, Moscow, Russia*

^c *Institut für Physik, FB Theoretische Physik, Universität Graz, Universitätsplatz 5, A-8010, Graz, Austria*

Received 10 June 2016; received in revised form 31 October 2016; accepted 1 November 2016

Available online 9 November 2016

Editor: Hubert Saleur

Abstract

The standard perturbative weak-coupling expansions in lattice models are asymptotic. The reason for this is hidden in the incorrect interchange of the summation and integration. However, substituting the Gaussian initial approximation of the perturbative expansions by a certain interacting model or regularizing original lattice integrals, one can construct desired convergent series. In this paper we develop methods, which are based on the joint and separate utilization of the regularization and new initial approximation. We prove, that the convergent series exist and can be expressed as re-summed standard perturbation theory for any model on the finite lattice with the polynomial interaction of even degree. We discuss properties of such series and study their applicability to practical computations on the example of the lattice ϕ^4 -model. We calculate $\langle \phi_n^2 \rangle$ expectation value using the convergent series, the comparison of the results with the Borel re-summation and Monte Carlo simulations shows a good agreement between all these methods.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The main objects of studies in quantum field theory are the Green's functions. They can be naturally expressed in terms of the path integral. However, the path integral itself gives only a

* Corresponding author.

E-mail addresses: ivanov.as@physics.msu.ru (A.S. Ivanov), vasily.sazonov@uni-graz.at (V.K. Sazonov).

formal solution and for the practical computations one has to find some efficient method of its evaluation. One of the possibility is to use a finite lattice approximation of the physical space and then investigate the truncated system using the Monte Carlo method. Such approach was successfully applied in many areas of statistical, condensed matter and elementary particle physics for a wide range of physical parameters, *e.g.* coupling constants. The Monte Carlo method is based on the probabilistic interpretation of the lattice integrals measure and, consequently, it can not be applied, when the action of the system is complex. For instance, this complex action problem, or in other words sign problem, impedes the computations in lattice QCD at finite chemical potentials [1]. Another traditional way of the path and lattice integrals computations is the utilization of standard perturbation theory with the Gaussian initial approximation (SPT). Usually, the series of SPT are asymptotic [2,3] and applicable only for the small coupling constants (parameters of the expansion). For example, the most precise results within standard perturbation theory were obtained in QED [4,5], where the expansion parameter is extremely small: $\alpha \simeq 1/137$. At the same time SPT fails to describe the low energy physics of graphene, which is effectively determined by QED with large coupling constant [6].

The SPT series are asymptotic because of the incorrect interchange of the summation and integration during the construction of the perturbative expansion. Nevertheless, the problem of the interchanging can be avoided. For instance, it can be done by an appropriate regularization of the original integral. The regularization by cutting off the large fluctuations of the fields in the lattice models was suggested in [7,8]. In [9–13] a special integral transform of the interaction part of the action and consequent regularization were applied to construct convergent perturbation theory for lattice models and path integrals with Gaussian measure defined by trace-class operators. The method, based on the modification of the interaction power due to the application of the intermediate field representation and on the consequent utilization of the forests/trees formula [14,15] was developed in works [16–20]. Alas, the computations with all these methods are highly complicated.

An alternative approach to the construction of the convergent perturbative series in scalar field theories, based on changing of the initial Gaussian approximation to a certain interacting theory, was proposed in [21–23]. Later, independently developed and similar ideas become a basis of the variational perturbation theory methods [24–26]. In [27] the approach [21–23] was extended to the construction of the strong coupling expansion for the anharmonic oscillator. In [28] the RG-equations consistent with the method [21] were derived. The critical indices of the ϕ^4 -model obtained within the latter RG-equations are in close agreement with the numerical results and corresponding experimental measurements for the liquid–gas transition, He₄ and binary mixtures [28,29]. However, a rigorous mathematical proof of the convergence of expansions [21–23, 28] is still missing. The main assumption in [21–23,25,26,28] is related to the applicability of the dimensional regularization [30], which is used to handle the limit of the infinite number of degrees of freedom during the scaling transformations of the fields. Nevertheless, the restriction to a finite amount of degrees of freedom allows one to carry out the derivation of the convergent series similar to [22,23] mathematically rigorously. A first step in this direction was done [31], where the one-dimensional lattice ϕ^4 -model was considered. The numerical computations within the convergent series in [31] revealed a perfect agreement with the results obtained by the Monte Carlo simulations for the lattice with the volume $V = 2$ and demonstrated slow convergence to the correct answers even for the slightly bigger lattices (with $V = 4$ and $V = 8$ lattice sites).

Here we continue the investigation of the convergent series method and present a solution of the slow convergence problem. We carry out main derivations on the example of the lattice ϕ^4 -model and perform generalizations, when it is necessary. We prove, that the convergent series

(CS) can be rigorously constructed for any finite lattice model with the polynomial interaction of even degree and that the series is expressed as a re-summation of SPT. We show, that up to all loops of standard perturbation theory the convergent series method has an internal symmetry, providing the possibility to introduce a variational parameter. The freedom in a choice of the variational parameter allows one to obtain numerical results, which are in agreement with the Monte Carlo data. Though, the convergence and non-perturbative correctness of CS, modified by the variational parameter, is not *a priori* evident. In the following we denote the latter series as variational (VS). To investigate the convergence of the variational series, we consider two regularizations of the lattice ϕ^4 . The first one, η -regularization, is a natural extension of the variational series construction and it gives suggestive arguments about the convergence of VS. The second, γ -regularization, is also based on the mathematical structures, used in VS. We prove, that γ -regularized model approximates original ϕ^4 -model with any arbitrary precision and that the Green's functions of the γ -regularized model can be calculated with the variational series, which is convergent in this case. We demonstrate the non-perturbative independence on the variational parameter of VS, when the γ -regularization is removed. Using this independence property, we propose a way for the computations in the infinite volume limit. Summarizing the properties of CS and VS with regularizations, we conjecture the convergence of the variational series. We study the applicability of the CS and VS for different lattice volumes and investigate the dependence on the variational and regularization parameters, computing the operator $\langle \phi_n^2 \rangle$ of the lattice ϕ^4 -model. The results are compared with the Monte Carlo simulations and Borel re-summation.

The paper is organized as follows. In Section 2 the convergent series for lattice models with polynomial interactions is constructed. We introduce variational series and discuss its perturbative properties in Section 3. To investigate non-perturbative aspects of VS we study regularized lattice models in Section 4. The results of the numerical computations are presented in Section 5. We conclude in Section 6.

2. Construction of the convergent series

We start with the construction of the convergent series for the lattice ϕ^4 -model. The model is defined by the action

$$S[\phi] = \frac{1}{2} \sum_{m,n=0}^{V-1} \phi_m K_{mn} \phi_n + \frac{\lambda}{4!} \sum_{n=0}^{V-1} \phi_n^4, \quad (1)$$

$$\frac{1}{2} \sum_{m,n=0}^{V-1} \phi_m K_{mn} \phi_n = \sum_{n=0}^{V-1} \left[\frac{1}{2} M^2 \phi_n^2 - \frac{1}{2} \sum_{\mu=1}^d (\phi_{n+\hat{\mu}} + \phi_{n-\hat{\mu}} - 2\phi_n) \phi_n \right], \quad (2)$$

where M is a mass parameter, λ is a coupling constant, V is the volume of the lattice, indices m and n label the lattice sites, $d = 1, 2$ is a dimension of the lattice, index μ runs over all spatial dimensions and $\hat{\mu}$ stands for the unit vector in the corresponding direction. The periodical boundary conditions in all possible directions are assumed.

Without loss of generality, as an example of an arbitrary Green's function, we consider the two point Green's function (propagator). The normalized to the free theory propagator is defined as

$$\langle \phi_i \phi_j \rangle = \int [d\phi] \phi_i \phi_j \exp\{-S[\phi]\}, \quad (3)$$

where $\int [d\phi] = \frac{1}{Z_0} \prod_n \int d\phi_n$ and

$$Z_0 = \prod_n \int d\phi_n e^{-\frac{1}{2} \sum_{l,m} \phi_l K_{lm} \phi_m} \quad (4)$$

is the partition function of the free theory. Following [22,23], we split the action into the new non-perturbed part $N[\phi]$ and perturbation: $S[\phi] = N[\phi] + (S[\phi] - N[\phi])$. For the following it is convenient to modify the latter expression as

$$S_\eta[\phi] = N[\phi] + \eta(S[\phi] - N[\phi]), \quad (5)$$

where $\eta \leq 1$ is a parameter labeling the order of the new perturbative expansion. The parameter η will be also used in Section 4.1 for the regularization. In the current section we derive formulas containing η , keeping in mind, that the initial model corresponds to $\eta = 1$. Then, the propagator (3) can be written as

$$\langle \phi_i \phi_j \rangle = \int [d\phi] \phi_i \phi_j e^{-N[\phi]} \sum_{l=0}^{\infty} \frac{\eta^l (N[\phi] - S[\phi])^l}{l!}. \quad (6)$$

When

$$N[\phi] \geq S[\phi], \quad (7)$$

the interchanging of the summation and integration in (6) leads to an absolutely convergent series

$$\langle \phi_i \phi_j \rangle = \sum_{l=0}^{\infty} \langle \phi_i \phi_j \rangle_l \quad (8)$$

with terms given by

$$\langle \phi_i \phi_j \rangle_l = \frac{\eta^l}{l!} \int [d\phi] \phi_i \phi_j (N[\phi] - S[\phi])^l e^{-N[\phi]}. \quad (9)$$

Indeed, it is easy to see, that

$$\begin{aligned} \left| \sum_{l=0}^{\infty} \langle \phi_i \phi_j \rangle_l \right| &\leq \sum_{l=0}^{\infty} |\langle \phi_i \phi_j \rangle_l| \leq \sum_{l=0}^{\infty} \frac{\eta^l}{l!} \int [d\phi] |\phi_i \phi_j| (N[\phi] - S[\phi])^l e^{-N[\phi]} \\ &= \int [d\phi] |\phi_i \phi_j| e^{-S_\eta[\phi]} < \infty, \quad \text{for } \eta \leq 1. \end{aligned} \quad (10)$$

There are many possibilities to choose the new initial approximation $N[\phi]$, satisfying inequality (7), however it should correspond to a solvable model. Here we take

$$N[\phi] = \sum_{n,m} \frac{1}{2} \phi_n K_{nm} \phi_m + \sigma \left(\sum_{n,m} \frac{1}{2} \phi_n K_{nm} \phi_m \right)^2, \quad (11)$$

where σ is an unknown positive parameter, which is determined by the substitution of (11) into (7):

$$\sigma \geq \frac{\lambda}{6M^4}. \quad (12)$$

The functions (9) can be calculated in the following way. Introducing an auxiliary integration, we change $\|\phi\| \equiv (\frac{1}{2} \phi_n K_{nm} \phi_m)^{\frac{1}{2}}$ to the one-dimensional variable t

$$\langle \phi_i \phi_j \rangle_l = \frac{\eta^l}{l!} \int [d\phi] \phi_i \phi_j \int_0^\infty dt e^{-t^2 - \sigma t^4} \delta(t - \|\phi\|) \left(\sigma t^4 - \frac{\lambda}{4!} \sum_n \phi_n^4 \right)^l. \tag{13}$$

Rescaling the field variables as $\phi_n^{old} = t \phi_n$, we get

$$\langle \phi_i \phi_j \rangle_l = J_\eta(V, l) \int [d\phi] \phi_i \phi_j \delta(1 - \|\phi\|) \left(\sigma - \frac{\lambda}{4!} \sum_n \phi_n^4 \right)^l, \tag{14}$$

where

$$[ll]J_\eta(V, l) = \frac{\eta^l}{l!} \int_0^\infty dt e^{-t^2 - \sigma t^4} t^{V+4l+1}. \tag{15}$$

The factor t^{V+4l-1} in the integrand in $J_\eta(V, l)$ is obtained by rescaling of the fields measure, fields and delta function. Now the multi-dimensional (lattice) part of the integral is factorized from the auxiliary integration. Applying the binomial expansion to the brackets $(\dots)^l$, we rewrite (14) as

$$\langle \phi_i \phi_j \rangle_l = J_\eta(V, l) \int [d\phi] \phi_i \phi_j \delta(1 - \|\phi\|) \sum_{k=0}^l C_l^k \sigma^{l-k} \left(-\frac{\lambda}{4!} \sum_n \phi_n^4 \right)^k. \tag{16}$$

To solve (16), we use the following equality¹

$$\int [d\phi] \phi_{n_1} \dots \phi_{n_Q} e^{-\|\phi\|^2} = \frac{1}{2} \Gamma\left(\frac{V+Q}{2}\right) \int [d\phi] \phi_{n_1} \dots \phi_{n_Q} \delta(1 - \|\phi\|). \tag{17}$$

By substituting (17) in (16), we obtain

$$\langle \phi_i \phi_j \rangle_l = J_\eta(V, l) \sum_{k=0}^l C_l^k \sigma^{l-k} \frac{2}{\Gamma\left(\frac{V+4k+2}{2}\right)} \int [d\phi] \phi_i \phi_j e^{-\|\phi\|^2} \left(-\frac{\lambda}{4!} \sum_n \phi_n^4 \right)^k. \tag{18}$$

Denoting the k -th order of standard perturbation theory as f_k , we rewrite (18) as

$$\langle \phi_i \phi_j \rangle_l = J_\eta(V, l) \left[\sum_{k=0}^l C_l^k \sigma^{l-k} \frac{2k! f_k}{\Gamma\left(\frac{V+4k+2}{2}\right)} \right]. \tag{19}$$

Therefore, each certain order l of the convergent series is expressed as a linear combination of first l orders of SPT with coefficients given by the one-dimensional analytically calculable t -depending integrals. However, the latter fact does not mean that the convergent series loses non-perturbative contributions from non-analytical functions such as $e^{-\frac{1}{\lambda}}$. Being an expansion with the non-Gaussian initial approximation, it automatically takes non-perturbative contributions into account in a similar way, as the function $e^{-\frac{1}{\lambda}}$ for positive λ can be reproduced by its Taylor series around $\lambda = 1$.

The lattice ϕ^4 is a Borel summable model, however, it is possible to generalize the results of the current section to a wider class of models, which include Borel non-summable cases.

¹ The r.h.s. of the identity (17) is obtained from the left one by the transformations analogous to (13), (14).

Proposition 1. Consider a model on the finite lattice, determined by the polynomial action $S[\phi] = P[\phi]$ with an even $\text{deg}(P)$, then it is always possible to construct a convergent series for this model with the terms which can be expressed as a linear combinations of terms of standard perturbation theory.

The proof follows from the construction presented above and from the fact, that each polynomial can be bounded as

$$|P[\phi]| \leq \text{const} (1 + \|\phi\|^{\text{deg}(P)}). \tag{20}$$

Moreover, it is possible to show, that convergent series is a re-summation of SPT, for details see Section 3.

The latter proposition demonstrates, that the whole non-perturbative physics of the lattice models with the polynomial actions can be encoded by the coefficients of standard perturbation theory. However, it is important to note, that this does not mean, that the non-perturbative information can be obtained only from the standard perturbative expansion. An additional input, needed for the construction of the convergent series is received from the model itself. In some sense it is similar to the resurgence program [32], where for the recovering of the non-perturbative contributions from the perturbative series, it is assumed that the solution is a resurgent function.

3. Variational series

The previous studies of the convergent series application to the lattice ϕ^4 -model [31] demonstrated a critical slow down of the convergence rate with the increasing of the lattice volume V (see Section 5). However, this problem can be resolved by the following observation. When $\eta = 1$, the explicit dependence on the lattice volume V in the sum (8) of functions (19) can be substituted by $\tau = V + \alpha$, i.e.,

$$\langle \phi_i \phi_j \rangle = \sum_{l=0}^{\infty} J_{\eta=1}(\tau, l) \left[\sum_{k=0}^l C_l^k \sigma^{l-k} \frac{2k! f_k}{\Gamma(\frac{\tau+4k+2}{2})} \right], \tag{21}$$

where $\tau > -2$, not to generate singularities in the integrals $J_{\eta}(\tau, l)$. The latter inequality is special for the propagator, for an arbitrary n -fields Green’s function it has to be substituted by $\tau > -n$.

Let us first consider only a perturbative proof of (21) (the non-perturbative aspects are considered in the Section 4). It can be obtained by changing the order of summations in (21)

$$\langle \phi_i \phi_j \rangle \approx \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} J_{\eta=1}(\tau, l) C_l^k \sigma^{l-k} \frac{2k! f_k}{\Gamma(\frac{\tau+4k+2}{2})}, \tag{22}$$

where the sign ‘ \approx ’ stands to indicate only perturbative equivalence between left and right parts of the expression. The summands in the latter expression are equal to zero, when $l < k$. Changing the summation index to $y = l - k$, we obtain

$$\begin{aligned} \langle \phi_i \phi_j \rangle &\approx \sum_{k=0}^{\infty} \frac{2 f_k \int_0^{\infty} dt t^{\tau+4k+1} e^{-t^2 - \sigma t^4} \sum_{y=0}^{\infty} \frac{C_{(y+k)}^k k! \sigma^y t^{4y}}{(y+k)!}}{\Gamma(\frac{\tau+4k+2}{2})} \\ &= \sum_{k=0}^{\infty} \frac{2 f_k \int_0^{\infty} dt t^{\tau+4k+1} e^{-t^2 - \sigma t^4} \sum_{y=0}^{\infty} \frac{\sigma^y t^{4y}}{y!}}{\Gamma(\frac{\tau+4k+2}{2})} \end{aligned}$$

$$= \sum_{k=0}^{\infty} \frac{2 f_k \int_0^{\infty} dt t^{\tau+4k+1} e^{-t^2}}{\Gamma\left(\frac{\tau+4k+2}{2}\right)} = \sum_{k=0}^{\infty} f_k. \tag{23}$$

We end up with the series of standard perturbation theory. There are two important consequences from this fact (the non-perturbative analogues of these statements are derived in the following sections).

- The whole sum of the series over l in (21) does not depend on τ . Therefore, τ is a variational parameter and can be taken arbitrary to optimize the convergence of the series.
- According to (23), the convergent series is a re-summation method. Thus, for the computation of the connected Green’s functions (including the normalized to the full partition sum propagator, which is a subject of our numerical studies) one can use the fact, that in standard perturbation theory connected functions are obtained from the full Green’s functions by throwing away disconnected Feynman diagrams from the expansions.

4. Convergence of the variational series

In the last section we have introduced a variational parameter τ , to improve the convergence rate of CS (see Section 5). However, the derivations of the Section 2 are not applicable when $\tau \neq V$. The proof of the convergence of the series (8) is based on the positivity of the brackets $(N[\phi] - S[\phi])^l$. When $\tau \neq V$, each summand of the binomial expansion of $(N[\phi] - S[\phi])^l$ transforms differently under the change of τ and the positivity of the brackets $(N[\phi] - S[\phi])^l$ can be lost. For instance, it is the case for the one-site lattice ϕ^4 -integral at $\tau = 0$. In Section 3 we have proved the perturbative independence on τ of the total sum of the series (21). One can not *a priori* exclude the possibility, that some non-analytical dependence on τ at $\eta = 1$, which gives zero contribution to the SPT series, still persist in (21). The prescription for the evaluation of the connected Green’s functions, suggested in the previous section, is based also only on the standard perturbation theory arguments.

Here we study all these issues from the non-perturbative point of view. For this we investigate the convergence properties of the variational expansions constructed for the lattice ϕ^4 with two different regularizations. The first one is the η -regularization and it is achieved by considering $0 < \eta < 1$ in (5). The second, γ -regularization, is defined in Section 4.2 by introducing an additional term proportional to $\|\phi\|^6$ into the action (1).

4.1. Convergence of the variational series depending on η and τ

Let us study the convergence of the series for the full propagator, obtained from (21), extending it to $\eta \leq 1$

$$\langle \phi_i \phi_j \rangle = \sum_{l=0}^{\infty} J_{\eta}(\tau, l) \left[\sum_{k=0}^l C_l^k \sigma^{l-k} \frac{2k! f_k}{\Gamma\left(\frac{\tau+4k+2}{2}\right)} \right]. \tag{24}$$

The asymptotic of large orders of the perturbation theory in quantum field theories and lattice models for the connected and full correlation functions have similar form [3,33,34]

$$f_k \sim (-1)^k \sqrt{2\pi} e \left(\frac{a}{e}\right)^k k^{k+b_0+1/2}, \tag{25}$$

where $a, b_0 \in \mathbb{R}$ are some constants. The upper bound for the series (24) can be obtained as




Fig. 1. The convergence of the series (24) in the half-band $-\eta_* < \eta < \eta_*$, $\tau > -2$ is provided by the bound (26). The convergence at $\tau = V$ and $\eta \leq 1$ follows from the estimate (10).

$$\begin{aligned}
 |(\phi_i \phi_j)| &\leq \sum_{l=0}^{\infty} |J_{\eta}(\tau, l)| \left[\sum_{k=0}^l C_l^k \sigma^{l-k} \frac{2k! |f_k|}{\Gamma(\frac{\tau+4k+2}{2})} \right] \\
 &= \sum_{k=0}^{\infty} \frac{2 \int_0^{\infty} dt t^{\tau+4k+1} e^{-t^2 - \sigma(1-|\eta|)t^4}}{\Gamma(\frac{\tau+4k+2}{2})} |f_k| |\eta|^k.
 \end{aligned}
 \tag{26}$$

The coefficients in front of $|f_k|$ in (26) at large k behave as

$$2^{1/2-2\beta} \sigma_{\eta}^{-\beta} k^{-k} \left(\frac{e|\eta|}{4\sigma_{\eta}} \right)^k,
 \tag{27}$$

where $\beta = (\tau + 2)/4$ and $\sigma_{\eta} = \sigma(1 - |\eta|)$. Consequently, the bound (26) is convergent, when $|\eta| < \eta_* = \frac{4\sigma}{|a|+4\sigma}$, independently on the value of τ . When $\tau = V$, the series (24) converges for $\eta \leq 1$ due to the estimate (10). The bound (26) has finite radius of the convergence in terms of η for any $\tau > -2$, including, for instance $\tau = 0$. In Fig. 1 we show the area of the parameters τ and η for which the convergence of the series (24) is guaranteed by (26) and (10).

The convergence of the series (24) is better than the convergence of (26), because of the cancellations in the internal sums over k in (24). Therefore, it might be, that the series (24) is convergent for $\tau > -2$ and $\eta = 1$. To support such possibility, let us consider an example of the series with an asymptotic of the type (25)

$$h_k = (-1)^k \frac{\Gamma(\frac{\tau+4k+2}{2})}{2\Gamma(k+1)} u^k, \quad u > 0.
 \tag{28}$$

At $\eta = 1$ the substitution of these coefficients into the expansion (26) produces a divergent series. In opposite, the expansion (24) in this case can be bounded by

$$\int_0^{\infty} dt t^{\tau+1} e^{-t^2 - \sigma t^4 + |\sigma-u|t^4} < \infty.
 \tag{29}$$

Therefore, the series (28), re-summed in accordance to (24), is convergent.

4.2. γ -Regularization

The η -regularization enforce an additional decay of the lattice Boltzmann weight for the large fluctuations of the fields ϕ_n . This results in the dumping of the coefficients of standard perturbation theory and in the consequent convergence of the series (24) for $|\eta| < \eta_*$. However, the additional decay is not sufficiently sharp to provide convergence for $\eta_* \leq \eta < 1$, which is

desired, since $\eta = 1$ corresponds to the original theory. Here we introduce alternative regularization, giving sharper vanishing of the large fluctuations of fields. We consider the regularized lattice ϕ^4 -model, defined by the action

$$S_\gamma[\phi] = S[\phi] + \gamma \|\phi\|^6. \tag{30}$$

When $\gamma = 0$ the action S_γ coincides with the action of the ϕ^4 -model (1). For the model defined by (30) one can construct two related expansions. As always, we demonstrate them on the example of the propagator. The first one is similar to standard perturbation theory and is obtained by expanding the interaction part of the original ϕ^4 -model into the Taylor series

$$\langle \phi_i \phi_j \rangle_\gamma = \sum_k \int [d\phi] \phi_i \phi_j e^{-\|\phi\|^2 - \gamma \|\phi\|^6} \frac{(-\frac{\lambda}{4!} \sum_n \phi_n^4)^k}{k!}. \tag{31}$$

Rewriting (31) in terms of Gaussian integrals analogously to Section 2 and introducing the dependence on τ , we get

$$\langle \phi_i \phi_j \rangle_\gamma = \sum_{k=0}^\infty \frac{2 \int_0^\infty dt t^{\tau+4k+1} e^{-t^2 - \gamma t^6}}{\Gamma(\frac{\tau+4k+2}{2})} f_k. \tag{32}$$

The second expansion is the variational series similar to (24). To derive it, we split the action as

$$S_\gamma[\phi] = (N[\phi] + \gamma \|\phi\|^6) + (S[\phi] - N[\phi]), \tag{33}$$

where $(N[\phi] + \gamma \|\phi\|^6)$ is treated as the initial approximation. Then, analogous to previous derivations, we have

$$\langle \phi_i \phi_j \rangle_\gamma = \sum_{l=0}^\infty J_\gamma(\tau, l) \left[\sum_{k=0}^l C_l^k \sigma^{l-k} \frac{2k! f_k}{\Gamma(\frac{\tau+4k+2}{2})} \right], \tag{34}$$

where

$$J_\gamma(\tau, l) = \frac{1}{l!} \int_0^\infty dt e^{-t^2 - \sigma t^4 - \gamma t^6} t^{\tau+4l-1}. \tag{35}$$

The series (32) can be obtained from the series (34) by changing the order of summations. Consequently, if both of these series converge absolutely, they converge to the same sum. The series (32) and (34) can be bounded by

$$\begin{aligned} |\langle \phi_i \phi_j \rangle_\gamma| &\leq \sum_{l=0}^\infty |J_\gamma(\tau, l)| \left[\sum_{k=0}^l C_l^k \sigma^{l-k} \frac{2k! |f_k|}{\Gamma(\frac{\tau+4k+2}{2})} \right] \\ &= \sum_{k=0}^\infty \frac{2 \int_0^\infty dt t^{\tau+4k+1} e^{-t^2 - \gamma t^6}}{\Gamma(\frac{\tau+4k+2}{2})} |f_k|. \end{aligned} \tag{36}$$

At large k the coefficients in front $|f_k|$ in the leading order are determined by

$$3(2\sqrt{3})^{-1-4\beta/3} \gamma^{-2\beta/3} k^{-4\beta/3} (12e^2\gamma)^{-2k/3} k^{-4k/3} \tag{37}$$

with $\beta = (\tau + 2)/4$. The bound (36) converges for any $\tau > -2$, when $\gamma > 0$. In Fig. 2 we present the area of parameters γ and τ for which the series (32) and (34) are convergent.

Similar to the η -regularization case the convergence of (34) is better than of the series (32) and bound (36).




Fig. 2. In the marked area the convergence of both series (32) and (34) is obtained due to the non-zero γ . At $\gamma = 0$ and $\tau = V$, as it follows from the estimate (10), only the series (34) is convergent.

4.3. Non-perturbative independence on τ and continuity of the γ -regularization

Here we show, that for any $\tau > -2$, there are such $\gamma_* \in \mathbb{R}_+$ and $K(\gamma_*) \in \mathbb{N}$, that it is possible to construct convergent series (32), which approximates the propagator of the lattice ϕ^4 -model with an arbitrary precision $\delta > 0$:

$$\left| \sum_{k=0}^{K(\gamma_*)} \frac{2 \int_0^\infty dt t^{\tau+4k+1} e^{-t^2-\gamma_* t^6}}{\Gamma(\frac{\tau+4k+2}{2})} f_k - \langle \phi_i \phi_j \rangle \right| < \delta. \tag{38}$$

The proof of (38) also demonstrates, that in the limit $\gamma \rightarrow 0$ the sum of the series (32) is independent on τ non-perturbatively, i.e., including all possible non-analytical contributions.

Since for $\gamma > 0$ the series (32) is absolutely convergent one can interchange in it the summation and integration (the SPT-coefficients f_k are integrals), this gives

$$\langle \phi_i \phi_j \rangle_\gamma = \int [d\phi] \phi_i \phi_j e^{-\|\phi\|^2} \sum_{k=0}^\infty h_k(\phi_n, \gamma, \tau), \tag{39}$$

where

$$h_k(\phi_n, \gamma, \tau) = \frac{(-\frac{\lambda}{4!} \sum_n \phi_n^4)^k}{k!} \frac{2 \int_0^\infty dt t^{\tau+4k+1} e^{-t^2-\gamma t^6}}{\Gamma(\frac{\tau+4k+2}{2})}. \tag{40}$$

The series $\sum_{k=0}^\infty h_k(\phi_n, \gamma, \tau)$ converges faster than the series for the exponent and, consequently, is uniformly convergent for each compact subset of the region of parameters $\bar{\mathcal{A}} = \{\phi_n \in \mathbb{R}, \gamma \geq 0, \tau > -2\}$. In region $\bar{\mathcal{A}}$ the functions $h_k(\phi_n, \gamma, \tau)$ are continuous in all parameters, therefore the sum of them is also continuous. At $\gamma = 0$ each function $h_k(\phi_n, \gamma, \tau)$ is independent on τ and the propagator (39) coincides with $\langle \phi_i \phi_j \rangle$, defined by (3). Hence, (39) is finite at $\gamma = 0$. The finiteness of (39) for $\gamma > 0, \tau > -2$ follows from the convergence of the series (32). Then, because $\sum_{k=0}^\infty h_k(\phi_n, \gamma, \tau)$ is continuous in $\bar{\mathcal{A}}$ and (39) is finite for $\gamma \geq 0, \tau > -2$, it follows, that for any $\tilde{\delta} > 0$ there is such $\tilde{\gamma}$, that

$$|\langle \phi_i \phi_j \rangle_{\tilde{\gamma}} - \langle \phi_i \phi_j \rangle| < \tilde{\delta}. \tag{41}$$

From the latter inequality and from the convergence of the series (39) we obtain (38).

As it was discussed above series (32) and (34) converge to the same quantities, therefore, all stated here is valid also for (34).

4.4. Evaluation of the connected Green's functions

In previous parts of the paper we predominantly discussed constructions of the convergent series for the full Green's functions on the example of the propagator. The generating functional of the full functions is the partition function $Z[A]$, which is a functional of the external field A . However, the thermodynamical quantities are naturally expressed in terms of the connected Green's functions, given by derivatives of the $\log(Z[A])$ with respect to the external field A . Within the framework of standard perturbation theory it is shown, that connected functions are obtained from the full functions by throwing away all disconnected diagrams of the perturbative expansion, see, for instance, [29]. As it is known from the theory of the combinatorial species [35] this relation is much more general. If the weights of a generating function for a combinatorial weighted species factorize among the connected components of the species, then the logarithm of that function is given by the sum over the connected associated species. In the convergent series (32) as in SPT the disconnected diagrams are the products of the connected ones, consequently, for the computation of the connected functions using the convergent series one has to substitute the contributions f_k of all diagrams in each order of SPT by the contributions only from connected diagrams \tilde{f}_k . The resulting series is convergent, since the asymptotic of the high orders of standard perturbation theory has similar form (25) for both full and connected Green's functions.

Let us change f_k by \tilde{f}_k in (34). Analogously to the relation between (32) and (34), the interchange of the summations in the series (34) with \tilde{f}_k gives the series (32) with f_k , changed by \tilde{f}_k . Therefore, the series (34) with \tilde{f}_k instead of f_k is also convergent.

5. Computations and numerical results

Here we present the results of the $\langle \phi_n^2 \rangle$ operator computations within the convergent and variational series and compare them with the Borel re-summation and Monte Carlo simulations. We show the numerical dependences on the variational and regularization parameters τ and γ . The propagator between all lattice sites $\langle \phi_i \phi_j \rangle$ is computed, but we present only the part of the resulting data representing $\langle \phi_n^2 \rangle$, for greater clarity. All calculations are performed at unit mass $M = 1$ and for the coupling constants λ in the region $[0, 10]$.

The computations within the convergent/variational series methods contain two main steps. The first one is the evaluation of the coefficients of standard perturbation theory for the propagator $\langle \phi_i \phi_j \rangle$. For this the connected diagrams of the ϕ^4 -theory are generated using the system 'GRACE' [36]. The free propagator in diagrams is obtained by the numerical inversion of the K_{mn} matrix (2). The best current results within the ϵ -expansion in the continuum ϕ^4 -theory are obtained in the 6-loops approximation [37]. We also compute 6 orders of the lattice SPT, since it is reasonable to test the efficiency of the convergent series within the same order of the perturbation theory. The second step of the calculations using the CS/VS methods is the re-summation of the perturbative results in accordance to the formulas (8), (19), (24), (34).

To perform the Borel re-summation procedure, we use the conformal mapping for the analytical continuation in the Borel plane. The conformal mapping can be done if the parameter a from the asymptotic of the high orders of the perturbation theory (25) is known. We estimate a using the values presented in [38] for the continuous one- and two-dimensional ϕ^4 -model

$$d = 1, \quad a = 1/8, \tag{42}$$

$$d = 2, \quad a = 1/35.102 \dots \tag{43}$$




Fig. 3. Dependence of $\langle \phi_n^2 \rangle$ on the order of the perturbation theory for the lattice volume $V = 4$ at $\lambda = 0.1$.




Fig. 4. Dependence of $\langle \phi_n^2 \rangle$ on the order of the perturbation theory for the lattice volume $V = 4, \tau = 0$ at $\lambda = 0.1$.

The results obtained on the lattice with $V = 4$ sites are the following. In Figs. 3 and 5 we present the operator $\langle \phi_n^2 \rangle$ computed with CS and other methods depending on the number of known orders of the perturbation theory (from 0 to 6 loops) at coupling constants $\lambda = 0.1$ and $\lambda = 10$ respectively. At small value of the coupling constant $\lambda = 0.1$ one observes an agreement between all methods including standard perturbation theory. When $\lambda = 10$, the result is qualitatively different. The convergent series is out of the Monte Carlo error bars, but the Borel re-summation method is in the agreement with the Monte Carlo data. The values of standard perturbation theory series are omitted because of the strong divergence.

In Figs. 4 and 6 we show the results obtained with the variational series and other methods at coupling constants $\lambda = 0.1$ and $\lambda = 10$ respectively. The variational series exhibits remarkable agreement with the Monte Carlo data and converges even faster than the Borel re-summation.

In Figs. 7 and 8 we demonstrate the dependence of $\langle \phi_n^2 \rangle$ on the coupling constant λ for different methods with $V = 4$. The computations carried according to VS agree with the Monte Carlo results with the precision of the one standard deviation.

The results obtained on the one-dimensional lattice with $V = 64$ are qualitatively similar. In Fig. 9 we show the computations performed with the convergent series method in comparison with other methods. Even at small coupling constants the convergent series does not agree with




Fig. 5. Dependence of $\langle \phi_n^2 \rangle$ on the order of the perturbation theory for the lattice volume $V = 4$ at $\lambda = 10$.




Fig. 6. Dependence of $\langle \phi_n^2 \rangle$ on the order of the perturbation theory for the lattice volume $V = 4$, $\tau = 0$ at $\lambda = 10$.




Fig. 7. Dependence of the operator $\langle \phi_n^2 \rangle$ on coupling constant for the lattice volume $V = 4$.




Fig. 8. Dependence of the operator $\langle \phi_n^2 \rangle$ on coupling constant for the lattice volume $V = 4, \tau = 0$.




Fig. 9. Dependence of the operator $\langle \phi_n^2 \rangle$ on coupling constant for the lattice volume $V = 64$.

the Monte Carlo. The Borel re-summation is in agreement with the Monte Carlo data in the whole region $\lambda \in [0, 10]$. However, the results obtained within VS are significantly different from the results of the simple convergent series. Corresponding computations are presented in Fig. 10. The variational series matches the Monte Carlo data with the precision of the one standard deviation again!

In Fig. 11 we present the dependence of $\langle \phi_n^2 \rangle$ on λ for the two-dimensional lattice with $V = 8 \times 8$, calculated with the VS method, $\tau = 0$. The VS method gives the results which are close to the Monte Carlo simulations. The strong deviation of the Borel re-summation can be caused by the not precise estimate of the parameter a in the asymptotic of the high orders of the perturbation theory (43).

In Section 4.3 we have proved the continuity of the series (34) with respect to the parameter $\gamma \geq 0$, in Fig. 12 we demonstrate it for $\lambda = 1$.

In the limit $\gamma \rightarrow 0$ the dependence on τ of the sum (34) has to disappear. In the real calculations one always has only a finite amount of terms of (34). In Figs. 13 and 14 we present the dependence on τ of the 6-th order approximation of the variational series (24) for the coupling constants $\lambda = 1$ and $\lambda = 5$ respectively. The optimal values of the parameter τ , giving the matching with the Monte Carlo mean value, are $\tau_{\lambda=1} \simeq -0.1, \tau_{\lambda=5} \simeq \{-1.62; -0.38\}$. Without




Fig. 10. Dependence of the operator $\langle \phi_n^2 \rangle$ on coupling constant for the lattice volume $V = 64, \tau = 0$.




Fig. 11. Dependence of the operator $\langle \phi_n^2 \rangle$ on coupling constant in the two-dimensional case $V = 8 \times 8, \tau = 0$.




Fig. 12. Dependence of the operator $\langle \phi_n^2 \rangle$, computed by the variational series, on the regularization parameter $\gamma, \lambda = 1, V = 64, \tau = 0$.




Fig. 13. Dependence of the operator $\langle \phi_n^2 \rangle$ on the variational parameter τ , $\lambda = 1$ for the lattice volume $V = 64$.




Fig. 14. Dependence of the operator $\langle \phi_n^2 \rangle$ on the variational parameter τ , $\lambda = 5$ for the lattice volume $V = 64$.

comparison with the Monte Carlo simulations one can apply the principle of the smallest contribution of the last term of the series, it gives uniform optimal $\tau_\lambda \simeq -0.086$. In the main part of our computations we use $\tau = 0$, what corresponds in the continuum limit to the utilization of the dimensional regularization [30].

6. Conclusions

We have presented the construction of the convergent series for the lattice models with the even degree polynomial interaction and investigated it in details on the example of the lattice ϕ^4 -model. We have proved, that CS is a re-summation method. The latter fact supports the resurgence idea, that the non-perturbative physics can be expressed via the coefficients of standard perturbation theory. The initial approximation of the convergent series is a non-Gaussian interacting non-local model, hence, CS automatically takes into account such non-analytical contributions as $e^{-\frac{1}{\lambda}}$.

We have observed an internal symmetry of the CS method and, using it, developed the variational series. The numerical values of the operator $\langle \phi_n^2 \rangle$ calculated using the CS and VS methods with 6 orders of standard perturbation theory were compared with the Borel re-summation and

Monte Carlo method. The comparison revealed, that for the small lattices and for the small coupling constants the convergent series exhibits the agreement with the Monte Carlo and Borel re-summation, but deviates from them for larger lattice volumes and/or for larger coupling constants. The variational series method, applied to the one- and two-dimensional cases, agrees with other methods for the wide range of lattice volumes and coupling constants and converges even faster to the Monte Carlo results than the Borel re-summation.

To study the convergence and correctness of the variational series from the analytical point of view, we have considered two regularizations of the lattice ϕ^4 -model. In both cases we have constructed for the variational series the upper bounds with finite radii of convergence in terms of the regularization parameters. For the γ -regularization the border of the convergence region approaches $\gamma = 0$, which corresponds to the non-regularized model. We also have shown, that the original model can be approximated by the γ -regularization with any arbitrary precision and it is always possible to construct two related convergent expansions for this regularization. Therefore, the γ -regularized series can be used itself for the non-perturbative computations, without the utilization of the variational series (21). The convergence properties of VS, depending on regularization parameters are summarized in Figs. 1 and 2. The series (34) and (24) are the regularizations of (21). Using this information and the fact, that the convergence of the series (34) and (24) should be better than the convergence of their bounds, we conjecture that the series (21) is convergent for any $\tau > -2$.

The convergence of (21) independently on values of τ and the independence on τ of its sum at $\gamma \rightarrow 0$ allows one to consider any finite $\tau > -2$ even for very large (infinite) volumes V . The diagrams of standard perturbation theory can be easily computed in the infinite volume. This provides a way for taking the infinite volume limit within the CS/VS methods. The computations at $\tau = 0$ work equally well for lattice volumes more than 10 times different, see Figs. 8 and 10, this supports the statement above and gives a numerical evidence of the possibility to perform the infinite volume limit.

It is important to note, that the applicability of the convergent/variational series is not based on any criteria of the kind of the Borel summability of the original perturbation theory. The presented construction can be generalized to the fermionic lattice models by employing bosonization. For instance, an application of the method [9–13] to the bosonized fermionic model, a model of lattice QED, was proposed in [39]. The bosonization of the complex actions has been recently suggested in [40]. Therefore, the convergent series, which is based only on the perturbative computations, can provide a way for the bypassing the sign problem in bosonic and fermionic models.

Acknowledgements

We acknowledge Vladimir V. Belokurov, Eugeny T. Shavgulidze, Vincent Rivasseau, Vladimir Yu. Lotoreychik for the discussions and Nikolay M. Gulitskiy for reading the manuscript. The work of A.S. Ivanov was supported by the Russian Science Foundation Grant No. 14-22-00161. V.K. Sazonov acknowledges the Austrian Science Fund (FWF), Grant No. I 1452-N27.

References

- [1] P. de Forcrand, Simulating QCD at finite density, PoS LAT2009 (2009) 010, arXiv:1005.0539.
- [2] F.J. Dyson, Divergence of perturbation theory in quantum electrodynamics, Phys. Rev. 85 (1952) 631–632, <http://dx.doi.org/10.1103/PhysRev.85.631>, <http://link.aps.org/doi/10.1103/PhysRev.85.631>.

- [3] L. Lipatov, Divergence of the perturbation theory series and the quasi-classical theory, *Sov. Phys. JETP* 45 (1977) 216.
- [4] S. Laporta, E. Remiddi, The analytical value of the electron ($g - 2$) at order α^3 in QED, *Phys. Lett. B* 379 (1–4) (1996) 283–291, [http://dx.doi.org/10.1016/0370-2693\(96\)00439-X](http://dx.doi.org/10.1016/0370-2693(96)00439-X), <http://www.sciencedirect.com/science/article/pii/037026939600439X>.
- [5] T. Aoyama, M. Hayakawa, T. Kinoshita, M. Nio, Tenth-order QED contribution to the electron $g - 2$ and an improved value of the fine structure constant, *Phys. Rev. Lett.* 109 (2012) 111807, <http://dx.doi.org/10.1103/PhysRevLett.109.111807>, <http://link.aps.org/doi/10.1103/PhysRevLett.109.111807>.
- [6] E.B. Kolomeisky, Optimal number of terms in QED series and its consequence in condensed-matter implementations of QED, *Phys. Rev. A* 92 (2015) 012113, <http://dx.doi.org/10.1103/PhysRevA.92.012113>, <http://link.aps.org/doi/10.1103/PhysRevA.92.012113>.
- [7] Y. Meurice, Simple method to make asymptotic series of Feynman diagrams converge, *Phys. Rev. Lett.* 88 (2002) 141601, <http://dx.doi.org/10.1103/PhysRevLett.88.141601>, <http://link.aps.org/doi/10.1103/PhysRevLett.88.141601>.
- [8] B. Kessler, L. Li, Y. Meurice, New optimization methods for converging perturbative series with a field cutoff, *Phys. Rev. D* 69 (2004) 045014, <http://dx.doi.org/10.1103/PhysRevD.69.045014>, <http://link.aps.org/doi/10.1103/PhysRevD.69.045014>.
- [9] V. Belokurov, Y. Solov'ev, E. Shavgulidze, Method of approximate evaluation of path integrals using perturbation theory with convergent series I, *Theor. Math. Phys.* 109 (1) (1996) 1287–1293, <http://dx.doi.org/10.1007/BF02069887>, <http://dx.doi.org/10.1007/BF02069887>.
- [10] V. Belokurov, Y. Solov'ev, E. Shavgulidze, Method for approximate evaluation of path integrals using perturbation theory with convergent series. II. Euclidean quantum field theory, *Theor. Math. Phys.* 109 (1) (1996) 1294–1301, <http://dx.doi.org/10.1007/BF02069888>, <http://dx.doi.org/10.1007/BF02069888>.
- [11] V. Belokurov, Y. Solov'ev, E. Shavgulidze, Perturbation theory with convergent series for functional integrals with respect to the Feynman measure, *Russ. Math. Surv.* 52 (2) (1997) 392.
- [12] V. Belokurov, Y. Solov'ev, E. Shavgulidze, Evaluation of the functional integrals with convergent series, *Fundam. Prikl. Mat.* 3 (3) (1997) 693–713 (In Russian).
- [13] V. Belokurov, Y. Solov'ev, E. Shavgulidze, General approach to the evaluation of the functional integral and summation of divergent series, *Fundam. Prikl. Mat.* 5 (2) (1999) 363–383 (in Russian).
- [14] D.C. Brydges, T. Kennedy, Mayer expansions and the Hamilton–Jacobi equation, *J. Stat. Phys.* 48 (1) (1995) 19–49, <http://dx.doi.org/10.1007/BF01010398>, <http://dx.doi.org/10.1007/BF01010398>.
- [15] A. Abdesselam, V. Rivasseau, Trees, forests and jungles: a botanical garden for cluster expansions, *Lect. Notes Phys.* 446 (1995) 7, http://dx.doi.org/10.1007/3-540-59190-7_20, arXiv:hep-th/9409094.
- [16] J. Magnen, V. Rivasseau, Constructive ϕ^4 field theory without tears, *Ann. Henri Poincaré* 9 (2008) 403–424, <http://dx.doi.org/10.1007/s00023-008-0360-1>, arXiv:0706.2457.
- [17] V. Rivasseau, Constructive field theory in zero dimension, *Adv. Math. Phys.* 2010 (2010) 180159, <http://dx.doi.org/10.1155/2009/180159>, arXiv:0906.3524.
- [18] V. Rivasseau, Z. Wang, Loop vertex expansion for ϕ^{2K} theory in zero dimension, *J. Math. Phys.* 51 (2010) 092304, <http://dx.doi.org/10.1063/1.3460320>, arXiv:1003.1037.
- [19] V. Rivasseau, Z. Wang, How to resum Feynman graphs, *Ann. Henri Poincaré* 15 (11) (2014) 2069–2083, <http://dx.doi.org/10.1007/s00023-013-0299-8>, arXiv:1304.5913.
- [20] V. Rivasseau, Z. Wang, Corrected loop vertex expansion for ϕ_2^4 theory, *J. Math. Phys.* 56 (6) (2015) 062301, <http://dx.doi.org/10.1063/1.4922116>, arXiv:1406.7428.
- [21] A.G. Ushveridze, Converging perturbational scheme for the field theory, *Yad. Fiz.* 38 (1983) 798–809 (in Russian).
- [22] B. Shaverdyan, A. Ushveridze, Convergent perturbation theory for the scalar ϕ^{2p} field theories; the Gell–Mann–Low function, *Phys. Lett. B* 123 (5) (1983) 316–318, [http://dx.doi.org/10.1016/0370-2693\(83\)91208-X](http://dx.doi.org/10.1016/0370-2693(83)91208-X), <http://www.sciencedirect.com/science/article/pii/037026938391208X>.
- [23] A. Ushveridze, Superconvergent perturbation theory for euclidean scalar field theories, *Phys. Lett. B* 142 (5–6) (1984) 403–406, [http://dx.doi.org/10.1016/0370-2693\(84\)91350-9](http://dx.doi.org/10.1016/0370-2693(84)91350-9), <http://www.sciencedirect.com/science/article/pii/0370269384913509>.
- [24] R.P. Feynman, H. Kleinert, Effective classical partition functions, *Phys. Rev. A* 34 (1986) 5080–5084, <http://dx.doi.org/10.1103/PhysRevA.34.5080>, <http://link.aps.org/doi/10.1103/PhysRevA.34.5080>.
- [25] A.N. Sisakian, I.L. Solovtsov, Variational perturbation theory: anharmonic oscillator, *Z. Phys. C* 54 (1992) 263–271, <http://dx.doi.org/10.1007/BF01566655>.
- [26] A.N. Sisakian, O.Yu. Shevchenko, I.L. Solovtsov, Convergent series in variational perturbation theory, *Phys. Lett. B* 297 (1992) 305–308, [http://dx.doi.org/10.1016/0370-2693\(92\)91267-D](http://dx.doi.org/10.1016/0370-2693(92)91267-D).

- [27] A.V. Turbiner, A.G. Ushveridze, Anharmonic oscillator: constructing the strong coupling expansions, *J. Math. Phys.* 29 (1988) 2053.
- [28] J. Honkonen, M. Nalimov, Convergent expansion for critical exponents in the $o(n)$ -symmetric φ^4 model for large ε , *Phys. Lett. B* 459 (4) (1999) 582–588, [http://dx.doi.org/10.1016/S0370-2693\(99\)00704-2](http://dx.doi.org/10.1016/S0370-2693(99)00704-2), <http://www.sciencedirect.com/science/article/pii/S0370269399007042>.
- [29] A.N. Vasil'ev, *The Field Theoretic Renormalization Group in Critical Behavior Theory and Stochastic Dynamics*, Chapman and Hall, ISBN 9780415310024, 2004.
- [30] G. Leibbrandt, Introduction to the technique of dimensional regularization, *Rev. Mod. Phys.* 47 (1975) 849–876, <http://dx.doi.org/10.1103/RevModPhys.47.849>.
- [31] V. Belokurov, A. Ivanov, V.K. Sazonov, E. Shavgulidze, Convergent perturbation theory for the lattice ϕ^4 -model, *PoS Lattice2015* (2016) 275, arXiv:1511.05959.
- [32] D. Dorigoni, An introduction to resurgence, trans-series and alien calculus, arXiv:1411.3585.
- [33] T. Spencer, The Lipatov argument, *Commun. Math. Phys.* 74 (3) (1980) 273–280.
- [34] C. Itzykson, G. Parisi, J.B. Zuber, Asymptotic estimates in scalar electrodynamics, *Phys. Rev. Lett.* 38 (7) (1977) 306–310.
- [35] P. Leroux, F. Bergeron, G. Labelle, *Combinatorial Species and Tree-like Structures*, *Encyclopedia of Mathematics and Its Applications*, vol. 67, Cambridge University Press, Cambridge, 1998.
- [36] F. Yuasa, et al., Automatic computation of cross-sections in HEP: Status of GRACE system, *Prog. Theor. Phys. Suppl.* 138 (2000) 18–23, <http://dx.doi.org/10.1143/PTPS.138.18>, arXiv:hep-ph/0007053.
- [37] D. Batkovich, K. Chetyrkin, M. Kompaniets, Six loop analytical calculation of the field anomalous dimension and the critical exponent η in $o(n)$ -symmetric φ^4 model, *Nucl. Phys. B* 906 (2016) 147–167, <http://dx.doi.org/10.1016/j.nuclphysb.2016.03.009>, <http://www.sciencedirect.com/science/article/pii/S0550321316000912>.
- [38] J. Zinn-Justin, U.D. Jentschura, Order-dependent mappings: strong coupling behaviour from weak coupling expansions in non-Hermitian theories, *J. Math. Phys.* 51 (2010) 072106, <http://dx.doi.org/10.1063/1.3451104>, arXiv:1006.4748.
- [39] V.K. Sazonov, Convergent perturbation theory for lattice models with fermions, *Int. J. Mod. Phys. A* 31 (13) (2016) 1650072, <http://dx.doi.org/10.1142/S0217751X1650072X>.
- [40] V.K. Sazonov, Non-hermitian bosonization, arXiv:1411.5046.