

## Differences between *EcoRI* Nonspecific and “Star” Sequence Complexes Revealed by Osmotic Stress

Nina Y. Sidorova and Donald C. Rau

Laboratory of Physical and Structural Biology, National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, Maryland 20892

**ABSTRACT** The binding of the restriction endonuclease *EcoRI* to DNA is exceptionally specific. Even a single basepair change (“star” sequence) from the recognition sequence, GAATTC, decreases the binding free energy of *EcoRI* to values nearly indistinguishable from nonspecific binding. The difference in the number of waters sequestered by the protein-DNA complexes of the “star” sequences TAATTC and CAATTC and by the specific sequence complex determined from the dependence of binding free energy on water activity is also practically indistinguishable at low osmotic pressures from the 110 water molecules sequestered by nonspecific sequence complexes. Novel measurements of the dissociation rates of noncognate sequence complexes and competition equilibrium show that sequestered water can be removed from “star” sequence complexes by high osmotic pressure, but not from a nonspecific complex. By 5 Osm, the TAATTC “star” sequence complex has lost almost 90 of the ~110 waters initially present. It is more difficult to remove water from the CAATTC “star” sequence complex. The sequence dependence of water loss correlates with the known sequence dependence of “star” cleavage activity.

### INTRODUCTION

It is generally thought that hydration water likely plays an important role in sequence recognition reactions, particularly in specific sequence DNA-protein binding (Hard and Lundback, 1996; Jen-Jacobson, 1997; Schwabe, 1997; Szwajkajzer and Carey, 1997). We have focused on measuring differences in hydration that distinguish specific and nonspecific DNA-protein interactions particularly of the restriction endonuclease *EcoRI*, an exceptionally stringent sequence specific DNA binding protein. Changes in hydration linked to binding reactions can be assessed using an experimental approach termed the osmotic stress technique (Parsegian et al., 1995, 2000). Differences in the numbers of solute excluding water molecules associated with specific and nonspecific DNA-protein complexes can be measured from the dependence of the relative free energy difference on osmotic pressure (or water chemical potential), using the same thermodynamics that allows differences in proton or salt binding accompanying macromolecular reactions to be probed by the sensitivity of the reaction to pH or salt activity. The osmotic stress technique has been widely used to measure the changes in water binding accompanying the DNA binding of several proteins: *Escherichia coli gal* (Garner and Rau, 1995), *lac* (Fried et al., 2002), and *tyr* (Poon et al., 1997) repressors, *E. coli* CAP protein (Vossen et al., 1997), Hin recombinase (Robinson and Sligar, 1996), ultrathorax and deformed homeodomains (Li and Matthews, 1997), the restriction endonucleases *EcoRI* (Robinson and Sligar, 1998; Sidorova

and Rau, 1996, 2001), *BamHI* (Lynch and Sligar, 2000), and *EcoRV* (Wenner and Bloomfield, 1999), HhaI methyl transferase (Swaminathan et al., 2002), Sso7d protein (Lundback et al., 1998), and the TATA binding protein (TBP) (Khrapunov and Brenowitz, 2004; Wu et al., 2001).

We have previously measured differences in water release for the binding of the *EcoRI* to specific and nonspecific DNA sequences (Sidorova and Rau, 1996). Several very different solutes, ranging from glycine and glycerol to triethylene glycol, were used to vary the water chemical potential. The free energy difference between specific and nonspecific DNA-*EcoRI* complexes was linearly dependent on the water chemical potential. The observed osmotic dependence indicated that the nonspecific complex retains some 110 waters more than the specific complex with the recognition sequence. All six osmolytes gave the same difference in waters within 15% experimental error. This slight sensitivity of the difference in the number of water molecules retained by specific and nonspecific complexes to the solute identity is much less than expected for a change in exposed surface area (Courtenay et al., 2000; Davis-Searles et al., 1998; Preisler et al., 1995; Timasheff, 1993). This led us to conclude that this water is sequestered in a space that is sterically inaccessible to solutes. Because the x-ray structure of the *EcoRI*-DNA specific complex (Rosenberg, 1991; Kim et al., 1990) shows that interface of this complex is essentially dry, the 110 waters are likely at the protein-DNA interface of the nonspecific complex. A structure for the nonspecific complex of *EcoRI* is not available, but x-ray structures for both specific sequence and noncognate sequence complexes of a closely related type II restriction endonuclease, *BamHI*, have been solved (Newman et al., 1995; Viadiu and Aggarwal, 2000). Unlike the extensive, direct protein-DNA contacts seen in the specific complex structure, the

Submitted March 5, 2004, and accepted for publication July 26, 2004.

Address reprint requests to Donald C. Rau, Laboratory of Physical and Structural Biology, National Institutes of Child Health and Human Development, National Institutes of Health, Building 9, Room 1E114, Bethesda, MD 20892-0924. Tel.: 301-402-4698; Fax: 301-496-9462; E-mail: raud@mail.nih.gov.

© 2004 by the Biophysical Society

0006-3495/04/10/2564/13 \$2.00

doi: 10.1529/biophysj.104.042390

nonspecific complex structure shows a gap between the *Bam*HI and DNA major groove surfaces that is large enough to hold ~150 waters. The large difference in sequestered water between nonspecific and specific complexes of *EcoRI* is in marked contrast to other solution components. Although salt and pH strongly affect the binding of *EcoRI* to the specific recognition sequence, there is practically no difference in their effect on the relative binding to specific and nonspecific sequences (Sidorova and Rau, 2001).

The observed osmotic sensitivity of the specific-nonspecific equilibrium binding constant is also reflected in the dissociation rate constant of the recognition sequence complex (Sidorova and Rau, 2000, 2001). For DNA-binding proteins that can efficiently locate their target sequences by initially binding nonspecifically and then diffusing along the DNA as *EcoRI*, the dissociation of nonspecifically bound protein is typically the rate-limiting reaction step. We showed that the observed osmotic dependence of the specific complex dissociation rate is dominated by the difference of 110 waters between the specific and nonspecific binding modes of the enzyme. The dissociation of nonspecifically bound protein from the DNA is coupled to an uptake of ~10–45 additional waters depending on the osmolyte used to set water activity. This dependence on solute nature is typical for a reaction that results in an increased exposed surface area of the complex.

At low pressures, the osmotic stress technique can be used as a convenient tool for probing differences in water release accompanying nonspecific versus specific DNA-protein binding without perturbing the system. A linear dependence of binding free energy on solute osmotic pressure indicates that the difference in retained water between complexes remains constant. In principle, however, any sequestered water can be removed by applying high enough osmotic stress. Essentially, the pressure-volume work,  $\Pi\Delta V$ , done in removing water from a complex or in stabilizing an alternate conformation with fewer sequestered waters is balanced by the resulting unfavorable interactions and conformational changes of the complex. A differential loss of water from a noncognate DNA-protein complex would be seen as a nonlinear dependence of the free energy difference between specific and noncognate complexes on osmotic stress at high pressures. Control experiments are necessary to ensure the solute is still acting osmotically at these high osmotic pressures.

We observed previously (Sidorova and Rau, 1999) that at low osmotic pressures an *EcoRI* complex with a DNA sequence that only differs from the recognition sequence by only a single basepair (a “star” sequence) sequesters the same amount of water as a complex with a nonspecific sequence oligonucleotide. Because the enzyme will cleave “star” sequences with low activity, but not sequences with two or more wrong basepairs (Gardner et al., 1982; Goodman et al., 1977; Hsu and Berg, 1978; Lesser et al., 1990; Malyguine et al., 1980; Polisky et al., 1975; Rosenberg and

Greene, 1982; Tikchonenko et al., 1978), “star” sequence complexes cannot be indistinguishable from nonspecific complexes. Additionally, it has been shown that *EcoRI* distinctly pauses at “star” sites while diffusing along DNA (Jeltsch et al., 1994). We extend our previous work (Sidorova and Rau, 1999) and measure to high stresses (~6 Osm) the osmotic dependence of the *EcoRI* dissociation rate from complexes with oligonucleotides containing either the TAATTC or CAATTC “star” sequence, both of which differ from the recognition sequence, GAATTC, by a single first basepair substitution, or an inverted recognition sequence, CTTAAG. We also measure the osmotic dependence of the relative binding free energies from equilibrium competition experiments at lower stresses (<2 Osm). Kinetic measurements allow use of much higher osmotic pressures than is practical for equilibrium studies. We measure the dissociation rate of “star” and inverted sequence complexes from the kinetics of appearance of specific complex after the enzyme is added to a mixture of specific and noncognate sequence DNA. We develop the equations necessary to describe the kinetics.

At low pressures (<1.2 Osm), the osmotic dependence of binding free energies relative to the specific complex indicates that *EcoRI* complexes with oligonucleotides containing “star” sequences and the inverted sequence all sequester about the same number of waters. The relative binding free energy of the inverted sequence oligonucleotide complex continues to vary linearly with osmotic pressure through 2 Osm. Both “star” sequence complexes, however, begin to show nonlinearity at the higher pressures, consistent with a loss of water from the complex or the stabilization of an alternate, dehydrated complex. The osmotic dependence of the dissociation rate from the inverted sequence complex shows no loss of water through 6 Osm (~150 atmospheres), the highest osmotic pressure used. Betaine glycine continues to act osmotically on this *EcoRI* complex even at these high stresses. Much of the water in the TAATTC “star” sequence complex, however, is removed at high osmotic pressures. Somewhat less water, but still a significant amount, is removed from the CAATTC “star” sequence complex. We estimate that a work of ~3 kcal/mol is needed to remove substantially all the water from the initial fully hydrated TAATTC “star” sequence complex and ~6 kcal/mol for the CAATTC complex. These energies are consistent with the observed enzymatic “star” activities for these sequences (Lesser et al., 1990).

## MATERIALS AND METHODS

### Materials

A DNA fragment containing a single *EcoRI* recognition sequence was isolated from the plasmid derived from pNEB193 (New England Biolabs, Beverly, MA) using standard techniques. The 360-bp fragment was purified from a PvuII (New England Biolabs) digestion of the plasmid. The

double-stranded 30-bp-long oligonucleotides used in competition and kinetic experiments were:

Specific sequence GAATTC oligo: ACGACGGCCAGT**GAATTC**GAGCTCGGTACC

TAATTC sequence oligo: ACGACGGCCAGT**TAATTC**GAGCTCGGTACC

CAATTC sequence oligo: ACGACGGCCAGT**CAATTC**GAGCTCGGTACC

CTTAAG sequence oligo: ACGACGGCCAGT**CTTAAG**GAGCTCGGTACC.

These oligonucleotides only differ in the central six basepairs shown in bold letters. The specific sequence oligonucleotide contains the *EcoRI* cognate recognition site, GAATTC. The TAATTC and CAATTC oligonucleotides contain different first basepair substitutions of the recognition sequence and are commonly termed “star” sites (Gardner et al., 1982; Goodman et al., 1977; Hsu and Berg, 1978; Lesser et al., 1990; Malyguine et al., 1980; Polisky et al., 1975; Rosenberg and Greene, 1982; Tikchonenko et al., 1978). The CTTAAG oligonucleotide contains an inverted specific sequence or a nonspecific site with all six basepairs wrong. Sequence differences from the recognition sequence in this central six basepair region are underlined. The oligonucleotides shown above and their complements were purchased from Gibco BRL, and dissolved in STE buffer (100 mM NaCl, 10 mM TrisCl (pH 7.5), 1 mM EDTA). Complementary strands were mixed in 1:1 proportion, heated to 92°C, and annealed by slow cooling to 25°C. Small molecular mass impurities were removed using P6 Bio-Spin columns (Bio-Rad Laboratories, Hercules, CA) at room temperature. Double-stranded oligonucleotides were then additionally purified using high-performance liquid chromatography (Waters (Milford, MA) model 2690, Protein Pak Q ion exchange column), ethanol precipitated, and dissolved in TE buffer (10 mM TrisCl (pH 7.5), 1 mM EDTA). The purity of the double-stranded oligonucleotides was confirmed by polyacrylamide gel electrophoresis. The concentrations of the DNA fragment and oligonucleotides were determined spectrophotometrically, using an extinction coefficient of  $0.013 (\mu\text{M basepairs})^{-1}$  at 260 nm. Absorption spectra were obtained with a Perkin Elmer (Wellesley, MA) Lambda 800 ultraviolet-visible spectrophotometer.

DNA-binding experiments were performed with highly purified *EcoRI* restriction endonuclease (kind gift of Dr. L. Jen-Jacobson). Titration of the *EcoRI* sample with known concentration with specific DNA fragment (1–10-nM concentration range) confirmed that stoichiometry of binding was one *EcoRI* dimer per 360-bp fragment with a single *EcoRI* specific recognition sequence (data not shown). Active protein concentrations for the individual binding experiments were determined by direct titration of the *EcoRI* with the 360-bp DNA fragment under conditions of stoichiometric binding as described previously (Sidorova and Rau, 1996).

Betaine glycine was purchased from United States Biochemical (Cleveland, OH) and used without further purification. Osmolal concentrations of betaine were determined by direct measurement using a vapor pressure osmometer operating at room temperature (Wescor, Logan, UT; model 5520XR). Changes in water chemical potentials are linearly proportional to solute osmolal concentrations, i.e.,  $\Delta\mu_w = \mu_w - \mu_w^{\text{ref}} = -RT[\text{Osm}]/55.6$ , where  $\mu_w$  and  $\mu_w^{\text{ref}}$  are the water chemical potentials of the solutions with and without added osmolyte. We confirmed previously that betaine glycine acts on *EcoRI*-DNA binding osmotically up to very high concentrations (at least up to 4 Osm) (Sidorova and Rau, 1996, 1999). The high solubility of betaine glycine coupled with its low viscosity makes it an excellent solute for experiments requiring high osmotic pressures.

## Equilibrium competition experiments

Specific-nonspecific equilibrium competition experiments were performed as described previously (Sidorova and Rau, 1996). Briefly, mixtures of *EcoRI* (~1 nM), the specific site 360-bp fragment (~2 nM), and the

oligonucleotide competitor (between 0 and ~50  $\mu\text{M}$  in oligonucleotide or 0 and ~1.5 mM in bp), were incubated at 20°C for 90 min. The loss of specific site binding as the concentration of nonspecific competitor DNA increased was determined by the gel mobility shift assay (Garner and Revzin, 1981; Fried and Crothers, 1981). In the experiments reported here all samples contained 20 mM imidazole (pH 7.5), 2 mM EDTA, 1 mM DTT, 0.1 mg/ml bovine serum albumin, and 2.5% ficoll. The total reaction volume was 30  $\mu\text{l}$ . Because the specific sequence complex dissociation rate is strongly dependent on osmotic pressure (Sidorova and Rau, 2000, 2001), the salt concentration was adjusted for different osmotic stress regimes such that the slowest dissociation half-life time is ~18 min, to ensure the reaction reaches equilibrium within 90 min. NaCl concentrations were 120 mM and 160 mM in the 0–1 and 1–2 Osm betaine glycine ranges, respectively. We have shown previously (Sidorova and Rau, 2001) that water activity and salt are acting as independent thermodynamic parameters. To prevent redistribution of *EcoRI* between specific site DNA fragment and competitor oligonucleotide between the time that the sample is loaded onto the gel and when the complex enters the gel, a quench reaction was used that “freezes” the equilibrium fraction of specifically bound *EcoRI*. After the 90-min incubation, 10  $\mu\text{l}$  of 0.5  $\mu\text{M}$  specific sequence oligonucleotide in 65 mM imidazole (pH 7.0), 10 mM NaCl, and 6.5 Osm betaine glycine was added to the 30- $\mu\text{l}$  reaction volume. The specific sequence oligonucleotide was added to bind *EcoRI* dissociating from nonspecific complexes to prevent binding to the specific DNA fragment. The combination of increased osmotic pressure and decreased salt and pH lengthens the dissociation half-life time of the specific sequence complex to several hours or more, depending on the initial salt and osmotic pressure conditions. Control experiments with stoichiometrically bound enzyme show no loss of *EcoRI* from specific site DNA fragment to specific site oligonucleotide under these conditions. In the absence of  $\text{Mg}^{2+}$ , we observed no measurable cleavage of the DNA.

## Dissociation kinetics



To measure the dissociation kinetics of noncognate complexes *EcoRI* was added to a mixture of the 360-bp specific sequence DNA fragment and nonspecific oligonucleotide. The final concentrations of *EcoRI* and the specific site 360-bp fragment were 1.5 nM and 3 nM, respectively. The ratio of oligonucleotide and specific sequence fragment concentrations was varied between 50 and 1000. The mixture was incubated for various times, quenched, and the appearance of specific site binding measured using the gel mobility shift assay.

Solution conditions for the kinetic experiments were 20 mM imidazole (pH 7.0), 90 mM NaCl, 2 mM EDTA, 1 mM DTT, and 0.1 mg/ml BSA, in 30  $\mu\text{l}$  volume. Samples were incubated at 20°C. The reaction was quenched by adding a 100-fold excess of specific sequence oligonucleotide and betaine glycine to a final concentration of at least 6 Osm. The samples were sufficiently dense with the added betaine glycine that ficoll was not necessary. The final salt concentration after quenching was 75 mM NaCl. The total reaction volume after quenching was 60  $\mu\text{l}$ .

The ratio of association rates between the specific recognition sequence 360-bp fragment and the specific sequence oligonucleotide was determined from the initial partitioning of *EcoRI* binding. *EcoRI* was added to a mixture of the DNA fragment and the specific sequence oligonucleotide and incubated for 10 min at 20°C in the same buffer as used for the dissociation kinetics experiments. The DNA fragment and protein concentrations were ~3 nM and ~1.5 nM, respectively. The ratio of oligonucleotide and DNA fragment concentrations was varied between 0.5 to 3 and the osmotic pressure between 1.5 and 5 Osm. After incubation the reaction was quenched by adding an equal volume of 2 molal betaine glycine, 20 mM imidazole (pH 7.0), and 60 mM NaCl. The distribution of the *EcoRI* between specific fragment and specific oligonucleotide was measured using the gel mobility shift assay. Under these conditions, binding is complete within 5 min incubation and the dissociation half-life time of the specific sequence complex is >6 h even at the lowest osmotic pressure (Sidorova and Rau, 2001). Thus, the distribution of *EcoRI* between the specific DNA fragment

and specific oligonucleotide on the 10-min timescale of the experiment is determined only by association rates.

The coupled reaction scheme we consider is:



If the ratio of specific site concentrations,  $[DNA_{sp}^{oligo}]/[DNA_{sp}^{fragment}]$ , is  $C$  and the fraction of specific sequence fragment DNA complex measured after the 10-min incubation is  $f$  and  $f^*$  in the presence and absence of specific sequence oligonucleotide, respectively, then the ratio of association rates,  $k_1/k_3$ , is given by,

$$\frac{k_1}{k_3} = \frac{\ln\left(\frac{C - f^* + f}{C}\right)}{\ln(1 - f)}. \quad (1)$$

### Gel mobility-shift experiments

Reaction mixtures from equilibrium and kinetic experiments were electrophoresed in a 10% polyacrylamide gel, TAE (45 mM Tris, 22.5 mM acetic acid, 1 mM EDTA, pH 8.3) buffer. Samples were loaded on a gel at 150 V, and gel was run for 40 min at this voltage. The voltage was then reduced to 60 V and the gel was run overnight at 20°C to separate free DNA fragments and *EcoRI*-bound complexes. *EcoRI*-specific DNA fragment complexes are remarkably stable in the polyacrylamide gels, no change in fractions of bound fragment was observed between 8- and 16-h runs.

Electrophoretic bands containing free DNA and DNA-protein complex were stained with the fluorescent dye SYBR Green I (Molecular Probes, Eugene, OR). The summed intensities of the two bands are constant within 5% over a wide range of protein binding (from no added protein to 70% of the DNA present as complex), indicating there is no significant difference in staining efficiency of free and *EcoRI*-bound DNA. The gels were imaged with a luminescent image analyzer LAS-1000 plus (Fuji Film, Valhalla, NY) that includes a 1.3-megapixel cooled charge-coupled device camera, epi-illumination at 470 nm (light-emitting diode), and a dichroic optical filter suitable for SYBR Green I. The LAS-1000 plus was interfaced to a Pentium (Intel, Santa Clara, CA) PC. Band intensities were quantified using ImageGauge (v.3.122) for Windows. The linearity of DNA fluorescent staining over the range of DNA concentrations studied was confirmed using pBR322 DNA fragments generated by *MspI* digestion.

### Equilibrium competition data analysis

As was developed previously (Sidorova and Rau, 1996), the ratio of specific (sp) and nonspecific (nonsp) association binding constants ( $K_{sp}/K_{nonsp}$ ) can be determined from the loss of specifically bound complex as the concentration of a nonspecific oligonucleotide competitor DNA is increased. If  $f_b$  and  $f_b^0$  are the fractions of *EcoRI*-bound specific sequence fragment with and without added oligonucleotide competitor, then under conditions of virtually stoichiometric protein binding and for much weaker nonspecific than specific binding ( $K_{nonsp} \ll K_{sp}$ ) the change in specific sequence binding is given by,

$$f_b = f_b^0 - \frac{K_{nonsp}}{K_{sp}} \frac{f_b}{1 - f_b} \frac{[DNA_{nonsp}]_{total}}{[DNA_{sp}]_{total}}, \quad (2)$$

where  $[DNA_{nonsp}]_{total}$  and  $[DNA_{sp}]_{total}$  are the molar concentrations of the competing oligonucleotide and the specific sequence fragment, respectively.

Relative binding constants,  $K_{sp}/K_{nonsp}$ , were straightforwardly calculated from the linear dependence of  $f_b$  on  $\frac{f_b}{1-f_b}[DNA_{nonsp}]_{total}$ , measured at constant specific sequence DNA and protein concentrations. The difference in the numbers of solute excluding waters between specifically and nonspecifically bound protein is calculated from the dependence of  $K_{sp}/K_{nonsp}$  on the solute osmolal concentration (Parsegian et al., 1995, 2000) by



$$\frac{d \ln(K_{sp}/K_{nonsp})}{d[\text{osmolal}]} = -\frac{\Delta N_{w,nonsp-sp}}{55.6}, \quad (3)$$

where  $\Delta N_{w,nonsp-sp} = N_{w,sp} - N_{w,nonsp}$ , the difference in the numbers of waters associated with specific and nonspecific complexes that exclude solute.

### THEORY

The dissociation rate from noncognate DNA sequence complexes can be determined by adding *EcoRI* to a mixture of noncognate oligonucleotide and specific sequence DNA fragment and measuring the time course for specific binding. Under conditions of fast association, the rate of appearance of specific complex depends on three factors: 1), the ratio of specific and nonspecific association rate constants; 2), the ratio of specific sequence fragment and nonspecific oligonucleotide concentrations; and 3), the nonspecific complex dissociation rate.

The coupled reaction scheme we consider is:



The rate constants are DNA length dependent. The dissociation rate constants from specific and nonspecific sites are very different. For proteins as *EcoRI* that initially bind nonspecifically then diffuse along the DNA in search of recognition sequences (Ehbrecht et al., 1985; Jack et al., 1982; Terry et al., 1985; Wright et al., 1999), association rate constants for specific and nonspecific sites are not expected to differ significantly for DNA of the same length.

We solve the kinetic equations with two assumptions. First, virtually all protein is bound to DNA over the time course of the experiment. For the experimental conditions used here, this means that the specific site concentration should be at least 1 nM. This is also equivalent to stating that the association rates,  $k_1[DNA_{sp}]_{tot}$  and  $k_3[DNA_{nonsp}]_{tot}$ , are fast compared to the experimental timescale. Kinetic measurements of protein association in the absence of oligonucleotide show that binding is complete within 5 min (data not shown). Secondly, virtually all protein is bound to the specific site fragment at equilibrium. This is equivalent to stating that the dissociation half-life time from the specific site fragment is much longer than the timescale of the

experiment. Our extensive measurements of specific site dissociation rates confirm this (Sidorova and Rau, 2001).

With these assumptions,

$$\ln\left(\frac{(f_\infty - f)}{(1 - f)}\right) - \ln\left(\frac{(f_\infty - f_0)}{(1 - f_0)}\right) = -k_4(1 - f_\infty) \frac{k_1 [DNA_{sp}]_{total}}{k_3 [DNA_{nonsp}]_{total}} t, \quad (4)$$

where  $f$  is the fraction of specific site DNA fragment with bound protein at time  $t$ ,  $f_0 = f$  at  $t = 0$ , and  $f_\infty = f$  at equilibrium. Under the conditions of the experiment,  $f_\infty$ , can be measured from protein binding in the absence of oligonucleotide and  $f_0$  can be calculated from the transcendental equation,

$$f_\infty - f_0 = -\frac{k_3 [DNA_{nonsp}]_{total}}{k_1 [DNA_{sp}]_{total}} \ln(1 - f_0). \quad (5)$$

For  $[DNA_{nonsp}]_{total} \gg [DNA_{sp}]_{total}$ ,  $f_0$  is small and can be approximated by,

$$f_0 \approx \frac{k_1 [DNA_{sp}]_{total}}{k_3 [DNA_{nonsp}]_{total}} f_\infty. \quad (6)$$

Equation 4 for the rate of appearance of DNA fragment with bound protein can be rewritten as,

$$f(t) = \frac{\left(f_\infty - \frac{(f_\infty - f_0)}{(1 - f_0)} e^{-k't}\right)}{\left(1 - \frac{(f_\infty - f_0)}{(1 - f_0)} e^{-k't}\right)}, \quad (7)$$

where

$$k' = (1 - f_\infty) k_4 \frac{k_1 [DNA_{sp}]_{total}}{k_3 [DNA_{nonsp}]_{total}}. \quad (8)$$

These expressions are somewhat more complicated than most dissociation rate equations because the second-order nature of the reaction of protein with specific sequence fragment is included. The apparent rate constant,  $k'$ , depends on the ratio of DNA concentrations because after dissociation of nonspecifically bound protein the probability of associating with the specific sequence DNA fragment rather than with another oligonucleotide is approximately proportional to  $k_1 [DNA_{sp}]_{total} / k_3 [DNA_{nonsp}]_{total}$ . The ratio of association constants  $k_1/k_3$  can be well estimated from the initial partitioning of *EcoRI* between the 360-bp specific site DNA fragment and specific recognition sequence oligonucleotide, i.e., under conditions such that the association time is much faster and the dissociation rate much slower than the

incubation time. We find that  $k_1/k_3 = 3.7 \pm 0.6$  for the pressure range 1 and 5 Osm as calculated from Eq. 1.

## RESULTS

### Equilibrium competition

The general strategy is the same as described earlier (Sidorova and Rau, 1996, 1999). *EcoRI* binding free energies of different DNA sequences relative to the specific sequence were measured using a competition assay. The ratio of binding constants for the specific fragment and competing oligonucleotide were extracted from the decrease in binding to the specific sequence DNA fragment due to the presence of the competitor. Changes in specific *EcoRI* binding were determined using the gel mobility shift assay. Fig. 1 *a* shows a typical gel image illustrating the competition assay. The fluorescently stained bands show the variation in free DNA fragment and protein-DNA complex as betaine glycine osmotic pressure is increased at constant competitor oligonucleotide concentration. *EcoRI* binding to the specific sequence fragment is enhanced with higher pressures. Fig. 1 *b* shows the osmotic stress dependence of the relative binding free energies for the TAATTC ‘‘star’’ sequence that differ from the canonical site by a single incorrect basepair and the CTTAAG inverted sequence competitor oligonucleotides calculated with Eq. 2. In the absence of added osmolyte, the association binding constant to the inverted sequence oligonucleotide is  $\sim 1.2 \times 10^4$  times smaller than the binding constant to the specific site. The TAATTC ‘‘star’’ sequence oligonucleotide binds *EcoRI*  $\sim 2.3$ -fold more strongly than the inverted sequence oligonucleotide. The dependence of  $\ln(K_{sp}/K_{nonsp})$  on osmotic pressure for the inverted sequence oligonucleotide-specific site competition is linear throughout the entire 2-Osm pressure range for both betaine glycine and  $\alpha$ -methyl glucoside including the no-added solute,  $\Pi = 0$ , limit. This linearity along with our previous observation of the linearity of  $\ln(k_d)$  vs.  $\Pi$  for specific complex dissociation (Sidorova and Rau, 2001) again including the  $\Pi = 0$  limit indicates that either solutes do not bind to the specific or nonspecific complexes or that solute binding is inconsequential for DNA-*EcoRI* binding. The slope translates into  $\Delta N_w = 108 \pm 5$  water molecules in a good agreement with previous results (Sidorova and Rau, 1996, 2001). Relative binding energies for competition between the TAATTC ‘‘star’’ sequence oligonucleotide and the specific site fragment are linear only through  $\sim 1$ – $1.5$  Osm. The slope in this region corresponds to a difference of  $105 \pm 7$  waters. The twofold stronger binding to this ‘‘star’’ sequence oligonucleotide compared to the inverted sequence oligonucleotide means that at least half the protein bound to the ‘‘star’’ sequence oligonucleotide at low stresses is directly associated with the TAATTC sequence. Thus, half of the osmotic dependence of  $\ln(K_{sp}/K_{nonsp})$  is the contribution from the ‘‘star’’ sequence complex itself. There is, therefore, no significant difference




FIGURE 1 (a) A polyacrylamide gel image illustrating that osmotic pressure favors specific sequence *EcoRI* binding over nonspecific binding in competition experiments. The gel mobility shift assay is used to monitor *EcoRI* binding to a 360-bp fragment containing its recognition sequence. The concentrations of protein, specific site fragment, and inverted sequence oligonucleotide competitor are constant as the concentration of the osmolyte betaine glycine is increased. The fraction of specific sequence complex is determined from fluorescent intensities of the SYBR Green I stained DNA bands. The short 30-bp competitor oligonucleotide is not seen on the gel. The particular conditions were: 120 mM NaCl, 20 mM imidazole pH 7.6, 20°C. Competitor oligonucleotide concentration is 14  $\mu$ M; *EcoRI* and specific fragment concentrations are 0.75 and 1.5 nM, respectively. (b) The dependence of  $\ln(K_{sp}/K_{nonsp})$  on betaine glycine or methyl glucoside osmolal concentration is shown for two noncognate oligonucleotides. The ratio of association binding constants for specific and nonspecific sequences,  $K_{sp}/K_{nonsp}$ , was calculated from experiments as shown in Fig. 1 *a* using Eq. 2. The nonspecific competitors are the TAATTC “star” sequence oligonucleotide (●, betaine; ○, methyl glucoside) and the nonspecific CTTAAG inverted sequence oligonucleotide (■, betaine; □, methyl glucoside). Except for the central six basepairs, the two 30-bp oligonucleotides are otherwise identical; full sequences are given in Methods and Materials. The error bars (shown for betaine) are calculated from three to four experiments at each osmotic pressure. Experimental conditions are given in Methods and Materials. The slopes at low pressures correspond to 105–110 more waters sequestered by the nonspecific sequence complexes compared with the specific site complex. The linear dependence on osmotic pressure for competition with the inverted sequence oligonucleotide indicates there is no change in the number of water sequestered by this nonspecific complex throughout the entire pressure range. The TAATTC sequence complex, however, apparently begins to lose water at osmotic pressures higher than 1–1.5 Osm.

between water sequestered by a nonspecific complex and the TAATTC “star” sequence complex at low stresses.

At higher stresses the slope of the plot for the TAATTC “star” sequence oligonucleotide complex decreases. This nonlinearity indicates either a differential loss of water from the “star” sequence complex or osmolyte binding. The solute binding explanation demands that both betaine glycine and  $\alpha$ -methyl glucoside bind not only with very similar energies and extents but also specifically to the “star” sequence complex, because both solutes continue to act osmotically on the competition between the inverted and specific sequence complexes. We believe this unlikely and consider probable that the nonlinearity reflects a loss of water from the “star” sequence complex. The average slope in the 1–2 osmolal range corresponds to  $\sim$ 85 waters. Competition with the CAATTC “star” site oligonucleotide was 1.6-fold stronger than with the inverted sequence oligonucleotide and showed similar nonlinear behavior at higher osmotic stresses (data not shown).

### Nonspecific complex dissociation kinetics

Measurement of competitive equilibrium binding at even higher pressures is difficult due to the very slow dissociation rates of the specific sequence complex and the impractically high oligonucleotide DNA concentrations needed for competition. Dissociation rates of *EcoRI* complexes, however, can be measured at very high osmotic pressures and give nearly the same osmotic stress information as equilibrium measurements. The dissociation rate of noncognate complexes can be measured using the gel mobility shift assay from the kinetics of the specific sequence complex formation after *EcoRI* is added to a mixture of the specific sequence fragment and noncognate oligonucleotide. Fig. 2 *a* shows a gel image illustrating the time course of specific complex appearance in the presence of 250-fold excess of the TAATTC oligonucleotide at 3.6 Osm betaine glycine. Fig. 2 *b* shows the dependence of the specific complex fraction on time and the fit to Eq. 7, allowing the initial and final fractions of *EcoRI* bound DNA fragment,  $f_0$  and  $f_\infty$ , respectively, and the apparent rate constant for specific site binding,  $k'$ , to vary. Setting  $f_0 = 0$  affected the fit negligibly. Enough protein was added such that  $f_\infty$  is  $\sim$ 0.5. The nonspecific dissociation rate constant,  $k_4$ , calculated from Eq. 8 is  $2.4 \text{ min}^{-1}$ . Under the same conditions, no dissociation from the specific recognition sequence complex can be observed within the time course of the experiment. If the rate is estimated by extrapolating our previous data (Sidorova and Rau, 2001),  $k_2 \sim 8 \times 10^{-6} \text{ min}^{-1}$ .

Equation 4 predicts a dependence of the rate of appearance of specific complex on the ratio of specific sequence fragment and oligonucleotide concentrations because this determines the probability for a free protein to encounter the specific sequence fragment or noncognate oligonucleotide. We have explicitly confirmed this DNA concentration dependence. Fig. 3 *a* shows a gel image illustrating the




FIGURE 2 (a) A polyacrylamide gel illustrating the kinetics of specific *Eco*RI binding to the 360-bp DNA fragment in the presence of the TAATTC “star” sequence oligonucleotide. In the absence of oligonucleotide, specific sequence binding is complete within 5 min. The dissociation rate of *Eco*RI from the TAATTC oligonucleotide complex is determined from the time dependence. The gel mobility shift assay is used to monitor the gain of specific *Eco*RI-DNA complex. *Eco*RI (1.5 nM) was incubated for different times with a mixture of specific sequence DNA fragment (3 nM) and the TAATTC sequence oligonucleotide. The particular conditions used in this experiment were 90 mM NaCl, 3.95 Osm betaine glycine, 20 mM imidazole (pH 7.0), and 20°C. The TAATTC sequence oligonucleotide was present in 250-fold molar excess over the specific sequence fragment. (b) The fraction of the specific *Eco*RI-DNA fragment complex is shown plotted against time for the gel shown in panel a. The solid line is the best fit of Eq. 7 to the data. The dissociation rate constant  $k_4$  for this complex was calculated as  $2.4 \text{ min}^{-1}$ .

decrease in specific fragment complex formed after 1 h incubation at 3.6 Osm betaine glycine as the concentration of the TAATTC oligonucleotide increases. Fig. 3 b shows the dependence of the specific complex fraction on oligonucleotide concentration. In Fig. 3 c the data is plotted using Eq. 4. The observed linearity over a wide range of concentrations for this “star” sequence oligonucleotide is consistent with the kinetic scheme. This DNA concentration dependence was confirmed for all three noncognate oligonucleotides at betaine glycine concentrations varying from 1.5 to 5 Osm and for  $[DNA_{\text{nonsp}}]_{\text{total}}/[DNA_{\text{sp}}]_{\text{total}}$  ratios ranging from 50 to 1000.

### Osmotic stress dependence of the nonspecific dissociation rate

A clear difference is seen in the effect of osmotic pressure on the dissociation rates of *Eco*RI from the three oligonucleo-




FIGURE 3 (a) The fraction of specific sequence complex formed after one hour incubation decreases as the concentration of noncognate oligonucleotide increases. *Eco*RI (1.5 nM) was incubated with specific sequence DNA fragment (3 nM) and varying concentrations of noncognate oligonucleotides in 75 mM NaCl, 20 mM imidazole (pH 7.0), 3.6 Osm betaine, and at 20°C. The gel mobility shift assay was used to monitor specific complex formation. (b) The dependence of the specific *Eco*RI-DNA complex fraction on TAATTC sequence oligonucleotide concentration for the gel shown in panel a. (c) The linear dependence of  $\ln((f-f)/(1-f))$  on  $[DNA_{\text{sp}}]_{\text{total}}/[DNA_{\text{nonsp}}]_{\text{total}}$  confirms the concentration dependence predicted by Eq. 4. The dissociation rate constant,  $k_4$ , of the TAATTC “star” sequence oligonucleotide complex can be calculated from the slope. For this particular experiment,  $k_4$  is  $0.86 \text{ min}^{-1}$ .

tides that differ only in the central six basepairs. Fig. 4 shows kinetic curves for the appearance of the *Eco*RI-specific 360-bp fragment complex in the presence of the three noncognate oligonucleotides at two osmotic pressures. Fig. 4 a shows the kinetics at 2.0 Osm betaine glycine with a




FIGURE 4 Noncognate complex dissociation rates are strongly dependent on osmotic stress and the oligonucleotide sequence. Dissociation of *EcoRI* from the noncognate oligonucleotides is monitored by the increase in protein binding to the specific 360-bp DNA fragment with the incubation time. (a) In 2 Osm betaine glycine, the kinetic curves are similar for the three oligonucleotide complexes. All three noncognate oligonucleotides are present at 1000-fold molar excess over the specific DNA fragment. The salt and pH conditions used in this experiment were 90 mM NaCl, 20 mM imidazole (pH 7.0), 20°C. (b) At 4.3 Osm betaine glycine, the dissociation rates are clearly different for the three oligonucleotides. Both of the “star” sequence complexes dissociate substantially more slowly than the nonspecific inverted sequence complex. Of the “star” sequence complexes, the dissociation rate of the TAATTC oligonucleotide complex is significantly slower than that of the CAATTC oligonucleotide complex. All three noncognate oligonucleotides are present in 100-fold molar excess over specific DNA fragment.

1000-fold molar excess of each oligonucleotide over specific site fragment. The dissociation rates for the two “star” sequence oligonucleotide complexes are ~50% slower than for the inverted sequence oligonucleotide complex. Under the same conditions, dissociation from the specific recognition site is too slow to be observed. We estimate from our previous work an expected reaction half-life time of ~60 h. Fig. 4b shows that at 4.3 Osm betaine glycine and with a 100-fold molar excess of oligonucleotide over specific sequence fragment, the dissociation rate of the TAATTC oligonucleotide complex is now ~10-fold slower than for the inverted

sequence oligonucleotide complex. The dissociation rate of the CAATTC star sequence oligonucleotide complex is intermediate.

The osmotic sensitivities of the *EcoRI* dissociation rate constant for both “star” sequence complexes and for the inverted sequence complex are shown in Fig. 5. Each rate constant shown was determined from full kinetic curve as shown in Figs. 2 and 4 and is the average of two to four measurements. Error bars are shown for the inverted sequence and TAATTC “star” sequence oligonucleotides. The *EcoRI* dissociation rate of the nonspecific oligonucleotide with the central inverted six-basepair sequence depends linearly on osmotic pressure over the entire range of stresses with a slope corresponding to 15 waters. The extrapolated dissociation rate of the inverted sequence oligonucleotide complex at 0 Osm is 40 min<sup>-1</sup>.

At low osmotic pressures, the dissociation rates for the two “star” sequence oligonucleotides are somewhat smaller than for the nonspecific, inverted sequence oligonucleotide





FIGURE 5 The dependence of  $\ln(k_4)$  for *EcoRI* dissociation on betaine glycine osmolar pressure is shown for three noncognate complexes, with the TAATTC and CAATTC “star” sequence oligonucleotides and the nonspecific inverted CTTAAG sequence oligonucleotide. Error bars are given for the TAATTC and inverted sequence oligonucleotide complexes. The solid lines show the best linear fit to the nonspecific inverted sequence complex data, quadratic fit to the CAATTC complex data, and cubic fit to the TAATTC complex data. The nonspecific inverted sequence complex shows a linear osmotic dependence over the entire range of pressures translating in a constant uptake of ~15 waters typical for a normally hydrated nonspecific complex. The nonlinearity for the “star” sequence complexes indicates an apparent loss of water from these complexes or stabilization of alternate conformations that sequester fewer waters. At 5 Osm the osmotic sensitivity of the  $k_4$  translates into a loss of ~90 waters from the amount initially present in the TAATTC complex, leaving only ~20 waters sequestered in the complex. The dashed line shows the best fit to the TAATTC complex data using Eq. 9 for a two-state model. The particular salt and pH conditions used in these experiments were 90 mM NaCl, 20 mM imidazole (pH 7.0), 20°C. The ratio of specific sequence DNA fragment and noncognate oligonucleotide was adjusted to give a convenient timescale for each experiment. Data points for  $k_4$  and associated error bars were extracted from two to four full kinetic curves measured at a fixed betaine glycine concentration.



complex, but the osmotic pressure dependence is similar. At 2 Osm the dissociation rate of the TAATTC oligonucleotide complex is  $\sim 1.6$ -fold smaller than the dissociation rate of the nonspecific oligonucleotide complex compared with the ratio of 2.3 for binding constants seen in Fig. 1 *b*. A clear deviation from linearity is seen for the TAATTC oligonucleotide complex at high osmotic pressures. Although not as dramatic, the CAATTC oligonucleotide also shows substantial nonlinearity at high osmotic pressures. Once again this nonlinearity could either reflect a change in the number of water molecules associated with the complex or betaine glycine binding. Because this solute continues to act osmotically on the dissociation rate of the inverted sequence complex through the entire osmotic pressure range and because we have previously observed that it also continues to act osmotically on the specific sequence complex through the highest pressure measured ( $\sim 3$  Osm) (Sidorova and Rau, 2001), we consider it likely that betaine glycine continues to act osmotically on the “star” sequence complexes.

The solid lines in Fig. 5 show cubic and quadratic fits to the TAATTC and CAATTC “star” sequence data, respectively. The slopes calculated from these fits at 1.5 Osm give 30 and 20 waters coupled with the dissociation of protein from the TAATTC and CAATTC oligonucleotides, respectively. At 5 Osm, the slopes calculated from the polynomial fits correspond to an uptake of some 105 waters coupled with the dissociation from the TAATTC oligonucleotide complex and 45 waters for the CAATTC oligonucleotide complex. Of these waters 15 can be attributed to water coupled to nonspecific complex dissociation. At 5 osmolal, therefore, the TAATTC “star” sequence complex has lost an apparent 90 water molecules compared with the inverted sequence complex. This is comparable to the difference of 105–110 water molecules between specific and nonspecific *EcoRI* binding seen from competition equilibrium experiments at low osmotic pressures (Fig. 1 *b*). The CAATTC “star” sequence complex has lost  $\sim 30$  water molecules compared with the inverted sequence complex at 5 Osm.

The data for the TAATTC oligonucleotide complex can also be fit to a two-state model. We consider in particular a scheme in which a hydrated “star” sequence complex is in equilibrium with a complex that is dehydrated,



If the difference in associated waters between the two complexes is  $\Delta N_w^*$  waters, then, at an osmotic pressure  $\Pi$  Osm,

$$\ln(k_4) = \ln(k_4^0) - \frac{15}{55.6} \Pi + \ln \left( \frac{K^* e^{-\Delta N_w^* \Pi / 55.6}}{1 + K^* e^{-\Delta N_w^* \Pi / 55.6}} \right), \quad (9)$$

where  $k_4^0$  is  $k_4$  at 0 Osm. The 15 waters linked to the dissociation of protein from the hydrated complex have been

explicitly incorporated. The dashed line in Fig. 5 shows the best fit of this equation to the TAATTC “star” sequence complex data. The best-fitting parameters are  $K^* = 210 \pm 80$  and  $\Delta N_w^* = 105 \pm 25$  waters. From the equilibrium constant, the alternate, dehydrated complex is 3.2 kcal/mol less stable than the hydrated state within this model. Not enough water is lost from the CAATTC oligonucleotide complex to give a fit to a two-state model with  $< 50\%$  standard errors and reasonable dependencies.

## DISCUSSION

The binding of the restriction nuclease *EcoRI* to varying DNA sequences is exceptionally stringent. A change of even a single basepair from the recognition sequence, GAATTC, is sufficient to lower the binding free energy to values essentially characteristic of nonspecific complexes (Lesser et al., 1990). This is unlike many other sequence specific DNA-binding proteins, as the *lac* (Frank et al., 1997) and  $\lambda$ -Cro repressors (Takeda et al., 1989, 1992), that show a more gradual decrease in binding energy as the consensus recognition sequence is altered. As seen in Fig. 1, the binding competition at 20°C between a DNA fragment containing the specific recognition sequence and a 30-bp nonspecific inverted sequence CTTAAG oligonucleotide gives an association binding constant ratio of  $K_{\text{sp}}/K_{\text{nonsp}} = 1.2 \times 10^4$ . The same oligonucleotide but with a central TAATTC “star” sequence binds *EcoRI* only  $\sim 2.3$ -fold stronger with no osmolyte. Another oligonucleotide with a central CAATTC “star” sequence binds  $\sim 1.6$ -fold more tightly than the inverted sequence oligonucleotide. These 30-bp oligonucleotides, of course, contain several nonspecific binding sites. The *EcoRI* binding constants to these oligonucleotides relative to the specific sequence ( $K_{\text{sp}}/K_{\text{nonsp}}$ ) reflect binding competition for the whole oligonucleotide, not just the central six basepairs that we use to identify the oligonucleotide. Differences in binding constants and osmotic stress sensitivities among the oligonucleotides, however, do reflect difference in the binding properties of these central sequences. The approximate twofold increase in binding constant for the “star” sequence oligonucleotide complexes indicates that about half of the protein at low stresses is bound to the “star” sequence itself.

Accompanying the precipitous loss of binding energy with even a single basepair change is a large increase in the number of waters associated with the complex. The *EcoRI* complexes with all three oligonucleotides initially include 105–110 more waters than the specific complex as measurements at low osmotic pressures show (Fig. 1 *b*). The absence of any intermediate hydration state for the “star” sequence complexes would suggest there are two stable and distinct binding modes of the protein to DNA, perhaps analogous to the structures of the specific and nonspecific DNA complexes of *BamHI* (Newman et al., 1995; Viadiu and Aggarwal, 2000).

The equilibrium measurements of the binding competition between specific site and noncognate sequences in Fig. 1, however, reveal a difference between “star” sequence and nonspecific complexes at higher osmotic pressures. In contrast to the inverted sequence oligonucleotide complex that shows linear behavior through 2 Osm implying a constant number of sequestered waters, the TAATTC “star” sequence complex shows a decreasing slope at higher stresses suggesting an apparent loss of water by the complex or that an alternate, dehydrated conformation is stabilized. This apparent loss of water could be connected to other differences between *EcoRI* complexes with nonspecific and “star” sequences. It has long been known that as with many other type II restriction endonucleases *EcoRI* is capable of cleaving “star” sequences with low activity (Polisky et al., 1975; Goodman et al., 1977; Tikchonenko et al., 1978; Hsu and Berg, 1978; Malyguine et al., 1980; Gardner et al., 1982; Rosenberg and Greene, 1982; Lesser et al., 1990). This “star” activity can be promoted by modifying reaction conditions, in particular, by the addition of neutral solutes to the reaction (Goodman et al., 1977; Robinson and Sligar, 1993). Robinson and Sligar (1993) further showed that the increased cleavage of “star” sequences by *EcoRI* associated with neutral osmolytes directly correlates with water activity. One possible explanation for this effect is that osmotic stress modulates the equilibrium between the inactive binding mode of the enzyme and an energetically unfavorable, but enzymatically active mode of *EcoRI* binding to “star” sequences.

The apparent numbers of water molecules lost by the “star” sequence complexes, however, are quite modest at 2 Osm,  $\sim 20$  for the TAATTC sequence complex. The equilibrium experiments are quite difficult at higher osmotic stresses because dissociation rates of the protein from specific sequence become very slow and the concentrations of oligonucleotide DNA needed for competition become problematically large. The osmotic stress dependence of noncognate complex dissociation rates, however, can in principle give the same information about differences in associated water as equilibrium measurements, but can be measured to very high osmotic stresses. The osmotic sensitivity of the dissociation rate is determined by the difference in the number of waters between the initial binding state and the final dissociation transition state. The rate-limiting step for many protein-DNA complexes including *EcoRI* is the release of protein from a nonspecific complex. As we previously reported (Sidorova and Rau, 2001) the number of waters coupled with specific sequence dissociation is comprised of the 110 waters coupled to the difference between specific and nonspecific complexes. Another  $\sim 10$  waters for betaine glycine are linked to the dissociation of nonspecifically bound *EcoRI*, that is, there is a difference of  $\sim 10$  waters between the nonspecific complex and the dissociation transition state.

Dissociation rates of nonspecific complexes are not commonly measured. Typically, these rates are too fast for convenient measurement and can be monitored directly only with difficulty. The method we use might be of general interest and application. Dissociation rates of noncognate complexes can be extracted from the kinetics of appearance of specific site complex after *EcoRI* is added to a mixture of specific sequence fragment DNA and noncognate oligonucleotide under conditions of fast association rates. Three criteria indicate that we indeed are measuring nonspecific complex dissociation rates. The predicted dependence of the observed dissociation rate on noncognate sequence oligonucleotide concentration was confirmed (Fig. 3). The observed osmotic sensitivity of the inverted sequence oligonucleotide dissociation rate corresponding to  $\sim 15$  waters (Fig. 5) is consistent with  $\Delta N_w^{\ddagger} \sim 10$  waters for betaine glycine extracted previously (Sidorova and Rau, 2001) from specific site off-rates for the release of protein from a nonspecific complex. Lastly and most importantly, the dissociation rate constant for an *EcoRI* specific site complex is determined by the nonspecific dissociation rate,  $k_4$ , and the ratio of specific and nonspecific dissociation equilibrium constants (Sidorova and Rau, 2001),

$$k_d = \frac{k_4}{1 - f_{\infty}} \frac{K_{\text{nonsp}}}{K_{\text{sp}}} \quad (10)$$

The ratio of equilibrium constants can be calculated, therefore, from the dissociation rate constant for the inverted sequence oligonucleotide complex extrapolated to 0 Osm,  $k_4 \sim 4.1 \times 10^1 \text{ min}^{-1}$ , and for the specific site dissociation rate at the same salt, pH, temperature, and osmotic pressure measured previously and corrected for the difference in length,  $k_d(1 - f_{\infty}) \sim 2.9 \times 10^{-3} \text{ min}^{-1}$ . The ratio of equilibrium constants calculated from dissociation rates,  $1.4 \times 10^4$ , is in excellent agreement with the direct equilibrium measurement of  $1.2 \times 10^4$  shown in Fig. 1.

We have used here a single osmolyte, betaine glycine, at high osmotic pressures to vary water activity. Betaine glycine is an excellent solute for these high osmotic pressure studies because of its high solubility and strong exclusion from proteins (Courtenay et al., 2000; Timasheff, 1993) coupled with its low viscosity. In our previous work (Sidorova and Rau, 1996) we investigated many solutes differing by size and nature and found that the action of all those examined on *EcoRI*-DNA binding is osmotic at least up to 1.6 Osm. Robinson and Sligar (1993, 1998) examined still other solutes and reported an osmotic effect on *EcoRI* binding and cleavage through at least 2 Osm. The nonlinear dependences of  $\ln(K_{\text{sp}}/K_{\text{nonsp}})$  (Fig. 1 *b*) and of  $\ln(k_4)$  (Fig. 5) on osmotic pressure seen for “star” sequence complexes signifies that these complexes are either binding solute or losing water. We argue for the continued osmotic action of betaine glycine for several reasons. First, betaine glycine continues to act osmotically (linearly) on the nonspecific sequence complex

competition with the specific complex at the same pressures that the “star” sequence complex competition begins showing nonlinearity (Fig. 1 *b*). A second solute,  $\alpha$ -methyl glucoside shows the same nonlinearity at higher pressures for the “star” sequence complex in the competition assay as betaine glycine (Fig. 1 *b*). It is unlikely these two solutes would bind identically. We have previously seen that both  $\alpha$ -methyl glucoside and betaine glycine continue to act osmotically on the dissociation of the specific sequence complex at least through 3 Osm (Sidorova and Rau, 2001). Betaine glycine continues to act osmotically on the dissociation of the nonspecific, inverted sequence oligonucleotide complex through the entire 6 Osm pressure range. It is improbable that betaine glycine binds differently to the “star” sequence oligonucleotide complexes than to specific or inverted sequence oligonucleotide complexes. The data indicate that betaine glycine very likely continues to act osmotically on the “star” sequence complex through high pressures and that the nonlinearity of the plots in Figs. 1 *b* and 5 indicates the “star” sequence complexes are losing water.

At low osmotic pressures (<2 Osm), the dissociation rate of the TAATTC “star” sequence complex is  $\sim 1.6$ -fold slower than the nonspecific complex, comparable to the twofold change seen in the ratio of specific-noncognate equilibrium constants. Approximately 30 water molecules are seen coupled with the dissociation of this “star” sequence complex at low osmotic pressure, slightly larger than  $\sim 15$  waters seen for the inverted sequence oligonucleotide complex. The apparent loss of 15 more waters by the TAATTC complex is comparable to the decrease of  $\sim 20$  waters (from 105 to 85 waters) estimated for this complex from the osmotic dependence of the equilibrium measurements for the same pressure range (Fig. 1 *b*). Because the dissociation transition state remains unchanged as is indicated by the continued linearity of the inverted sequence dissociation rate data, the initial TAATTC complex is losing water with increasing osmotic pressure. A cubic fit to the data indicates that at 5-Osm pressure the TAATTC complex has lost some 90 waters compared with the inverted sequence oligonucleotide complex. The osmotic work,  $W$ , associated with the loss of  $\Delta N_w$  waters over the osmotic pressure range 0– $\Pi$  Osm is given by:

$$W = RT \left( \ln \left( \frac{k_4(\Pi)}{k_4(0)} \right) + \frac{\Pi \Delta N_w(\Pi)}{55.6} \right), \quad (11)$$

where the thermal factor  $RT = 0.6$  kcal/mol. The work needed to remove these 90 waters at 5 Osm from the TAATTC oligonucleotide complex is  $\sim 3.5$  kcal/mol. The osmotic energy gained through the release of sequestered water to the bulk solution is balanced by the resulting unfavorable interactions between DNA and protein surfaces.

The TAATTC oligonucleotide complex data can be well fit to a simple two-state model assuming equilibrium between the two states with different numbers of associated waters. The best fit gives a difference of  $\sim 105$  waters between the two states. Although we do not know where this water is located, it seems likely it is the same excess water in the nonspecific complex that is lost upon specific binding. The dehydrated state would then resemble the specific recognition sequence complex. The best fit also gives a free energy for the alternate dehydrated complex in the absence of solute that is  $\sim 3.2$  kcal/mol less favorable than the hydrated state. A more complicated two-state model that accounts for the fraction of protein actually bound to the “star” sequence itself changes the fitting parameters only slightly,  $\Delta G = -2.8$  kcal/mol and  $\Delta N_w = 90$ .

It is substantially more difficult to remove water from the CAATTC “star” sequence oligonucleotide complex. From a quadratic fit to the data, only  $\sim 25$  waters are apparently lost at 5 Osm from the hydrated complex initially present at low pressures. The osmotic work necessary to remove these 25 waters,  $\sim 0.9$  kcal/mol, compared with  $\sim 0.45$  kcal/mol required to remove the first 25 waters from the TAATTC “star” sequence complex. Although not enough water is lost from the CAATTC oligonucleotide complex to give a meaningful two-state model fit, an energy difference between the two states of  $\sim 6.2$  kcal/mol can be estimated if  $\Delta N_w^*$  is fixed at 105 waters as for the TAATTC oligonucleotide complex.

Of the “star” sequences, the TAATTC is the most readily cleaved by *EcoRI* and CAATTC the most cleavage resistant of the first basepair substitutions. Lesser et al. (1990) estimated energy differences for *EcoRI* binding to the cleavage transition state between the recognition sequence and various “star” sequences at 20°C. In particular, a free energy of 6.6 kcal/mol was reported for a TAATTC “star” sequence and 9.5 kcal/mol for a CAATTC sequence. We can compare these values with the energy differences we estimate for the dehydrated complexes of the TAATTC and CAATTC “star” sequences relative to specific sequence complex. This comparison is strictly valid only if the activation energy differences between the initial complexes with specific and “star” sequences and the cleavage transition state are small. The oligonucleotides used here and by Lesser et al. (1990) have different flanking sequences that can strongly affect activity (Jen-Jacobson, 1997) and also have different lengths and consequently different numbers of possible nonspecific sites that will affect relative binding energies. Nonetheless, from the competition experiments, with no osmolyte both TAATTC and CAATTC oligonucleotide complexes are  $\sim 5$  kcal/mol less stable than the specific recognition sequence complex. The dehydrated, TAATTC complex is an additional  $\sim 3$  kcal/mol less stable than the hydrated binding mode complex, or, therefore,  $\sim 8$  kcal/mol less stable than the specific recognition sequence complex. Similarly, we estimate that the

dehydrated CAATTC complex is  $\sim 11$  kcal/mol less stable than the specific sequence complex. The agreement between our results and Lesser et al. (1990) is reasonable enough to suggest that the dehydrated "star" sequence complex is an enzymatically active form. In particular, both methods give  $\sim 3$  kcal/mol for the difference between TAATTC and CAATTA complexes. This difference should be much less sensitive to cleavage activation energies and the effect of flanking sequences and DNA length.

## CONCLUSIONS

On the basis of the osmotic stress measurements, the nonspecific complex for *EcoRI* binding seems to be a well-defined structure. All noncognate sequence complexes we have investigated, even sequences that differ by only a single first basepair from the recognition sequence, sequester 105–110 waters more than the specific sequence complex. In analogy with the nonspecific complex of *BamHI* (Viadiu and Aggarwal, 2000), the protein likely contacts the sugar-phosphate backbone of DNA rather than bases in the major groove. The small differences in noncognate binding constants among sequences, e.g., the approximate twofold increase in binding for "star" sequences seen here to compare with inverted sequence could simply result from an indirect readout of sequence dependent helical parameters. A difference between "star" sequence and nonspecific complexes, however, is revealed at higher osmotic pressures. "Star" sequence complexes appear to lose water as the higher osmotic pressures or an alternate conformation with fewer sequestered waters is stabilized. If analyzed as two states, the alternate complex is almost as dehydrated as the specific sequence complex. Osmotic stress can be used to modulate the energy difference between a nonspecific complex and this specific-like binding mode. The work needed to remove virtually all the water from the "star" sequence complex, indicates that this dehydrated first basepair "star" sequence complex is destabilized by an estimated 8–11 kcal/mol from the specific sequence complex, depending on the "star" sequence. This large energy difference is consistent with a rigid and interconnected protein recognition surface that is unable to adapt to recognize mismatches. In this way the x-ray structure of the specific *EcoRI* complex (Kim et al., 1990; McClarin et al., 1986; Rosenberg, 1991) is different from *lac* (Lewis et al., 1996) or  $\lambda$ -Cro repressor-DNA (Albright and Matthews, 1998) structures that do not show the extensive DNA-protein and protein-protein bonding interconnections seen for *EcoRI* binding. The loss of water from the "star" sequence complexes provides a structural basis for the enzymatic "star" activity of *EcoRI* and a natural link between "star" activity and osmotic pressure as was anticipated by Pingoud and Jeltsch (1997). We cannot unambiguously discriminate between a gradual loss of water from the star sequence complexes and the two-state model. The calculated energies to remove water will

depend somewhat on the actual mechanism, but only slightly. More importantly, the key component that differentiates *EcoRI* binding to various sequences is found in the water sequestered by the complexes.

We are deeply grateful to V. A. Parsegian for the valuable discussion and support, to L. Jen-Jacobson for the kind gift of highly purified *EcoRI*, and to Shakir Muradymov for his enthusiastic help with the experiments.

## REFERENCES

- Albright, R. A., and B. W. Matthews. 1998. Crystal structure of lambda-Cro bound to a consensus operator at 3.0 Å resolution. *J. Mol. Biol.* 280:137–151.
- Courtenay, E. S., M. W. Capp, C. F. Anderson, and M. T. Record, Jr. 2000. Vapor pressure osmometry studies of osmolyte-protein interactions: implications for the action of osmoprotectants in vivo and for the interpretation of osmotic stress experiments in vitro. *Biochemistry.* 39:4455–4471.
- Davis-Searles, P. R., A. S. Morar, A. J. Saunders, D. A. Erie, and G. J. Pielak. 1998. Sugar-induced molten-globule model. *Biochemistry.* 37:17048–17053.
- Ehbrecht, H. J., A. Pingoud, C. Urbanke, G. Maass, and C. Gualerzi. 1985. Linear diffusion of restriction endonucleases on DNA. *J. Biol. Chem.* 260:6160–6166.
- Frank, D. E., R. M. Saecker, J. P. Bond, M. W. Capp, O. V. Tsodikov, S. E. Melcher, M. M. Levandovski, and M. T. Record, Jr. 1997. Thermodynamic of the interactions of *lac* repressor with variants of the symmetric *lac* operator: effects of converting a consensus site to a non-specific site. *J. Mol. Biol.* 267:1186–1206.
- Fried, M. G., and D. M. Crothers. 1981. Equilibria and kinetics of *lac* repressor-operator interactions by polyacrylamide gel electrophoresis. *Nucleic Acids Res.* 9:6505–6525.
- Fried, M. G., D. F. Stickle, K. V. Smirnakis, C. Adams, D. MacDonald, and P. Lu. 2002. Role of hydration in the binding of *lac* repressor to DNA. *J. Biol. Chem.* 277:50676–50682.
- Gardner, R. C., A. J. Howarth, J. Messing, and R. J. Shepherd. 1982. Cloning and sequencing of restriction fragments generated by *EcoRI*\*. *DNA.* 1:109–115.
- Garner, M. M., and D. C. Rau. 1995. Water release associated with specific binding of *gal* repressor. *EMBO J.* 14:1257–1263.
- Garner, M. M., and A. Revzin. 1981. A gel electrophoresis method for quantifying the binding of proteins to specific DNA regions: application to components of the *Escherichia coli* lactose operon regulatory system. *Nucl. Acids Res.* 9:3047–3060.
- Goodman, H. M., P. J. Green, D. E. Garfin, and H. W. Boyer. 1977. DNA site recognition by the *EcoRI* restriction endonuclease and modification methylase. In *Nucleic Acid: Protein Recognition*. H. J. Vogel, editor. Academic Press, New York. 329–359.
- Hard, T., and T. Lundback. 1996. Thermodynamics of sequence-specific protein-DNA interactions. *Biophys. Chem.* 62:121–139.
- Hsu, M., and P. Berg. 1978. Altering the specificity of restriction endonuclease: effect of replacing  $Mg^{2+}$  with  $Mn^{2+}$ . *Biochemistry.* 17:131–138.
- Jack, W. E., B. J. Terry, and P. Modrich. 1982. Involvement of outside DNA sequences in the major kinetic path by which *EcoRI* endonuclease locates and leaves its recognition sequence. *Proc. Natl. Acad. Sci. USA.* 79:4010–4014.
- Jeltsch, A., J. Alves, H. Wolfes, G. Maas, and A. Pingoud. 1994. Pausing of the restriction endonuclease *EcoRI* during linear diffusion on DNA. *Biochemistry.* 33:10215–10219.
- Jen-Jacobson, L. 1997. Protein-DNA recognition complexes: conservation of structure and binding energy in the transition state. *Biopolymers.* 44:153–180.

- Khrapunov, S., and M. Brenowitz. 2004. Comparison of the effect of water release on the interaction of the *Saccharomyces cerevisiae* TATA binding protein (TBP) with TATA box sequences composed of adenosine or inosine. *Biophys. J.* 86:371–383.
- Kim, Y., J. C. Grable, R. Love, P. J. Greene, and J. M. Rosenberg. 1990. Refinement of EcoRI endonuclease crystal structure: a revised protein chain tracing. *Science.* 249:1307–1309.
- Lesser, D. R., M. R. Kurpiewski, and L. Jen-Jacobson. 1990. The energetic basis of specificity in the EcoRI endonuclease-DNA interaction. *Science.* 250:776–786.
- Lewis, M., G. Chang, N. C. Hortin, M. A. Kercher, H. C. Pace, M. A. Schumacher, R. G. Brennan, and P. Lu. 1996. Crystal structure of the lactose operon repressor and its complexes with DNA and inducer. *Science.* 271:1247–1254.
- Li, L., and K. S. Matthews. 1997. Differences in water release with DNA binding by ultrathin and deformed homeodomains. *Biochemistry.* 36:7003–7011.
- Lundback, T., H. Hansson, S. Knapp, R. Ladenstein, and T. Hard. 1998. Thermodynamic characterization of non-sequence-specific DNA-binding by the Sso7d protein from *Sulfolobus solfataricus*. *J. Mol. Biol.* 276:775–786.
- Lynch, T. W., and S. G. Sligar. 2000. Macromolecular hydration changes associated with BamHI binding and catalysis. *J. Biol. Chem.* 275:30561–30565.
- Malyguine, E., P. Vannier, and P. Yot. 1980. Alteration of the specificity of restriction endonucleases in the presence of organic solvents. *Gene.* 8:163–177.
- McClarín, J. A., C. A. Frederick, B.-C. Wang, H. W. Boyer, J. Grable, and J. M. Rosenberg. 1986. Structure of the DNA-EcoRI endonuclease recognition complex at 3 Å resolution. *Science.* 234:1526–1541.
- Newman, M., T. Strzelecka, L. F. Dorner, I. Schildkraut, and A. K. Aggarwal. 1995. Structure of BamHI endonuclease bound to DNA: partial folding and unfolding on DNA binding. *Science.* 269:656–663.
- Parsegian, V. A., R. P. Rand, and D. C. Rau. 1995. Macromolecules and water: probing with osmotic stress. *Methods Enzymol.* 259:43–94.
- Parsegian, V. A., R. P. Rand, and D. C. Rau. 2000. Osmotic stress, crowding, preferential hydration, and binding: a comparison of perspectives. *Proc. Natl. Acad. Sci. USA.* 97:3987–3992.
- Pingoud, A., and A. Jeltsch. 1997. Recognition and cleavage of DNA by type-II restriction endonucleases. *Eur. J. Biochem.* 246:1–22.
- Polisky, B., P. Greene, D. E. Garfin, B. J. McCarthy, H. M. Goodman, and H. W. Boyer. 1975. Specificity of substrate recognition by the EcoRI restriction endonuclease. *Proc. Natl. Acad. Sci. USA.* 72:3310–3314.
- Poon, J., M. Bailey, D. J. Winzor, B. E. Davidson, and W. H. Sawyer. 1997. Effects of molecular crowding on the interaction between DNA and the *Escherichia coli* regulatory protein TyrR. *Biophys. J.* 73:3257–3264.
- Preisler, R. S., H.-H. Chen, M. F. Colombo, Y. Choe, B. J. Short, Jr., and D. C. Rau. 1995. The B-form to Z-form transition of poly(dG-m5dC) is sensitive to neutral solutes through an osmotic stress. *Biochemistry.* 34:14400–14407.
- Robinson, C. R., and S. G. Sligar. 1993. Molecular recognition mediated by bound water: a mechanism for star activity of the restriction endonuclease EcoRI. *J. Mol. Biol.* 234:302–306.
- Robinson, C. R., and S. G. Sligar. 1996. Participation of water in Hin recombinase-DNA recognition. *Protein Sci.* 5:2119–2124.
- Robinson, C. R., and S. G. Sligar. 1998. Changes in solvation during DNA binding and cleavage are critical to altered specificity of the EcoRI endonuclease. *Proc. Natl. Acad. Sci. USA.* 95:2186–2191.
- Rosenberg, J. M. 1991. Structure and function of restriction endonucleases. *Curr. Opin. Struct. Biol.* 1:104–113.
- Rosenberg, J. M., and P. Greene. 1982. EcoRI\* specificity and hydrogen bonding. *DNA.* 1:117–124.
- Schwabe, J. W. R. 1997. The role of water in protein-DNA interactions. *Curr. Opin. Struct. Biol.* 7:124–134.
- Sidorova, N. Y., and D. C. Rau. 1996. Differences in water release for the binding of EcoRI to specific and nonspecific DNA sequences. *Proc. Natl. Acad. Sci. USA.* 93:12272–12277.
- Sidorova, N. Y., and D. C. Rau. 1999. Removing water from an EcoRI-noncognate DNA complex with osmotic stress. *J. Biomol. Struct. and Dynam.* 17:19–31.
- Sidorova, N. Y., and D. C. Rau. 2000. The dissociation rate of the EcoRI-DNA specific complex is linked to water activity. *Biopolymers.* 53:363–368.
- Sidorova, N. Y., and D. C. Rau. 2001. Linkage of EcoRI dissociation from its specific DNA recognition site to water activity, salt concentration, and pH: separating their roles in specific and non-specific binding. *J. Mol. Biol.* 310:801–816.
- Swaminathan, C. P., U. T. Sankpal, D. N. Rao, and A. Suroliya. 2002. Water-assisted dual mode cofactor recognition by HhaI DNA methyltransferase. *J. Biol. Chem.* 277:4042–4049.
- Szwajkajzer, D., and J. Carey. 1997. Molecular and biological constraints on ligand-binding affinity and specificity. *Biopolymers.* 44:181–198.
- Takeda, Y., P. D. Ross, and C. P. Mudd. 1992. Thermodynamics of Cro protein-DNA interactions. *Proc. Natl. Acad. Sci. USA.* 89:8180–8184.
- Takeda, Y., A. Sarai, and V. M. Rivera. 1989. Analysis of the sequence-specific interactions between Cro repressor and operator DNA by systematic base substitution experiments. *Proc. Natl. Acad. Sci. USA.* 86:439–443.
- Terry, B. J., W. E. Jack, and P. Modrich. 1985. Facilitated diffusion during catalysis by EcoRI endonuclease. Nonspecific interactions in EcoRI catalysis. *J. Biol. Chem.* 260:13130–13137.
- Tikhonchenko, T. I., E. V. Karamov, B. A. Zavizion, and B. S. Naroditsky. 1978. EcoRI\* activity: enzyme modification of activation of accompanying endonuclease. *Gene.* 4:195–212.
- Timasheff, S. N. 1993. The control of protein stability and association by weak interactions with water: how do solvents affect these processes? *Annu. Rev. Biophys. Biomol. Struct.* 22:27–65.
- Viadiu, H., and A. K. Aggarwal. 2000. Structure of BamHI bound to nonspecific DNA: a model for DNA sliding. *Mol. Cell.* 5:889–895.
- Vossen, K. M., R. Wolz, M. A. Daugherty, and M. G. Fried. 1997. Role of macromolecular hydration in the binding of the *Escherichia coli* cyclic AMP receptor to DNA. *Biochemistry.* 36:11640–11647.
- Wenner, J. R., and V. Bloomfield. 1999. Osmotic pressure effects on EcoRV cleavage and binding. *J. Biomol. Struct. Dyn.* 17:461–471.
- Wright, D. J., W. E. Jack, and P. Modrich. 1999. The kinetic mechanism of EcoRI endonuclease. *J. Biol. Chem.* 274:31896–31902.
- Wu, J., K. M. Parkhurst, R. M. Powell, M. Brenowitz, and L. J. Parkhurst. 2001. DNA bends in TATA-binding protein-TATA complexes in solution are DNA sequence-dependent. *J. Biol. Chem.* 276:14614–14622.