



## MEETING ABSTRACT

## Open Access

# Mild cognitive disorder and depression: treatment with combination of galantamine and escitalopram

Julio Zarra<sup>1\*</sup>, Luisa Schmidt<sup>2</sup>

From 1<sup>st</sup> International Congress on Neurobiology and Clinical Psychopharmacology and European Psychiatric Association Conference on Treatment Guidance Thessaloniki, Greece. 19-22 November 2009

## Background

To evaluate the efficacy of galantamine-escitalopram combination in patients with Mild Cognitive Disorder and Depression. So there is a possible relation between the deficit of cerebral oxygenation and depression or relation between the serotonin system and cholinergic system in relation with disease comorbidity cognitive-depression.

## Objective

To evaluate the therapeutic response in patients with comorbidity between Mild Cognitive Disorder and Depression in treatment with Galantamine, Escitalopram and the two drugs in combination.

## Materials and methods

A group of 300 patients with symptoms of Mild Cognitive Disorder and Depression (DSM IV-R criteria) were separated in 3 groups of 100 patients. Each group received different treatment in an 8 months period:

Group 1: Galantamine 16 mg/day.

Group 2: Escitalopram 20 mg/day.

Group 3: both drugs, same dose.

## Results

The therapeutic response evaluated in Hamilton Scale for Depression (HAM-D), Montgomery and Åsberg Depression Rating Scale (M.A.D.R.S.), Mini Mental State Examination (M.M.S.E.) and Global Clinical Impression (G.C.I.) scores during 8 months. In the third group who received the two drugs associated, had much better response than the others and "brain enhancer".

## Conclusions

The group who received the combination of the nootropic agent Galantamine with antidepressant (SSRIs) Escitalopram had a relevant satisfactory therapeutic response (the best result), so there is a possible relation between the deficit in cholinergic systems and depression. Could be cerebral cholinergic systems deficit a generator of Depressive Disorder?

Attention and memory functions are closely tied to the cholinergic neurotransmitter system. The cholinergic system is one of the neurotransmitter systems implicated in the pathophysiology of mood disorders. Evidence suggests that during major depressive episodes, the cholinergic system is hypersensitive to acetylcholine.

## Author details

<sup>1</sup>Servicio de Psiquiatría, Hospital Italiano de La Plata, Buenos Aires, Argentina.

<sup>2</sup>Servicio de Salud Mental, Hospital Centenario, Gualaguaychú, Argentina.

Published: 22 April 2010

## Reference

1. Blesa R: Galantamine: Therapeutic effects beyond Cognition. *Dement Geriatr Cogn Disord* 2000, **11**(suppl 1):28-34.

doi:10.1186/1744-859X-9-S1-S100

**Cite this article as:** Zarra and Schmidt: Mild cognitive disorder and depression: treatment with combination of galantamine and escitalopram. *Annals of General Psychiatry* 2010 **9**(Suppl 1):S100.

<sup>1</sup>Servicio de Psiquiatría, Hospital Italiano de La Plata, Buenos Aires, Argentina