

Available online at www.sciencedirect.com

ScienceDirect

Nuclear Physics B 886 (2014) 312–327

NUCLEAR
PHYSICS Bwww.elsevier.com/locate/nuclphysb

Quantifying the reheating temperature of the universe

Anupam Mazumdar^a, Bryan Zaldivar^{b,*}^a *Consortium for Fundamental Physics, Lancaster University, Lancaster LA1 4YB, UK*^b *Instituto de Fisica Teorica, IFT-UAM/CSIC, 28049 Madrid, Spain*

Received 11 April 2014; received in revised form 30 June 2014; accepted 1 July 2014

Available online 7 July 2014

Editor: Hong-Jian He

Abstract

The aim of this paper is to determine an exact definition of the reheat temperature for a generic perturbative decay of the inflaton. In order to estimate the reheat temperature, there are two important conditions one needs to satisfy: (a) the decay products of the inflaton must dominate the energy density of the universe, i.e. the universe becomes completely radiation dominated, and (b) the decay products of the inflaton have attained local thermodynamical equilibrium. For some choices of parameters, the latter is a more stringent condition, such that the decay products may thermalise much after the beginning of radiation–domination. Consequently, we have obtained that the reheat temperature can be much lower than the standard lore estimation. In this paper we describe under what conditions our universe could have *efficient* or *inefficient* thermalisation, and quantify the reheat temperature for both the scenarios. This result has an immediate impact on many applications which rely on the thermal history of the universe, in particular gravitino abundance.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP³.

1. Introduction

The transition from a cold inflating universe to a hot thermal universe depends *solely* on the inflaton mass, m_ϕ , its coupling α_ϕ to the relevant degrees of freedom (d.o.f.), and the dominant coupling between the decay products. In the case of Standard Model (SM) particles, it is predominantly the strong interaction, $\alpha_s \sim 1/30$. This epoch is known as reheating [1], or preheating [2] (for a review see [3]). In this paper we will mostly concentrate on the case where the inflaton

* Corresponding author.

has a small Yukawa coupling to the relevant d.o.f., which would typically yield a perturbative decay of the inflaton to its almost massless quarks, leptons and gluons. This is well justified for an SM *gauge singlet* inflaton, since the SM quarks and leptons are *chiral* in nature, and therefore the lowest order couplings are determined by the dimensional 5 operators in the potential, see [4]. Inflation could be driven by many independent sectors [5], but what matters is the last field which is responsible for finally reheating the universe in our patch for the success of Big Bang Nucleosynthesis (BBN) [6].

Especially, an SM gauge singlet inflaton could also couple to the SM Higgs with a 4-dimensional coupling, but through quartic coupling the inflaton never decays unless ϕ develops a VEV (vacuum expectation value): it rather leads to $\phi\phi \leftrightarrow HH$ scatterings, where ϕ is the inflaton and H denotes the SM Higgs. In order to deplete the inflaton quanta it is still important to rely on the perturbative decay of the inflaton [7].¹

Typically, the reheating process is assumed to be instantaneous, with an efficient energy density conversion from the inflaton to the relativistic plasma. Within this framework the concept of reheating temperature T_{rh} has been defined, see [1,11], ultimately relying on the assumption of the presence of local thermal equilibrium (LTE) at the very instant of conversion from the initial coherent oscillations of the inflaton domination to the radiation domination.

The aim of this work is to determine a proper definition of the reheat temperature of the universe keeping in mind when the LTE is established along with the fact that the inflaton has completely decayed into radiation. When and how should we evaluate the reheat temperature is an important question for a number of applications ranging from evaluating the baryonic asymmetry, dark matter abundance and the success of BBN [11]. In this paper we shall put down the criteria of estimating the reheat temperature, based on when the inflaton decay products attain their thermalisation. Depending on whether the decay products of the inflaton thermalise before or after the radiation has dominated the universe, the reheat temperature will be very different. In either situation the notion of reheat temperature only makes sense when the universe is completely dominated by the radiation bath.

If thermalisation of the ambient plasma occurs during the coherent oscillations of the inflaton, one may be able to associate a maximum temperature with the relativistic species [11,12], but if the thermalisation time scale is longer than that of the inflaton-to-radiation domination transition time scale, the notion of temperature does not make sense until the universe reaches its full LTE. In this respect there could be three regimes of interest which we will discuss in this paper:

1. Instant thermalisation: when the inflaton decay products instantly thermalise upon decay.
2. Efficient thermalisation: when the inflaton decay products thermalise right at the instant when radiation epoch starts dominating the universe.
3. Delayed thermalisation: when the inflaton decay products thermalise deep inside the radiation dominated epoch after the transition from inflaton-to-radiation domination had occurred.

This paper is organised as follows. In Section 2 we set the stage and write down the relevant equations for our analysis. The standard lore about the reheating epoch is briefly commented in Section 3. Section 4 is devoted to present our analysis, in which we study the conditions under

¹ Our treatment is very general and it can be applicable to supersymmetric theories. However there is a word of caution on how the inflaton couples to the supersymmetric Standard Model degrees of freedom, which depends very much on the origin of the inflaton. If inflaton is SM gauge singlet, see [8], if inflaton is SM gauge invariant field, such as one belonging to the supersymmetric flat directions of squarks and sleptons [9], see [10].

which the plasma attains thermalisation. Later on, in Section 5 we discuss the concept of reheat temperature such as to properly capture the issues of thermalisation. Finally, we conclude in Section 6.

2. Key assumptions and equations

For the sake of simplicity, we will assume *universal* inflaton coupling, α_ϕ , to all its decay products, determined by the number of relativistic d.o.f. g_* . Since the decay products of the inflaton are light, just from kinematics, they will typically have an initial momentum roughly given by: $m_\phi/2$ for two-body decay, or $m_\phi/3$ for a three-body decay processes. The inflaton is assumed here to be an SM gauge singlet – it will decay universally to all its decay products, i.e. all the relativistic species g_* would be excited.

Once the decay products are all excited there are two important processes which lead to thermalisation of all the d.o.f., or establish an LTE. Whereas a detailed thermalisation analysis of the plasma is out of the scope of this paper, some of its features are essential to our analysis, see Refs. [13,14]:

1. *Kinetic equilibrium*: Redistribution of the momentum between different decay particles. This can be achieved by number conserving $2 \rightarrow 2$ scatterings with gauge boson exchange in the t -channel [13,14].
2. *Chemical equilibrium*: Number violating $2 \rightarrow 3$ scatterings via t -channel are required to establish the chemical equilibrium [13,14]. Higher order process are suppressed by further powers of the gauge coupling. Typically $2 \rightarrow 3$ interaction rate is higher than that of $2 \rightarrow 2$.

The inelastic cross section for $2 \rightarrow 3$ processes are roughly estimated by [14]:

$$\sigma \sim \frac{\alpha_s^3}{p(t)^2} \log\left(\frac{m_\phi^2}{p(t)^2}\right), \quad (1)$$

where $\alpha_s \sim 1/30$ is the typical strong gauge coupling of the SM, and $p(t)$ is the 3-momentum transferred in the scattering process.

There are two interesting regimes which we will discuss below:

1. **t -channel enhancement**: If the scatterings $2 \rightarrow 3$ processes via t -channel are mediated by light or massless gauge bosons, the cross section in question has an infrared divergence, which can be reasonably cut off by the Debye length, given by the inverse of the average separation between the two quanta, i.e. $\bar{r} \sim n^{-1/3}$, where n is the number density of the particles in the plasma. In this case the scattering rate is extremely fast due to the infrared divergence and would yield an efficient thermalisation of the plasma, as discussed in Ref. [18].
2. **t -channel suppression**: As noted in [8,18] this singularity is absent if, for example, the scattering happens via exchange of massive gauge boson. There, the thermalisation process may be considerably delayed due to suppression in the scattering rate. Such examples have been investigated in Ref. [8], in presence of supersymmetric flat directions developing VEV or finite temperature effects, which naturally gives rise to massive gauge boson. In those cases, the most important processes for thermalisation are either $2 \rightarrow 3$ scatterings with scalar boson exchanges, or s -channel resonant gauge boson exchange. In either case, the infrared divergences disappear.

In this work we are going to discuss both the possibilities, although we will concentrate more on the delayed scenario, since the situation with enhanced cross sections has been extensively discussed in the literature [18].

On the other hand, the evolution of the inflaton, and the relativistic decay product’s energy densities during the reheating period is described by the coupled set of Boltzmann equations, see [11]:

$$\begin{cases} \dot{\rho}_\phi + 3H(t)\rho_\phi = -\Gamma_\phi\rho_\phi \\ \dot{\rho}_R + 4H(t)\rho_R = \Gamma_\phi\rho_\phi + \Gamma_{th}(\rho_R - \rho_R^{eq}), \end{cases} \quad (2)$$

where the dots denote derivatives w.r.t. the physical time, $\rho_\phi(\rho_R)$ is the energy density of inflaton (radiation), being ρ_R^{eq} the equilibrium one; $H(t)$ is the Hubble parameter accounting for the expansion of the universe; $\Gamma_\phi \equiv \alpha_\phi m_\phi$ is the inflaton decay rate,² and Γ_{th} is the reaction rate responsible for thermalisation of the radiation plasma. Of course, once LTE is attained, the direct and inverse interactions among relativistic species counterbalance each other and the evolution of ρ_R is dictated solely by the inflaton source and the Hubble expansion.

3. Assuming LTE is established soon after inflaton decay

Previous works which are relevant to our study have assumed LTE while studying the evolution of the relativistic species during the reheating period [11], see however [12–14,18] for emphasising the importance of acquiring LTE. At any epoch during reheating, as long as there is a relativistic bath in thermal equilibrium, we can extract an instantaneous temperature as:

$$T(t) = \left[\frac{30}{\pi^2} \rho_R(t) / g_*(t) \right]^{1/4} \quad (3)$$

For a constant g_* during the whole period, the evolution of the temperature according to Eq. (3) is such that it has a maximum T_{max} [11,12], which can be estimated as:

$$T_{max} \simeq \left[\frac{1.57}{\pi^3 g_*} \right]^{1/4} \sqrt{M_P} (\Gamma_\phi H_I)^{1/4}, \quad (4)$$

being H_I the initial Hubble rate. Indeed, T_{max} can be potentially much larger than the reheating temperature, T_{rh} . The latter is usually defined as the temperature of the plasma assuming an instantaneous conversion of the inflaton’s energy density into radiation, at the time when $H(t) \approx \Gamma_\phi$, such that:

$$T_{rh} = \left(\frac{90}{8\pi^3 g_*} \right)^{1/4} \sqrt{\Gamma_\phi M_P}. \quad (5)$$

4. When is LTE attained?

However, LTE has to be attained and should not be taken for granted from the onset of the inflaton decay. In our analysis *we do not assume LTE* as a given condition for the relativistic species. Instead, we evaluate when and for which region of the inflaton parameters, m_ϕ and α_ϕ for a fixed $\alpha_s = 1/30$, this condition is achieved.

² See [19] for a recent very detailed analysis on Γ_ϕ .




Fig. 1. Radiation energy density (blue line), ρ_R , and inflaton energy density (red line), ρ_ϕ , as a function of the scale factor, for $m_\phi = 10^{13}$ GeV and $\alpha_\phi = 10^{-11}$. The power laws indicate the behaviour of ρ_R in the different regimes. The region in grey represents the reheating epoch, which by the *standard lore* finishes when radiation dominates the expansion (see text for details). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

There are (as justified later) two regions of the parameter space (α_ϕ, m_ϕ), for which Eq. (2) can be simplified such that the term $\Gamma_{th}(\rho_R - \rho_R^{eq})$ can be safely discarded:

1. Very small α_ϕ and very large m_ϕ , for which Γ_{th} is very small:

$$\Gamma_{th} \ll \Gamma_\phi \cdot \left(\frac{\rho_\phi}{\rho_R}\right), \quad \Gamma_{th} \ll H \tag{6}$$

2. Very large α_ϕ and very small m_ϕ , for which $\rho_R \approx \rho_R^{eq}$:

$$\Gamma_{th} \gg \Gamma_\phi \cdot \left(\frac{\rho_\phi}{\rho_R}\right), \quad \Gamma_{th} \gg H \tag{7}$$

We will justify the notion of very small and very large below. For these two cases, Eq. (2) simplifies to (working with a comoving coordinate, $x \equiv a(t) \times m_\phi$, where $a(t)$ is the scale factor) [12]:

$$\begin{cases} \frac{d\Phi}{dx} = -\left(\sqrt{\frac{3}{8\pi}} \frac{M_P}{m_\phi} \alpha_\phi\right) \frac{x\Phi}{\sqrt{R+x\Phi}} \\ \frac{dR}{dx} = \left(\sqrt{\frac{3}{8\pi}} \frac{M_P}{m_\phi} \alpha_\phi\right) \frac{x^2\Phi}{\sqrt{R+x\Phi}} \end{cases} \tag{8}$$

with

$$\Phi \equiv \rho_\phi m_\phi^{-4} x^3, \quad R \equiv \rho_R m_\phi^{-4} x^4. \tag{9}$$

The initial condition is:

$$R(x_I) = 0, \quad \Phi_I \equiv \Phi(x_I) = \frac{H_I^2 M_P^2}{8\pi/3} \cdot m_\phi^{-4} x_I^3, \tag{10}$$

where the subindex I refers to initial values. In many inflationary scenarios it is a good approximation to take $H_I \sim m_\phi$.

We have solved Eq. (8) numerically, and the result is shown in Fig. 1, where for illustration we have taken $m_\phi = 10^{13}$ GeV and $\alpha_\phi = 10^{-11}$. We can infer that the radiation energy density (blue

line) peaks very fast, around $x = x_{\max} \sim 1.5x_I$, followed by a dilution due to the expansion. The position of the maximum is independent of the inflaton parameters. We also show for reference the inflaton energy density (red line), which as we can see completely dominates the expansion of the universe until the end of the reheating epoch. Analytically, during the inflaton–dominated period the radiation energy density goes like:

$$\begin{aligned} \rho_R^{id}(x) &\approx \frac{2}{5} \sqrt{\frac{3}{8\pi}} \Gamma_\phi m_\phi^2 M_P \sqrt{\Phi_I} x^{-3/2}, \quad x_I \ll x < x_{rh} \\ &\approx \frac{0.15}{\pi} M_P^2 m_\phi^2 \alpha_\phi \left(\frac{x_I}{x}\right)^{3/2} \end{aligned} \tag{11}$$

whereas for radiation–domination the expected x^{-4} -law is recovered:

$$\rho_R^{rd}(x) \approx \rho_R^{id}(x_{rh}) \left(\frac{x_{rh}}{x}\right)^4, \quad x_{rh} < x. \tag{12}$$

Here x_{rh} (to be computed below) encodes the moment at which reheating ends. The super-indices (*id*) and (*rd*) stem for (inflaton–domination) and (radiation–domination), respectively.

The condition under which the plasma enters in thermal equilibrium can be naively estimated by the requirement

$$\Gamma_{\text{th}} = n_R(x) \langle \sigma(x)v \rangle > H(x), \tag{13}$$

where we approximate the cross-section σ by Eq. (1), $v \approx c$ for relativistic species, and $n_R(x)$ is the relativistic number density. The latter can be directly extracted by solving Eq. (8) in terms of number densities instead of energy densities. Assuming 2-body decays of the inflaton (our results will not be affected much if we assume 3-body decay of the inflaton), see also [14]:

$$n_R(x) \approx 2n_\phi^I \left[1 - e^{(-\Gamma_\phi \int_{x_0}^x \frac{d\tilde{x}}{\tilde{x}H(\tilde{x})})} \right] \left(\frac{x_I}{x}\right)^3 \tag{14}$$

where the initial inflaton number density, $n_\phi^I \sim \rho_\phi^I/m_\phi$, as well as $H(x)$, are computed according to the our numerical solution of Eq. (8). Analytical estimations of Eq. (14) can be obtained, as for the case of ρ_R , in two regimes.

1. **During inflaton–domination:** In this case, $R(x)$ gives a negligible contribution to the Hubble rate, whereas Φ remains approximately constant, $\Phi \approx \Phi_I$. In this case, it is straightforward to obtain:

$$n_R^{id}(x) \simeq 2n_\phi^I (1 - e^{-\kappa x^{3/2}}) \left(\frac{x_I}{x}\right)^3 \approx 2n_\phi^I \kappa x_I^3 x^{-3/2} \simeq \frac{0.5}{\pi} M_P^2 m_\phi \alpha_\phi \left(\frac{x_I}{x}\right)^{3/2}, \tag{15}$$

with $\kappa = (2/3)\alpha_\phi/x_I^{3/2}$. The superscript ‘*id*’ denotes *inflaton–domination*, since the inflaton oscillations are dominating over the relativistic species.

2. **During radiation–domination:** On the other hand, for *radiation–domination*, denoted below by the superscript ‘*rd*’, we clearly have:

$$n_R^{rd}(x) \simeq 2n_\phi^I \left(\frac{x_I}{x}\right)^3 = \frac{3}{4\pi} M_P^2 m_\phi \left(\frac{x_I}{x}\right)^3. \tag{16}$$




Fig. 2. Radiation number density as a function of the scale factor, for $m_\phi = 10^{13}$ GeV and $\alpha_\phi = 10^{-11}$. The solid red line is the solution of Eq. (14), where $H(x)$ is computed numerically from Eq. (8). Dashed black line is the solution in Eq. (15), whereas the dotted black line is the solution in Eq. (16). The solid black vertical lines is the value of x_{rh} according to Eq. (17). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

The value x_{rh} at which the regime changes could be computed in several ways, one of which is demanding $n_R^{id}(x_{rh}) = n_R^{rd}(x_{rh})$, resulting in:

$$x_{rh} = \kappa^{-2/3} \simeq \frac{1.3x_I}{\alpha_\phi^{2/3}}. \tag{17}$$

Note that this value is independent of m_ϕ – heavier inflaton would have a shorter lifetime, but at the same time they would cause a faster expansion rates at early times.

We have shown in Fig. 2 the perfect agreement of the analytical estimations made in Eqs. (15)–(17) w.r.t. the numerical solution in Eq. (14).

4.1. Evolution of the momenta of relativistic particles

Coming back to the thermalisation analysis, since we cannot rely on an equilibrium distribution at this point, we take the typical momentum $\bar{p}(x)$ in Eq. (1) to be:

$$\bar{p}(x) = \frac{d\rho_R(x)}{dn_R(x)} = \frac{d\rho_R(x)}{dx} \cdot \left[\frac{dn_R(x)}{dx} \right]^{-1}. \tag{18}$$

This expression directly follows from the definitions of n_R and ρ_R , *without* assuming any particular shape of the distribution function $f(p)$. We would like to emphasise here that Eq. (3), in the absence of LTE, should not even have an interpretation of mean kinetic energy, since its functional shape incorporates the assumption of LTE-like $f(p)$.

Taking then Eq. (18) as a measure of the mean kinetic energy \bar{E} of particles in the plasma, we compare $\bar{E}(x)$ with the temperature $T(x)$, extracted from Eq. (3) under the assumption of thermal equilibrium. This is shown in Fig. 3. As can be observed, \bar{E} is constant over almost the whole reheating period, whereas after reheating its evolution follows the same law as for $T(x)$, i.e. the well-known $T \propto x^{-1}$ behaviour of the radiation-dominated universe, resulting in:

$$\bar{E}^{id} \simeq \frac{m_\phi}{3}, \quad \bar{E}(x)^{rd} \simeq \frac{0.5m_\phi}{\alpha_\phi^{2/3}} \frac{x_I}{x}. \tag{19}$$




Fig. 3. Comparison of the mean kinetic energy as computed according to Eq. (18), and the temperature assuming LTE from the very onset of the inflaton decay as in Eq. (3), for $m_\phi = 10^{13}$ GeV, $\alpha_\phi = 10^{-11}$. One can see the obvious distinction and the importance of understanding when one should associate a temperature to the decay products of the inflaton.

Physically it makes sense: during inflaton–domination the plasma (containing the relativistic species from the inflaton decay) is getting constantly reheated by the inflaton decay, and it turns out that it does so at a rate which is equal to the cooling rate due to the expansion. Afterwards, when the inflaton has decayed completely and only radiation remains, the energy of the relativistic species gets only redshifted by the expansion of the universe.

While this estimation for a typical momentum is reasonable in the scenario of delayed LTE, in the pure SM for example the emitted soft particles (out of the $2 \rightarrow 3$ inelastic processes) may have momenta as low as:

$$\bar{p}_{\text{cut}} \sim n_R^{1/3} \tag{20}$$

where n_R is given approximately by Eq. (15) or (16) depending on the period of energy density domination. Comparing \bar{p}_{cut} with \bar{E} for the two regimes, we have that $\bar{p}_{\text{cut}} < \bar{E}$ for:

$$m_\phi \gtrsim 2M_P \alpha_\phi^{1/2} \left(\frac{x_I}{x}\right)^{3/4}, \quad (\text{id-epoch}) \tag{21}$$

$$m_\phi \gtrsim 1.4M_P \alpha_\phi, \quad (\text{rd-epoch}).$$

This means that in scenarios where the infrared enhancement is accessible, the thermalisation is much faster than in the delayed scenario, mainly for larger inflaton masses and smaller couplings.

4.2. Evaluating the thermalisation time

As for the thermalisation condition is concerned, depending on the value of (α_ϕ, m_ϕ) , in the delayed scenario the LTE can be attained during inflaton–domination or afterwards, during radiation–domination. We should evaluate Γ_{th} by making use of the mean energies, \bar{E} , instead of the *temperature*, since as we pointed out above – we cannot rely at this point on thermal distribution. In the case of efficient thermalisation we use \bar{p}_{cut} instead. We then compare Γ_{th} , according to the case, with:

$$H(x) \approx \begin{cases} 2.9m_\phi \left| \frac{0.6x_I^{3/2}}{\sqrt{\pi}x^{3/2}} - 0.1\alpha_\phi \right|, & (\text{id-epoch}) \\ 1.6 \frac{m_\phi}{\sqrt{\pi}\alpha_\phi^{1/3}} \frac{x_I^2}{x^2}, & (\text{rd-epoch}) \end{cases} \tag{22}$$




Fig. 4. Reaction rate Γ as a function of x for the delayed (efficient) scenario in red (black), as compared to the Hubble parameter (blue). Break in the slopes determine the transition from inflaton–domination to the radiation–domination. Note that thermalisation time scale in the delayed scenario is larger than the matter-to-radiation transition scale. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

where in inflaton–domination, the Hubble rate is approximately given by: $H \propto (\rho_\phi^{id})^{1/2}$, whereas in radiation–domination case, we have: $H \propto (\rho_R^{rd})^{1/2}$.

We have shown in Fig. 4 the comparison of Γ_{th} and H from the numerical solution of Eq. (8), using $\alpha_\phi = 10^{-11}$ and $m_\phi = 10^{13}$ GeV, for the sake of illustration, in the two scenarios: delayed and efficient thermalisation.

For the efficient scenario, the thermalisation happens much before the beginning of radiation–domination epoch, as expected.

On the other hand, for the delayed scenario, the evolution of Γ_{th} is parallel to that of H for nearly the whole reheating period. Indeed, in this region $\sigma(x)$ is nearly constant (because \bar{E} is) and thus Γ_{th} scales as $n_R(x)$, the latter evolving as $\rho_R(x)$ as was already deduced above (see Eqs. (11) and (15)). On the other hand the Hubble rate, even if dominated by the inflaton oscillations, also evolves as $\rho_R(x)$.³ It is only after the inflaton population decreases substantially that the universe starts being radiation–dominated, thus the thermalisation processes become faster than the expansion rate and thermal equilibrium is achieved. The numerical solution for the thermalisation time, x_{th} , is around $x_{th} \sim 10^{10} x_I$ for this choice of parameters.

Analytically it is possible to obtain the value of x at which the thermalisation occurs, $\Gamma_{th}(x_{th}) = H(x_{th})$. We just need to build up Γ_{th} from Eqs. (16) and (1), whereas the Hubble rate is approximated by Eq. (22). As we are considering a delayed LTE scenario, we evaluate our cross section using (19) as explained above.

For the sake of illustration, assuming a total thermalisation cross-section which goes like $\sigma_{th} = \alpha_s^3/E^2$, see Eq. (1), we obtain the following solution for x_{th} :

$$\frac{x_{th}^{rd}}{x_I} \approx \frac{m_\phi^2}{\alpha_s^3 M_P^2 \alpha_\phi^{5/3}} \quad (23)$$

for the case of a radiation–dominated thermalisation. However when including the log contribution, see Eq. (1), it is not possible to obtain an analytical solution of x_{th} . In this more accurate case, the solution is numerical and the thermalisation time x_{th} is between 10 and 100 times smaller than what Eq. (23) predicts.

³ This can be deduced from Eq. (10) under the assumption of $\Phi \approx \text{const}$.

On the other corner of the parameter space, for large α_ϕ and small m_ϕ , it usually happens that thermalisation happens very fast, $x_{th}^{id} \lesssim x_{max}$, when the inflaton still dominates the expansion.

This is one of the main results of our analysis: for some choices of the pair (α_ϕ, m_ϕ) , the plasma does not reach thermalisation at the time when the universe becomes radiation-dominated, but later. This happens for:

$$\alpha_\phi \lesssim (0.01-0.1) \times \left(\frac{1}{\alpha_s^3}\right) \left(\frac{m_\phi}{M_p}\right)^2, \tag{24}$$

where in the RHS we have corrected for the fact that a realistic x_{th} may be 10^{-2} smaller than that of (23).

As an example for illustration, for a heavy mass, $m_\phi = 10^{14}$ GeV, thermalisation reactions driven by $2 \rightarrow 3$ processes of strong gauge coupling (as in Eq. (1)), the relativistic species reaches thermal equilibrium later than the beginning of the radiation-domination era as long as $\alpha_\phi \lesssim 10^{-8}$.

Physically speaking this can be understood as follows: even when the universe starts to become dominated by the radiation energy density, the thermalisation reaction rates may still be inefficient because of the very large typical energies of the interacting particles, $\bar{E} \lesssim \mathcal{O}(m_\phi)$, inherited from the inflaton decays and almost unaffected otherwise (see Fig. 3, for an illustrative point). These large energies penalise the cross-sections, until the redshift is important enough as for the scattering process to become efficient enough, such that $\Gamma_{th} > H$ and LTE is finally attained. Of course this works as long as the population of soft particles is not large enough as for affecting noticeably the rate of scattering processes.

5. Definition of reheating temperature

Now let us define the reheating temperature, T_{rh} , as computed according to energy density (cf. Eq. (3)), provided the radiation has just thermalised, and dominates the Hubble expansion rate of the universe.

$$T_{rh} = T(x), \quad x = \max(x_{th}, x_{rh}). \tag{25}$$

There are three cases of interest:

1. **Instant thermalisation** – ($x_{rh} \ll x_{th}$): Thermalisation of relativistic species is attained almost instantaneously (usually even around x_{max}), already during the coherent oscillations of the inflaton, and they maintained LTE throughout reheating and also at the time when the universe becomes radiation dominated. Following our prescription in Eq. (25), in this case the reheating temperature is determined by:

$$T_{rh}(x_{th} \ll x_{rd}) \approx \frac{0.6}{g_*^{1/4}} \sqrt{\alpha_\phi m_\phi M_p}. \tag{26}$$

A couple of points to note: we see that $T_{rh}(x_{th} \ll x_{rd})$ behaves exactly as the usual T_{rh} of instant-reheating scenario, see Eq. (5), with an $\mathcal{O}(1)$ -difference in a prefactor. Indeed $T_{rh}(x_{th} < x_{rd})$ is a bit smaller than the usual definition of reheating case, as assumed in Eq. (5), since the latter corresponds to a maximal thermalisation-efficiency by definition. In our case, the lower efficiency translates into a bit smaller reheating temperature (see Fig. 5 below). On the other hand, in this scenario, it is indeed possible to define a maximum temperature of the relativistic species, $T_{max} \equiv T(x_{max}) > T_{rh}$.




Fig. 5. Reheating temperature computed numerically by solving Eq. (8). This is represented with dots joined by solid lines for different values of α_ϕ : 10^{-13} (blue), 10^{-11} (red), 10^{-9} (green) and 10^{-7} (cyan), for $\alpha_s = 1/30$. Delayed scenario is shown in top panel, whereas efficient scenario is shown in bottom panel. The blue dashed-line is an incorrect depiction of reheating temperature (as in Eq. (5)), corresponding to $\alpha_\phi = 10^{-13}$. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

2. **Efficient thermalisation.** This may be the case of the SM for example, where due to the presence of infrared divergences in Eq. (1), cured by a cut-off given in (20), we get $x_{\max} \ll x_{th} \leq x_{rh}$. In this case the estimation of T_{rh} is again as in Eq. (26) following the recipe Eq. (25), and thermalisation happens within the inflaton–domination era.
3. **Delayed thermalisation** – ($x_{th} \gg x_{rh}$): Thermalisation happens deep inside the radiation dominated era, such that the reheat temperature is determined by:

$$T_{rh}(x_{th} \gg x_{rd}) \approx (7-70) \times \frac{\alpha_s^3 \alpha_\phi^{3/2} M_P^{5/2}}{g_*^{1/4} m_\phi^{3/2}}. \tag{27}$$

Note that $T_{rh}(x_{th} \gg x_{rd})$ has an opposite behaviour with respect to m_ϕ . This is the most important result of our work – in some region of the parameter space (α_ϕ, m_ϕ), where thermalisation happens after radiation starts dominating, the reheating temperature actually decreases with the inflaton mass. Physically this is due to the following. For larger m_ϕ , larger is the mean energy \bar{E} of the relativistic species. This penalises the cross-sections for the thermalisation reactions which occur at the beginning, when soft processes are still unimportant,

thus rendering the thermalisation rate less efficient at the end of the day, which is attained later. Consequently this lowers down $T_{rh}(x_{th} \gg x_{rd})$.

For scenarios beyond SM where large VEVs of scalar fields prevent the appearance of infrared divergences in (1)⁴ we have shown in Fig. 5 (top) the reheating temperature T_{rh} as a function of m_ϕ for different values of α_ϕ , computed numerically by solving Eq. (8) and represented with coloured dots joined by full lines. As commented above there are two regimes: one for which the thermalisation happens at inflaton–domination, where $T_{rh}(x_{th} < x_{rd})$ grows with m_ϕ and follows closely to $T_{rh} = (90/8\pi^3 g_*)^{1/4} \sqrt{\Gamma_\phi} M_P$ (see dashed blue line in Fig. 5); and a second regime for which the thermalisation happens deep inside radiation–domination era, where $T_{rh}(x_{th} > x_{rd})$ decreases with m_ϕ . Essentially, for the largest α_ϕ and the smallest m_ϕ , we are in the former regime, whereas for the smallest α_ϕ and the largest m_ϕ , we are in the latter regime.

Note that, for example, for $\alpha_\phi = 10^{-13}$ and $m_\phi = 10^{13}$ GeV the usual T_{rh} (as in Eq. (5)) largely overestimates the (more realistic) reheating temperature we have obtained in our analysis. For the second regime, where Eq. (27) applies, we obtain a prediction in the correct ballpark for the numerical solution shown in Fig. 5. On the other hand the parametric dependence of Eq. (27) is verified.

A closer inspection of Fig. 5 reveals some values of m_ϕ and α_ϕ for which the numerical results are not shown. These “holes” in the scan are due to the limited validity of our numerical solution of Eq. (8). As discussed above this expression, the $\Gamma_{th}(\rho_R - \rho_R^{eq})$ term is important when Γ_{th} becomes essentially comparable in size to the Hubble expansion and the inflaton source. Otherwise, either radiation-to-radiation terms are very inefficient (such that they do not play a role in the ρ_R -evolution), or if they are too efficient (such that production and annihilation balance each other in an equilibrium distribution), Eq. (8) is a reasonable simplification of the original Boltzmann set of equations Eq. (2).

We have also obtained T_{rh} numerically in the “efficient thermalisation” case, for all the parameter space (see Fig. 5 (bottom)). There, T_{rh} grows monotonically as approximated by Eq. (26) even for the largest inflaton masses, contrary to the delayed scenario where T_{rh} decrease with m_ϕ .

6. Phenomenological implications of delayed thermalisation

The scenario of delayed thermalisation may have important implications for phenomenology and model building. We next briefly discuss some of the most direct ones:

6.1. Leptogenesis

In this scenario,⁵ the existence of right-handed (RH) neutrinos give rise to the observed baryon asymmetry by sphaleron conversion processes. The lightest RH neutrino N_1 has to be massive enough as for producing sufficient CP asymmetry. Thus, since the thermal plasma needs to produce enough number density of those N_1 for a successful mechanism, a lower bound on the reheating temperature is imposed. Here we quote $T_{rh} \gtrsim 2 \times 10^9$ GeV [17]. By direct inspection of Fig. 5, we can see that the leptogenesis bound forbids couplings $\alpha_\phi \lesssim 10^{-11}$ for whatever value of the inflaton mass.

⁴ I.e. where the “efficient thermalisation” described above does not apply.

⁵ For a review see [15].




Fig. 6. Upper bounds on the coupling α_ϕ coming from BBN constraints. This have been obtained by convoluting one of the bounds presented in [16]. Upper (blue) line is for an inflaton mass of $M_\phi = 10^{13}$ GeV, whereas the lower (red) line uses $M_\phi = 10^{10}$ GeV. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

6.2. Gravitinos

Certainly one of the most important scenarios sensitive to the very early universe, the gravitino over-production poses a serious cosmological problem. The gravitinos are produced mostly at the reheating period out of the thermal bath and, if unstable, their late decays could potentially spoil the mechanisms leading to Big Bang Nucleosynthesis (BBN). On the other hand, if they are stable, they can over-close the Universe as dark matter candidates if the reheating temperature is large enough. Thus, in both cases we are able to place upper bounds on the reheating temperature.

6.2.1. Unstable gravitinos

In this scenario the bounds on T_{rh} come from the abundance of light elements. Given a situation of delayed thermalisation as the one we have discussed in this work, we could translate these upper bounds on T_{rh} from BBN to bounds on the coupling α_ϕ for given inflaton masses, as a function of the gravitino mass. The result is shown in Fig. 6. Based on [16], we have, for each gravitino mass $m_{3/2}$, maximum allowed values of T_{rh} coming from the abundances of D, ^3He , ^4He , ^6Li and ^7Li elements. We take for each $m_{3/2}$ the strongest (i.e. the minimum) of these bounds, such that at the end of the day we have a convolution of these upper bounds as a function of $m_{3/2}$. Then, in Fig. 6 we have translated the obtained universal upper bound on T_{rh} , to upper bounds on α_ϕ , given two fixed inflaton masses: 10^{10} and 10^{13} GeV, by making use of our result (27).

We see that in general the bounds from BBN greatly constrain the couplings α_ϕ : even for inflaton masses as large as 10^{13} GeV, couplings larger than 10^{-12} – 10^{-10} are forbidden for a large range of gravitino masses. This bound goes in the opposite direction as the one from leptogenesis. Only if the gravitino is very heavy (larger than 10^4 GeV) the two constraints are compatible, since then the BBN bound becomes loose.

6.2.2. Stable gravitinos

If the gravitinos were stable instead, thus dark matter candidates, their number density would freeze shortly after the reheating period. There are distinct ways in which gravitinos can be




Fig. 7. Values of the coupling α_ϕ for two inflaton masses which give rise to the gravitino relic abundance as full dark matter candidate. Colour code is the same as in previous figure. See text for more details.

produced, for example, from direct perturbative decays of the inflaton [20]; from scatterings of the inflaton decay products (i.e. relativistic species of supersymmetric Standard Model), or from thermal processes once the radiation bath has attained thermal equilibrium, see [17]. However, as it is known the gravitino production before the thermalisation is attained gets diluted by the entropy release. Thus, it's final population is in very good approximation given by the thermal production yield, at the time where $T = T_{rh}$, the resulting relic abundance being⁶:

$$\Omega_{3/2} h^2 \approx 1.7 \times 10^{-3} \left(\frac{m_{3/2}}{\text{GeV}} \right) \left(\frac{T_{rh}}{10^{10} \text{ GeV}} \right) \left(\frac{\gamma(T)}{T^6/M_P^2} \right)_{T=T_{rh}} \tag{28}$$

where $\gamma(T)$ is the total gravitino production rate, typically proportional to T^6/M_P^2 , and dependent on the supersymmetric spectrum, particularly the gaugino masses. In Fig. 7 we show the prediction for a simplified scenario where the gaugino masses are degenerate and equal to 1 TeV at GUT scale. By fixing the gravitino relic abundance to $\Omega_{3/2} h^2 = 0.12$, in [17] they have obtained a prediction for $T_{rh}(m_{3/2})$, which we then translate to a prediction for the coupling α_ϕ according to (27), in a similar fashion as above. Amusingly, the resulting couplings are in the same ballpark as the upper bounds obtained above from a completely independent analysis.

7. Conclusions and discussions

In this work we have studied inflationary reheating, in particular revisiting the study of the thermalisation of the inflaton decay products from both an analytical and numerical point of view, by analysing the dominant thermalisation process of the relativistic plasma as a whole. We have solved the coupled set of Boltzmann equations in two clearly defined regimes: a) The $2 \rightarrow 3$ processes leading to thermalisation are too inefficient to affect the global evolution of the radiation energy-density itself, as a result the universe could be radiation dominated, but still not in local thermodynamical equilibrium (LTE), and b) when thermalisation process is very quick, at much larger rates compared to the Hubble expansion and the inflaton decay rate, in such a way that LTE of the decay products is attained very fast. In both regimes the Boltzmann equations are simplified in a similar fashion.

⁶ We have computed the evolution of the gravitino yield in this case. Our result is in agreement with [17].

We have obtained the following important results:

- For sufficiently small α_ϕ and sufficiently large inflaton-mass m_ϕ , the relativistic plasma does not thermalise at the time where radiation–domination era begins, but (in some cases, much) later. When α_ϕ is very small there are not enough relativistic species at the matter-to-radiation transition to immediately thermalise, whereas for very large m_ϕ , the species are too energetic as for the relevant scattering processes to be efficient enough. Although, this requires significant suppression in t -channel scattering rate, which may happen for a massive gauge boson mediated interactions due to VEV or finite temperature effects giving mass to the gauge bosons. If there is a t -channel enhancement due to massless gauge boson mediation, then the scattering rate is enhanced due to infrared effect and the Debye cut-off is determined by the number density of relativistic species present in the plasma. In this case thermalisation occurs during the inflaton oscillations dominating the universe.
- We have determined a proper definition of the reheat temperature, in a generic scenario of perturbative decays of the inflaton. Essentially, two necessary conditions have to be eventually fulfilled: the plasma have to attain LTE, and it must dominate the expansion rate of the universe. This is such that for some region of the inflaton parameters (precisely the one commented in the first point), the reheat temperature turns out to be much smaller than the standard estimations.

Finally, we have discussed some connection with phenomenology by presenting implications on the gravitino cosmology. In general, for unstable or stable gravitinos, the predicted inflaton couplings need to be very small, order $\alpha_\phi \sim 10^{-12}$ – 10^{-10} for an inflaton mass of 10^{13} GeV.

Other phenomenological implications may be obtained in the context of Affleck–Dine baryogenesis (see e.g. [9]) and dark matter creation during reheating [18]. More recently in the context of freeze-in mechanism through heavy portals [21–23], we have other examples of DM which are sensitive to the reheat temperature. Some of this topical issues might as well have important implications for the inflaton mass and couplings, as for the gravitino case discussed above.

Acknowledgements

We thank E. Fernandez-Martinez and J. Rubio for very useful discussions. A.M. is supported by the Lancaster–Manchester–Sheffield Consortium for Fundamental Physics under STFC grant ST/J000418/1. B.Z. acknowledges the Consolider–Ingenio PAU CSD2007-00060, CPAN CSD2007-00042, under the contract FPA2010-17747; the Comunidad Autonoma de Madrid through the project HEPHACOS P-ESP-00346, and the European Commission under contract PITN-GA-2009-237920, as well as the support of the Spanish MINECOs “Centro de Excelencia Severo Ochoa” Programme under grant SEV-2012-0249.

References

- [1] A. Albrecht, P.J. Steinhardt, M.S. Turner, F. Wilczek, *Phys. Rev. Lett.* 48 (1982) 1437.
- [2] J.H. Traschen, R.H. Brandenberger, *Phys. Rev. D* 42 (1990) 2491;
Y. Shtanov, J.H. Traschen, R.H. Brandenberger, *Phys. Rev. D* 51 (1995) 5438, arXiv:hep-ph/9407247;
L. Kofman, A.D. Linde, A.A. Starobinsky, *Phys. Rev. Lett.* 73 (1994) 3195, arXiv:hep-th/9405187;
L. Kofman, A.D. Linde, A.A. Starobinsky, *Phys. Rev. D* 56 (1997) 3258, arXiv:hep-ph/9704452.
- [3] R. Allahverdi, et al., *Annu. Rev. Nucl. Part. Sci.* 60 (2010) 27, arXiv:1001.2600 [hep-th].
- [4] A. Mazumdar, J. Rocher, *Phys. Rep.* 497 (2011) 85, arXiv:1001.0993 [hep-ph].
- [5] A.R. Liddle, A. Mazumdar, F.E. Schunck, *Phys. Rev. D* 58 (1998) 061301, arXiv:astro-ph/9804177.
- [6] J. Beringer, et al., Particle Data Group Collaboration, *Phys. Rev. D* 86 (2012) 010001

- [7] J. Garcia-Bellido, D.G. Figueroa, J. Rubio, Phys. Rev. D 79 (2009) 063531, arXiv:0812.4624 [hep-ph].
- [8] R. Allahverdi, A. Mazumdar, Phys. Rev. D 76 (2007) 103526, arXiv:hep-ph/0603244;
R. Allahverdi, A. Mazumdar, J. Cosmol. Astropart. Phys. 0610 (2006) 008, arXiv:hep-ph/0512227.
- [9] K. Enqvist, A. Mazumdar, Phys. Rep. 380 (2003) 99, arXiv:hep-ph/0209244.
- [10] R. Allahverdi, et al., Phys. Rev. D 83 (2011) 123507, arXiv:1103.2123 [hep-ph].
- [11] E.W. Kolb, M.S. Turner, Front. Phys. 69 (1990) 1.
- [12] D.J.H. Chung, E.W. Kolb, A. Riotto, Phys. Rev. D 60 (1999) 063504, arXiv:hep-ph/9809453;
D.J.H. Chung, E.W. Kolb, A. Riotto, Phys. Rev. D 59 (1999) 023501, arXiv:hep-ph/9802238.
- [13] K. Enqvist, K.J. Eskola, Mod. Phys. Lett. A 5 (1990) 1919;
K. Enqvist, J. Sirkka, Phys. Lett. B 314 (1993) 298, arXiv:hep-ph/9304273.
- [14] S. Davidson, S. Sarkar, J. High Energy Phys. 0011 (2000) 012, arXiv:hep-ph/0009078.
- [15] S. Davidson, E. Nardi, Y. Nir, Phys. Rep. 466 (2008) 105, arXiv:0802.2962 [hep-ph].
- [16] M. Kawasaki, K. Kohri, T. Moroi, A. Yotsuyanagi, Phys. Rev. D 78 (2008) 065011, arXiv:0804.3745 [hep-ph].
- [17] V.S. Rychkov, A. Strumia, Phys. Rev. D 75 (2007) 075011, arXiv:hep-ph/0701104.
- [18] R. Allahverdi, M. Drees, Phys. Rev. D 66 (2002) 063513, arXiv:hep-ph/0205246;
R. Allahverdi, M. Drees, Phys. Rev. Lett. 89 (2002) 091302, arXiv:hep-ph/0203118.
- [19] M. Drewes, J.U. Kang, Nucl. Phys. B 875 (2013) 315, arXiv:1305.0267 [hep-ph].
- [20] A.L. Maroto, A. Mazumdar, Phys. Rev. Lett. 84 (2000) 1655, arXiv:hep-ph/9904206.
- [21] Y. Mambrini, K.A. Olive, J. Quevillon, B. Zaldivar, Phys. Rev. Lett. 110 (2013) 241306, arXiv:1302.4438 [hep-ph].
- [22] M. Blennow, E. Fernandez-Martinez, B. Zaldivar, arXiv:1309.7348 [hep-ph].
- [23] X. Chu, Y. Mambrini, Jérémie Quevillon, B. Zaldivar, arXiv:1306.4677 [hep-ph].