

Phenomenological analysis of the decay $B^\pm \rightarrow K^\pm p \bar{p}$

E. Di Salvo^{1,2,3,a}, F. Fontanelli^{1,2,b}¹ Dipartimento di Fisica, Via Dodecaneso 33, 16146 Genoa, Italy² INFN-Sez. di Genova, Genoa, Italy³ Laboratoire de Physique Corpusculaire de Clermont-Ferrand, IN2P3/CNRS Université Blaise Pascal, 63177 Aubière Cedex, France

Received: 31 May 2015 / Accepted: 8 October 2015 / Published online: 20 November 2015

© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract We propose a parametrization for interpreting some of the presently available data of the $B^\pm \rightarrow K^\pm p \bar{p}$ decay, in particular those by the LHCb and Belle Collaborations. The model is inspired by the well-known current and transition contributions, usually assumed in this kind of decay. However, in the light of considerations as regards the dominant diagrams and final state interactions, we modify some parameters of the model, determining them by means of a best fit to the data. We show the results, which we discuss in some detail. Moreover, we give some predictions on other observables relative to the decays.

1 Introduction

The physics of the B -meson has opened a new door in the sector of hadronic weak interactions. In particular, numerous B decays, analyzed during more than a decade, have confirmed the CKM mechanism for CP violation [1]; yet, it is generally believed that some particular decays of this kind could reveal new physics beyond the standard model (SM). Therefore several experiments in this sense have been suggested or even realized. In particular, data of $B^\pm \rightarrow K^\pm p \bar{p}$ decay have been published recently by the LHCb Collaboration [2–4].

Three-body baryon-antibaryon B decays were detected for the first time by ARGUS in 1987 [5]. Although immediately ruled out by CLEO [6], that claim gave rise to a lively theoretical interest in the subject for several years: see Ref. [7] for a complete and thorough review. More recently, this interest was revived [8–12] and new experimental results were

found by the CLEO [13], Belle [14,15] and BaBar [16,17] Collaborations.

A three-body decay presents some advantages over a two-body one. For example, if one chooses a three-body charged B decay such that all final particles are charged, its detection is favored. Moreover, an amplitude analysis in different intervals of the Dalitz plot allows for direct measurement of the relative strong phases [18].

Baryonic three-body decays of charged B are especially useful for studying the strong dynamics and, in principle, for analyzing the observables sensitive to new physics [8]. Indeed, it is customary to assume that final state interactions [19] (FSI) are much smaller [2] than those of three-meson decays, characterized by large re-scattering effects [18–20]. This fact would allow for the study of FSI by comparison between the two different kinds of decays [2] and therefore could help in determining the weak phase, which generally presents serious difficulties [21]. However, as we shall see in the present paper, FSI might play a surprisingly important role also in baryonic decays.

Besides, such decays are somewhat advantageous with respect to the baryonic two-body ones (e.g., $B^0 \rightarrow p \bar{p}$), in that they give rise to greater decay rates than two-body baryonic decays. This is connected with the threshold baryon-antibaryon peaks, predicted by Hou and Soni [8] as a consequence of the increasing difficulty of a quark to hadronize to a baryon at higher energies. In particular, if the final state includes the proton-antiproton system, the effect is attributed to re-scattering [22]. Moreover, the decays considered exhibit intriguing forward-backward (FB) and Dalitz plot asymmetries [2,15,17].

This kind of decays is generally described by assuming two factorizable amplitudes of the type [7,9]

$$\mathcal{T} = \langle M | (\bar{q}_3 q_2) | 0 \rangle \langle \mathcal{B}_1 \bar{\mathcal{B}}_2 | (\bar{q}_1 b) | B^- \rangle \quad \text{and} \\ \mathcal{I} = \langle \mathcal{B}_1 \bar{\mathcal{B}}_2 | (\bar{q}_1 q_2) | 0 \rangle \langle M | (\bar{q}_3 b) | B^- \rangle,$$

^a e-mail: disalvo@ge.infn.it; Elvio.Disalvo@ge.infn.it^b e-mail: Flavio.Fontanelli@ge.infn.it

called, respectively, transition and current terms. Here M , \mathcal{B}_1 , and $\bar{\mathcal{B}}_2$ are the final meson, baryon and antibaryon respectively, while b and q (\bar{q}) are the destruction (creation) operators of the active (light) quarks involved in the decay.

Turning to the decay $B^\pm \rightarrow K^\pm p \bar{p}$ (to be named K decay in the following), it presents two main differences with respect to the $B^\pm \rightarrow \pi^\pm p \bar{p}$ (π decay from now on). First, it has a wider $p \bar{p}$ peak, which may be explained as an effect of the interference between the transition and the current contributions [9]: indeed, while suppressed in the π decay, the current term contributes considerably to the K decay and decreases more slowly than the transition term at increasing $p \bar{p}$ effective mass. Second, the sign of the FB asymmetry (A_{FB}) in the π decay is opposite to that in the K decay: while the behavior of the π asymmetry can be explained in the framework of quark dynamics [7], the other one is more difficult to interpret.

An important feature of the K decay is that it derives contributions from two different amplitudes, typically tree and penguin, endowed with different weak and strong phases. Therefore, we expect that data exhibit a sizable direct CP asymmetry (A_{CP}), to be compared with the SM predictions. Indeed, this observable has been measured recently to high precision by the LHCb Collaboration [2]. Unfortunately, as recalled above, comparison with theory is particularly difficult in weak decays, owing to the problem of disentangling real CP-violation effects from strong FSI. Therefore, in the interest of understanding and parametrizing as precisely as possible the effects of non-perturbative QCD in hadronic weak decays, we perform a best fit to some available data, like the differential branching fraction and the asymmetries A_{FB} and A_{CP} . More precisely, on the one hand, we adopt for the decay amplitude the parametrization by Ref. [9]; on the other hand, however, we modify some parameters by means of a best fit, in view of various considerations. As we shall see, our analysis, based on the data by Belle [15] and on the most recent data of LHCb [2], leads to conclusions in contrast with previous ones; in particular, we give a different answer to one of the main questions illustrated above, that is, the origin of the FB asymmetry. Lastly, we make some predictions about other observables measurable in principle.

Section 2 is devoted to the definition of some observables, either presently available or measurable in the future, and to the theoretical function used for interpreting them. In Sect. 3, we present our phenomenological analysis, both assuming factorization and relaxing this assumption; moreover, we show some predictions of our model. Lastly, in Sect. 4, we discuss the results obtained and draw some conclusions.

2 Observables and theoretical function

2.1 Observables

We fit the data relative to the the differential branching fraction of the decay $B^\pm \rightarrow K^\pm p \bar{p}$ [2, 15], i.e.,

$$\gamma = \frac{dBf_{Kp\bar{p}}}{dm_{p\bar{p}}}. \quad (1)$$

The LHCb data [2] are re-scaled in such a way that the integral over all of the $p \bar{p} K$ spectrum equals the total branching fraction [23], i.e.,

$$Bf(B^\pm \rightarrow K^\pm p \bar{p}) = (5.9 \pm 0.5) \cdot 10^{-6}. \quad (2)$$

On the other hand, we calculate the CP asymmetry and the FB asymmetry, whose experimental values are, respectively [2],

$$A_{\text{CP}} = -0.022 \pm 0.031 \pm 0.007 \quad (3)$$

and

$$A_{\text{FB}} = 0.370 \pm 0.018 \pm 0.016. \quad (4)$$

The latter is defined as

$$A_{\text{FB}} = \frac{N^+ - N^-}{N^+ + N^-} \quad (5)$$

and N^\pm is the number of events for which $\cos\theta_p$ is positive (negative), θ_p being the angle between the meson and the opposite-sign baryon in the $p \bar{p}$ rest frame. This definition holds as well for the π decay.

2.2 Theoretical function

2.2.1 Matrix element of the decay

We parametrize the decay amplitude for $B^\pm \rightarrow K^\pm p \bar{p}$, according to Chua et al. [9] (see also Refs. [7, 11]), i. e.,

$$\mathcal{M} = \frac{G_F}{\sqrt{2}}(\mathcal{I} + \mathcal{T}); \quad (6)$$

here G_F is, as usual, the Fermi constant of weak interactions and \mathcal{I} and \mathcal{T} are respectively the current and transition terms, according to the definition given in the introduction. As regards the current term, we have

$$\mathcal{I} = \mathcal{I}_1 + \mathcal{I}_2, \quad (7)$$

$$\mathcal{I}_1 = \langle K^- | L_{sb}^\mu | B^- \rangle \left(\sum_{i=1}^4 \alpha_i \langle p \bar{p} | L_\mu^i | 0 \rangle + \alpha_5 \langle p \bar{p} | R_\mu | 0 \rangle \right), \tag{8}$$

$$\mathcal{I}_2 = \langle | K^- | \tilde{L}_{sb} | B^- \rangle \alpha_6 \langle p \bar{p} | \tilde{R}_{ss} | 0 \rangle. \tag{9}$$

On the other hand, the transition term reads

$$\mathcal{T} = \langle K^- | \tilde{L}_{us} | 0 \rangle (\alpha_7 \langle p \bar{p} | S_{ub} | B^- \rangle + \alpha_8 \langle p \bar{p} | P_{ub} | B^- \rangle). \tag{10}$$

Here we have set

$$L_{sb}^\mu = \bar{s}_L \gamma^\mu b_L, \quad L_\mu^1 = \bar{u}_L \gamma_\mu u_L, \tag{11}$$

$$L_\mu^2 = \bar{u}_L \gamma_\mu u_L + \bar{d}_L \gamma_\mu d_L + \bar{s}_L \gamma_\mu s_L,$$

$$L_\mu^3 = \bar{s}_L \gamma_\mu s_L, \quad L_\mu^4 = e_u \bar{u}_L \gamma_\mu u_L + e_d \bar{d}_L \gamma_\mu d_L + e_s \bar{s}_L \gamma_\mu s_L, \tag{12}$$

$$R_\mu = \bar{u}_R \gamma_\mu u_R + \bar{d}_R \gamma_\mu d_R + \bar{s}_R \gamma_\mu s_R, \quad \tilde{L}_{sb} = \bar{s}_L b_L, \tag{13}$$

$$\tilde{R}_{ss} = \bar{s}_R s_R, \quad \tilde{L}_{us} = \bar{s}_L u_L, \quad S_{ub} = \bar{u} b, \quad P_{ub} = \bar{u} \gamma_5 b. \tag{14}$$

Moreover, the α are connected to the CKM matrix elements and to the Wilson coefficients. Focusing on the B^- decay, we have

$$\alpha_1 = V_{ub} V_{us}^* a_2, \quad \alpha_2 = -V_{tb} V_{ts}^* a_3, \quad \alpha_3 = -V_{tb} V_{ts}^* a_4, \tag{15}$$

$$\alpha_4 = -\frac{3}{2} V_{tb} V_{ts}^* a_9, \quad \alpha_5 = -V_{tb} V_{ts}^* a_5, \tag{16}$$

$$\alpha_6 = 2V_{tb} V_{ts}^* a_6,$$

$$\alpha_{7,8} = V_{ub} V_{us}^* a_1 - V_{tb} V_{ts}^* a_{\pm}, \tag{17}$$

$$a_{\pm} = a_4 \pm a_6 \frac{2m_K^2}{m_b(m_s + m_u)}.$$

The numerical values of the a and of the V are listed in the appendix. The formulas of the B^+ decay are obtained by taking the complex conjugates of the CKM matrix elements. Since V_{ub} is the only complex CKM matrix element in the equations above, only α_1 , α_7 , and α_8 are involved in the CP asymmetry.

The various (non-perturbative) matrix elements of the quark operators are connected to some form factors, as shown in the appendix. As a result we get

$$\mathcal{M} = \frac{G_F}{\sqrt{2}} \frac{1}{2m_p} \sum_{l=1}^4 \beta_l \bar{u}(p_p) O_l v(p_{\bar{p}}). \tag{18}$$

Here p_p and $p_{\bar{p}}$ are the four-momenta of the proton and of the antiproton respectively; u and v are their standard Dirac spinors, normalized as $\bar{u}u = -\bar{v}v = 2m_p$, where m_p is the proton mass. Moreover,

$$O_1 = \not{p}_B, \quad O_2 = \not{p}_B \gamma_5, \quad O_3 = I, \quad O_4 = \gamma_5, \tag{19}$$

with p_B being the four-momentum of the decaying resonance B . Lastly, the coefficients β are

$$\beta_1 = 2F_1 \sum_{i=1}^4 \Phi^i \xi_i - f_K m_b \alpha_8 F_{V5}, \tag{20}$$

$$\beta_2 = 2F_1 \sum_{i=1}^4 g_A^i \eta_i - f_K m_b \alpha_7 F_A, \tag{21}$$

$$\beta_3 = \frac{\ell \cdot \delta}{2m_p} F_1 \sum_{i=1}^4 \xi_i \hat{\Phi}^i - f_K m_b \alpha_8 F_P, \tag{22}$$

$$\beta_4 = F_0 \left[\frac{r p_K^2}{2m_p} \sum_{i=1}^4 \eta_i h_A^i + \alpha_6 \frac{m_B^2 - m_K^2}{m_b - m_s} \right. \\ \left. \times \left(\frac{m_p}{m_s} g_A^3 + \frac{p_K^2}{4m_p m_s} h_A^3 \right) \right] + \\ - f_K m_b \alpha_7 F_P - 2(r+1)m_p F_1 \sum_{i=1}^4 g_A^i \eta_i \\ + 2m_p \left(r F_0 \sum_{i=1}^4 g_A^i \eta_i + f_K m_b \alpha_7 F_A \right). \tag{23}$$

Here

$$\xi_i(\eta_i) = \pm \alpha_i + \alpha_5 \delta_{i,2}, \quad (i = 1, 2, 3, 4), \\ r = \frac{m_B^2 - m_K^2}{(p_{\bar{p}} + p_p)^2}, \tag{24}$$

$$l = 2p_B - (1+r)(p_{\bar{p}} + p_p), \quad \delta = p_{\bar{p}} - p_p, \tag{25}$$

p_K is the four-momentum of the K -meson, $m_{b(s)}$ the masses of the quarks; m_K and m_B are the meson masses. Moreover, f_K is the decay constant of the K -meson and $F_0, F_1, F_A, F_P, F_{V5}, \Phi_i, \hat{\Phi}_i, g_A^i$, and h_A^i ($i = 1$ to 4) are form factors. In particular, $\hat{\Phi}_i$ – related to the time-like nucleon form factors, as well as the Φ_i – have been set equal to zero, according to Ref. [9]; we shall discuss this assumption in Sect. 4. In this connection, it is worth noting that $F_{0,1}$, as well as the corresponding proportionality factors $S_{0,1}$, defined in the appendix, have the dimensions of an energy, unlike assumed in Refs. [9,24].

Lastly, as we shall see in the next section, our analysis involves also the π decay, for which some variations have to be performed: first of all, one has to replace $s_{L(R)}$ with $d_{L(R)}$, f_K with f_π , m_s with m_d and V_{us}, V_{ts} with V_{ud}, V_{td} ; second, the values of some a_i and parameters in the form factors have to be changed, as shown in the appendix.

2.2.2 Calculations of observables

The modulus squared of the matrix element (18) reads

$$|\mathcal{M}|^2 = \frac{G_F^2}{2} \frac{1}{4m_p^2} \left\{ \sum_{l=1}^4 |\beta_l|^2 \Omega_{ll} + 2 [\Omega_{13} \Re(\beta_1 \beta_3^*) + \Omega_{24} \Re(\beta_2 \beta_4^*)] \right\}. \tag{26}$$

Here

$$\Omega_{11} = 4[2p_B \cdot p_p p_B \cdot p_{\bar{p}} - m_B^2(m_p^2 + p_p \cdot p_{\bar{p}})], \tag{27}$$

$$\Omega_{22} = 4[2p_B \cdot p_p p_B \cdot p_{\bar{p}} + m_B^2(m_p^2 - p_p \cdot p_{\bar{p}})], \tag{28}$$

$$\Omega_{33} = 4(-m_p^2 + p_p \cdot p_{\bar{p}}), \tag{29}$$

$$\Omega_{44} = 4(m_p^2 + p_p \cdot p_{\bar{p}}), \tag{30}$$

$$\Omega_{13} = 4m_p p_B \cdot (p_{\bar{p}} - p_p), \tag{31}$$

$$\Omega_{24} = 4m_p p_B \cdot (p_{\bar{p}} + p_p). \tag{32}$$

Then the differential decay width is

$$d\Gamma = \frac{(2\pi)^4}{2m_B} |\mathcal{M}|^2 \prod_{i=1}^3 \frac{4m_p^2 d^3 p_i}{(2\pi)^3 2E_i} \delta\left(m_B - \sum_{i=1}^3 E_i\right) \delta^3\left(\sum_{i=1}^3 \mathbf{p}_i\right), \tag{33}$$

where E_i and \mathbf{p}_i ($i = 1$ to 3) are the energies and the momenta, respectively, of the proton, of the antiproton and of the K -meson in the B rest frame. By integrating over all variables but the momentum of the proton, we get

$$d\Gamma = \frac{m_p^2}{4(2\pi)^4 m_B} |\mathcal{M}|^2 p_3 \frac{d^3 p_1}{E_1 E_2}, \tag{34}$$

where $p_3 = |\mathbf{p}_3|$. In order to calculate the FB (or helicity [2]) asymmetry, it is convenient to express energies and momenta as functions of kinematic quantities in the $p\bar{p}$ rest frame. To this end, we perform a Lorentz boost from that frame to the B rest frame. Moreover, we integrate over the azimuthal angle of the proton: since $|\mathcal{M}|^2$ does not depend on it, we get

$$d\Gamma = \frac{J m_p^2 p_3 \pi_p^2}{2^4 \pi^3 m_B} |\mathcal{M}|^2 \left(\frac{1}{4} E_0^2 - \frac{p_3^2}{t} \pi_z^2\right)^{-1} d\pi_p d\cos\theta_p. \tag{35}$$

Here $\pi_z = \pi_p \cos\theta_p$, π_p is the modulus of the momentum of the proton and of the antiproton in the $p\bar{p}$ frame and θ_p the helicity angle [2], between the meson and the opposite-sign baryon in the same frame; moreover, $t = 4(m_p^2 + \pi_p^2)$ is the effective mass squared of the $p\bar{p}$ system, $E_0 = \sqrt{t + p_3^2}$ and

$$J = \frac{E_0}{\sqrt{t}} - 4\pi_z \frac{dp_3}{dt} + \frac{8\pi_z^2}{\sqrt{t}} \left[\frac{1}{2E_0} \left(1 + \frac{dp_3^2}{dt}\right) - \frac{E_0}{2t} \right]. \tag{36}$$

Taking into account energy-momentum conservation, we have

$$p_3 = \frac{1}{2m_B} \left[(m_B^2 - m_K^2)^2 - 2t(m_B^2 - m_K^2) + t^2 \right]^{1/2}. \tag{37}$$

Our analysis requires the formulas for the differential decay width and for the differential FB difference, as functions of the $p\bar{p}$ invariant mass $m_{p\bar{p}} = \sqrt{t}$. To this end, we perform the appropriate integrations over $\cos\theta_p$ – respectively, \int_{-1}^1 and $\int_0^1 - \int_{-1}^0$ – expressing the Lorentz invariant coefficients Ω_{ij} in terms of the kinematic variables just defined. The results are

$$\frac{d\Gamma}{dm_{p\bar{p}}} \left(\frac{d\Delta\Gamma}{dm_{p\bar{p}}} \right) = \frac{G_F^2}{2} \frac{m_{p\bar{p}} \pi_p p_3}{4^4 \pi^3 m_B} I(\pi_p) [\Delta I(\pi_p)]. \tag{38}$$

Here

$$I(\pi_p) = \sum_{i=1}^3 \rho_i G_i, \quad \Delta I(\pi_p) = \sum_{i=1}^3 \rho_i \Delta G_i, \tag{39}$$

with

$$\rho_1 = 8m_B^2 (|\beta_1|^2 + |\beta_2|^2), \tag{40}$$

$$\rho_2 = -8 \left\{ m_B^2 [(m_p^2 + \pi_p^2) |\beta_1|^2 + \pi_p^2 (|\beta_2|^2) - \pi_p^2 (|\beta_3|^2 + [(m_p^2 + \pi_p^2) |\beta_4|^2 - m_p m_B E_0 \Re(\beta_2 \beta_4^*)]) \right\}, \tag{41}$$

$$\rho_3 = 16m_p m_B p_3 t^{-1/2} \Re(\beta_1 \beta_3^*), \tag{42}$$

$$G_1 = 2 \left(A + \frac{1}{3} C \right),$$

$$G_2 = \frac{1}{h} \left[\left(\frac{A}{a} + aC \right) \ln \frac{a+1}{|a-1|} - 2C \right] \tag{43}$$

$$G_3 = \frac{\pi_p B}{h} \left(2 - a \ln \frac{a+1}{|a-1|} \right), \tag{44}$$

$$\Delta G_1 = -B, \quad \Delta G_2 = \frac{B}{h} \ln \frac{|a^2 - 1|}{a^2}, \tag{45}$$

$$\Delta G_3 = -\frac{\pi_p}{h} \left[(A + Ca^2) \ln \frac{|a^2 - 1|}{a^2} + C \right] \tag{46}$$

and

$$A = \frac{E_0}{\sqrt{t}}, \quad B = 4\pi_p \frac{dp_3}{dt},$$

$$C = \frac{8\pi_p^2}{\sqrt{t}} \left[\frac{1}{2E_0} \left(1 + \frac{dp_3^2}{dt} \right) - \frac{E_0}{2t} \right], \tag{47}$$

$$a = \frac{E_0 \sqrt{t}}{2\pi_p p_3}, \quad h = \frac{p_3^2 \pi_p^2}{t}. \tag{48}$$

Equation (38) allow one to calculate several observables, to be compared with data. In particular, we are interested in the quantities defined in Sect. 2.1 – that is, the

differential and total branching fractions and the overall FB and CP asymmetries – and in the differential CP asymmetry,

$$A_{CP} = \frac{\gamma^- - \gamma^+}{\gamma^- + \gamma^+}, \tag{49}$$

where γ is defined by Eq. (1).

3 Phenomenological analysis

Here we compare the theoretical function just written with the available observables listed in Sect. 2.1, that is, the differential branching fraction and the overall CP and FB asymmetries. In particular, as regards the first observable, we exclude the contribution of the two charmonium bands, around 3.1 GeV and around 3.6 GeV. Concerning the form factors involved in Eq. (38), first we adopt the factorization assumption, like Chua et al. [9]. However, as we shall see, this disagrees with data [2, 15]. Therefore we shall propose a modification of the model, in view of some considerations about the main contributing graphs.

3.1 Factorization assumption

As a first attempt, we assume factorization for the current and transition terms, according to Refs. [7, 9, 11, 12] and refs. therein. Indeed, this scheme generally describes the three-body baryonic B decays in a satisfactory way; in particular, it seemed to be supported [9] by the early data of the K and π decays [25]. Therefore a comparison with the most recent data of those decays [2, 15] is in order.

In the context of factorization, the transition term has the same proportionality coefficient for the K decay as for the π decay; as regards the current form factors, the proportionality coefficients have been calculated by Melikhov and Stech [24] (see also Ref. [9]).

In the case of the π decay, the penguin amplitude is subdominant with respect to the color-allowed tree diagram [17], even taking into account re-scattering effects to be discussed in Sect. 3.2. Therefore, the current term, which derives contributions almost exclusively from the penguin amplitude, is suppressed with respect to the transition term, which consists, instead, of tree and penguin contributions. Incidentally, in that decay, also a subdominant, color-suppressed tree diagram has to be accounted for. It corresponds, in the case of the B^- decay, to the quark subprocess $b \rightarrow u\bar{u}d$, followed by recombination of the d -quark with the spectator \bar{u} to form the π^- , and by the fragmentation of the active $u\bar{u}$ pair into $p\bar{p}$; the \bar{p} is next to the π^- in rapidity space and can resonate with it as a Δ^{--} , giving a negative contribution to the FB asymmetry.




Fig. 1 Diagram of annihilation of a virtual γ into $p\bar{p}$

The above evaluation about the dominating diagrams agrees qualitatively with the numerical results of the current form factors given in Ref. [24]; indeed, the current term turns out to be negligibly small for the π decay [9]. Therefore, in a simplifying assumption, we neglect the current term; we impose as well the functions F_{V5} and F_P to vanish, according to Ref. [9]. Then our expression of the branching fraction depends only on F_A , whose proportionality coefficient is C_A ; imposing the branching fraction to be equal the experimental value [23],

$$Bf(B^\pm \rightarrow \pi^\pm p\bar{p}) = (1.62 \pm 0.20) \cdot 10^{-6}, \tag{50}$$

yields

$$|C_A| = \left(42.1_{-2.6}^{+2.4}\right) \text{ GeV}^5. \tag{51}$$

Factorization implies that this contribution – up to the sign of C_A , and up to some constants, as explained at Sect. 2.2.1 – should be present also in the K decay. Here, however, also the current term is important [9], since, in this case, it receives contributions exclusively from the penguin amplitude, which is dominant over the tree amplitude. This fact is confirmed by the sizes of the proportionality coefficients of the current form factors for that decay [24]. Therefore, in order to calculate the current term, we insert into Eqs. (8) and (9) the formulas of the electromagnetic nucleon form factors in the time-like region [9, 24], shown diagrammatically in Fig. 1; such formulas are given in the appendix, while the proportionality coefficients of the form factors, S_0 and S_1 , are listed in Table 1 [9].

As regards the transition term, we assume the value (51) for C_A , with the two possible signs. The choice of the – sign yields

$$Bf(B^\pm \rightarrow K^\pm p\bar{p}) = (2.96 \pm 0.27) \cdot 10^{-6}, \tag{52}$$

Table 1 Parameters of the theoretical function. F1 refers to factorization, F2 shows the results of the best fit. C_A and C_{V5} are expressed in GeV^5 , C_P in GeV^8 , S_0 and S_1 in GeV

Fit	C_A	C_{V5}	C_P	S_0	S_1
F1	$-42.1^{+2.4}_{-2.6}$	–	–	0.36	0.36
F2	$-66.0^{+24.8}_{-9.1}$	–	–	$187.3^{+48.2}_{-126.9}$	$-1.25^{+0.05}_{-0.03}$




Fig. 2 Main diagrams in the decays $B^\pm \rightarrow K^\pm p \bar{p}$

which is considerably smaller than the experimental value (2). The choice of the + sign for C_A yields even a lower value. Also data [2, 15] of the differential branching fraction are in disagreement with the factorization assumption, as shown by the dashed line in Fig. 3; therefore we conclude that this assumption is inadequate. The situation is somewhat analogous to the one illustrated in Ref. [26] about the decays of B to $K\chi_0$, to $D^{(*)0}\pi^0$ and to $\pi^0\pi^0$, which are surprisingly enhanced with respect to the factorization assumption and demand both non-factorizing terms and re-scattering effects.

3.2 Remarks on the main contributing graphs

After the result just shown above, a criticism of the assumptions made is in order. Therefore we analyze in detail the main graphs contributing to the decay considered, which we illustrate in Fig. 2.

Figure 2a, b represent, respectively, the color-allowed and the penguin diagram. According to the former one, the meson is by no means correlated to the baryons, since the quark hadronization to the meson occurs just after the weak (very short-ranged) interaction; therefore factorization could be of course assumed for this diagram, which contributes to the transition term. But as told, in the K decay, this amplitude is subdominant with respect to the penguin one. This latter consists of several components contributing to the K and/or π decay, described and represented in detail in Refs. [27] and [7]: some of them may be regarded as factorizable, but not all. The evident failure of factorization leads us to conclude

that non-factorizing diagrams play quite an important role in the K decay. In particular, we point out a contribution fed by the quark subprocess $b \rightarrow c\bar{c}s$. It may give rise to a $p\bar{p}$ pair through re-scattering from $\Lambda_c\bar{\Lambda}_c$ pair formation, which can occur either by $c\bar{c}$ fragmentation, or via intermediate states like $D^{(*)}\bar{D}_s^{(*)}$, $\Xi_c\bar{\Lambda}_c$ [7]; this causes an anomalously large enhancement of the decay rate with respect to the naive factorization [7]. In the π decay, the corresponding amplitude is suppressed by a factor $|V_{cd}/V_{cs}|$ with respect to the K decay. Since the current term interferes only weakly with the transition term [9], this process gives a negligibly small contribution to the differential decay width, as anticipated in Sect. 3.1. On the contrary, it might explain the prevalent role of the current term in the K decay, as we shall establish in Sect. 3.3.

The failure of factorization is also confirmed by the numerical results presented by Chua et al. [9], since the current contribution depends crucially on the effective number of colors, N_c (see also Ref. [12]). On the contrary, the fit to the π decay, dominated by the tree diagram, is stable versus N_c ; this is reflected also on the transition term of the K decay [9], for which those authors assume the same parametrization as for π . The fit to $B^\pm \rightarrow \pi^\pm \Lambda \bar{p}$, which depends essentially on a factorizable penguin diagram [27], is also reasonably stable versus N_c .

Aside from that, as stated before, the long-range FSI might be not so negligible, somewhat analogously to the case of B decays to mesons [26], although on a smaller scale. This may contribute to the failure of factorization and may give rise to further effects, to be discussed in Sect. 4.

3.3 An alternative parametrization

All of the above considerations suggest to relax the factorization assumption. Indeed, we have to take into account both non-factorizable terms and FSI; the latter are typically non-perturbative and therefore cannot be deduced from QCD principles. To this end, we regard C_A , C_{V5} , C_P , S_0 , and S_1 as free parameters, which we determine by minimizing the χ^2 in a fit to the data, that is, as stated before, the differential branching fraction [2, 15] and the asymmetries A_{FB} and A_{CP} [2]. In order to avoid strong correlations between the various parameters, we set $C_{V5} = C_P = 0$, like Chua et al. [9]. The parameters of the fit are shown in Table 1, where they are compared to those of factorization. Furthermore the fit to the $m_{p\bar{p}}$ distribution is exhibited in Fig. 3, continuous line.

3.4 Results and model predictions

Table 2 shows the experimental values of the branching fraction and of the CP and FB asymmetries and the theoretical results about such quantities, both according to the factorization assumption and to our best fit. Furthermore, Figs. 4 and




Fig. 3 $B^\pm \rightarrow K^\pm p\bar{p}$ decay: differential branching fraction vs $m_{p\bar{p}}$, charge averaged case. LHCb and Belle data are represented, respectively, by *rhombuses* and *triangles*. The continuous (*dashed*) line refers to the best fit (factorization assumption)




Fig. 4 Differential branching fractions vs $m_{p\bar{p}}$ for $B^\pm \rightarrow K^\pm p\bar{p}$ decays separately: the continuous (*dashed*) line refers to the B^\pm decay

Table 2 Branching fraction and CP and FB asymmetries: experimental data (Ex) are compared to results of the factorization assumption (F1) and of our fit (F2). Bf is multiplied by 10^6 , A_{CP} and A_{FB} by 10^2 . Here the experimental errors of the asymmetries are summed quadratically (see Eqs. (3) and (4))

Fit	Bf	A_{CP}	A_{FB}
Ex	5.9 ± 0.5	-2.2 ± 3.2	37.0 ± 2.4
F1	2.96 ± 0.27	$8.0^{+0.24}_{-0.29}$	7.6 ± 0.1
F2	$6.21^{+2.40}_{-0.49}$	$11.7^{+9.5}_{-7.1}$	$11.4^{+0.1}_{-1.5}$

5 represent, respectively, the differential branching fractions of the K^+ and K^- decays separately and the differential CP asymmetry, A_{CP} , Eq. (49), which could be measured in the future.

Two comments are in order about the results given in Tables 1 and 2. First of all, the parameters C_A and S_0 , as determined by our best fit (see line F2 of Table 1), are affected by large, asymmetric errors; however, while the value of C_A is compatible with factorization (compare with line F1), S_0 is much greater, which suggests that the current term derives a large contribution from the effects described in Sect. 3.2. As regards Table 2, the theoretical value of A_{CP} is affected by a large error, while the one of A_{FB} is quite small. This is connected to the experimental errors and to the definition itself of these asymmetries. A_{CP} is proportional to the difference between two large and almost equal quantities and is known experimentally with a relative error of 141 %; therefore, its dependence on the fit parameters is quite mild. The opposite happens for A_{FB} , whose experimental value is not so close




Fig. 5 Differential CP asymmetry vs $m_{p\bar{p}}$

to the theoretical one and is affected by a quite small relative error [2].

4 Discussion and conclusions

Here we comment on our main results.

- (a) As regards the current contribution, we have adopted the parametrization suggested by the pole model [24], like Ref. [9]. As already observed in Sect. 3.4, by comparing the lines L1 and L2 of Table 1, we conclude that,

according to our fit, the current term is much greater than expected from factorization. Moreover, we see from line F2 that $|S_0| \gg |S_1|$, therefore the term corresponding to the angular momentum $J = 0$ prevails neatly over the one with $J = 1$, in agreement with the low-mass character of the $p\bar{p}$ peak. However, the term with $J = 1$ gives an essential contribution to the fit to the Dalitz plot distribution, as expected from C and P violation in weak decays [22].

- (b) The FB asymmetry (A_{FB}) – confirmed by the Dalitz plot distribution [2, 17] – is caused by the linear terms in $\cos \theta_p$, which appear in the differential decay width (38), precisely,
 - in the Jacobian (36), owing to the Lorentz boost from the $p\bar{p}$ frame to the B rest frame;
 - in the interference term between the pseudoscalar and the vector amplitude, whose coefficient is Ω_{13} , Eq. (31).

In our model – for which, incidentally, the second cause of asymmetry shown above is ruled out, owing to the absence of the pseudoscalar amplitude – A_{FB} is positive for the K decay, not only as a result of our fit, but also according to the factorization assumption, as shown in Table 2. In this connection, it is worth observing that, in the model proposed, the sign of A_{FB} is not a priori determined; it depends on the moduli and phases of the amplitudes involved in the decay, according to the β coefficients (20) to (23). Furthermore, this is not typical of the K decay. Indeed, also in the case of π decay, the parametrization by Ref. [9] yields a positive A_{FB} , whereas experiments find a negative value [2, 15, 17]. Therefore, our picture seems to contradict the pole model conclusions and naive quark model expectations [7]. Rather, as observed in Sect. 3.1, the negative FB asymmetry of the π decay could be explained by means of the color-suppressed tree diagram.

- (c) In the K decay, the gap between the experimental value of A_{FB} and the result of our fit (see Table 2) could be filled by the first term of Eq. (22), concerning the electric form factor of the proton, neglected also in Ref. [9], which is linear in $\cos \theta_p$. This explanation of the large asymmetry could be alternative to the one proposed by other authors [11].
- (d) Lastly, the CP asymmetry (A_{CP}) is found to be positive according to our parametrization, which does not differ so much from the prediction of the factorization assumption (see also Ref. [11]); on the contrary, experiments seem to indicate a negative value, although compatible with a positive asymmetry within errors. Therefore, if a more accurate determination of this observable could be desirable, also a deeper analysis of the model is needed. Indeed, although less important than in meson B decays,

the FSI could produce a relative phase between the current term and the transition amplitude, so as to change the sign of A_{CP} [26, 28]. In other words, the difference of sign between data and model predictions does not necessarily imply new physics.

Acknowledgments The authors are grateful to Prof. Chua for his very kind help. Moreover, they thank their friends C. Patrignani and P. Saracco for useful and stimulating discussions and for technical support.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Funded by SCOAP³.

Appendix

Here we give the numerical values of the CKM matrix elements and of the Wilson coefficients involved in the decays considered, as well as the expressions of the matrix elements that appear in the expressions of the current and of the transition terms, according to Refs. [9, 23].

CKM matrix elements

We give the CKM matrix elements [23] which enter the expressions of the decay amplitudes for $B^- \rightarrow K^-(\pi^-)p\bar{p}$:

$$V_{ub} = A\lambda^3(\rho - i\eta), \quad V_{tb} = 1, \quad V_{us}^* = \lambda, \quad (A. 1)$$

$$V_{ts}^* = -A\lambda^2, \quad V_{ud}^* = 1 - \frac{1}{2}\lambda^2, \quad (A. 2)$$

$$V_{td}^* = A\lambda^3(1 - \rho + i\eta),$$

with $A = 0.814$, $\lambda = 0.226$, $\rho = 0.117$ and $\eta = 0.353$. For the CP-conjugated decays one has to take the complex conjugated elements.

Wilson coefficients

For the $B^\pm \rightarrow K^\pm p\bar{p}$ decay, we have [9]

$$\begin{aligned} a_1 &= 1.05, & a_2 &= 0.02, & a_3 &= (72.7 - 0.3i) \cdot 10^{-4}, \\ a_4 &= -(3.873 + 1.21i) \cdot 10^{-2}, & a_5 &= -(66 + 0.3i) \cdot 10^{-4}, \\ a_6 &= -(5.553 + 1.21i) \cdot 10^{-2}, & a_9 &= -(92.6 + 2.7i) \cdot 10^{-4}. \end{aligned} \quad (A. 3)$$

As for the $B^\pm \rightarrow \pi^\pm p\bar{p}$ decay, the coefficients read

$$\begin{aligned} a_1 &= 1.05, & a_2 &= 0.02, & a_3 &= (73 + 0.3i) \cdot 10^{-4}, \\ a_4 &= -(3.757 + 1.083i) \cdot 10^{-2}, & a_5 &= (-66 + 0.3i) \cdot 10^{-4}, \\ a_6 &= -(5.447 + 1.083i) \cdot 10^{-2}, & a_9 &= -(92.4 + 2.5i) \cdot 10^{-4}. \end{aligned} \quad (A. 4)$$

Non-perturbative matrix elements

(A) The matrix element $\langle K^- | L_{sb}^\mu | B^- \rangle$, which appears in the current term (8), reads [9]

$$\langle K^- | L_{sb}^\mu | B^- \rangle = k^\mu r F_0 + l^\mu F_1, \tag{A. 5}$$

with

$$k = p_p + p_{\bar{p}}, \quad l = 2p_B - (1+r)k, \quad r = \frac{m_B^2 - m_K^2}{k^2} \tag{A. 6}$$

and

$$F_0 = \frac{S_0}{1 - u_1\tau + u_2\tau^2}, \quad F_1 = \frac{S_1}{(1 - \tau)(1 - w\tau)}, \tag{A. 7}$$

$$\tau = \frac{t}{m_V^2}.$$

In order to get the expression of $\langle \pi^- | L_{sb}^\mu | B^- \rangle$, we have to replace m_K by m_π in r . Moreover, the numerical values of the parameters which appear in Eq. (A. 7) are

$$m_V = 5.42 \text{ (5.32) GeV}, \quad u_1 = 0.70 \text{ (0.76)}, \tag{A. 8}$$

$$u_2 = 0.27 \text{ (0.28)}, \quad w = 0.43 \text{ (0.48)},$$

the values outside (inside) parentheses referring to the $K(\pi)$ decay. Lastly, S_0 and S_1 , which have the dimension of an energy, are treated as free parameters and determined according to the two possible choices described in the text; see Table 1.

(B) Now we give the expressions of other matrix elements involved in the current terms in K decay [9]:

$$\langle K | \tilde{L}_{sb} | B \rangle = \frac{m_B^2 - m_K^2}{m_b - m_s} F_0, \tag{A. 9}$$

$$\langle p\bar{p} | L_\mu^i | 0 \rangle = \bar{u}(p_p) \left[\Phi^i \gamma_\mu + \hat{\Phi}^i \frac{\delta_\mu}{2m_p} - \left(g_A^i \gamma_\mu \gamma_5 + \gamma_5 \frac{h_A^i}{2m_p} k_\mu \right) \right] v(p_{\bar{p}}), \tag{A. 10}$$

$$\langle p\bar{p} | R_\mu | 0 \rangle = \bar{u}(p_p) \left[\Phi^2 \gamma_\mu + \hat{\Phi}^2 \frac{\delta_\mu}{2m_p} - \left(g_A^2 \gamma_\mu \gamma_5 + \gamma_5 \frac{h_A^2}{2m_p} k_\mu \right) \right] v(p_{\bar{p}}), \tag{A. 11}$$

$$\langle p\bar{p} | \tilde{R}_{ss} | 0 \rangle = \left(\frac{m_p}{m_s} g_A^3 + \frac{k^2}{4m_p m_s} h_A^3 \right) \bar{u}(p_p) \gamma_5 v(p_{\bar{p}}). \tag{A. 12}$$

Here the first and last formulas are deduced from the equations of motion; in particular, the first one exploits also Eq. (A. 5), contracted with k_μ , and the relation $k \cdot l = 0$. Moreover

$$\delta = p_{\bar{p}} - p_p, \tag{A. 13}$$

$$\Phi^1 = G_M^p - G_M^n + S_V, \quad \Phi^2 = 3(S_V - G_M^n), \tag{A. 14}$$

$$\Phi^3 = S_V - G_M^p - 2G_M^n, \quad \Phi^4 = G_M^p,$$

$$G_M^p = \lambda(t) \sum_{n=1}^5 \frac{x_n}{t^{n+1}}, \quad G_M^n = -\lambda(t) \sum_{n=1}^2 \frac{y_n}{t^{n+1}}, \tag{A. 15}$$

$$S_V = -\lambda(t) \frac{y_3}{t^2},$$

$$\lambda(t) = [ln(t/\Lambda_0^2)]^\gamma, \quad \Lambda_0 = 0.3 \text{ GeV}, \quad \gamma = -2.148, \tag{A. 16}$$

$$x_1 = 420.96 \text{ GeV}^4, \quad x_2 = -10485.50 \text{ GeV}^6, \tag{A. 17}$$

$$x_3 = 106390.97 \text{ GeV}^8, \quad x_4 = -433916.61 \text{ GeV}^{10},$$

$$x_5 = 613780.15 \text{ GeV}^{12}, \tag{A. 18}$$

$$y_1 = 236.69 \text{ GeV}^4, \quad y_2 = -579.51 \text{ GeV}^6, \tag{A. 19}$$

$$y_3 = -52.42 \text{ GeV}^4$$

and

$$g_A^1 = f_A + d_A + s_A, \quad g_A^2 = 2d_A + 3s_A, \tag{A. 20}$$

$$g_A^3 = -f_A + d_A + s_A, \quad g_A^4 = f_A + 1/3d_A,$$

$$f_A = \lambda(t) \sum_{n=1}^2 \frac{w_n}{t^{n+1}}, \quad d_A = \lambda(t) \sum_{n=1}^2 \frac{z_n}{t^{n+1}}, \tag{A. 21}$$

$$s_A = \lambda(t) \frac{z_3}{t^2},$$

$$w_1 = 399 \text{ GeV}^4, \quad w_2 = -1055 \text{ GeV}^6, \tag{A. 22}$$

$$z_1 = 65.93 \text{ GeV}^4, \quad z_2 = -1055 \text{ GeV}^6, \tag{A. 23}$$

$$z_3 = 333.06 \text{ GeV}^4.$$

Lastly, we have set

$$\hat{\Phi}^i = 0 \quad \text{and} \quad h_A^i = \frac{-4m_p^2}{t - m_{\pi_0}} g_A^i, \quad i = 1 \text{ to } 4. \tag{A. 24}$$

The corresponding expressions for the π decay are obtained by replacing m_K by m_π , m_s by m_d , Φ^3 by S_V , and g_A^3 by s_A .

(C) The matrix elements involved in the transition term (10) are [9]

$$\langle K | \tilde{L}_{us} | 0 \rangle \langle p\bar{p} | S^{ub} | 0 \rangle = -f_K m_b [F_A \bar{u}(p_p) \not{p}_K \gamma_5 v(p_{\bar{p}}) + F_P \bar{u}(p_p) \gamma_5 v(p_{\bar{p}})], \tag{A. 25}$$

$$\langle K | \tilde{L}_{us} | 0 \rangle \langle p\bar{p} | P^{ub} | 0 \rangle = -f_K m_b [F_V \bar{u}(p_p) \not{p}_K v(p_{\bar{p}}) + F_P \bar{u}(p_p) v(p_{\bar{p}})]. \tag{A. 26}$$

Here $f_K = 0.158$ GeV and

$$F_A = \frac{C_A}{t^3}, \quad F_{V5} = \frac{C_{V5}}{t^3}, \quad F_P = \frac{C_P}{t^4}, \quad (\text{A. 27})$$

C_A , C_{V5} , and C_P being assumed, again, as free parameters and determined as described in the text.

In the case of the π decay, one has to replace f_K by $f_\pi = 0.133$ GeV.

References

1. I.I. Bigi and A.I. Sanda: "CP Violation", Second Edition, Cambridge University Press, Cambridge CB2 8R, UK, May 2009
2. R. Aaij et al., LHCb Coll. Phys. Rev. D **88**, 052015 (2013)
3. R. Aaij et al., LHCb Coll. Phys. Rev. Lett. **111**, 101801 (2013)
4. R. Aaij et al., LHCb Coll. Phys. Rev. Lett. **113**, 141801 (2014)
5. H. Albrecht et al., ARGUS Coll. Phys. Lett. B **209**, 119 (1988)
6. C. Bebek et al., CLEO Coll. Phys. Rev. Lett. **62**, 2436 (1989)
7. H.-Y. Cheng, Int. Jou. Mod. Phys. A **21**, 4209 (2006)
8. W.-S. Hou, A. Soni, Phys. Rev. Lett. **86**, 4247 (2001)
9. C.-K. Chua, W.-S. Hou, S.-Y. Tsai, Phys. Rev. D **66**, 054004 (2002)
10. J. Rosner, Phys. Rev. D **68**, 014004 (2003)
11. C.Q. Geng, Y.K. Hsiao, Phys. Rev. D **74**, 094023 (2006)
12. C.Q. Geng, Y.K. Hsiao, J.N. Ng, Phys. Rev. Lett. **98**, 011810 (2007)
13. A. Bornheim et al., CLEO Coll. Phys. Rev. D **68**, 052002 (2003)
14. M.C. Chang et al., Belle Coll. Phys. Rev. D **71**, 072007 (2005)
15. J. T. Wei et al., Belle Coll.: Phys Lett. B **659**, 80 (2008)
16. B. Aubert et al., BaBar Coll. Phys. Rev. D **69**, 091503 (2004)
17. B. Aubert et al., BaBar Coll. Phys. Rev. D **72**, 051101 (2005)
18. T. Latham: DPF 2013 - The Meeting of the APS - Division of Particles and Fields, Santa Cruz, California, August 13-17, 2013
19. R. Marshak, Riazuddin and C, *Theory of Weak Interactions in Particle Physics* (Wiley-Interscience, New York, NY, 1969)
20. L. Wolfenstein, Phys. Rev. D **43**, 151 (1991)
21. Z.J. Ajaltouni, E. Di Salvo, Int. Jou. Mod. Phys. E **22**, 1330006 (2013)
22. J. Haidenbauer et al., Phys. Rev. D **74**, 017501 (2006)
23. K.A. Olive et al., Particle Data Group. Chin. Phys. C **38**, 090001 (2014)
24. D. Melikhov, B. Stech, Phys. Rev. D **62**, 014006 (2000)
25. K. Abe et al., Belle Coll. Phys. Rev. Lett. **88**, 181803 (2002)
26. H.-Y. Cheng, C.-K. Chua, A. Soni, Phys. Rev. D **71**, 014030 (2005)
27. H.-Y. Cheng, K.-C. Yang, Phys. Rev. D **66**, 014020 (2002)
28. H.-Y. Cheng, C.-K. Chua, A. Soni, Phys. Rev. D **72**, 014006 (2005)