



RESEARCH

Open Access

The effects of dietary fiber level on nutrient digestibility in growing pigs

Wenjuan Zhang, Defa Li, Ling Liu, Jianjun Zang, Qiwu Duan, Wenjun Yang and Liying Zhang*

Abstract

The objective of this study was to investigate the effects of total dietary fiber level on nutrient digestibility and the relationship between apparent total tract digestibility of total dietary fiber, and soluble dietary fiber, insoluble dietary fiber and available energy. Sugar beet pulp was as the only fiber source. The experiment was designed as a 6×6 Latin square with an adaptation period of 7 d followed by a 5-d total collection of feces and urine. Feed intake tended to decrease ($P = 0.10$) as total dietary fiber level increased. The apparent total tract digestibility of dry matter, crude protein and gross energy decreased ($P < 0.01$) when total dietary fiber increased but the digestibility of soluble dietary fiber and insoluble dietary fiber increased ($P < 0.01$). The digestible energy and metabolizable energy content of diets decreased ($P < 0.01$) as the total dietary fiber increased.

Keywords: Growing pigs, Insoluble dietary fiber, Nutrient digestibility, Soluble dietary fiber, Total dietary fiber

Background

Total dietary fiber (TDF) is the sum of the dietary carbohydrates that are resistant to digestion by mammalian enzymes in the small intestine but can be partially or completely fermented in the hindgut [1]. According to its solubility, TDF can be divided into soluble dietary fiber (SDF) and insoluble dietary fiber (IDF) [2]. Dietary fiber is a key factor determining nutrient utilization in the diet and more emphasis should be given to routine techniques that identify the nutritional and physiological "quality" of dietary fiber [3].

Serena et al. (2008) reported that SDF has a high water holding capacity, delays gastric emptying, slows the rate of nutrient absorption [4]. Most of SDF and partial of IDF are degraded by bacteria in either the small or large intestine [4-6]. Energy produced by microflora in the hindgut can satisfy up to 30% of the maintenance energy requirements of the pig [7]. In addition, IDF was found to decrease intestinal transit time, binds organic compounds and increases fecal bulk [8].

A minimum level of fiber has to be included in pig diets to support normal physiological activity in the digestive tract [9]. Mateos et al. (2006) suggested that young pigs may have a minimum requirement for a fiber level

of 6% [10]. However, diets or ingredients with a high fiber content may negatively affect voluntary feed intake and nutrient digestibility in young pigs [11,12]. Understanding the effects of TDF level in the diet on nutrient digestibility and feed intake is critical for optimal swine production.

The effect of TDF on the digestibility of nutrients in the diet is controversial. Wilfart et al. (2007) added 0, 20 and 40% wheat bran to a wheat-barley-soybean meal diet and found that an increase of TDF significantly decreased the apparent total tract digestibility (ATTD) of dry matter, organic matter, crude protein and gross energy, but the ATTD of TDF was unaffected [12]. However, Urriola and Stein (2010) reported that the digestibility of dry matter, gross energy and TDF in diet with 30% distillers dried grains with soluble (DDGS) was significantly lower than a corn-soybean meal control diet fed to growing pigs [13]. Additionally, Bindelle et al. (2009) reported that when growing pigs were fed corn-soybean meal diets supplemented with sugar beet pulp at levels of 0, 10, 20 and 30%, the TDF level increased from 9.6 to 25.4% while the ATTD of dry matter, organic matter and crude protein linearly decreased but the ATTD of NDF was linearly increased [14]. Some of these differences may be due to differences in ATTD of TDF between the basal diet and the fiber sources.

* Correspondence: zhangliying01@sina.com

State Key Laboratory of Animal Nutrition, China Agricultural University, Beijing 100193, P. R. China

The impact of fiber level on digestibility may differ with the properties of the fiber (i.e. soluble vs. insoluble) [15]. Sugar beet pulp is characterized by a high content of soluble fiber such as pectins and glucans [16]. However, there are no reports in the literature about the relationship between the type of fiber and its effects on nutrient digestibility. Therefore, in the present experiment, sugar beet pulp was added to diets to determine the effect of fiber type on nutrient digestibility and to study the relationship between the apparent total tract digestibility of TDF and SDF or IDF.

Materials and methods

The Institutional Animal Care and Use Committee at China Agricultural University (Beijing, China) reviewed and approved the protocols used in this study.

Experiment design and housing

Six healthy crossbred (Duroc × Landrace × Large White) barrows were allotted to a 6 × 6 Latin square design. The pigs (average initial body weight of 30.0 ± 1.8 kg) were individually housed in 1.2 m × 0.7 m × 0.96 m stainless steel metabolism cages in an environmentally controlled room (22 ± 2°C).

Diets and feeding

Table 1 shows the nutrient content of the main ingredients used in this experiment. Six diets were formulated by replacing the basal diet with 0, 15, 25, 35, 45 or 55% sugar beet pulp (Table 2). The sugar beet pulp was ground to pass through a > 3.5 mm mesh screen before mixing into the diets. The protein level of diets was maintained at the same level by adjusting the amount of casein. The mineral, vitamin and amino acid premix was added to the diets at a level sufficient to meet or exceed the nutrient requirements of the National Research Council [17] for pigs weighing 20 to 50 kg.

The daily feed allowance was equivalent to 4% of body weight at the beginning of each period [18]. The allowance was divided into two equal parts and fed at 08:00 and 17:00 h. The diets were mixed with water in a ratio of 1:1 (Wt/Wt) before feeding. Water was available *ad libitum* through a drinking nipple. The pigs were weighed individually at the beginning of each period and the amount of feed supplied each period was recorded, as well

as any feed refusals. Each experimental period consisted of a 7-d adaptation period followed by a 5-d collection of feces and urine. The collected urine was weighed and 10% of the daily urine volume was stored at -20°C. The collection of feces and urine were conducted according to the methods described by Song et al. (2003) [19]. Feces were collected immediately when the feces appeared in the metabolism cages, kept in plastic bags and stored at -20°C. Urine was collected into urine collection buckets that were placed under the metabolism cages. The buckets were emptied each afternoon and 50 mL of 6 mol/L HCl was added. At the end of the experiment, feces and urine samples were thawed and mixed within animal and diet, and a subsample was collected for chemical analysis. Fecal samples were dried in a forced air oven and ground through a 1-mm screen, and thoroughly mixed before a subsample was collected for chemical analysis.

Chemical analyses

Diets and feces were analyzed for dry matter (AOAC method 930.15) [20] and crude protein (AOAC method 990.03) [20]. TDF and IDF were also determined (AOAC method 985.29) [20]. The concentration of SDF in the diets was calculated as the difference between TDF and IDF. In addition, the diets were analyzed for calcium and total phosphorus (AOAC method 985.01) [20]. The gross energy in diets, feces, and urine were analyzed using an adiabatic oxygen bomb calorimeter (Parr Instruments, Moline, IL). The content of nitrogen in the urine was also analyzed (AOAC method 990.03) [20]. Amino acids in the feeds were determined by hydrolyzing the feed with 6 mol/L HCl for 24 h at 110°C (AOAC method 982.30 E) [20] and analyzed using a Hitachi L-8900 Amino Acid Analyzer (Tokyo, Japan). Methionine was determined as methionine sulfone after cold performic acid oxidation over night and hydrolyzing with 7.5 mol/L HCl for 24 h at 110°C.

Statistical analysis

The data for effects of dietary sugar beet pulp on the apparent total tract digestibility (ATTD) of fiber, dry matter, protein, energy and the available energy of diets in growing pigs were subjected to an Analysis of Variance using PROC GLM of SAS (Statistical Analysis System 9.1, SAS Institute, Cary, NC, USA). Orthogonal polynomial

Table 1 Chemical composition of rice starch, casein and sugar beet pulp (as-fed basis)

	Moisture	Gross energy [MJ/kg]	Nutrient value ^a , %			
			Crude protein	TDF	SDF	IDF
Rice starch	12.6	14.7				
Casein	8.2	21.6	82.4			
Sugar beet pulp	13.1	15.9	9.6	69.1	22.0	47.1

^aAnalyzed values. TDF = Total dietary fiber; SDF = Soluble dietary fiber; IDF = Insoluble dietary fiber.

Table 2 Ingredient composition, energy and nutrient levels of the diets (% as fed)

	Sugar beet pulp, %					
	0.0	15.0	25.0	35.0	45.0	55.0
Ingredients, %						
Rice starch	76.0	62.8	54.0	45.2	36.3	27.5
Casein	20.0	18.2	17.0	15.8	14.7	13.5
Sugar beet pulp	0.0	15.0	25.0	35.0	45.0	55.0
Vitamin and mineral premix ^a	4.0	4.0	4.0	4.0	4.0	4.0
Energy and nutrient values ^b						
Digestible energy, MJ/kg	14.7	14.8	14.2	14.0	13.6	13.1
Crude protein, %	17.5	17.3	17.3	17.4	17.3	17.3
Total dietary fiber, %	0.0	12.1	18.8	24.8	32.0	38.9
Soluble dietary fiber, %	0.0	4.2	6.0	8.0	10.1	12.2
Insoluble dietary fiber, %	0.0	7.9	12.8	16.8	21.9	26.7
Calcium, %	0.77	0.79	0.81	0.81	0.80	0.82
Phosphorus, %	0.50	0.55	0.54	0.57	0.54	0.55
	Sugar beet pulp, %					
	0.0	15.0	25.0	35.0	45.0	55.0
Amino acids, %						
Aspartic acid	1.28	1.26	1.13	1.26	1.14	1.34
Threonine	0.64	0.70	0.69	0.71	0.71	0.78
Serine	0.81	0.88	0.88	0.83	0.76	0.87
Glutamic acid	3.85	3.96	3.86	3.65	3.53	3.48
Proline	1.61	1.69	1.59	1.40	1.61	1.48
Glycine	0.42	0.37	0.35	0.42	0.41	0.50
Alanine	0.65	0.61	0.66	0.64	0.65	0.68
Valine	1.08	1.14	1.02	1.11	1.00	1.14
Isoleucine	0.89	0.90	0.85	0.86	0.82	0.85
Leucine	1.56	1.63	1.59	1.52	1.49	1.50
Tyrosine	0.82	0.80	0.72	0.76	0.72	0.77
Phenylalanine	0.92	0.92	0.90	0.88	0.85	0.84
Histidine	0.47	0.48	0.54	0.49	0.50	0.50
Lysine	1.55	1.60	1.54	1.58	1.46	1.58
Arginine	0.41	0.55	0.56	0.56	0.55	0.63
Methionine	0.44	0.46	0.47	0.47	0.47	0.48
Tryptophan	0.21	0.22	0.21	0.23	0.22	0.21

^a Supplied per kilogram of diet: vitamin A; 6.0 KIU; vitamin D₃; 2.4 KIU; vitamin E, 21.6 IU; vitamin K, 2.0 mg; thiamine 1.0 mg; riboflavin 5.2 mg; pyridoxine 2.0 mg; vitamin B₁₂ 0.01 mg; D-pantothenic acid 11.2 mg; niacin 22 mg; biotin 40 µg; folic acid 0.4 mg; 120 mg of Fe; 120 mg of Zn; 40.0 mg of Mn; 80 mg of Cu; 400 µg of I; and 240 µg of Se; 8.0 g of calcium; 0.4 g of phosphorus.

^b The nutrient levels are analyzed values. Digestible energy were calculated values.

contrasts were used to determine linear and quadratic effects of the TDF level on ATTD of energy, nutrients and the available energy of diets in growing pigs. Pig and period were random effects, and TDF level was considered a fixed effect. The PROC CORR and GLM of SAS were

used to analyze the relationship between the ATTD of TDF and the ATTD of IDF or SDF. The model included dietary treatment and the residual mean square error was used as the error term. The means were separated using Duncan's new multiple range test. The individual pig (n = 6 pigs/treatment group) served as the experimental unit. Results are reported as means plus standard errors with $P < 0.05$ defined as significant and $P < 0.10$ as indicative of a trend.

Results

The effects of sugar beet pulp on feed intake, fecal output, and the relationship between digestibility of TDF and IDF or SDF

The effect of TDF on the digestibility of fibrous components is shown in Table 3. The feed intake of diets tended to decrease ($P = 0.10$) as the TDF increased. As expected, the intake of TDF, SDF and IDF and the excretion of TDF and IDF increased ($P < 0.01$) as TDF increased, but the excretion of SDF was unaffected by TDF. The output of feces increased ($P < 0.01$) as the TDF increased. The ATTD of TDF, SDF and IDF increased ($P < 0.01$) when TDF increased. The relationship between ATTD of SDF, IDF and TDF is shown in Figure 1. There was a good relationship between ATTD of TDF and IDF ($r^2 = 0.93$), but a poor relationship between the ATTD of TDF and SDF ($r^2 = 0.28$).

The effects of sugar beet pulp on the digestibility of dry matter, crude protein and energy

The effect of TDF on the digestibility of dry matter, crude protein and energy is shown in Table 4. The gross energy intake tended to decrease ($P = 0.10$) as TDF increased. In addition, the amounts of nitrogen and energy excreted from feces were increased ($P < 0.01$) as dietary fiber level increased, while the amounts of energy excreted from urine was decreased ($P < 0.05$). The ratio of urinary nitrogen to fecal nitrogen tended to decrease ($P = 0.10$) as TDF increased although the amount of nitrogen excreted from urine was not affected ($P > 0.1$) by the dietary fiber level. The digestibility of dry matter, crude protein and energy decreased ($P < 0.01$) as TDF increased.

The effects of sugar beet pulp level on the available energy in the diet

The digestible energy (DE) and metabolizable energy (ME) content of the diets decreased ($P < 0.01$) when the level of sugar beet pulp increased from 15.0 to 55.0% (Table 4). The correlation between the content of TDF and DE, ME of diets is shown in Table 5. The content of TDF had a negative correlation with the DE and ME content of the diet.

Table 3 Effects of dietary sugar beet pulp on the apparent total tract digestibility (ATTD) of fiber in growing pigs

Items	Sugar beet pulp, %						P Values		
	0.0	15.0	25.0	35.0	45.0	55.0	SEM	Linear	Quadratic
Intake, g/d									
Feed intake	1,190.0	1,193.0	1,193.0	1,205.0	1,107.0	1,062.0	64.15	0.09	0.16
Total dietary fiber	0.0	124.0	206.4	292.8	344.3	404.6	23.86	<0.01	<0.01
Soluble dietary fiber	0.0	39.4	65.6	92.8	109.6	128.5	7.58	<0.01	<0.01
Insoluble dietary fiber	0.0	84.7	140.8	200.0	234.7	276.1	16.28	<0.01	<0.01
Excretion, g/d									
Feces, wet basis	135.0	194.9	317.0	409.7	415.2	571.5	19.60	<0.01	<0.01
Feces, dry matter basis	79.3	90.2	127.8	145.3	124.9	159.8	9.00	<0.01	<0.01
Total dietary fiber	0.0	29.9	29.3	35.6	36.4	43.2	3.34	<0.01	<0.01
Soluble dietary fiber	0.0	3.5	3.7	5.5	4.1	4.7	0.67	0.22	0.38
Insoluble dietary fiber	0.0	16.4	25.6	30.1	32.3	38.4	2.86	<0.01	<0.01
ATTD of fiber, %									
Total dietary fiber	0.0	86.2	86.9	88.1	89.7	89.5	0.45	<0.01	<0.01
Soluble dietary fiber	0.0	92.9	94.9	94.3	96.3	96.3	0.49	<0.01	<0.01
Insoluble dietary fiber	0.0	82.6	83.2	85.1	86.7	86.4	0.80	<0.01	<0.01

Discussion

The level of fiber in a pig’s diet is considered an important factor affecting palatability and feed intake although pigs can tolerate relatively high levels of fiber [21]. High fiber levels in diets can decrease the voluntary feed intake of the animals as a consequence of gut fill, compromising the energy intake of pigs [22]. In the present study, the voluntary feed intake of pigs was lower than the full allowance during the trial period when the inclusion of sugar beet pulp was higher than 35.0% or TDF was higher than 24.8%. Likewise, Anguita et al. (2007) also reported that the inclusion of sugar beet pulp decreased voluntary feed intake of pigs more than other less digestible ingredients [23], probably as a result of

the higher amount of digesta and its water retention capacity when the diet contained sugar beet pulp.

Fiber-containing diets could increase fecal output [24]. Wilfart et al. (2007) reported that the output of fecal dry matter increased as TDF increased [12]. In the current study, the output of feces increased as the TDF level increased. The main reason for this included two aspects. One reason is mainly due to the fact that the moisture content of feces increased as the TDF level increased. Another reason is the TDF excretion increased because 50 to 60% of the dry matter excretion at the rectum was TDF [12]. Generally speaking, the increase in fecal output as TDF level increased in the diet was related to the water holding capacity of SDF and increases in fecal bulk of IDF [4].




Figure 1 Relationship between the apparent total tract digestibility (ATTD) of total dietary fiber (TDF) and the ATTD of soluble fiber (SDF) (a) and insoluble dietary fiber (IDF) (b) in sugar beet pulp fed to growing pigs (n = 30).

Table 4 Effects of dietary sugar beet pulp on the apparent total tract digestibility (ATTD) of dry matter, protein, energy and the available energy of diets in growing pigs

Items	Sugar beet pulp, %						P Values		
	0.0	15.0	25.0	35.0	45.0	55.0	SEM	Linear	Quadratic
Intake									
Dry matter intake, g/d	1,088.0	1,095.0	1,091.8	1,083.0	1,022.3	968.3	56.96	0.46	0.26
Nitrogen intake, g/d	33.4	33.6	34.2	33.6	29.8	29.6	1.69	0.40	0.28
Gross energy intake, MJ/d	17.9	18.7	18.5	18.8	17.1	16.3	0.58	0.10	0.07
Excretion									
Fecal nitrogen, g/d	1.9	3.3	4.8	5.2	4.9	6.1	0.44	<0.01	<0.01
Urinary nitrogen, g/d	9.3	7.6	7.4	7.0	6.8	6.2	1.00	0.27	0.54
Urinary:fecal nitrogen ratio	4.9	2.3	1.6	1.4	2.0	1.1	0.13	0.10	0.17
Fecal energy, MJ/d	0.4	1.1	1.6	1.9	2.0	2.3	0.13	<0.01	<0.01
Urinary energy, MJ/d	0.9	0.7	0.6	0.6	0.4	0.4	0.11	<0.01	0.02
ATTD, %									
Dry matter	95.4	91.8	88.3	86.6	87.8	83.5	0.32	<0.01	<0.01
Nitrogen	94.6	90.0	85.9	84.4	83.7	80.0	1.00	<0.01	<0.01
Energy	97.8	94.5	91.4	89.9	88.4	86.3	0.47	<0.01	<0.01
Energy content of diets, MJ/kg									
Digestible energy	14.7	14.8	14.2	14.0	13.6	13.1	0.07	<0.01	0.06
Metabolizable energy	14.0	14.2	13.7	13.5	13.2	12.8	0.10	<0.01	0.71

Soluble fiber is usually susceptible to microbial degradation, thus increasing bacteria growth in the lower gut [25]. The greater the amount of fiber in the diet, the greater the disappearance of fiber (g of fiber disappeared/kg of feed DM) after fermentation in growing pigs [26]. Bindelle et al. (2009) added sugar beet pulp at levels of 0, 10, 20 and 30% to a corn-soybean meal basal diet fed to growing pigs and found a linear increase in the digestibility of neutral detergent fiber [14]. In agreement with the report of Bindelle et al. (2009) [14], the digestibility of TDF, SDF and IDF increased as the TDF level increased in the present study. One reason for this observation may be that there are components in endogenous secretions that are analyzed as TDF (although they are not TDF). The influence of these components is reduced as more TDF is included in the diets, which is the reason for the increased values for ATTD as TDF concentrations in the diets. But this result is not in line with the report of Wilfart et al. (2007) [12], who reported that the digestibility of TDF is unaffected by TDF level of diets fed to growing pigs. The occurrence of different results about the fiber digestibility may be due to different fiber sources used in the two trials (wheat bran vs. sugar beet pulp).

The strong relationship between the ATTD of TDF and the ATTD of IDF but poor relationship between SDF in the current study is in agreement with Urriola et al. (2010) [5], who reported that there was a strong relationship between the ATTD of TDF and the ATTD of IDF in distillers dried grains with solubles but a poor

relationship between the ATTD of TDF and the ATTD of SDF. The poor relationship between the ATTD of TDF and the ATTD of SDF is due to the fact that most of the fiber in distillers dried grains with solubles is insoluble [27]. In this study, although the main component of sugar beet pulp is fiber, and the fiber is about 1/3 soluble and 2/3 insoluble (Table 1), the ATTD of SDF was higher than 92.0%. In other words, most of SDF in sugar beet pulp was fermented in the hindgut. Therefore, there is a poor relationship between the ATTD of TDF and the ATTD of SDF in the present study.

Fiber may improve intestinal health because it is necessary for the stimulation of the intestinal compartments [28], and it is usually associated with a reduction of potentially harmful products from protein fermentation [29]. However, the inclusion of fiber in the diet offered to pigs results in reductions in foregut and whole-tract digestibility of dry matter [27] leading to a lower absorption of nutrients and energy. The digestibility of nutrients in pig diets has been shown to be related

Table 5 Correlation coefficients between dietary fiber and available energy of diets

Items	TDF	DE	ME
Total dietary fiber ^a , %	1.00		
Digestible energy, MJ/kg	- 0.92	1.00	
Metabolizable energy, MJ/kg	- 0.83	0.97	1.00

^a TDF = Total dietary fiber, DE = digestible energy, ME = metabolizable energy.

to the origin and content of dietary fiber [30]. In the present study, inclusion of sugar beet pulp in a casein-rice starch basal diet increased the concentration of TDF, SDF and IDF in the diet. The ATTD of dry matter, crude protein and energy were negatively correlated as sugar beet pulp level increased from 15.0 to 55.0%. The lower ATTD of crude protein can be explained by increased endogenous secretions, or by decreased hydrolysis and absorption of nutrients, or both [12]. Part of the endogenous nitrogen loss was the bacterial nitrogen in the feces. It has been reported that 60 to 90% of fecal nitrogen was of bacterial origin [31]. Bindelle et al. (2009) examined the effect of dietary fiber on bacterial protein synthesis and reported a linear increase of bacterial nitrogen incorporation with graded levels of sugar beet pulp at levels of 10, 20, and 30%, respectively [14]. Similar to the report of Bindelle et al. (2009) [14], the excretion of fecal nitrogen increased when the TDF level increased. This may be the main reason for the decrease in digestibility of crude protein. Just et al. (1984) reported that dietary fiber concentration could account for about 70% of the variation in energy digestibility in diets [32]. Previous studies showed that the digestibility of gross energy decreased with an increase of TDF in the diet [12-14]. In agreement with previous reports, it was found that the digestibility of gross energy decreased when TDF level increased in the diet in this study. Castiglia-Delavaud et al. (1998) reported that about 35.0% of the fermented sugar beet non-starch polysaccharide energy appeared as fecal bacteria energy [33].

Evaluation of the available energy content of pig feeds is usually based on their DE or ME content [3]. It was found that a high fiber content is responsible for adverse effects on the digestible energy content of feeds for pigs [34]. A similar result was found in the present study that the content of TDF in the diet was negatively related to the DE and ME content of the diet. In addition, in agreement with the work of Noblet (2006) [3], who reported that the ratio of ME to DE of complete feeds is approximately 0.96 while in this study the ratio of ME to DE was 0.97.

Conclusions

It is concluded that the digestibility of dry matter, gross energy, and crude protein in diets were negatively affected by the level of sugar beet pulp, which ranged from 15.0 to 55.0%, but the digestibility of SDF and IDF increased with the increase of TDF. There was a strong relationship between the ATTD of TDF and the ATTD of IDE, but the relationship between ATTD of TDF and the ATTD of SDF was poor.

Abbreviations

TDF: Total dietary fiber; SDF: Soluble dietary fiber; IDF: Insoluble dietary fiber; ATTD: Apparent total tract digestibility; DDGS: Distillers dried grains with soluble; DE: Digestible energy; ME: Metabolizable energy.

Competing interests

The authors declare that they have no competing interests.

Author's contributions

WJZ carried out the experiment trial, performed the statistics and drafted the manuscript. DFL and LL participated in design of the study. JJZ and QWD participated in animal trial. WJY participated sample analysis. LYZ conceived the study, and participated in its design and coordination. All authors read and approved the final manuscript.

Acknowledgements

This research was financially supported by the National Key Technology R & D Program in the 11th Five Yr Plan of China (2006BAD12B01) and Special Public Sector Fund in Agriculture (200903006). Thanks to Xianjiang Wen, Zijuan Fan and Lingfeng Xue for their practical support.

Received: 4 February 2013 Accepted: 11 April 2013

Published: 15 April 2013

References

1. Institute of Medicine: *Dietary, functional, and total dietary fiber*. Washington, DC: National Academic Press; 2006:340-421.
2. Prosky L, Asp NG, Schweizer TF, de Vries JW, Furda I: **Determination of insoluble and soluble dietary fiber in foods and food products: Collaborative study**. *J Assoc Off Anal Chem* 1988, **71**:1017-1023.
3. Noblet J: **Energy evaluation of feeds for pigs: consequences on diet formulation and environment protection**. *Lohmann Inf* 2006, **41**:38-51.
4. Serena A, Jørgensen H, Bach Knudsen KE: **Digestion of carbohydrates and utilization of energy in sows fed diets with contrasting levels and physicochemical properties of dietary fiber**. *J Anim Sci* 2008, **86**:2208-2216.
5. Urriola PE, Shurson GC, Stein HH: **Digestibility of dietary fiber in distillers coproducts fed to growing pigs**. *J Anim Sci* 2010, **88**:2373-2381.
6. Urriola PE, Stein HH: **Comparative digestibility of energy and nutrients in fibrous feed ingredients fed to Meishan and Yorkshire pigs**. *J Anim Sci* 2012, **90**:802-812.
7. Adeshinwa AOK: **Energy and protein requirements of pigs and the utilization of fibrous feedstuffs in Nigeria: A review**. *Afr J Biotech* 2008, **7**:4798-4806.
8. Renteria-Flores JA: *Effect of soluble and insoluble fiber on diet digestibility and sow performance*, Doctoral Thesis. University of Minnesota: St. Paul; 2006:161.
9. Wenk C: **The role of dietary fibre in the digestive physiology of the pig**. *Anim Feed Sci Technol* 2001, **90**:21-33.
10. Mateos GG, Martin F, Latorre MA, Vicente B, Lazaro R: **Inclusion of oat hulls in diets for young pigs based on cooked maize or cooked rice**. *J Anim Sci* 2006, **82**:57-63.
11. Kyriazakis I, Emmans GC: **The voluntary feed intake of pigs given feeds based on wheat bran, dried citrus pulp and grass meal, in relation to measurements of bulk**. *Br J Nutr* 1995, **73**:191-207.
12. Wilfart A, Montagne L, Simmins H, Noblet J, Van Milgen J: **Effect of fibre content in the diet on the mean retention time in different segments of the digestive tract in growing pigs**. *Livest Sci* 2007, **109**:27-29.
13. Urriola PE, Stein HH: **Effects of distillers dried grains with solubles on amino acid, energy, and fiber digestibility and on hindgut fermentation of dietary fiber in a corn-soybean meal diet fed to growing pigs**. *J Anim Sci* 2010, **88**:1454-1462.
14. Bindelle J, Buldgen A, Delacollette M, Wavreille J, Agneessens R, Destain JP, Leterme P: **Influence of source and concentrations of dietary fiber on in vivo nitrogen excretion pathways in pigs as reflected by in vitro fermentation and nitrogen incorporation by fecal bacteria**. *J Anim Sci* 2009, **87**:583-593.
15. Högberg A, Lindberg JE: **Influence of cereal non-starch polysaccharides and enzyme supplementation on digestion site and gut environment in weaned piglets**. *Anim Feed Sci Technol* 2004, **116**:113-128.
16. Fadel JG, DePeters EJ, Arosemena A: **Composition and digestibility of beet pulp with and without molasses and dried using three methods**. *Anim Feed Sci Technol* 2000, **85**:121-129.
17. NRC: *Nutrient Requirements of Swine, 10th rev. ed.* Washington, DC: Natl. Acad. Press; 1998.
18. Adeola O: *Digestion and balance techniques in pigs*. New York: CRC Press; 2001:903-916.

19. Song GL, Li DF, Piao XS, Chi F, Yang WJ: **Apparent ileal digestibility of amino acids and the digestible and metabolizable energy content of high-oil corn varieties and its effects on growth performance of pigs.** *Arch Anim Nutr* 2003, **57**:297–306.
20. AOAC: *Official methods of analysis*. 18th ed. Gaithersburg, MD, USA: Association of Official Analytical Chemists; 2007.
21. Solà-Oriol D, Roura E, Torrallardona D: **Feed preference in pigs: Effect of cereal sources at different inclusion rates.** *J Anim Sci* 2009, **87**:562–570.
22. da Silva CS, van den Borne JJ, Gerrits WG, Kemp B, Bolhuis JE: **Effects of dietary fibers with different physicochemical properties on feeding motivation in adult female pigs.** *Physiol Behav* 2012, **107**:218–230.
23. Anguita M, Gasa J, Nofrarias M, Martín-Orúe SM, Pérez JF: **Effect of coarse ground corn, sugar beet pulp and wheat bran on the voluntary intake and physicochemical characteristics of digesta of growing pigs.** *Livest Sci* 2007, **107**:182–191.
24. Hansen MJ, Chwalibog A, Tauson AH: **Influence of different fibre sources in diets for growing pigs on chemical composition of faeces and slurry and ammonia emission from slurry.** *Anim Feed Sci Technol* 2007, **134**:326–336.
25. He GF: *Improving pig performance and nutrient utilization by manipulating dietary protein level and using fermentable fiber ingredients*, Doctoral Thesis. University of Minnesota: St. Paul; 2004:220.
26. Anguita M, Canibe N, Pérez JF, Jensen BB: **Influence of the amount of dietary fiber on the available energy from hindgut fermentation in growing pigs: Use of cannulated pigs and in vitro fermentation.** *J Anim Sci* 2006, **84**:2766–2778.
27. Bach Knudsen KE, Hansen I: **Gastrointestinal implications in pigs of wheat and oat fractions 1. Digestibility and bulking properties of polysaccharides and other major constituents.** *Br J Nutr* 1991, **65**:217–232.
28. Longland AC, Carruthers J, Low AG: **The ability of piglets 4 to 8 wks old to digest and perform on diets containing two contrasting sources of non starch polysaccharide.** *Anim Prod* 1994, **58**:405–410.
29. McBurney MI, Van Soest PJ, Jeraci JL: **Colonic carcano genesis: The microbial feast or famine mechanism.** *Nutr Cancer* 1987, **10**:23–28.
30. de Vriesa S, Pustjensb AM, Scholsb HA, Hendriksa WH, Gerrits JJ: **Improving digestive utilization of fiber-rich feedstuffs in pigs and poultry by processing and enzyme technologies: A review.** *Anim Feed Sci Technol* 2012, **178**:123–138.
31. Luis AR: **Carbohydrates digestibility and faecal nitrogen excretion in rats fed rawor germinated faba bean (*Vicia faba*) and chickpea (*Cicerarietinum*) based diets.** *Br J Nutr* 2003, **90**:301–309.
32. Just A, Jørgensen H, Fernández JA: **Prediction of metabolizable energy for pigs on the basis of crude nutrients in the feeds.** *Livest Prod Sci* 1984, **11**:105–128.
33. Castiglia-Delavaud C, Verdier E, Besle JM, Vernet J, Boirie Y, Beaufre're B, De Baynast R, Vermorel M: **Netenergy value of non-starch polysaccharide isolates (sugar beet fibre and commercial inulin) and their impact on nutrient digestive utilization in healthy human subjects.** *Br J Nutr* 1998, **80**:343–352.
34. Noblet J, Perez JM: **Prediction of digestibility of nutrients and energy values of pig diets from chemical analysis.** *J Anim Sci* 1993, **71**:3389–3398.

doi:10.1186/2049-1891-4-17

Cite this article as: Zhang et al.: The effects of dietary fiber level on nutrient digestibility in growing pigs. *Journal of Animal Science and Biotechnology* 2013 **4**:17.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

