

## ORIGINAL RESEARCH

## Open Access

# Hemolymph osmotic, ionic status, and branchial $\text{Na}^+/\text{K}^+$ -ATPase activity under varying environmental conditions in the intertidal grapsid crab, *Gaetice depressus*

Takeshi Nanba<sup>1</sup>, Hideya Takahashi<sup>1,2</sup>, Tsukasa Abe<sup>1,3</sup>, Waichirou Godo<sup>1</sup>, Maho Ogoshi<sup>1</sup>, Hirotaka Sakamoto<sup>1</sup>, Naoaki Tsutsui<sup>1</sup> and Tatsuya Sakamoto<sup>1\*</sup>

\* Correspondence: [ryu@uml.okayama-u.ac.jp](mailto:ryu@uml.okayama-u.ac.jp)  
<sup>1</sup>Ushimado Marine Institute, Faculty of Science, Okayama University, Ushimado, Setouchi 701-4303, Japan  
Full list of author information is available at the end of the article

## Abstract

Osmo- and ionoregulatory abilities were examined in the intertidal grapsid crab, *Gaetice depressus*, transferred from normal seawater (30 ppt) to low (10 ppt) or high (50 ppt) salinities for 2 and 10 days, in addition to animals kept out of water for 2 days. The results of the hemolymph osmotic and ionic status indicate that *G. depressus* is able to adapt for more than 10 days in these salinities and for 2 days under terrestrial conditions. Especially, the free  $\text{Ca}^{2+}$  concentration was relatively maintained compared with concentrations of monovalent ions and osmolality values in 10 and 50 ppt, partly using the complexed calcium (total minus free calcium) as an internal reserve in the hemolymph. In 10 ppt, complexed calcium disappeared from the hemolymph after 10 days, indicating that all the hemolymph calcium was ionized. In 50 ppt, free  $\text{Ca}^{2+}$  was regulated to lower levels than concentrations in the medium, while total calcium increased to higher levels after 2 days. Examination of  $\text{Na}^+/\text{K}^+$ -ATPase activity, which has been implicated in ion transport in many crustaceans, revealed that induction of high  $\text{Na}^+/\text{K}^+$ -ATPase activity varies among the posterior gills in response to salinities. Ten-ppt salinity induces activity in two of the posterior gills (gill numbers 6 and 7, eight in total), albeit with differing degrees of response. In contrast, 50-ppt salinity stimulates the activity primarily in gill number 8, suggesting that this gill may be associated specifically with ion excretion in *G. depressus*. As a euryhaline amphibious crab, this abundant species around Japan will serve as a model to study the osmotic/ionic regulatory mechanisms which operate in crustaceans.

**Keywords:** Crustacean, Osmoregulation, Salinity

## Background

Osmoregulation of decapod crustaceans after hyposaline exposure has been extensively studied. In crabs, the primary site of regulation is the posterior gills which have the highest specific activity of  $\text{Na}^+/\text{K}^+$ -ATPase, the enzyme thought to provide the major driving force for salt uptake, although the underlying cellular components of active ion uptake also include other transport proteins and transport-related enzymes such as a  $\text{Na}^+/\text{H}^+$  antiporter, a  $\text{Na}^+/\text{K}^+/\text{2Cl}^-$  co-transporter, V-ATPases, and carbonic anhydrases

(Bianchini et al. 2008; Lucu and Towle 2003; Pequeux 1995; Henry et al. 2002; Towle et al. 2011, Towle et al. 1997; Ahearn et al. 1999; Serrano and Henry 2008; Towle and Weihrauch 2001). Many terrestrial crabs are capable of directing their urine from their nephropores into their branchial chambers (Morris 2002), and the salt uptake from the urine is also regulated by branchial  $\text{Na}^+/\text{K}^+$ -ATPase (Morris 2001, 2002). Therefore, this branchial  $\text{Na}^+/\text{K}^+$ -ATPase appears to be one of the central players in osmotic and ionic regulation generally in crabs. However, information about the roles of this enzyme at high salinity is limited to some species (the fiddler crab, burrowing crab, and green crab) (Bianchini et al. 2008; Dorazio and Holliday 1985; Lin et al. 2002; Zanders and Rojas 1996; Jillette et al. 2011; Freire et al. 2008). Furthermore,  $\text{Na}^+/\text{Cl}^-$  regulation has been focused on, and little attention has been given to  $\text{Ca}^{2+}$  handling during the adaptation to different salinities (Freire et al. 2008; Charmantier et al. 2009), while  $\text{Ca}^{2+}$  homeostasis in the molting cycle of crustaceans has been examined extensively (Wheatly et al. 2002; Ahearn et al. 2004). For example, only the total calcium concentrations in the hemolymph (complexed (protein bound) plus free (unbound) calcium) have been reported as the  $\text{Ca}^{2+}$  concentration in most of the studies on the responses to different salinities, although it has been known that about 20% of calcium in the hemolymph of intermolt crabs is protein bound unlike other ions (Pequeux 1995; Wheatly 1999; Robertson 1960) and the free moiety which is more relevant can now be measured (Neufeld and Cameron 1992; Wilder et al. 1998).

*G. depressus*, a grapsid crab, is found in subtropical regions and is one of the most abundant northwestern-Pacific species (Kikuchi et al. 1981). This species occurs in intertidal cobble areas and tide pools where the salinities fluctuate considerably due to the effects of rainfall, evaporation, and influx of groundwater (Lohrer et al. 2000; Kawane et al. 2008). Therefore, this amphibious and euryhaline *G. depressus* provides a model for studies on osmotic/ionic responses to various environmental conditions. In this study, we examined changes in the hemolymph osmotic and ionic concentrations and in the activity of branchial  $\text{Na}^+/\text{K}^+$ -ATPase after exposure to high and low salinities as well as after water deprivation in *G. depressus*.

## Methods

### Collection and maintenance of animals

Adult male individuals of grapsid crab, *G. depressus*, weighing 3 to 6 g were collected from June to September along the shore near the Ushimado Marine Institute in Okayama Prefecture, Honshu, Japan. Crabs were transferred to the Marine Institute and held in undiluted natural seawater at a practical salinity of 30 ppt (448 mM  $\text{Na}^+$ , 506 mM  $\text{Cl}^-$ , 9.7 mM  $\text{Ca}^{2+}$ , 9.7 mM  $\text{K}^+$ , 994 mOsm  $\text{kg}^{-1}$ ) and a temperature of  $24 \pm 1^\circ\text{C}$ . A 12:12-h light/dark photoperiod was maintained. Small rocks were placed in each tank to allow animals the opportunity to hide and come out of the water by climbing on them. Animals were acclimated to laboratory conditions for 1 month prior to experimentation and fed a commercial diet ad libitum daily but were not fed for a minimum of 48 h prior to use in the experimentation. Crabs were killed following anesthesia on ice. All procedures were conducted in accordance with the Guidelines for Animal Experimentation established by the Okayama University.

### Experimental design

Each crab in 30-ppt seawater was transferred to an individual 2-L aquarium with 10-ppt salinity (seawater diluted with dechlorinated fresh water), 30-ppt salinity (control), or 50-ppt salinity (seawater supplemented with artificial sea salt, GEX, Osaka, Japan) and submerged in these media. The salinity was checked with a refractometer, and its osmolality later confirmed with a vapor pressure osmometer (Wescor Inc. 5500, Logan, UT, USA). The water in the tanks was replaced daily. The eight gills were dissected out from the crabs acclimated for 10 days for the measurement of  $\text{Na}^+/\text{K}^+$ -ATPase activity. The gills were blotted and placed in ice-cold SEI buffer (250 mM sucrose, 10 mM di-sodium EDTA, 50 mM imidazole, pH 7.3) and frozen immediately at  $-80^\circ\text{C}$ . A separate group of crabs was transferred from 30 ppt to the different salinities or to aquaria without water (terrestrial condition), and subsamples of this group were sampled on day 2 and day 10. Due to a high mortality rate (>50%) 3 days after water deprivation, crabs could only be acclimated to the terrestrial condition for 2 days, and mortality was less than 5% in all groups sampled. Hemolymph (0.2 ml) was withdrawn from the arthrodistal membrane of the walking legs, immediately centrifuged (5 min at  $6,000\times g$ ), and the supernatant was analyzed. The posterior gills (G6 to G8) where  $\text{Na}^+/\text{K}^+$ -ATPase activity was high in 10- to 50-ppt salinity (see Results) were used for  $\text{Na}^+/\text{K}^+$ -ATPase analysis. To assess the effects of osmotic conditions on body mass, the crabs were weighed with a Shimadzu AB54 balance (Shimadzu Corporation, Nakagyo-ku, Kyoto, Japan; accuracy  $\pm 1$  mg) after the visible water had been removed with adsorbent tissue. Only specimens in intermolt stage C were retained for analysis (Drach and Tchernigovtzeff 1967; Fukui 1993; Drach 1939).

### Determination of osmolality and ionic concentrations

The osmolality and  $\text{Cl}^-$  concentration were measured on 5- $\mu\text{l}$  samples (diluted 1:1 with deionized water) using the vapor pressure osmometer and a digital chloride meter (Buchler, Lenexa, KS, USA), respectively. Determinations of  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Ca}^{2+}$  were made on 50- $\mu\text{l}$  samples (diluted 1:3 with deionized water) using ion-specific electrodes of an electrolyte analyzer (AVL 984-S, Graz, Austria). In the case of calcium, total (free plus complexed) calcium was also analyzed since about 20% of calcium in the hemolymph of intermolt crabs is a complexed moiety unlike other ions (Wheatly 1999; Robertson 1960). A 5- $\mu\text{l}$  aliquot of hemolymph was diluted 1:400 in deionized  $\text{H}_2\text{O}$ , and total calcium concentrations were determined by an atomic absorption spectrophotometer (Hitachi Z5300, Tokyo, Japan).

### Assay of $\text{Na}^+/\text{K}^+$ -ATPase enzyme activity

The  $\text{Na}^+/\text{K}^+$ -ATPase activity was determined with a linked pyruvate kinase/lactate dehydrogenase-NADH assay (McCormick 1993). Gill tissue was homogenized in ice-cold 0.1%-deoxycholate SEI buffer (1:9 *w/v*) and centrifuged at  $5,000\times g$ . The resulting supernatant was diluted and assayed for  $\text{Na}^+/\text{K}^+$ -ATPase activity. Each sample of gill homogenate was plated in quadruplicates of 10  $\mu\text{l}$ , two contained 2.8 mM ouabain and two did not. Fifty microliters of salt solution (50 mM imidazole, 189 mM NaCl, 10.5 mM  $\text{MgCl}_2$ , and 42 mM KCl) and 150  $\mu\text{l}$  of assay mixture (50 mM imidazole, 2 mM phosphoenolpyruvate, 0.16 mM nicotinamide adenine dinucleotide, 0.5 mM adenosine

triphosphate, 3.3 U/ml lactic dehydrogenase, and 3.6 U/ml pyruvate kinase) were added to each well. The kinetic assay was read at a wavelength of 340 nm at 24°C with a run time of 10 min and intervals of 10 s. The difference between the kinetic reading with and without ouabain is the Na<sup>+</sup>/K<sup>+</sup>-ATPase activity and is expressed as micromoles ADP per milligram protein per hour. Total protein in homogenates was measured using a BCA Protein Assay kit (Pierce Chemical Co., Rockford, IL, USA). Assays were run on a microplate reader (Multiskan Ascent, Thermo Electron Corporation, Vaanta, Finland).

For validation of this system, standard conditions described above were employed, varying one factor while keeping all the other parameters constant. Inhibition by ouabain corresponding to the actual measurement of Na<sup>+</sup>/K<sup>+</sup>-ATPase activity, as a function of ouabain concentration in the reaction mixture, was first examined. Under the standard conditions, final concentrations of ouabain in the reaction mixture varied from 0, 0.5, 1.4, 2.8, and 5.0 mM in wells, and maximal inhibition was observed at 2.8 mM ouabain, and thus this concentration was fixed in the examination of other parameters in the remainder of the validation. Optimal conditions for actual analyses of response to changing environmental condition were set according to such results. In examinations of the effects of gill protein concentration on the enzymatic activity, it was seen that in a sample consisting of 0.1 mg protein/10 µl of 0.1%-deoxycholate SEI buffer diluted from 2- to 16-fold, activity decreased linearly in proportion to the protein quantity. Therefore, measurements were valid for samples diluted at least twofold, but samples were usually diluted tenfold in this investigation.

### Statistical analyses

Statistics were performed using Statview 4.11 (Abacus Concept). Since there was a significant interaction between the treatment (environmental condition) and time by two-way ANOVA, data for day 2 and day 10 were analyzed separately by the appropriate *post hoc* test to determine the differences between the control (30 ppt) and treatments (different environmental conditions). All data were checked for normality and equal variances. Where assumptions of normality or equal variances were not satisfied, equivalent nonparametric tests were used.

## Results

### Body mass

Exposure of *G. depressus* to salinities of 10, 30, or 50 ppt did not result in any significant change in body mass ( $P > 0.05$ ), while the loss of 10% in the crabs after exposure to terrestrial conditions was significant ( $P < 0.001$ , Figure 1a).

### Hemolymph osmotic and ionic status

In crabs exposed to 10-ppt salinity (331 mOsm), hemolymph osmolality decreased after 2 days to its new acclimation level at 580 mOsm ( $P < 0.001$ , Figure 1b) and then did not change significantly ( $P > 0.05$ ) thereafter. After exposure to 50 ppt (1,660 mOsm) and terrestrial conditions, hemolymph osmolality increased ( $P < 0.001$ ), although the




Figure 1 (See legend on next page.)

(See figure on previous page.)

**Figure 1 The wet body weight (a) as well as hemolymph osmotic and ionic concentrations (b to f) of *G. depressus*.** Change in the wet body weight (%) as well as osmotic and ionic concentrations in the hemolymph of *G. depressus* under various environmental conditions for 2 and 10 days after being transferred from 30-ppt seawater. Dotted line indicates osmotic and ionic concentrations of the media. Mean  $\pm$  SE ( $N = 4$  to 10) is indicated. Means with asterisks are significantly different from the means of the control (30 ppt) on the same day (asterisk indicates  $P < 0.05$ ; double asterisk,  $P < 0.01$ ; triple asterisk,  $P < 0.001$ ).

levels in 50-ppt seawater also stabilized by day 10 to new acclimated values which were about 350 mOsm hypoosmotic to the medium. The osmoregulatory performance of crabs acclimated for 10 days in various salinity media is shown in Figure 2a.


At 30 ppt, hemolymph was isoionic to the medium in the case of the three ions,  $\text{Na}^+$ ,  $\text{Cl}^-$ , and  $\text{Ca}^{2+}$  (Figure 1c,d,e). A virtually identical pattern was seen for the major hemolymph ions,  $\text{Na}^+$  and  $\text{Cl}^-$ . The concentrations of these ions decreased and increased 2 days after being transferred to 10 and 50 ppt, respectively ( $P < 0.001$ ). Thereafter, they appeared to stabilize at new acclimated values, although the  $\text{Na}^+$  levels after transfer to 50 ppt decreased on day 10 compared to those on day 2 (Figure 1c,d). Changes in hemolymph  $\text{K}^+$  concentrations also paralleled to those of osmolality,  $\text{Na}^+$  and  $\text{Cl}^-$  (data not shown).

The changes in  $\text{K}^+$ ,  $\text{Na}^+$ , and  $\text{Cl}^-$  showed a relationship with changes in hemolymph osmolality, but free  $\text{Ca}^{2+}$  showed different patterns after exposure to 50 ppt and terrestrial conditions (Figure 1e). After being transferred to 50 ppt, both the free and total calcium increased significantly ( $P < 0.001$ ) on day 2, but returned to the levels similar to those at 30 ppt on day 10. In crabs kept out of the water, free  $\text{Ca}^{2+}$  decreased slightly but significantly ( $P < 0.05$ ). Furthermore, the complexed calcium (total minus free calcium) virtually disappeared from the hemolymph on day 10 in 10 and 50 ppt, indicating that all the hemolymph calcium was ionized, and the concentrations appeared to be maintained at acclimated values. The calcium regulatory performance of crabs acclimated for 10 days in various salinity media is shown in Figure 2b.

#### Branchial $\text{Na}^+/\text{K}^+$ -ATPase activity


Branchial  $\text{Na}^+/\text{K}^+$ -ATPase activity was heterogeneously distributed among the eight gill pairs in crabs acclimated to 30, 10, or 50 ppt for 10 days (Figure 3a). Activity was high in the posterior three gills, G6 to G8. This distribution and the absolute values of the activity reported here were similar to those reported in the literature (see the 'Discussion' section).

The responses of  $\text{Na}^+/\text{K}^+$ -ATPase activity after exposure to various environmental conditions were examined in the posterior gills (G6 to G8) since the above results (Figure 3a) demonstrated the high levels of  $\text{Na}^+/\text{K}^+$ -ATPase activity in these gills after acclimation to 30, 10, or 50 ppt. The  $\text{Na}^+/\text{K}^+$ -ATPase activity in G6 and G7 increased significantly during exposure to 10 ppt by day 10 ( $P < 0.05$ ). The  $\text{Na}^+/\text{K}^+$ -ATPase activity in G8 did not change in 10 ppt ( $P > 0.05$ ) but increased significantly 10 days after transfer to 50 ppt ( $P < 0.05$ ). In G6 to G8 of the crabs kept out of water, there was no significant difference in the  $\text{Na}^+/\text{K}^+$ -ATPase activity ( $P > 0.05$ ), although a threefold increase in the mean activity was observed in G6 (Figure 3b).



## Discussion

The results of the transfer to 10- and 50-ppt salinities indicate that *G. depressus* is able to osmoregulate and survive at least for 10 days in these salinities as a hyper-hypoosmoregulating marine crab, and are consistent with the previous reports (Charmantier et al. 1998, 2002) on hemolymph osmolality in the other grapsid crabs (*Armases miersii* and *Chasmagnathus granulata*) as a function of the ambient salinity. Together with the results of the terrestrial exposure showing that hemolymph osmolality was in the same physiological range after 2 days despite dehydration, *G. depressus* can be considered as an appropriate model to study the osmotic/ionic regulatory mechanisms supporting salinity/terrestrial acclimation. In this investigation, it was considered necessary to obtain more basic information concerning the ionic status in



**Figure 3 Na<sup>+</sup>/K<sup>+</sup>-ATPase activity in the gills of *G. depressus*.** Mean  $\pm$  SE ( $N = 4$  to  $9$ ) is indicated. (a) The activity in the gills of crabs acclimated for 10 days in 10-, 30-, or 50-ppt salinity after being transferred from the 30-ppt seawater. (b) The activity in the posterior gills (G6 to G8) of crabs under various environmental conditions on day 2 and day 10 after being transferred from 30 ppt; means with an asterisk are significantly different from the mean of the control (30 ppt) on the same day ( $P < 0.05$ ).



the hemolymph as well as the branchial  $\text{Na}^+/\text{K}^+$ -ATPase activity as a prerequisite for further studies.

Regarding ionic changes in the hemolymph, changes in  $\text{Na}^+$  and  $\text{Cl}^-$  generally paralleled those in osmotic concentrations in response to high and low salinity exposure as well as to water deprivation, and it is likely that altered osmolality of the hemolymph under varying environmental conditions was based for the most part on altered levels of  $\text{Na}^+$  and  $\text{Cl}^-$  (Wilder et al. 1998). An interesting finding of this study, however, is the pattern of hemolymph calcium levels, particularly for those of complexed calcium. Slightly lower levels of hemolymph free  $\text{Ca}^{2+}$  in crabs kept out of water seems to be related to the fasted condition since intermolt terrestrial crabs regulate hemolymph calcium by controlling intake of dietary calcium (Wheatly 1999; Zanotto and Wheatly 2002). In crabs acclimated to 10-ppt salinity, free  $\text{Ca}^{2+}$  was maintained at twofold higher values than concentrations of the medium, and complexed calcium disappeared from the hemolymph after 10 days. In crabs exposed to 50-ppt salinity, free  $\text{Ca}^{2+}$  was regulated to the lower levels than those in the medium through the experiment while total calcium increased to higher levels after 2 days. These responses may indicate that hemolymph complexed calcium could serve as an internal reserve for maintaining free  $\text{Ca}^{2+}$  levels in the hemolymph. On the other hand, the complexed calcium decreased dramatically and disappeared from the hemolymph after 10 days in 50 ppt, and we speculate that this represents a surplus (unnecessary) calcium reservoir in the hemolymph for prolonged period in higher  $\text{Ca}^{2+}$  environment. Calcium regulation in various environments has been studied in crustaceans, mostly with respect to the control of epithelial calcium transport (Freire et al. 2008; Ahearn et al. 2004; Wheatly 1999; Zanotto and Wheatly 2002), and the role of complexed calcium in the hemolymph as a reserve for free  $\text{Ca}^{2+}$  is unknown. Our findings, therefore, suggest a new control mechanism of hemolymph free  $\text{Ca}^{2+}$  and imply that hemolymph concentrations of both total and free calcium need to be analyzed. At any rate, it appears that it is necessary to regulate free  $\text{Ca}^{2+}$  to a specific range and that this control is separate from the osmoregulatory mechanisms.

One of the ion transporters that has received the most intensive studies in osmoregulating crustaceans is  $\text{Na}^+/\text{K}^+$ -ATPase (Bianchini et al. 2008; Lucu and Towle 2003; Pequeux 1995; Henry et al. 2002; Towle et al. 1997, 2011; Ahearn et al. 1999; Serrano and Henry 2008; Towle and Weihrauch 2001). In addition to the  $\text{Ca}^{2+}$  channel,  $\text{Ca}^{2+}$ -ATPase, and  $\text{Na}^+/\text{Ca}^{2+}$  exchanger, the transportation of  $\text{Ca}^{2+}$  in the hemolymph of crustaceans is also affected by the potential energy of the  $\text{Na}^+$  gradient, established by  $\text{Na}^+/\text{K}^+$  ATPase activity (Roer and Dillaman 1993). In this study, we observed higher  $\text{Na}^+/\text{K}^+$ -ATPase activity in the posterior gills when compared with the anterior gills (see Figure 3a), consistent with the molecular biological and physiological studies on many crab species (Bianchini et al. 2008; Lucu and Towle 2003; Pequeux 1995; Towle and Weihrauch 2001; Freire et al. 2008; Charmantier et al. 2009; Siebers et al. 1982; Onken and Putzenlechner 1995), which have designated the posterior gill epithelium, with its high abundance of  $\text{Na}^+/\text{K}^+$ -ATPase activity, as the principal site of osmoregulatory ion transport. These differences constitute the basis of the paradigm that anterior gills are structurally and functionally specialized for respiratory gas exchange, while the posterior gills have become specialized for active ion absorption counterbalancing passive losses in dilute media (Freire et al. 2008; Charmantier et al.

2009) as reflected in the significant increases in the  $\text{Na}^+/\text{K}^+$ -ATPase activity of G6 and G7 after acclimation to 10-ppt salinity (Figure 3b).

Following these observations, the present investigation provides an interesting finding that acclimation to 50-ppt salinity for 10 days, in which hypo-ionoregulation occurred, was accompanied by increased  $\text{Na}^+/\text{K}^+$ -ATPase activity exclusively in G8 (Figure 3b), suggesting that G8 may participate in ion excretion into the concentrated media. These different responses of the enzyme activity among the individual gills indicate a gill-specific pattern of the regulation and a higher degree of specialization in gill function in *G. depressus*. Together with the differences between the terrestrial condition and 50-ppt salinity, the increased  $\text{Na}^+/\text{K}^+$ -ATPase activity is not simply part of a cellular regulation since the cells were exposed to the similar osmo/ionic stresses. A study with the marble shore crab (*Pachygrapsus marmoratus*) also showed that the abundance of  $\text{Na}^+/\text{K}^+$ -ATPase mRNA induced in all nine gills following the transfer of crabs to low salinity but increased only in G7 after being transferred to high salinity (Jayasundara et al. 2007). Our enzyme activity results for *G. depressus* support the notion that individual gills do indeed play distinct osmoregulatory roles in euryhaline crustaceans. Other transport proteins and transport-related enzymes in gills, including a  $\text{Na}^+/\text{H}^+$  antiporter, carbonic anhydrase, and  $\text{Na}^+/\text{K}^+/\text{2Cl}^-$  cotransporters (Bianchini et al. 2008; Pequeux 1995; Henry et al. 2002; Towle et al. 1997, 2011; Serrano and Henry 2008; Jillette et al. 2011; Ahearn et al. 2004; Wheatly 1999), might be involved in the specificity of function. For example, a basolateral  $\text{Na}^+/\text{K}^+/\text{2Cl}^-$  cotransporter involved in NaCl excretion appears to be induced during acclimation to concentrated seawater (Freire et al. 2008; Luquet et al. 2005). To further develop *G. depressus* as a new model for the study of salinity acclimation in crabs, future investigations will examine the role of these transporters and possible ionocytes in the acclimation responses of this euryhaline species.

## Conclusions

The hemolymph osmotic and ionic status of *G. depressus* indicates that this intertidal grapsid crab is a hyper/hypo-ionoregulating amphibious species. Especially, the free  $\text{Ca}^{2+}$  concentration was well-maintained partly by the hemolymph complexed calcium as an internal reserve. Induction of  $\text{Na}^+/\text{K}^+$ -ATPase activity in response to salinities varies between the gills. This abundant species around Japan will serve as a model to study the crustacean osmotic/ionic regulation.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

TS conceived of the study, participated in its design and coordination, and drafted the manuscript. TN participated in the design of the study and performed the experiments and analyses. HT and TA participated in the design of the study, helped perform the experiments, and drafted the manuscript. WG helped perform the experiments. MO, HS, and NT helped draft the manuscript. All authors read and approved the final manuscript.

## Acknowledgments

We thank S. Hasegawa, S. Ogawa, T. Akada, and Y. Nishiyama for their help in this research. We also thank Dr. J. P. Breves for the critical reading of the manuscript. This research was funded in part by grants to Tatsuya Sakamoto (Grants-in-Aid for Scientific Research (C) nos. 17570049, 19570057, and 22570065 from Japan Society for the Promotion of Science) and was carried out in part as a joint research in Japanese Association for Marine Biology (JAMBIO).

#### Author details

<sup>1</sup>Ushimado Marine Institute, Faculty of Science, Okayama University, Ushimado, Setouchi 701-4303, Japan. <sup>2</sup>Department of Environmental Science, Faculty of Science, Niigata University, Ikarashi Niigata 950-2181, Japan. <sup>3</sup>Lago Co., Ltd., Taga, Oumihachiman, Shiga 523-0821, Japan.

Received: 27 August 2012 Accepted: 26 October 2012

Published: 9 November 2012

#### References

- Ahearn GA, Duerr JM, Zhuang Z, Brown RJ, Aslamkhan A, Killebrew DA (1999) Ion transport processes of crustacean epithelial cells. *Physiol Biochem Zool* 72(1):1–18
- Ahearn GA, Mandal PK, Mandal A (2004) Calcium regulation in crustaceans during the molt cycle: a review and update. *Comp Biochem Physiol A Mol Integr Physiol* 137(2):247–257
- Bianchini A, Lauer MM, Nery LE, Colares EP, Monserrat JM, Dos Santos Filho EA (2008) Biochemical and physiological adaptations in the estuarine crab *Neohelice granulata* during salinity acclimation. *Comp Biochem Physiol A Mol Integr Physiol* 151(3):423–436
- Charmantier G, Charmantier-Daures M, Anger K (1998) Ontogeny of osmoregulation in the grapsid crab *Armases miersii* (Crustacea, Decapoda). *Mar Ecol Prog Ser* 164:285–292
- Charmantier G, Gimenez L, Charmantier-Daures M, Anger K (2002) Ontogeny of osmoregulation, physiological plasticity and larval export strategy in the grapsid crab *Chasmagnathus granulata* (Crustacea, Decapoda). *Mar Ecol Prog Ser* 229:185–194
- Charmantier G, Charmantier-Daures M, Towle D (2009) Osmotic and ionic regulation in aquatic arthropods. In: Evans DH (ed) *Osmotic and ionic regulation: cells and animals*. CRC Press, New York, p 165
- Dorazio SE, Holliday CW (1985) Gill Na, K-ATPase and osmoregulation in the sand fiddler crab, *Uca pugilator*. *Physiol Zool* 58(4):364–373
- Drach P (1939) Mue et cycle d'intermue chez les Crustacés Décapodes. *Annales de l'Institut Oceanographique* 19(1):103–388
- Drach P, Tchernigovtzeff C (1967) Sur la méthode de détermination des stades d'intermue et son application générale aux Crustacés. *Vie et Milieu* 18559(3):595–609
- Freire CA, Onken H, McNamara JC (2008) A structure-function analysis of ion transport in crustacean gills and excretory organs. *Comp Biochem Physiol A Mol Integr Physiol* 151(3):272–304
- Fukui Y (1993) Timing of copulation in the molting and reproductive cycles in a grapsid crab, *Gaetice depressus* (Crustacea: Brachyura). *Mar Biol* 117(2):221–226
- Henry RP, Garrelts EE, McCarty MM, Towle DW (2002) Differential induction of branchial carbonic anhydrase and Na<sup>+</sup>/K<sup>+</sup> ATPase activity in the euryhaline crab, *Carcinus maenas*, in response to low salinity exposure. *J Exp Zool* 292(7):595–603
- Jayasundara N, Towle DW, Weihrauch D, Spanings-Pierrot C (2007) Gill-specific transcriptional regulation of Na<sup>+</sup>/K<sup>+</sup>–ATPase alpha-subunit in the euryhaline shore crab *Pachygrapsus marmoratus*: sequence variants and promoter structure. *J Exp Biol* 210(Pt 12):2070–2081
- Jillette N, Cammack L, Lowenstein M, Henry RP (2011) Down-regulation of activity and expression of three transport-related proteins in the gills of the euryhaline green crab, *Carcinus maenas*, in response to high salinity acclimation. *Comp Biochem Physiol A Mol Integr Physiol* 158(2):189–193
- Kawane M, Wada K, Watanabe K (2008) Comparisons of genetic population structures in four intertidal brachyuran species of contrasting habitat characteristics. *Mar Biol* 156(2):193–203
- Kikuchi T, Tanaka M, Nojima S, Takahashi T (1981) Ecological studies on the pebble crab, *Gaetice depressus* (de Haan). I. Ecological distribution of the crab and environmental conditions. Publication from the Amakusa Marine Biological Laboratory 61:23–34
- Lin HC, Su YC, Su SH (2002) A comparative study of osmoregulation in four fiddler crabs (Ocypodidae: Uca). *Zool Sci* 19(6):643–650
- Lohrer A, Fukui Y, Wada K, Whitlatch R (2000) Structural complexity and vertical zonation of intertidal crabs, with focus on habitat requirements of the invasive Asian shore crab, *Hemigrapsus sanguineus* (de Haan). *J Exp Mar Biol Ecol* 244(2):203–217
- Lucu C, Towle DW (2003) Na<sup>+</sup>+K<sup>+</sup>-ATPase in gills of aquatic crustacea. *Comp Biochem Physiol A Mol Integr Physiol* 135(2):195–214
- Luquet CM, Weihrauch D, Senek M, Towle DW (2005) Induction of branchial ion transporter mRNA expression during acclimation to salinity change in the euryhaline crab *Chasmagnathus granulatus*. *J Exp Biol* 208(Pt 19):3627–3636
- McCormick SD (1993) Methods for nonlethal gill biopsy and measurement of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity. *Can J Fish Aquat Sci* 50(3):656–658
- Morris S (2001) Neuroendocrine regulation of osmoregulation and the evolution of air-breathing in decapod crustaceans. *J Exp Biol* 204(Pt 5):979–989
- Morris S (2002) The ecophysiology of air-breathing in crabs with special reference to *Gecarcoidea natalis*. *Comp Biochem Physiol B Biochem Mol Biol* 131(4):559–570
- Neufeld DS, Cameron JN (1992) Postmoult uptake of calcium by the blue crab (*Callinectes sapidus*) in water of low Salinity. *J Exp Biol* 171:283–299
- Onken H, Putzenlechner M (1995) A V-ATPase drives active, electrogenic and Na<sup>+</sup>-independent Cl<sup>-</sup> absorption across the gills of *Eriocheir sinensis*. *J Exp Biol* 198(Pt 3):767–774
- Pequeux A (1995) Osmotic regulation in crustaceans. *J Crustac Biol* 15(1):1–60
- Robertson JD (1960) Ionic regulation in the crab *Carcinus maenas* (L.) in relation to the moulting cycle. *Comp Biochem Physiol* 1(3):183–212
- Roer RD, Dillaman RM (1993) Molt-related change in integumental structure and function. In: Horst MN, Freeman JA (ed) *The crustacean integument morphology and biochemistry*. CRC Press, Boca Raton, pp 1–37

- Serrano L, Henry RP (2008) Differential expression and induction of two carbonic anhydrase isoforms in the gills of the euryhaline green crab, *Carcinus maenas*, in response to low salinity. *Comp Biochem Physiol Part D Genomics Proteomics* 3(2):186–193
- Siebers D, Leweck K, Markus H, Winkler A (1982) Sodium regulation in the shore crab *Carcinus maenas* as related to ambient salinity. *Mar Biol* 69(1):37–43
- Towle DW, Weihrauch D (2001) Osmoregulation by gills of euryhaline crabs: molecular analysis of transporters. *Am Zool* 41(4):770–780
- Towle DW, Rushton ME, Heidysch D, Magnani JJ, Rose MJ, Amstutz A, Jordan MK, Shearer DW, Wu WS (1997) Sodium/proton antiporter in the euryhaline crab *Carcinus maenas*: molecular cloning, expression and tissue distribution. *J Exp Biol* 200(Pt 6):1003–1014
- Towle DW, Henry RP, Terwilliger NB (2011) Microarray-detected changes in gene expression in gills of green crabs (*Carcinus maenas*) upon dilution of environmental salinity. *Comp Biochem Physiol Part D Genomics Proteomics* 6(2):115–125
- Wheatly MG (1999) Calcium homeostasis in crustacea: the evolving role of branchial, renal, digestive and hypodermal epithelia. *J Exp Zool* 283(7):620–640
- Wheatly MG, Zanutto FP, Hubbard MG (2002) Calcium homeostasis in crustaceans: subcellular Ca dynamics. *Comp Biochem Physiol B Biochem Mol Biol* 132(1):163–178
- Wilder MN, Ikuta K, Atmomarsono M, Hatta T, Komuro K (1998) Changes in osmotic and ionic concentrations in the hemolymph of *Macrobrachium rosenbergii* exposed to varying salinities and correlation to ionic and crystalline composition of the cuticle. *Comp Biochem Phys A* 119(4):941–950
- Zanders IP, Rojas WE (1996) Transbranchial potentials and ion fluxes across isolated, perfused gills of *Uca rapax*. *Mar Biol* 125(2):307–314
- Zanutto FP, Wheatly MG (2002) Calcium balance in crustaceans: nutritional aspects of physiological regulation. *Comp Biochem Physiol A Mol Integr Physiol* 133(3):645–660

doi:10.1186/2008-6970-4-18

**Cite this article as:** Nanba et al.: Hemolymph osmotic, ionic status, and branchial Na<sup>+</sup>/K<sup>+</sup>-ATPase activity under varying environmental conditions in the intertidal grapsid crab, *Gaetice depressus*. *International Aquatic Research* 2012 4:18.

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

---