

Research Article

Sharpening the Becker-Stark Inequalities

Ling Zhu and Jiukun Hua

Department of Mathematics, Zhejiang Gongshang University, Hangzhou, Zhejiang 310018, China

Correspondence should be addressed to Ling Zhu, zhuling0571@163.com

Received 3 April 2009; Accepted 14 January 2010

Academic Editor: Sever Silvestru Dragomir

Copyright © 2010 L. Zhu and J. Hua. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In this paper, we establish a general refinement of the Becker-Stark inequalities by using the power series expansion of the tangent function via Bernoulli numbers and the property of a function involving Riemann's zeta one.

1. Introduction

Steckin [1] (or see Mitrinovic [2, 3.4.19, page 246]) gives us a result as follows.

Theorem 1.1 (see [1, Lemma 2.1]). *If $0 < x < \pi/2$, then*

$$\frac{4}{\pi} \frac{x}{\pi - 2x} < \tan x. \quad (1.1)$$

Later, Becker and Stark [3] (or see Kuang [4, 5.1.102, page 248]) obtain the following two-sided rational approximation for $(\tan x)/x$.

Theorem 1.2. *Let $0 < x < \pi/2$, then*

$$\frac{8}{\pi^2 - 4x^2} < \frac{\tan x}{x} < \frac{\pi^2}{\pi^2 - 4x^2}. \quad (1.2)$$

Furthermore, 8 and π^2 are the best constants in (1.2).

In fact, we can obtain the following further results.

Theorem 1.3. Let $0 < x < \pi/2$, then

$$\frac{\pi^2 + ((4(8 - \pi^2))/\pi^2)x^2}{\pi^2 - 4x^2} < \frac{\tan x}{x} < \frac{\pi^2 + (\pi^2/3 - 4)x^2}{\pi^2 - 4x^2}. \quad (1.3)$$

Furthermore, $\alpha = (4(8 - \pi^2))/\pi^2$ and $\beta = \pi^2/3 - 4$ are the best constants in (1.3).

In this paper, in the form of (1.2) and (1.3) we shall show a general refinement of the Becker-Stark inequalities as follows.

Theorem 1.4. Let $0 < x < \pi/2$, and let $N \geq 0$ be a natural number. Then

$$\frac{P_{2N}(x) + \alpha x^{2N+2}}{\pi^2 - 4x^2} < \frac{\tan x}{x} < \frac{P_{2N}(x) + \beta x^{2N+2}}{\pi^2 - 4x^2} \quad (1.4)$$

holds, where $P_{2N}(x) = a_0 + a_1x^2 + \dots + a_Nx^{2N}$, and

$$a_n = \frac{2^{2n+2}(2^{2n+2} - 1)\pi^2}{(2n + 2)!} |B_{2n+2}| - \frac{4 \cdot 2^{2n}(2^{2n} - 1)}{(2n)!} |B_{2n}|, \quad n = 0, 1, 2, \dots, \quad (1.5)$$

where B_{2n} are the even-indexed Bernoulli numbers.

Furthermore, $\alpha = (8 - a_0 - a_1(\pi/2)^2 - \dots - a_N(\pi/2)^{2N})/(\pi/2)^{2N+2}$ and $\beta = a_{N+1}$ are the best constants in (1.4).

2. Four Lemmas

Lemma 2.1. The function $(1 - 1/2^n)\zeta(n)$ ($n = 1, 2, \dots$) is decreasing, where $\zeta(n)$ is Riemann's zeta function.

Proof. $(1 - 1/2^n)\zeta(n) = \zeta(n) - \zeta(n)/2^n$ is equivalent to the function $\lambda(n) = \sum_{k=0}^{\infty} 1/(2k + 1)^n$, which is decreasing. \square

Lemma 2.2 (see [5, Theorem 3.4]). Let $\zeta(n)$ be Riemann's zeta function and B_{2n} the even-indexed Bernoulli numbers. Then

$$\zeta(2n) = \frac{(2\pi)^{2n}}{2(2n)!} |B_{2n}|, \quad n = 1, 2, \dots \quad (2.1)$$

Lemma 2.3 (see [6, 1.3.1.4 (1.3)]). Let $|x| < \pi/2$. Then

$$\tan x = \sum_{n=1}^{\infty} \frac{2^{2n}(2^{2n} - 1)}{(2n)!} (-1)^{n-1} B_{2n} x^{2n-1} = \sum_{n=1}^{\infty} \frac{2^{2n}(2^{2n} - 1)}{(2n)!} |B_{2n}| x^{2n-1}. \quad (2.2)$$

Lemma 2.4. Let $F(x) = (\pi^2 - 4x^2)(\tan x/x)$ and $|x| < \pi/2$. Then $F(x) = \pi^2 + \sum_{n=1}^{+\infty} a_n x^{2n}$, where

$$a_n = \frac{2^{2n+2}(2^{2n+2} - 1)\pi^2}{(2n + 2)!} |B_{2n+2}| - \frac{4 \cdot 2^{2n}(2^{2n} - 1)}{(2n)!} |B_{2n}| < 0, \quad n = 1, 2, \dots \tag{2.3}$$

Proof. By Lemma 2.3, we have

$$\begin{aligned} F(x) &= (\pi^2 - 4x^2) \sum_{n=1}^{\infty} \frac{2^{2n}(2^{2n} - 1)}{(2n)!} |B_{2n}| x^{2n-2} \\ &= \pi^2 + \sum_{n=1}^{+\infty} \left[\frac{2^{2n+2}(2^{2n+2} - 1)\pi^2}{(2n + 2)!} |B_{2n+2}| - \frac{4 \cdot 2^{2n}(2^{2n} - 1)}{(2n)!} |B_{2n}| \right] x^{2n} \\ &:= \pi^2 + \sum_{n=1}^{+\infty} a_n x^{2n}. \end{aligned} \tag{2.4}$$

Since $(1 - (1/2^{2n}))\zeta(2n)$ is decreasing by Lemma 2.1, it follows that

$$\frac{2^{2n+2} - 1}{4} \zeta(2n + 2) < (2^{2n} - 1) \zeta(2n). \tag{2.5}$$

From Lemma 2.2, we get

$$\frac{\pi^2(2^{2n+2} - 1)}{(2n + 2)!} |B_{2n+2}| < \frac{(2^{2n} - 1)}{(2n)!} |B_{2n}|, \tag{2.6}$$

which implies that $a_n < 0$ for $n = 1, 2, \dots$ □

3. Proofs of Theorems

Proof of Theorem 1.4. Let

$$G(x) = \frac{((\tan x)/x)(\pi^2 - 4x^2) - (a_0 + a_1x^2 + \dots + a_Nx^{2N})}{x^{2N+2}}. \tag{3.1}$$

Then

$$G(x) = \frac{F(x) - (a_0 + a_1x^2 + \dots + a_Nx^{2N})}{x^{2N+2}} = \frac{\sum_{n=N+1}^{+\infty} a_n x^{2n}}{x^{2N+2}} = \sum_{k=0}^{+\infty} a_{N+1+k} x^{2k}. \tag{3.2}$$

By Lemma 2.4, we have $a_n < 0$ for $n \in \mathbb{N}^+$, and $G(x)$ is decreasing on $(0, \pi/2)$.

At the same time, $\alpha = \lim_{x \rightarrow (\pi/2)^-} G(x) = (8 - a_0 - a_1(\pi/2)^2 - \dots - a_N(\pi/2)^{2N})/(\pi/2)^{2N+2}$ by (3.1), and $\beta = \lim_{x \rightarrow 0^+} G(x) = a_{N+1}$ by (3.2), so α and β are the best constants in (1.4). □

Proof of Theorem 1.3. Let $N = 0$ in Theorem 1.4; we obtain that $\alpha = (4(8 - \pi^2))/\pi^2$ and $\beta = \pi^2/3 - 4$. Then the proof of Theorem 1.3 is complete. \square

References

- [1] S. B. Steckin, "Some remarks on trigonometric polynomials," *Uspekhi Matematicheskikh Nauk*, vol. 10, no. 1(63), pp. 159–166, 1955 (Russian).
- [2] D. S. Mitrinovic, *Analytic Inequalities*, Springer, New York, NY, USA, 1970.
- [3] M. Becker and E. L. Strak, "On a hierarchy of quolynomial inequalities for $\tan x$," *University of Beograd Publikacije Elektrotehnicki Fakultet. Serija Matematika i fizika*, no. 602–633, pp. 133–138, 1978.
- [4] J. C. Kuang, *Applied Inequalities*, Shangdong Science and Technology Press, Jinan City, China, 3rd edition, 2004.
- [5] W. Scharlau and H. Opolka, *From Fermat to Minkowski*, Undergraduate Texts in Mathematics, Springer, New York, NY, USA, 1985.
- [6] A. Jeffrey, *Handbook of Mathematical Formulas and Integrals*, Elsevier Academic Press, San Diego, Calif, USA, 3rd edition, 2004.