

RESEARCH ARTICLE

Open Access



Effects of treatment with *Astragalus Membranaceus* on function of rat leydig cells

Xiaolong Jiang^{1†}, Xia Cao^{1†}, Yang Huang¹, Jianwei Chen¹, Xiaolei Yao¹, Miaomiao Zhao¹, Yan Liu¹, Jinzhu Meng¹, Pengfei Li², Zhiyan Li³, Jianbo Yao^{1,7}, George W Smith^{1,4,5,6} and Lihua Lv^{1*}

Abstract

Background: *Astragalus membranaceus* (AM) is a Chinese traditional herb which has been reported to have broad positive effects on many diseases, including hepatitis, heart disease, diabetes and skin disease. AM can promote cell proliferation, increase the activities of superoxide dismutase (SOD) and glutathione peroxidase (GPx), and inhibit apoptosis by regulating the transcription of proto-oncogenes controlling cell death. While AM is included in some commercially available “testosterone boosting supplements”, studies directly testing ability of AM to modulate testosterone production are lacking. In the present study, we examined the effects of AM on Leydig cell function *in vitro*.

Methods: Rat Leydig cells were purified and treated with AM at different concentrations (0 µg/mL, 10 µg/mL, 20 µg/mL, 50 µg/mL, 100 µg/mL and 150 µg/mL) and cell counting-8 (CCK-8) assay, Enzyme-linked immunosorbent assay, quantitative real time PCR and analysis of activities of SOD and GPx were done respectively.

Results: Treatment with 100 µg/mL ($P < 0.05$) and 150 µg/mL AM ($P < 0.01$) significantly increased Leydig cell numbers. Treatment with AM (20 µg/mL, 50 µg/mL and 100 µg/mL) significantly increased testosterone production ($P < 0.01$). In addition, increased Leydig cell SOD and GPx activities were observed in response to 20 µg/mL and 50 µg/mL AM treatment ($P < 0.01$). Furthermore, expression of *Bax* mRNA was significantly decreased ($P < 0.01$), and the ratio of *Bcl-2/Bax* mRNA was significantly increased in response to 20 µg/mL AM in the culture medium ($P < 0.05$).

Conclusions: Results supported a beneficial effect of AM on multiple aspects of rat Leydig cell function *in vitro* including testosterone production.

Keywords: *Astragalus membranaceus*, Leydig cell, SOD and GPx, *Bax*, *Bcl-2*, Testosterone

Background

Astragalus membranaceus (AM), a well-known Chinese medicinal plant, has been widely used as a traditional prescription medicine for treatment of various diseases, including hepatitis [1], diabetic nephropathy [2, 3], cardiovascular disorders [4, 5] and skin diseases [6]. AM has various favorable pharmacological effects including anti-fibrotic [1, 7–9], anti-oxidant [10], anti-apoptotic [11], anti-inflammatory [12] and immune-enhancing properties [13]. AM inhibits apoptosis mainly by regulating the expression of *Bax* and *Bcl-2*, members of the B cell leukemia protein (Bcl)-2 family that controls the intrinsic apoptotic

pathway [14]. A previous study has shown that glucose-induced podocyte apoptosis was inhibited by intravenous AM injection *in vivo* and AM treatment *in vitro* with effects linked to down regulation of *Bax* expression and up-regulation of *Bcl-2* expression [15]. Collectively, these observations indicate that AM has therapeutic effects in the context of various types of disease.

Although numerous studies support potential medicinal value of AM, its effects on the reproductive system have not been well investigated. Limited studies have demonstrated effects of AM on the motility of human spermatozoa *in vitro* [16] and amelioration of reproductive toxicity induced by cyclophosphamide in male mice [17] and sperm abnormalities in cadmium-treated rats [18]. Hong *et al.* tested effects of 18 types of Chinese herbs on semen parameters and only AM aqueous

* Correspondence: lihualvxau@126.com

†Equal contributors

¹College of Animal Science and Technology, Shanxi Agricultural University, No.1 Mingxian Nan Road, Taigu 030801, China

Full list of author information is available at the end of the article

extract showed a significant stimulatory effect on the motility of human spermatozoa [16].

Leydig cells are distributed in the loose connective tissue between the seminiferous tubules, accounting for 2 %-4 % of testicular cells. Their main physiological function is to produce testosterone. Adult Leydig cells secrete testosterone required for the onset and maintenance of spermatogenesis [19]. However, studies directly testing effects of AM on Leydig cells and ability of AM to modulate testosterone production are lacking.

In the present study, we examined the effects of AM injection on rat Leydig cells, which supported a stimulatory effect on multiple aspects of rat Leydig cell function *in vitro*.

Methods

Materials All materials were obtained from Huaxia Biotech (Beijing, China) unless stated otherwise.

Animals Healthy male rats ($n = 3$) with average weight of 280-300 grams at 50-60 days of age were obtained from the Central Animal Services, Fudan University. Animals were euthanized by CO₂ inhalation followed by cervical dislocation. All animal procedures were performed with approval of the Fudan University institutional animal care and use committee and in conformity with the guidelines for the care and use of laboratory animals formulated by the Ministry and Science and Technology of China (The Ministry and Science and Technology of the People's Republic of China, Beijing, China, 2006).

AM solution preparation AM injection was purchased from Wu Jing Hospital (Shanghai, China), and each was loaded 100 mL which was equivalent to 20 grams of raw herb. AM injection was diluted in dimethyl sulfoxide (DMSO) (Sigma, Shanghai, China) at 10 µg/mL, 20 µg/mL, 50 µg/mL, 100 µg/mL and 150 µg/mL respectively for cell culture.

Isolation and purification of rat Leydig cells Testes were excised sterilely from male rats ($n = 3$) following euthanasia and placed in PBS (pH 7.0-7.2). After washing 2 times with PBS, the testes were placed in DMEM. After removing the epididymis, fat and tunica albuginea, the testes were shredded into small pieces and transferred to a 50 mL centrifuge tube containing DMEM. Leydig cells were dispersed by pipetting followed by centrifugation at 800 rpm for 10 min at 4 °C. Pellet was resuspended in medium followed by centrifugation twice at 600 rpm and 4 °C for 2 min. Supernatant was collected and the cells were seeded into culture dishes for 12 h at 37 °C with 5 % CO₂. Culture medium was then replaced and cells cultured for another 12 h with Leydig cell purity assessed as described below.

3β-hydroxysteroid dehydrogenase (3β-HSD) staining The purity of cells was determined by 3β-HSD staining. After above described 24 h culture, media was removed and cells were removed from culture dish and placed in suspension. The Leydig cell suspension was incubated for 1 h

at 37 °C with 1 mg/mL NBT (nitroblue tetrazolium), 3 mg/mL NAD⁺ (Nicotinamide Adenine Dinucleotide), 2 mg/mL DHEA (dehydroepiandrosterone) and 1.6 mg/mL nicotinamide in 0.1 M PBS (Phosphate Buffered Saline). Stained cells were washed with PBS once and fixed in 10 % formaldehyde - 50 % ethanol (v/v) for 30 min, after which the cells were sedimented and washed two times. A drop of resuspended cell suspension was placed on a glass microscope slide. After drying, the percent of positive cells with a distinct blue reaction product were observed under a microscope (Olympus, Hong Kong, China).

Culture of Leydig cells and AM treatment Purified Leydig cells were seeded in either 96-well culture plates (3×10^3 cells/well) or 24-well culture plates (2×10^4 cells/well) and cultured in 90 % DMEM plus 10 % FBS and 1 % Penicillin-Streptomycin solution at 37 °C and 5 % CO₂. Treatments consisted of culture medium without (control group with equivalent DMSO instead of AM injection) or with different concentrations of AM injection (10 µg/mL, 20 µg/mL, 50 µg/mL, 100 µg/mL, 150 µg/mL). 48 h after treatment, the cells and/or culture medium were collected and used in various assays described below. All experiments were repeated 3 times using testes obtained from different rats on different days.

Cell proliferation assay Cell proliferation was detected using a CCK-8 (cell counting kit-8) assay kit according to the manufacturer's instructions (Dojindo, Shanghai, China). After above described 48 h culture, the culture medium was collected and Leydig cells were washed two times with 0.1 M PBS. Then 10 µL of CCK-8 reagent was added to each well and incubated at 37 °C for 2 h. The WST-8 (2 - (2 - methoxy - 4 - (phenyl) - 3 - (4 - (phenyl) - 5 - (2, 4 - sulpho benzene) - 2 H - tetrazolium monosodium salt) in the reagent can be reduced to orange-yellow formazan by dehydrogenase, which is proportional to the number of viable cells. Absorbance at 450 nm was recorded using a microplate reader (Thermo Scientific, Shanghai, China). A standard curve was designed using Leydig cell suspension with different dilution rates to calculate the viable cell numbers in each sample.

Testosterone Enzyme-Linked Immunosorbent assay (ELISA) Concentrations of testosterone in culture medium were detected using a Rat Free testosterone ELISA kit following the manufacturer's instructions (Zhongtianjingwei Science and Technology Co., LTD, Beijing, China). The ELISA plates were read with a microplate reader (Thermo Scientific, Shanghai, China) to record the optical densities and testosterone concentrations derived from standard curve.

Assays for SOD and GPx activities After removing the medium, Leydig cells were suspended in 10 mM PBS (pH 7.0-7.2, 100-200 µL/10⁶ cells) and cells lysed via homogenization. The mixture was centrifuged at 4,000 rpm for 10 min at 4 °C and supernatants (20-100 µg protein per sample according to the reagent specification)

used for measurement of activity of SOD and GPx using the Total Superoxide Dismutase Assay kit (Jiancheng Bioengineering Institute, Nanjing, China) with WST-8 and Cellular Glutathione Peroxidase Assay Kit respectively (Jiancheng Bioengineering Institute, Nanjing, China). The activity of SOD and GPx was adjusted according to the protein concentrations in different samples.

Quantitative real time PCR Total RNA was isolated from the Leydig cell lysates using the miRNeasy mini kit following the manufacturer's protocol (Qiagen, Shanghai, China). The integrity and concentration of total RNA were measured by agarose gel electrophoresis and Nanodrop-1000 spectrophotometer (Gene Company Limited, Hongkong, China), respectively. Total RNA (500 ng/sample) was then converted to cDNA using the iScript cDNA synthesis kit following the manufacturer's instructions (Thermo Scientific, Shanghai, China). cDNA was diluted using nuclease free water to a final volume of 40 μ L.

Quantitative real time PCR (qRT-PCR) was performed using a 20 μ L reaction volume containing 10 μ L of SYBR[®] *premix Ex Taq*[™] II (Ruian Biotech., Shanghai, China), 0.4 of ROX Reference Dye II, 0.8 μ L each of forward and reverse primer, 2 μ L of cDNA and 6 μ L of nuclease free water. Reactions were run on a 7500 Real Time PCR system (Thermo Scientific, Beijing, China) for 45 cycles of 95 $^{\circ}$ C for 15s followed by 60 $^{\circ}$ C for 1 min. *GAPDH* gene was used as the endogenous control. Primers were designed using Primer 3 (<http://primer3.ut.ee/>). The primers were as follows: *GAPDH*, 5'- TGGGTGTGAACCACGAGA -3' (forward) and 5'-GGCATGGACTGTGGTCATGA -3' (reverse); *Bax*, 5'- AGGATGCGTCCACCAAGAAGC -3' (forward) and 5'-CGGAAGAAGACCTCTCGGGG-3' (reverse); *Bcl-2*, 5'- GGAGCGTCAACAGGGAGATG-3' (forward) and 5'- CAGCCAGGAGAAATCAAACAGA -3' (reverse). The relative mRNA expression level of *Bax* and *Bcl-2* was calculated using the comparative $2^{-\Delta\Delta CT}$ method [20]. Here, we chose 20 μ g/mL *AM* as the treatment, because the results of preliminary test indicated that *AM* in concentration of 20 μ g/mL can significantly

improve the biological function of Leydig cells, and the concentration is more economical in a real application.

Statistical analysis All data were analyzed in one way ANOVA using SPSS computer software (IBM, USA). Effects of *AM* treatment (control versus 20 μ g/mL *AM*) on *Bcl-2* and *Bax* mRNA were analyzed by Student's *t*-test. Means were separated using Tukey's test. Data are presented as mean \pm SE.

Results

Leydig cell purity To assess the purity of cultured Leydig cells, 3β -HSD staining was performed. The results showed that the purity of cultured Leydig cells was > 95 % (Fig. 1).

***AM* treatment increased Leydig cell numbers** To investigate the effect of *AM* injection on Leydig cells, numbers of viable cells were determined by CCK-8 assay 48 h after *AM* treatment. Compared with the untreated control, numbers of viable cells increased within a certain range of concentration, particularly at the concentrations of 100 μ g/mL ($P < 0.05$) and 150 μ g/mL ($P < 0.01$) (Fig. 2). The result suggested that addition of *AM* increases numbers of viable Leydig cells 48 h after treatment.

***AM* treatment increased Leydig cell testosterone production** To determine potential effects of *AM* on the production of testosterone in Leydig cells, cells were treated with increasing concentrations of *AM*. Significantly higher concentrations ($P < 0.01$) of testosterone were observed in culture media of cells treated with 20 μ g/mL, 50 μ g/mL, and 100 μ g/mL of *AM* injection for 48 h (compared with untreated control; Fig. 3), suggesting certain stimulatory effects of *AM* injection on Leydig cell testosterone production. Intra- assay and inter- assay coefficients of variation were < 11% and < 9 %, respectively.

***AM* injection treatment increased SOD and GPx activities** To determine the effect of *AM* on the antioxidant defense system, the activities of SOD and GPx were detected in cultured Leydig cells 48 h after *AM* treatment. As shown in Fig. 4, the activities of SOD and GPx were significantly increased in Leydig cell cultures treated






Fig. 1 3β -HSD staining of Leydig cells. **a** Magnification, 100x; **b** Magnification, 400x. LC, Leydig cell



with 20 µg/mL and 50 µg/mL of AM relative to untreated controls ($P < 0.01$). Results indicated AM treatment could increase Leydig cell antioxidant activity within a certain range of concentration.

Effect of AM on Bax and Bcl-2 mRNA expression in AM treated Leydig cells To further elucidate the potential effects of AM on Leydig cell numbers, the expression of apoptosis-related genes *Bax* and *Bcl-2* in Leydig cells treated with 0 µg/mL or 20 µg/mL of AM injection for 48 h were analyzed using qRT-PCR. Results showed in Fig. 5 indicated the expression of *Bax* mRNA was significantly reduced in AM treated group ($P < 0.01$) compared with the untreated control, while the expression of *Bcl-2* mRNA had no obvious change between AM treated group and the untreated control ($P > 0.05$). However, the ratio of



Bcl-2/Bax mRNA was significantly higher in the AM treated group versus the untreated control ($P < 0.05$).

Discussion

Many studies have suggested a positive role for AM in the treatment of various diseases. In the present study, we investigated the effect of AM injection on function of rat Leydig cells. According to the instructions for AM injection and relevant reference [21], we at first used AM concentrations that ranged from 0-80 µg/mL to take the preliminary test. The result showed that there was no significant difference in cell number among these concentrations (the result was not provided). After several times of preliminary test and adjusting the concentration gradient, finally we chose 0 µg/mL, 10 µg/mL, 20 µg/mL, 50 µg/mL, 100 µg/mL, 150 µg/mL as the test concentration gradient.

Several studies have reported stimulatory effects of Astragaloside (main compound extracted from AM) [4, 22] on cell proliferation, which was also observed in this study. Leydig cells influence sexual maturation of males by secreting testosterone, which is influenced by cell number and rate of steroidogenesis. However, significant changes in cell proliferation and testosterone production did not occur at the same dosage of AM. The increase in testosterone production occurred at lower concentrations of AM than those demonstrated to increase Leydig cell numbers. Results suggested direct effects of AM on regulation of steroidogenesis independent of modulation of cell numbers.

Apoptosis is regulated by various factors, among which pro- and anti-apoptotic proteins of Bcl-2 family play an important role. Relative to the absolute expression of either gene alone, the ratio of *Bcl-2* to *Bax* is regarded as a better determinant to measure cell survival [23]. In the present study, we found significantly decreased *Bax* expression in the AM treated Leydig cells, which is consistent with a previous study demonstrating similar effects of AM on *Bax* expression in skin [11]. However, AM treatment did not increase Leydig cell *Bcl-2* expression in the current studies. Therefore, the increased ratio of *Bcl-2/Bax* in AM treated group may be attributed to the inhibitory effect of AM on *Bax* mRNA. A previous study demonstrated that intravenous injection of Astragaloside down-regulated *Bax* mRNA to reduce apoptosis in a rodent acute kidney injury model [3].

SOD catalyzes the dismutation of the superoxide anion, and GPx mediates the breakdown of hydrogen peroxide. These enzymes, as well as other antioxidant enzymes, are the main components of antioxidant defense system that possesses high potency to scavenge reactive oxygen species free radicals that are detrimental to cell survival by affecting cellular signaling pathways and gene expression. In the present study, we detected increased activities of SOD and GPx in the AM treated Leydig cells, supporting the




Fig. 4 Effect of AM treatment on the activities of SOD and GPx in Leydig cells. Cells were treated with 0 μg/mL, 10 μg/mL, 20 μg/mL, 50 μg/mL, 100 μg/mL and 150 μg/mL of AM. **a** SOD activity. **b** GPx activity. Results are depicted as mean +/- SE, * $P < 0.05$, ** $P < 0.01$

antioxidant role of AM. Previous studies linked the therapeutic effects of AM on myocardial ischemia [24], ischemic brain injury [25], hemorrhagic shock-reperfusion injury of intestinal mucosa [26] and the epithelial-to-mesenchymal transition in diabetic kidney disease [21] to increased activities of antioxidant enzymes such as GPx and SOD. A large body of evidence indicates antioxidants can suppress apoptosis. Hence, it is plausible that the increase in Leydig cell numbers is linked, at least in part, to stimulatory effects of AM on SOD and GPx activity resulting in reduced apoptotic cell death.

Cell proliferation is a process regulated by lots of factors. The SOD and GPx activity was just one aspect that could be considered to evaluate the growth of Leydig cells. Perhaps AM at the concentrations of 20 μg/mL and 50 μg/mL was optimal for the SOD and GPx activity of Leydig cells although it had no obvious promotion for cell proliferation. With the increase of concentration, the amount of AM was too much for the SOD and GPx, so the activity decreased, but the number of Leydig cells was significantly increased in the process. Perhaps all above lead to the result that the treated group in which there was

a significant difference of Leydig cell number was not consistent with that of SOD and GPx activity. Results suggested direct effects of AM on regulation of SOD and GPx activity independent of modulation of cell numbers.

Although the study preliminarily indicated that AM injection has a stimulatory effect on multiple aspects of rat Leydig cell function *in vitro*, it had some limitations, such as the design of AM injection gradient, and this was just the results of cell culture level. Therefore, lots of work and study should be done to elucidate the exact mechanism of AM promoting the function of Leydig cells.

Conclusions

In the present study, we investigated the influence of AM injection on cell numbers and testosterone production of rat Leydig cells cultured *in vitro*. The results suggested that AM injection has a favorable effect on the function of Leydig cells. The study provides a foundation for future studies to understand the mechanisms responsible for stimulatory effects of AM injection on Leydig cells and whether AM treatment *in vivo* can enhance indices of male fertility.




Fig. 5 Effect of AM treatment on the expression of Bax and Bcl-2 in Leydig cells. qRT-PCR was performed to determine the abundances of Bax (a) and Bcl-2 (b) mRNA in Leydig cells treated with 20 μg/mL of AM, and to further determine the effect of AM treatment on expression level of Bax and Bcl-2, we calculated the ratio of Bcl-2 to Bax (c). Expression of Bax and Bcl-2 mRNA was normalized relative to the abundance of GAPDH mRNA. Results are depicted as mean +/- SE (n = 3). * $P < 0.05$, ** $P < 0.01$

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

XLJ performed the statistical analyses and drafted the manuscript. XC participated in the design of the study and carried out the assays. XLY, JWC and MMZ assisted in the cell culture assays. YL and JZM assisted in the Quantitative real time PCR and ELSIA. ZYL participated in the design of the study and provided technical supports. YH and PFL participated in the statistical analyses and assisted in drafting the manuscript. JBY, GS and LHL provided conceptual advice and critically revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This study was Supported by Chinese Natural Science Foundation (No. 31172211), Shanxi Scholarship Council of China (2014-key 5), and Shanxi Talent Introduction Project and SXAU Research Achievement Development Project(71060003).

Author details

¹College of Animal Science and Technology, Shanxi Agricultural University, No.1 Mingxian Nan Road, Taigu 030801, China. ²College of Life Science, Shanxi Agricultural University, Taigu, Shanxi 030801, China. ³Pharmaceutical Analysis Laboratory of Fudan University, Shanghai 200000, China. ⁴Laboratory of Mammalian Reproductive Biology and Genomics, Michigan State University, East Lansing, MI 48824, USA. ⁵Departments of Animal Science, Michigan State University, East Lansing, MI 48824, USA. ⁶Departments of Animal Physiology, Michigan State University, East Lansing, MI 48824, USA. ⁷Division of Animal and Nutritional Sciences, West Virginia University, Morgantown, WV 26506, USA.

Received: 25 March 2015 Accepted: 13 July 2015

Published online: 01 August 2015

References

- Rato L, Alves MG, Socorro S, Duarte AI, Cavaco JE, Oliveira PF. Metabolic regulation is important for spermatogenesis. *Nat Rev Urol*. 2012;9:330–8.
- Zhang YW, Wu CY, Cheng JT. Merit of Astragalus polysaccharide in the improvement of early diabetic nephropathy with an effect on mRNA expressions of NF-kappaB and IkappaB in renal cortex of streptozotocin-induced diabetic rats. *J Ethnopharmacol*. 2007;114:387–92.
- Gui D, Huang J, Guo Y, Chen J, Chen Y, Xiao W, et al. Astragaloside IV ameliorates renal injury in streptozotocin-induced diabetic rats through inhibiting NF-kappaB-mediated inflammatory genes expression. *Cytokine*. 2013;61:970–7.
- Zhang Y, Hu G, Li S, Li ZH, Lam CO, Hong SJ, et al. Pro-angiogenic activity of astragaloside IV in HUVECs *in vitro* and zebrafish *in vivo*. *Mol Med Rep*. 2012;5:805–11.
- Zhang L, Liu Q, Lu L, Zhao X, Gao X, Wang Y. Astragaloside IV stimulates angiogenesis and increases hypoxia-inducible factor-1alpha accumulation via phosphatidylinositol 3-kinase/Akt pathway. *J Pharmacol Exp Ther*. 2011;338:485–91.
- Kim BH, Oh I, Kim JH, Jeon JE, Jeon B, Shin J, et al. Anti-inflammatory activity of compounds isolated from *Astragalus sinicus* L. in cytokine-induced keratinocytes and skin. *Exp Mol Med*. 2014;46:e87.
- Gao J, Feng LJ, Huang Y, Li P, Xu DJ, Li J, et al. Total glucosides of Danggui Buxue Tang attenuates bleomycin-induced pulmonary fibrosis via inhibition of extracellular matrix remodelling. *J Pharm Pharmacol*. 2012;64:811–20.
- Li Z, Zhang L, He W, Zhu C, Yang J, Sheng M. Astragalus membranaceus inhibits peritoneal fibrosis via monocyte chemoattractant protein (MCP)-1 and the transforming growth factor-beta1 (TGF-beta1) pathway in rats submitted to peritoneal dialysis. *Int J Mol Sci*. 2014;15:12959–71.
- Wei MG, Sun W, Xiong PH, Shao JD. Antifibrotic effect of the Chinese herbs Modified Danggui Buxue Decoction on adriamycin-induced nephropathy in rats. *Chin J Integr Med*. 2012;18:591–8.
- He Y, Du M, Gao Y, Liu H, Wang H, Wu X, et al. Astragaloside IV attenuates experimental autoimmune encephalomyelitis of mice by counteracting oxidative stress at multiple levels. *PLoS ONE*. 2013;8:e76495.
- Kim MH, Kim SH, Yang WM. Beneficial effects of Astragaloside IV for hair loss via inhibition of Fas/Fas L-mediated apoptotic signaling. *PLoS ONE*. 2014;9:e92984.
- Wang B, Chen MZ. Astragaloside IV possesses antiarthritic effect by preventing interleukin 1beta-induced joint inflammation and cartilage damage. *Arch Pharm Res*. 2014;37:793–802.
- Hong F, Xiao W, Ragupathi G, Lau CB, Leung PC, Yeung KS, et al. The known immunologically active components of *Astragalus* account for only a small proportion of the immunological adjuvant activity when combined with conjugate vaccines. *Planta Med*. 2011;77:817–24.
- Botchkareva NV, Ahluwalia G, Shander D. Apoptosis in the hair follicle. *J Invest Dermatol*. 2006;126:258–64.
- Gui D, Guo Y, Wang F, Liu W, Chen J, Chen Y, et al. Astragaloside IV, a novel antioxidant, prevents glucose-induced podocyte apoptosis *in vitro* and *in vivo*. *PLoS ONE*. 2012;7:e39824.
- Hong CY, Ku J, Wu P. Astragalus membranaceus stimulates human sperm motility *in vitro*. *Am J Chin Med*. 1992;20:289–94.
- Kim W, Kim SH, Park SK, Chang MS. Astragalus membranaceus ameliorates reproductive toxicity induced by cyclophosphamide in male mice. *Phytother Res*. 2012;26:1418–21.
- Liang P, Li H, Peng X, Xiao J, Liu J, Ye Z. [Effects of astragalus membranaceus injection on sperm abnormality in Cd-induced rats]. *Zhonghua Nan Ke Xue*. 2004;10:42–5. 48.
- Chen H, Ge RS, Zirkin BR. Leydig cells: from stem cells to aging. *Mol Cell Endocrinol*. 2009;306:9–16.
- Shi L, Zhao H, Ren Y, Yao X, Song R, Yue W. Effects of different levels of dietary selenium on the proliferation of spermatogonial stem cells and antioxidant status in testis of roosters. *Anim Reprod Sci*. 2014;149:266–72.
- Qi W, Niu J, Qin Q, Qiao Z, Gu Y. Astragaloside IV attenuates glycated albumin-induced epithelial-to-mesenchymal transition by inhibiting oxidative stress in renal proximal tubular cells. *Cell Stress Chaperones*. 2014;19:105–14.
- Junior AA, Oliveira LS, Assis Neto AC, Alves FR, Miglino MA, Carvalho MA. Spermatogenesis in goats with and without scrotum bipartition. *Anim Reprod Sci*. 2012;130:42–50.
- Tanaka K, Asanuma M, Ogawa N. Molecular basis of anti-apoptotic effect of immunophilin ligands on hydrogen peroxide-induced apoptosis in human glioma cells. *Neurochem Res*. 2004;29:1529–36.
- Zhang WD, Chen H, Zhang C, Liu RH, Li HL, Chen HZ. Astragaloside IV from *Astragalus membranaceus* shows cardioprotection during myocardial ischemia *in vivo* and *in vitro*. *Planta Med*. 2006;72:4–8.
- Luo Y, Qin Z, Hong Z, Zhang X, Ding D, Fu JH, et al. Astragaloside IV protects against ischemic brain injury in a murine model of transient focal ischemia. *Neurosci Lett*. 2004;363:218–23.
- Hei ZQ, Huang HQ, Zhang JJ, Chen BX, Li XY. Protective effect of *Astragalus membranaceus* on intestinal mucosa reperfusion injury after hemorrhagic shock in rats. *World J Gastroenterol*. 2005;11:4986–91.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

