Smorenberg et al. Perioperative Medicine 2014, 3:3 http://www.perioperativemedicinejournal.com/content/3/1/3



CORRECTION **Open Access**

Correction: Dose and type of crystalloid fluid therapy in adult hospitalized patients

Annemieke Smorenberg, Can Ince and A B Johan Groeneveld*

After the publication of this work [1] it was brought to our attention that there was an error in Table two (Table 1 here) of the article, in which the table columns were misaligned. The correct version of the table is included here:

Table 1 Composition of bodyl fluids (in mmol/L)

	Na ⁺	K ⁺	Ca ²⁺	Mg ²⁺	Cl ⁻	HCO ₃	HPO ₄	HSO ₄	Organic acid	Protein
Plasma	142	4	5	2	101	27	2	1	6	16
Plasma wa	ter									
	153	4.3	5.4	2.2	109	29	2.2	1	6.5	17
Interstitial	water									
	139	4	5	2	114	31	2	1	7	1
Intracellula	ar water									
	10	160	2	26	3	10	100	20		65

Received: 13 May 2014 Accepted: 14 May 2014

Published: 23 May 2014

Reference

Smorenberg A, Ince C, Groeneveld ABJ: Dose and type of crystalloid fluid therapy in adult hospitalized patients. Perioper Med 2013, 2:17.

doi:10.1186/2047-0525-3-3

Cite this article as: Smorenberg et al.: Correction: Dose and type of crystalloid fluid therapy in adult hospitalized patients. Perioperative Medicine 2014 3:3

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



^{*} Correspondence: a.b.j.groeneveld@erasmusmc.nl Department of Intensive Care, Erasmus Medical Centre, 's Gravendijkwal 230, 3015 CE Rotterdam, The Netherlands

