

Collect. Math. (2011) 62:275–296
 DOI 10.1007/s13348-011-0044-x

On the full, strongly exceptional collections on toric varieties with Picard number three

Michał Lason · Mateusz Michałek

Received: 23 February 2011 / Accepted: 11 May 2011 / Published online: 28 May 2011
 © The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract We investigate full strongly exceptional collections on smooth, complete toric varieties. We obtain explicit results for a large family of varieties with Picard number three, containing many of the families already known. We also describe the relations between the collections and the split of the push forward of the trivial line bundle by the toric Frobenius morphism.

Contents

1	Introduction	276
2	Preliminaries	277
	2.1 Full, strongly exceptional collections	277
	2.2 Toric varieties	277
3	First results and methods	279
	3.1 Bondal's construction and Thomsen's algorithm	279
	3.2 Techniques of counting homology	283
4	Main theorem	286
	4.1 Our setting	287
	4.2 Acyclicity of differences of line bundles from <i>Col</i>	288
	4.3 Generating the derived category	290

M. Lason is supported by the grant of Polish MNiSzW N N201 413139.

M. Michałek is supported by the grant of Polish MNiSzW N N201 413539.

M. Lason · M. Michałek

Institute of Mathematics of the Polish Academy of Sciences, Św. Tomasza 30, 31-027 Kraków, Poland

M. Lason (✉)

Theoretical Computer Science Department, Faculty of Mathematics and Computer Science,
 Jagiellonian University, S. Łojasiewicza 6, 30-348 Kraków, Poland
 e-mail: mlason@op.pl

M. Michałek

Institut Fourier, Université Joseph Fourier, 100 rue des Maths, BP 74, 38402 St Martin d'Hères, France
 e-mail: wajcha2@poczta.onet.pl

5	Bondal’s construction not containing a full, strongly exceptional collection	294
5.1	Example	294
5.2	Our case	295

1 Introduction

Let X be a smooth variety over an algebraically closed field \mathbb{K} of characteristic zero and let $D^b(X)$ be the derived category of bounded complexes of coherent sheaves of \mathcal{O}_X -modules. This category is an important algebraic invariant of X . In order to understand the derived category $D^b(X)$ one is interested in knowing a strongly exceptional collection of objects that generate $D^b(X)$, see also [4].

For a smooth, complete toric variety X there is a well known construction due to Bondal which gives a full collection of line bundles in $D^b(X)$. In some cases Bondal’s collection of line bundles is a strongly exceptional collection (see also [3]), but it is not true in general. Often one can find a subset of this collection and order it in such a way that it becomes strongly exceptional and remains full. This approach was well described in [8] for a class of toric varieties with Picard number three.

One of the first conjectures concerning this topic was made by King [21]:

Conjecture 1.1 King’s *For any smooth, complete toric variety X there exists a full, strongly exceptional collection of line bundles.*

Originally this conjecture was made in terms of existence of tilting bundles whose direct summands are line bundles, but it is easy to see that they are equivalent, see [9]. It was disproved by Hille and Perling [17]. They gave an example of a smooth, complete toric surface which does not have a full, strongly exceptional collection of line bundles. The conjecture was reformulated by Miró-Roig and Costa (stated also in [6]):

Conjecture 1.2 *For any smooth, complete Fano toric variety there exists a full, strongly exceptional collection of line bundles.*

This conjecture has an affirmative answer when the Picard number of X is less then or equal to two [9] or the dimension of X is at most three [2, 4, 6]. Recently it was disproved by Efimov [12]. In the same paper the author states the following conjecture, suggested by D. Orlov.

Conjecture 1.3 [12] *For any smooth projective toric DM stack Y , the derived category $D^b(Y)$ is generated by a strong exceptional collection.*

Here the assumption on the objects forming the collection is relaxed. We believe that one could possibly ask if the collection can be made from coherent sheaves or toric vector bundles. There is a well known result due to Kawamata in this direction [20].

Theorem 1 [20] *For any smooth projective toric DM stack Y , the derived category $D^b(Y)$ is generated by an exceptional collection of coherent sheaves.*

The goal of this paper is to investigate when it is possible to find a full, strongly exceptional collection and whether line bundles that come from Bondal’s construction contain such a collection. The examples in which such collections do or do not exist are now excessively studied, see for example [18, 23, 26]. We restrict our attention to smooth, complete toric varieties with Picard number three. There are some families among these varieties for which

Conjecture 1.2 is true [8, 11]. We state Theorem 4.24 for a much larger family of varieties containing both families already known. Namely for family having half of parameters fixed and the other half arbitrary, among toric varieties with Picard number three.

In Sect. 5 we also show that in general it is not possible to find a full, strongly exceptional collection among line bundles that come from Bondal’s construction, even in the Fano case.

To determine the image of Bondal’s construction we look at the image of the real torus in the Picard group of a toric variety. We also compare this with the result of Thomsen’s algorithm [27] that gives a decomposition of the push forward of a line bundle by a toric Frobenius morphism. This leads to some unexpected results like Corollary 3.5.

To prove that a given collection of line bundles is strongly exceptional we develop new, efficient methods of counting homologies of simplicial complexes given by primitive collections, that is minimal subsets of points that do not form a simplex. To do this we use the results of [24]. In particular this enables us to determine all acyclic simplicial complexes arising from complete toric varieties with Picard number three.

2 Preliminaries

2.1 Full, strongly exceptional collections

For an algebraic variety X let $D^b(X)$ be the derived category of coherent sheaves on X . For an introduction to derived categories the reader is advised to look in [7, 14, 19]. The structure and properties of the derived category of an arbitrary variety X can be very complicated and they are an object of many studies. One of the approaches to understand the derived category uses the notion of exceptional objects. Let us introduce the following definitions (see also [15]):

Definition 2.1

1. A coherent sheaf F on X is **exceptional** if $\text{Hom}(F, F) = \mathbb{K}$ and $\text{Ext}_{\mathcal{O}_X}^i(F, F) = 0$ for $i \geq 1$.
2. An ordered collection (F_0, F_1, \dots, F_m) of coherent sheaves on X is an **exceptional collection** if each sheaf F_i is exceptional and $\text{Ext}_{\mathcal{O}_X}^i(F_k, F_j) = 0$ for $j < k$ and $i \geq 0$.
3. An exceptional collection (F_0, F_1, \dots, F_m) of coherent sheaves on X is a **strongly exceptional collection** if $\text{Ext}_{\mathcal{O}_X}^i(F_j, F_k) = 0$ for $j \leq k$ and $i \geq 1$.
4. A (strongly) exceptional collection (F_0, F_1, \dots, F_m) of coherent sheaves on X is a **full, (strongly) exceptional collection** if it generates the bounded derived category $D^b(X)$ of X i.e. the smallest triangulated category containing $\{F_0, F_1, \dots, F_n\}$ is equivalent to $D^b(X)$.

For an exceptional collection (F_0, \dots, F_m) one may define an object $F = \bigoplus_{i=0}^m F_i$ and an algebra $A = \text{Hom}(F, F)$. Such an object gives us a functor G_F from $D^b(X)$ to the derived category $D^b(A\text{-mod})$ of right finite-dimensional modules over the algebra A . Bondal proved in [4], that if X is smooth and (F_i) is a full, strongly exceptional collection, then the functor G_F gives an equivalence of these categories. For further reading only the definition of the full strongly exceptional collection is necessary.

2.2 Toric varieties

A normal algebraic variety is called toric if it contains a dense torus $(\mathbb{C}^*)^n$ whose action on itself extends to the action on the whole variety. For a good introduction to toric varieties the

reader is advised to look in [10] or [13]. Varieties of this type form a sufficiently large class among normal varieties to test many hypothesis in algebraic geometry. Many invariants of a toric variety can be effectively computed using combinatorial description. Let us recall it.

Given an n dimensional torus T we may consider one parameter subgroups of T , that is morphisms $\mathbb{C}^* \rightarrow T$ and characters of T , that is morphisms $T \rightarrow \mathbb{C}^*$. One parameter subgroups form a lattice N and characters form a lattice M . These lattices are dual to each other and isomorphic to \mathbb{Z}^n .

A toric variety X is constructed from a fan Σ , that is a system of cones $\sigma_i \subset N$. This is done by gluing together affine schemes $\text{Spec}(\mathbb{C}[\sigma_i^*])$, where $\sigma_i^* \subset M$ is a cone dual to σ_i . One dimensional cones in Σ are called rays. The generators of these semigroups are called ray generators.

Many properties of the variety X can be described using the fan Σ . For example X is smooth if and only if for every cone σ_i the set of its ray generators can be extended to the basis of N . Moreover to each ray generator v we may associate a unique T invariant Weil divisor denoted by D_v . There is a well known exact sequence:

$$0 \rightarrow M \rightarrow Div_T \rightarrow Cl(X) \rightarrow 0, \tag{2.1}$$

where Div_T is the group of T invariant Weil divisors and $Cl(X)$ is the class group. The map $M \rightarrow Div_T$ is given by:

$$m \rightarrow \sum m(v_i)D_{v_i},$$

where the sum is taken over all ray generators v_i .

Smooth, complete toric varieties with Picard number three have been classified by Betyrev [1] according to their primitive relations. Let Σ be a fan in $N = \mathbb{Z}^n$ and let R be the set of rays of Σ .

Definition 2.2 We say that a subset $P \subset R$ is a primitive collection if it is a minimal subset of R which does not span a cone in Σ .

In other words a primitive collection is a subset of ray generators, such that all together they do not span a cone in Σ but if we remove any generator, then the rest spans a cone that belongs to Σ . To each primitive collection $P = \{x_1, \dots, x_k\}$ we associate a primitive relation. Let $w = \sum_{i=1}^k x_i$. Let $\sigma \in \Sigma$ be the cone of the smallest dimension that contains w and let y_1, \dots, y_s be the ray generators of this cone. The toric variety of Σ was assumed to be smooth, so there are unique positive integers n_1, \dots, n_s such that

$$w = \sum_{i=1}^s n_i y_i.$$

Definition 2.3 For each primitive collection $P = \{x_1, \dots, x_k\}$ let n_i and y_i be as described above. The linear relation:

$$x_1 + \dots + x_k - n_1 y_1 - \dots - n_s y_s = 0$$

is called the primitive relation (associated to P).

Using the results of [16,25] Batyrev proved in [1] that for any smooth, complete n dimensional fan with $n + 3$ generators its set of ray generators can be partitioned into l non-empty sets X_0, \dots, X_{l-1} in such a way that the primitive collections are exactly sums of $p + 1$ consecutive sets X_i (we use a circular numeration, that is we assume that $i \in \mathbb{Z}/l\mathbb{Z}$), where

$l = 2p + 3$. Moreover l is equal to 3 or 5. The number l is of course the number of primitive collections. In the case $l = 3$ the fan Σ is a splitting fan (that is any two primitive collections are disjoint). These varieties are well characterized, and we know much about full, strongly exceptional collections of line bundles on them. The case of five primitive collections is much more complicated and is our object of study. For $l = 5$ we have the following result of Batyrev [1, Theorem 6.6]:

Theorem 2.4 *Let $Y_i = X_i \cup X_{i+1}$, where $i \in \mathbb{Z}/5\mathbb{Z}$,*

$$\begin{aligned} X_0 &= \{v_1, \dots, v_{p_0}\}, & X_1 &= \{y_1, \dots, y_{p_1}\}, & X_2 &= \{z_1, \dots, z_{p_2}\}, \\ X_3 &= \{t_1, \dots, t_{p_3}\}, & X_4 &= \{u_1, \dots, u_{p_4}\}, \end{aligned}$$

where $p_0 + p_1 + p_2 + p_3 + p_4 = n + 3$. Then any n -dimensional fan Σ with the set of generators $\bigcup X_i$ and five primitive collections Y_i can be described up to a symmetry of the pentagon by the following primitive relations with nonnegative integral coefficients $c_2, \dots, c_{p_2}, b_1, \dots, b_{p_3}$:

$$\begin{aligned} v_1 + \dots + v_{p_0} + y_1 + \dots + y_{p_1} - c_2 z_2 - \dots - c_{p_2} z_{p_2} \\ - (b_1 + 1)t_1 - \dots - (b_{p_3} + 1)t_{p_3} &= 0, \\ y_1 + \dots + y_{p_1} + z_1 + \dots + z_{p_2} - u_1 - \dots - u_{p_4} &= 0, \\ z_1 + \dots + z_{p_2} + t_1 + \dots + t_{p_3} &= 0, \\ t_1 + \dots + t_{p_3} + u_1 + \dots + u_{p_4} - y_1 - \dots - y_{p_1} &= 0, \\ u_1 + \dots + u_{p_4} + v_1 + \dots + v_{p_0} - c_2 z_2 - \dots - c_{p_2} z_{p_2} - b_1 t_1 - \dots - b_{p_3} t_{p_3} &= 0. \end{aligned}$$

□

In this case we may assume that

$$v_1, \dots, v_{p_0}, \quad y_2, \dots, y_{p_1}, \quad z_2, \dots, z_{p_2}, \quad t_1, \dots, t_{p_3}, \quad u_2, \dots, u_{p_4}$$

form a basis of the lattice N . The other vectors are given by

$$\begin{aligned} z_1 &= -z_2 - \dots - z_{p_2} - t_1 - \dots - t_{p_3} \\ y_1 &= -y_2 - \dots - y_{p_1} - z_1 - \dots - z_{p_2} + u_1 + \dots + u_{p_4} \\ u_1 &= -u_2 - \dots - u_{p_4} - v_1 - \dots - v_{p_0} + c_2 z_2 + \dots + c_{p_2} z_{p_2} \\ &\quad + b_1 t_1 + \dots + b_{p_3} t_{p_3}. \end{aligned} \tag{2.2}$$

3 First results and methods

3.1 Bondal’s construction and Thomsen’s algorithm

We start this section by recalling Thomsen’s [27] algorit for computing the summands of the push forward of a line bundle by a Frobenius morphism. We do this because of two reasons.

First is that Thomsen in his paper assumes finite characteristic of the ground field and uses absolute Frobenius morphism. We claim that the arguments used apply also in case of geometric Frobenius morphism and characteristic zero.

Moreover by recalling all methods we are able to show that the results of Thomsen coincide with the results stated by Bondal [3]. Combining these both methods enables us to deduce some interesting facts about toric varieties.

Most of the results of this section are due to Bondal and Thomsen. We use the notation from [27]. Let $\Sigma \subset N$ be a fan such that the toric variety $X = X(\Sigma)$ is smooth. Let us denote by $\sigma_i \in \Sigma$ the cones of our fan and by T the torus of our variety. If we fix a basis (e_1, \dots, e_n) of the lattice N , then of course $T = \text{Spec } R$, where $R = k[X_{e_1}^{\pm 1}, \dots, X_{e_n}^{\pm 1}]$.

In characteristic p we have got two p th Frobenius morphisms $F : X \rightarrow X$. One of them is the absolute Frobenius morphism given as an identity on the underlying topological space and a p th power on sheaves. Notice that on the torus it is given by a map $R \rightarrow R$ that is simply a p th power map, hence it is not a morphism of k algebras (it is not an identity on k).

The other morphism is called the geometric Frobenius morphism and can be defined in any characteristic. Let us fix an integer m . Consider a morphism of tori $T \rightarrow T$ that associates t^m to a point t . This is a morphism of schemes over k that can be extended to the m th geometric Frobenius morphism $F : X \rightarrow X$. What is important is that both of these morphisms can be considered as endomorphisms of open affine subsets associated to cones of Σ . We claim that in both cases the Thomsen’s algorithm works.

We begin by recalling the algorithm from [27]. Let v_{i1}, \dots, v_{id_i} be the ray generators of the d_i dimensional cone σ_i . As the variety was assumed to be smooth we may extend this set to a basis of N . Let A_i be a square matrix whose rows are vectors v_{ij} in the fixed basis of N . Let $B_i = A_i^{-1}$ and let w_{ij} be the j th column of B_i . Of course the columns of B_i are ray generators (extended to a basis) of the dual cone $\sigma_i^* \subset M = N^*$.

Let us remind that $X(\Sigma)$ is covered by affine open subsets $U_{\sigma_i} = \text{Spec } R_i$, where $R_i = k[X^{w_{i1}}, \dots, X^{w_{id_i}}, X^{\pm w_{id_i+1}}, \dots, X^{\pm w_{in}}]$. Here we use the notation $X^v = X_{e_1}^{v_1} \cdots X_{e_n}^{v_n}$. Let also $X_{ij} = X^{w_{ij}}$. In this way the monomials X_{i1}, \dots, X_{in} should be considered as coordinates on the affine subset U_{σ_i} , so we are able to think about monomials on U_{σ_i} as vectors: a vector v corresponds to the monomial X_i^v . Of course all of these affine subsets contain T , that corresponds to the inclusions $R_i \subset R$.

Using the results of [13] we know that $U_{\sigma_i} \cap U_{\sigma_j} = U_{\sigma_i \cap \sigma_j}$ and this is a principal open subset of U_{σ_i} . This means that there is a monomial M_{ij} such that $U_{\sigma_i \cap \sigma_j} = \text{Spec}((R_i)_{M_{ij}})$.

We are interested in Picard divisors. A T invariant Picard divisor is given by a compatible collection $\{(U_{\sigma_i}, X_i^{u_i})\}_{\sigma_i \in \Sigma}$. Compatible means that the quotient of any two functions in the collection is invertible on the intersection of domains. This motivates the definition:

$$I_{ij} = \{v : X_i^v \text{ is invertible in } (R_i)_{M_{ij}}\}.$$

Given a monomial X_i^v , if we want to know how it looks in coordinates $X_{e_1}^*, \dots, X_{e_n}^*$ (obviously from the definition of X_i) we just have to multiply v by B_i : $X_i^v = X^{B_i v}$. We see that $X_i^v = X_j^{B_j^{-1} B_i v}$. That is why we define $C_{ij} = B_j^{-1} B_i$ and we think of C_{ij} as the matrices that translate the monomials in coordinates of one affine piece to another.

Now the compatibility in the definition of a Cartier divisor simply is equivalent to the condition $u_j - C_{ij} u_i \in I_{ji}$. We define $u_{ij} = u_j - C_{ij} u_i$ and think about them as transition maps. Of course a divisor is principal if and only if $u_{ij} = 0$ for all i, j (vector equal to 0 corresponds to a constant function equal to 1).

Let $P_m = \{v = (v_1, \dots, v_n) : 0 \leq v_i < m\}$. Later we will see that this set has got a description in terms of characters of the kernel of the Frobenius map between tori.

Using simple algebra Thomsen proves that the following functions are well defined (the only think to prove is that the image of h is in I_{ji}):

Let us fix $w \in I_{ji}$ and a positive integer m . We define the functions

$$\begin{aligned} h_{ijm}^w &: P_m \rightarrow I_{ji} \\ r_{ijm}^w &: P_m \rightarrow P_m, \end{aligned}$$

for any $v \in P_m$ by the equation

$$C_{ij}v + w = mh_{ijm}^w(v) + r_{ijm}^w(v).$$

This is a simple division by m with the rest. Moreover r_{ijm}^w is bijective.

Now if we have any $v \in P_m$, a T -Cartier divisor $D = \{(U_{\sigma_i}, X_i^{u_i})\}_{\sigma_i \in \Sigma}$ and a fixed $\sigma_l \in \Sigma$ then Thomsen defines $t_i = h_{lij}^{u_i}(v)$. He proves that the collection $\{(U_{\sigma_i}, X_i^{t_i})\}_{\sigma_i \in \Sigma}$ is a T -Cartier divisor D_v . This is of course independent on the representation of D up to linear equivalence. The choice of l corresponds to “normalizing” the representation of D on the affine subset U_{σ_l} . Although the definition of D_v may depend on l , the vector bundle $\bigoplus_{v \in P_m} \mathcal{O}(D_v)$ is independent on l . Moreover Thomsen proves that in case of p th absolute Frobenius morphism and characteristic $p > 0$ this vector bundle is a push forward of the line bundle $\mathcal{O}(D)$. The proof uses only the fact that the Frobenius morphism can be considered as a morphism of affine pieces U_{σ_i} , so can be extended to the case of geometric Frobenius morphism and arbitrary characteristic. One only has to notice that the basis of free modules obtained by Thomsen in [27, Section 5, Theorem 1] are exactly the same in all cases.

Now let us remind that there is an exact sequence 2.1:

$$0 \rightarrow M \rightarrow D_T \rightarrow Pic \rightarrow 0,$$

where D_T are T invariant divisors. Let (g_j) be the collection of ray generators of the fan Σ and D_{g_j} a divisor associated to the ray generator g_j . The morphism from M to D_T is given by $v \rightarrow \sum_j v(g_j)D_{g_j}$. Such a map may be extended to a map from $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$ by $f : v \rightarrow \sum_j [v(g_j)]D_{g_j}$. Notice that this is no longer a morphism, however if $a \in M$ and $b \in M_{\mathbb{R}}$, then $f(a + b) = f(a) + f(b)$. We obtain a map $\mathbf{T} := \frac{M_{\mathbb{R}}}{M} \rightarrow Pic$, where \mathbf{T} is a real torus (do not confuse with T). We also fix the notation for an \mathbb{R} -divisor $D = \sum_j a_j D_{g_j}$:

$$[D] := \sum_j [a_j]D_{g_j}.$$

Let G be the kernel of the m th geometric Frobenius morphism between the tori T . By acting with the functor $\text{Hom}(\cdot, \mathbb{C}^*)$ we obtain an exact sequence:

$$0 \rightarrow M \rightarrow M \rightarrow G^* \simeq \frac{M}{mM} \rightarrow 0.$$

We also have a morphism:

$$\frac{1}{m} : G^* \simeq \frac{M}{mM} \rightarrow \mathbf{T},$$

that simply divides the coordinates by m . By composing it with the morphism from $\mathbf{T} \rightarrow Pic$ we get a morphism from G^* to Pic . It can be also described as follows:

We fix $\chi \in G^*$ and arbitrarily lift it to an element $\chi_M \in M$. Now we use the morphism $M \rightarrow Div_T$ to obtain a T invariant principal divisor D_{χ} . The image of χ in Pic is simply equal to $[\frac{D_{\chi}}{m}]$. Of course for different lifts of χ to M we get linearly equivalent divisors. Now we prove one of the results stated by Bondal in [3]:

Proposition 3.1 *Let $L = \mathcal{O}(D)$ by any line bundle on a smooth toric variety X . The push forward $F_*(\mathcal{O}(D))$ is equal to $\bigoplus_{\chi \in G^*} \mathcal{O}\left(\lceil \frac{D+D_{\chi}}{m} \rceil\right)$.*

Remark 3.2 The characters of G play the role of $v \in P_m$ in Thomsen’s algorithm. Notice also that it is not clear that $\bigoplus_{\chi \in G^*} \mathcal{O}\left(\lceil \frac{D+D_{\chi}}{m} \rceil\right)$ is independent on the representation of L

by D . If we prove that this is equal to the push forward then this fact will follow, but in the proof we have to take any representation of L and we cannot change D with a linearly equivalent divisor.

Proof Let $D = \{(U_{\sigma_i}, X_i^{u_i})\}$ and let us fix $\chi \in G^*$. We have to prove that $\mathcal{O}\left(\left[\frac{D+D_\chi}{m}\right]\right)$ is one of $\mathcal{O}(D_v)$ for $v \in P_m$ and that this correspondence is one to one over all $\chi \in G^*$. We already know that $\left[\frac{D_\chi}{m}\right]$ is independent on the choice of the lift of χ , so we may take such a lift, that $v = \chi_M + u_l$ is in the P_m . Here l is an index of a cone, but we may assume that its ray generators form a standard basis of N , so $A_l = Id$. Of course such a matching between $\chi \in G^*$ and $v \in P_m$ is bijective.

Now let us compare the coefficients of $\left[\frac{D+D_\chi}{m}\right]$ and D_v . We fix a ray generator $r = (r_1, \dots, r_n) \in \sigma_j$. Let k be such that this ray generator is the k th row of matrix A_j . We compare coefficients of D_r . Let $\chi_M = (a_1, \dots, a_n)$. We see that:

$$\left[\frac{D + D_\chi}{m}\right] = \dots + \left[\frac{(u_j)_k + \sum_{w=1}^n a_w r_w}{m}\right] D_r + \dots$$

Here of course $(u_j)_k$ is not a transition map u_{jk} , but the k th entry of vector u_j that is of course the coefficient of D_r of the divisor D . Now from Thomsen’s algorithm described above we know that

$$C_{lj}(\chi + u_l) + u_{lj} = m t_j + r,$$

where $r \in P_m$. We see that

$$t_j = \left[\frac{C_{lj}(\chi + u_l) + u_{lj}}{m}\right].$$

Now $A_l = Id$ and from the definition of u_{lj} we have $C_{lj}u_l + u_{lj} = u_j$, so:

$$t_j = \left[\frac{A_j \chi + u_j}{m}\right].$$

This gives us:

$$D_v = \dots + \left[\frac{\sum_{w=1}^n a_w r_w + (u_j)_k}{m}\right] D_r + \dots$$

what completes the proof.

From [3] we know that the image B of \mathbf{T} in Pic is a full collection of line bundles. Of course B is a finite set (the coefficients of divisors associated to ray generators are bounded). Moreover the image of rational points of \mathbf{T} contains the whole image of \mathbf{T} (a set of equalities and inequalities with rational coefficients has got a solution in \mathbb{R} if and only if it has got a solution in \mathbb{Q}). This means that for sufficiently large m the split of the push forward of the trivial bundle by the m th Frobenius morphism coincides with the image of \mathbf{T} and hence is full.

Let us now consider an example of \mathbb{P}^2 . Let v_1, v_2 and $v_3 = -v_1 - v_2$ be the ray generators of the fan. We fix a basis (v_1, v_2) of N . The image of the torus \mathbf{T} is equal to the set of all divisors of the form $[a]D_{v_1} + [b]D_{v_2} + [-a - b]D_{v_3}$ for $0 \leq a, b < 1$. We see that the image of the torus \mathbf{T} is $\mathcal{O}, \mathcal{O}(-1), \mathcal{O}(-2)$. This is a full collection. Notice however that it is not true that if we have a line bundle L then there exists an integer m_0 such that the push forward of L by the m th Frobenius morphism for $m > m_0$ is a direct sum of line bundles from B .

For example the push forward of $\mathcal{O}(-3)$ always contains in the split $\mathcal{O}(-3)$ that is not an element of B . However, as we will see only minor differences from the set B are possible.

Definition 3.3 Let us fix a natural bijection between points of \mathbf{T} and elements of $M_{\mathbb{R}}$ with entries from $[0, 1)$ in some fixed basis. Now each element of B has got a natural representant in Div_T as sum of D_{g_j} with integer coefficients. Let $B_0 \subset Div_T$ be the set of these representatives. We define the set B' as the set of all divisors D in Pic for which there exists an element in $b \in B_0$, such that there exists a representation of D whose coefficients differ by at most one from the coefficients of b .

In other words we take (some fixed) representations of all elements of B , we take all other representations whose coefficients differ by at most one and we take the image in Pic to obtain B' .

Let us look once more at the example of \mathbb{P}^2 . With previous notation B is equal to $0, -D_{v_3}, -2D_{v_3}$. The set B' would be equal to $\pm D_{v_1} \pm D_{v_2} \pm D_{v_3}, \pm D_{v_1} \pm D_{v_2} \pm D_{v_3} - D_{v_3}, \pm D_{v_1} \pm D_{v_2} \pm D_{v_3} - 2D_{v_3}$. This gives us $\mathcal{O}(3), \mathcal{O}(2), \mathcal{O}(1), \mathcal{O}, \mathcal{O}(-1), \mathcal{O}(-2), \mathcal{O}(-3), \mathcal{O}(-4), \mathcal{O}(-5)$.

Proposition 3.4 For any smooth toric variety and any line bundle there exists an integer m_0 such that the push forward by the m th Frobenius morphism for any $m > m_0$ splits into the line bundles form B' .

Proof From 3.1 we know that the line bundles from the split are of the form $\left[\frac{D}{m} + \frac{D_{\chi}}{m}\right]$, where $L = \mathcal{O}(D)$ is a fixed representation of L . Of course for sufficiently large m all coefficients of $\frac{D}{m}$ belong to the interval $(-1, 1)$, so the coefficients of $\left[\frac{D}{m} + \frac{D_{\chi}}{m}\right]$ differ by at most one from the coefficients of $\left[\frac{D_{\chi}}{m}\right]$ that is in B , so in fact $\left[\frac{D}{m} + \frac{D_{\chi}}{m}\right] \in B'$.

This combined with the result of Thomsen [27] that the push forward and the line bundle are isomorphic as sheaves or abelian groups gives us the following result:

Corollary 3.5 There exists a finite set, namely B' , such that each line bundle is isomorphic as a sheaf of abelian groups to a direct sum of line bundles from B' . In particular their cohomologies agree.

3.2 Techniques of counting homology

Our aim will be to describe line bundles on toric varieties with vanishing higher cohomologies, that we call acyclic. Later, we will use this characterization to check if $Ext^i(L, M) = H^i(L^{\vee} \otimes M)$ is equal to zero for $i > 0$. We start with general remarks on cohomology of line bundles on smooth, complete toric varieties.

Let Σ be a fan in $N = \mathbb{Z}^n$ with rays x_1, \dots, x_m and let \mathbb{P}_{Σ} denote the variety constructed from the fan Σ . For $I \subset \{1, \dots, m\}$ let C_I be a simplicial complex generated by sets $J \subset I$ such that $\{x_i : i \in J\}$ generate a cone in Σ . For $r = (r_i : i = 1, \dots, m)$ let us define $Supp(r) := C_{\{i: r_i \geq 0\}}$.

The proof of the following well known fact can be found in the paper [6]:

Proposition 3.6 The cohomology $H^j(\mathbb{P}_{\Sigma}, L)$ is isomorphic to the direct sum over all $r = (r_i : i = 1, \dots, m)$ such that $\mathcal{O}(\sum_{i=1}^m r_i D_{x_i}) \cong L$ of the $(n - j)$ th reduced homology of the simplicial complex $Supp(r)$.

Definition 3.7 We call a line bundle L on \mathbb{P}_Σ acyclic if $H^i(\mathbb{P}_\Sigma, L) = 0$ for all $i \geq 1$.

Definition 3.8 For a fixed fan Σ we call a proper subset I of $\{1, \dots, m\}$ a forbidden set if the simplicial complex C_I has nontrivial reduced homology.

From Proposition 3.6 we have the following characterization of acyclic line bundles:

Proposition 3.9 *A line bundle L on \mathbb{P}_Σ is acyclic if it is not isomorphic to any of the following line bundles*

$$\mathcal{O} \left(\sum_{i \in I} r_i D_{x_i} - \sum_{i \notin I} (1 + r_i) D_{x_i} \right)$$

where $r_i \geq 0$ and I is a proper forbidden subset of $\{1, \dots, m\}$.

Hence to determine which bundles on \mathbb{P}_Σ are acyclic it is enough to know which sets I are forbidden.

In our case $C_I = \{J \subset I : \widehat{Y}_i := \{j : x_j \in Y_i\} \not\subseteq J \text{ for } i = 1, \dots, 5\}$, since Y_i are primitive collections. We call sets \widehat{Y}_i also primitive collections. The only difference between sets \widehat{Y}_i and Y_i is that the first one is the set of indices of rays in the second one, so in fact they could be even identified.

In case of a simplicial complex S on the set of vertices V we also define a primitive collection as a minimal subset of vertices that do not form a simplex. Complex S is determined by its primitive collections, namely it contains simplexes (subsets of V) that contain none of primitive collections.

We describe a very powerful method of counting homologies of simplicial complexes which are given by their primitive collections (as in our case). We use the result of Mrozek and Batko [24]:

Lemma 3.10 *Let X be a simplicial complex and let Z be a cycle in the chain complex whose boundary B is exactly one simplex. Then we can remove the pair (Z, B) from the chain complex without changing the homology.*

Definition 3.11 Let X be a simplicial complex defined by its set of primitive collections P on the set of vertices V . We say that simplicial complex X' on the set of vertices $V \setminus P$ is obtained from X by delating a primitive collection P if the set of primitive collections of X' is equal to the set of minimal sets in $\{Q \cap (X \setminus P) : Q \in P\}$.

Lemma 3.12 *Let X be a simplicial complex and suppose that there exists an element x which belongs to exactly one primitive collection P . Let $m = |P|$ and let X' be a simplicial complex obtained from X by delating P , then*

$$h^i(X) = h^{i-m+1}(X').$$

Proof Using Lemma 3.10 we will be removing subsequently on dimension reductive pairs (Z, B) such that $x \in Z$. We start from $(\{x\}, \emptyset)$. One can see that in each dimension we can take all $(Z, Z \setminus \{x\})$ for Z containing x as reductive pairs. Let us consider all simplexes of X that do not contain $P \setminus \{x\}$. One can prove by induction on dimension that we will remove all of them:

Let D be a simplex. If it contains x , than it will be removed as a first element of a reductive pair. If it does not, then $D \cup \{x\}$ is also a simplex of X and we will remove $(D \cup \{x\}, D)$.

We see that our simplicial complex can be reduced to a complex with simplexes containing $P \setminus \{x\}$. Now one immediately sees that such a complex is isomorphic to a complex X' (with a degree shifted by $|P \setminus \{x\}| = m - 1$).

The same method allows us to easily compute homologies when there are few primitive collections and many points. The idea is that we can glue together points that are in exactly the same primitive collections.

Definition 3.13 Let X be a simplicial complex defined by its set of primitive collections P on the set of vertices V . Suppose that there exist two points $x, y \in X$ such that they belong to the same primitive collections. We say that a simplicial complex X' on the set of vertices $V \setminus \{y\}$ is obtained from X by gluing points x and y if the set of primitive collections of X' is equal $\{Q \setminus \{y\} : Q \in P\}$. We can think of it like x was in fact two points x, y .

Proposition 3.14 Let X be a simplicial complex and suppose that there exist two points $x, y \in X$ such that they belong to the same primitive collections. Let X' be a simplicial complex obtained from X by gluing points x and y , then

$$h^i(X) = h^{i-1}(X').$$

Proof In both complexes we will be removing reductive pairs of the form (Z, B) with $x \in Z$ just as in Lemma 3.12. In both situations all that is left are simplexes that contain a set of a form $P \setminus \{x\}$, where P is a primitive collection containing x . In this situation all of simplexes of X that are left contain y and they can be identified with simplexes of X' that are left, the maps are exactly the same what finishes the proof.

Corollary 3.15 Let X be a simplicial complex on the set of vertices V . Let X' be a simplicial complex obtained from X by gluing equivalence classes of the relation \sim that identifies elements that are in exactly the same primitive collections. Suppose $|V| - |V/\sim| = m$, then

$$h^i(X) = h^{i-m}(X').$$

Proof We use Proposition 3.14 for pairs of points in the equivalence classes.

Corollary 3.16 In the situation of Lemma 3.12 and Corollary 3.15 X is acyclic if and only if X' is acyclic.

With these tools we are ready to determine forbidden subsets. In general we have got two following lemmas:

Lemma 3.17 If a nonempty subset I is not a sum of primitive collections, then it is not forbidden.

Proof There exists $a \in I$ such that a does not belong to any primitive collection which is contained in I . Using Lemma 3.10 we can remove subsequently on dimension reductive pairs (Z, B) such that $a \in Z$. We start from $(\{a\}, \emptyset)$. One can see that in this way we remove all of simplexes and as a consequence the chain complex is exact.

Lemma 3.18 A primitive collection is a forbidden subset.

Proof Using Lemma 3.12 we can remove this primitive collection and get a complex consisting of the empty set only that has nontrivial reduced homologies.

This can be also seen from the fact that the considered complex topologically is a sphere.

The following lemmas apply to the case when the Picard number is three and we have five primitive collections as in Batyrev’s classification. Let us remind that primitive collections of simplicial complex in this case are $\widehat{Y}_i := \{j : x_j \in Y_i\}$, for our convenience we define also $\widehat{X}_i := \{j : x_j \in X_i\}$.

Lemma 3.19 *A sum of two consecutive primitive collections is a forbidden subset.*

Proof Using Lemma 3.12 we remove one primitive collection and get a situation of Lemma 3.18.

Lemma 3.20 *A sum of three consecutive primitive collections $\widehat{Y}_i, \widehat{Y}_{i+1}, \widehat{Y}_{i+2}$ is not a forbidden subset.*

Proof First we can remove primitive collection \widehat{Y}_i . The image of \widehat{Y}_{i+2} contains the image of \widehat{Y}_{i+1} , so in fact we are left with just one primitive collection P which is an image of \widehat{Y}_{i+1} . We can remove P and obtain a nonempty full simplicial complex which is known to have trivial homologies.

Above lemmas match together to the following

Theorem 3.21 *The only forbidden subsets are primitive collections, their complements and the empty set.*

This gives us that in our situation

Corollary 3.22 *A line bundle L is acyclic if and only if it is not isomorphic to any of the following line bundles*

$$\mathcal{O}(\alpha_1^1 D_{v_1} + \dots + \alpha_2^1 D_{y_1} + \dots + \alpha_3^1 D_{z_1} + \dots + \alpha_4^1 D_{t_1} + \dots + \alpha_5^1 D_{u_1} + \dots)$$

where exactly 2, 3 or 5 consecutive $\alpha_i := (\alpha_i^1, \dots, \alpha_i^{p_i})$ are all less or equal to -1 and the rest is nonnegative.

Proof It is an immediate consequence of Proposition 3.9 and Theorem 3.21

Corollary 3.23 *If all of the coefficients b and c are zero in the primitive relations from Theorem 2.4 then a line bundle L is acyclic if and only if it is not isomorphic to any of the following line bundles*

$$\mathcal{O}(\alpha_1 D_v + \alpha_2 D_y + \alpha_3 D_z + \alpha_4 D_t + \alpha_5 D_u)$$

where exactly 2, 3 or 5 consecutive α_i are negative and if $\alpha_i < 0$ then $\alpha_i \leq -|X_i|$.

Proof Since all divisors corresponding to elements of the set X_i are linearly equivalent we match them together and as a consequence α_i is the sum of all of their coefficients.

4 Main theorem

This section contains the main, new result of this work. We give an explicit construction of a full, strongly exceptional collection of line bundles in the derived category $D^b(X)$ for a large family of smooth, complete toric varieties X with Picard number three. Namely for varieties X whose sets X_1, X_3 and X_4 from Batyrev’s classification presented in Theorem 2.4 have only one element. We will use results from Sect. 3.

4.1 Our setting

In this section we establish a family of varieties which we consider in this section and we also fix notation.

From now on for the whole Section let X be a smooth, complete toric variety with Picard number three, which using the notation from Theorem 2.4 has $|X_1| = |X_3| = |X_4| = 1$.

Let $r = |X_2|$. Then of course $|X_0| = n - r$. We allow arbitrary nonnegative integer parameters $b := b_1, c_2, \dots, c_r$. This family generalizes one considered in [11] (there, the case $r = 1$ was considered) and [8] (there the case $b = c_1 = \dots = c_r = 0$ was considered).

Remark 4.1 A variety of this type is Fano iff

$$n - r > \sum_{i=2}^r c_i + b.$$

In what follows we do not restrict to the Fano case.

Let e_1, \dots, e_n be a basis of the lattice N . Let us write what are the coordinates of the ray generators in the considered situation:

$$\begin{aligned} v_1 &= e_1, & v_2 &= e_2, \dots, & v_{n-r} &= e_{n-r} \\ y &= -e_1 - \dots - e_{n-r} + c_2 e_{n-r+2} + \dots + c_r e_n - (b + 1)(e_{n-r+1} + \dots + e_n) \\ z_1 &= e_{n-r+1}, \dots, z_r = e_n \\ t &= -e_{n-r+1} - \dots - e_n \\ u &= -e_1 - \dots - e_{n-r} + c_2 e_{n-r+2} + \dots + c_r e_n - b(e_{n-r+1} + \dots + e_n) \end{aligned} \tag{4.1}$$

Let D_w be the divisor associated to the ray generator w . One can easily see that the divisors $D_{v_1}, \dots, D_{v_{n-r}}$ are all linearly equivalent. Let D_v be any their representant in the Picard group. The other equivalence relations that generate all the relations in the Picard group are:

$$\begin{aligned} D_v &\simeq D_u + D_y \\ D_{z_i} &\simeq D_t + bD_u + (b + 1)D_y \\ D_{z_i} &\simeq D_t + (b - c_i)D_u + (b - c_i + 1)D_y, \quad 2 \leq i \leq r \end{aligned} \tag{4.2}$$

From these relations we can easily deduce:

Proposition 4.2 *The Picard group of the variety X is isomorphic to \mathbb{Z}^3 and is generated by D_t, D_y, D_v .*

We introduce two sets of divisors. We claim that these sets can be ordered in such a way that line bundles corresponding to divisors from these sets form a strongly exceptional collection.

$$Col_1 = \{-sD_t - sD_y + (-(n - r) - bs + q)D_v : 0 \leq s \leq r, 0 \leq q \leq n - r\} \tag{4.3}$$

$$Col_2 = \{-sD_t - (s - 1)D_y + (-(n - r) - bs + q)D_v : 1 \leq s \leq r, 0 \leq q \leq n - r - 1\}$$

Definition 4.3 Let $Col = Col_1 \cup Col_2$.

Remark 4.4 Let us notice that $|Col_1| = (r + 1)(n - r + 1)$ and $|Col_2| = r(n - r)$, so $|Col| = 2rn - 2r^2 + n + 1$.

We calculate the number of maximal cones in the fan defining the variety X . In order to obtain a maximal cone we have to choose n ray generators that do not contain a primitive collection. This is equivalent to removing three ray generators in such a way that the rest do not contain a primitive collection. First let us notice that we can remove at most one element from each group X_i because otherwise the rest would contain a primitive collection. We have the following possibilities:

- (1) We remove one element from X_0 and X_2 . Then we have to remove one element from X_3 or X_4 . We have got $2(n - r)r$ such possibilities.
- (2) We remove one element from X_0 and none from X_2 . We have got $n - r$ such possibilities.
- (3) We remove one element from X_2 and none from X_0 . We have got r such possibilities.
- (4) We do not remove any elements from X_0 and from X_2 . We have got 1 such possibility.

All together we see that we have $2rn - 2r^2 + n + 1$ maximal cones. From the general theory we know that the rank of the Grothendieck group is the same. Let us notice that from Remark 4.4 our set Col is of the same number of elements.

4.2 Acyclicity of differences of line bundles from Col

In this section we order the set Col and prove that line bundles corresponding to divisors from Col form a strongly exceptional collection.

Let us first check that $\text{Ext}_{\mathcal{O}_X}^i(\mathcal{O}(D_1), \mathcal{O}(D_2)) = 0$ for any divisors D_1, D_2 from the set Col and for any $i > 0$. We know that

$$\text{Ext}_{\mathcal{O}_X}^i(\mathcal{O}(D_1), \mathcal{O}(D_2)) = H^i(\mathcal{O}(D_1)^\vee \otimes \mathcal{O}(D_2)) = H^i(\mathcal{O}(D_2 - D_1)).$$

This means that we have to show that all line bundles associated to differences of divisors from Col are acyclic.

Definition 4.5 Let $Diff$ be the set of all divisors of the form $D_1 - D_2$, where $D_1, D_2 \in Col$.

Proposition 4.6 The set $Diff$ is the sum of sets $Diff_1, Diff_2, Diff_3$, where:

$$Diff_1 = \{sD_t + sD_y + (bs + q)D_v : -r \leq s \leq r, r - n \leq q \leq n - r\}$$

$$Diff_2 = \{sD_t + (s - 1)D_y + (bs + q)D_v : -r + 1 \leq s \leq r, r - n + 1 \leq q \leq n - r\}$$

$$Diff_3 = \{sD_t + (s + 1)D_y + (bs + q)D_v : -r \leq s \leq r - 1, r - n \leq q \leq n - r - 1\}.$$

Proof The set $Diff_1$ is equal to the set of all possible differences of two divisors from Col_1 and this set contains all possible differences of two divisors from Col_2 . The set $Diff_2$ is the set of all possible differences of the form $D_1 - D_2$, where $D_1 \in Col_1, D_2 \in Col_2$. The set $Diff_3$ is equal to $-Diff_2$ and so it is equal to the set of all differences of the form $D_2 - D_1$, where $D_1 \in Col_1, D_2 \in Col_2$. These are of course all possible differences of two elements from Col .

From the Corollary 3.22 we know that it is enough to prove that elements of $Diff$ are not of the form

$$\alpha_1 D_v + \alpha_2 D_y + \alpha_3^1 D_{z_1} + \alpha_3^2 D_{z_2} + \dots + \alpha_3^r D_{z_r} + \alpha_4 D_t + \alpha_5 D_u,$$

where exactly two, three or five consecutive α_i 's are negative (we call a number positive when it is nonnegative and consider only two signs positive and negative) and:

- (1) if $\alpha_1 < 0$, then $\alpha_1 \leq -(n - r)$ (α_1 is in fact sum of all the coefficients of D_{v_i} , which have to be of the same sign),
- (2) if any $\alpha_3^i < 0$ then $\alpha_3^j < 0$ (all parameters α_3^j are treated as one group and have the same sign).

From now on we assume that these conditions on α_i 's are satisfied.

Using the relations 4.2 we obtain:

$$\begin{aligned} & \alpha_1 D_v + \alpha_2 D_y + \alpha_3^1 D_{z_1} + \alpha_3^2 D_{z_2} + \dots + \alpha_3^r D_{z_r} + \alpha_4 D_t + \alpha_5 D_u \\ &= \left(\alpha_4 + \sum_{j=1}^r \alpha_3^j \right) D_t + \left(\alpha_2 - \alpha_5 + \sum_{j=1}^r \alpha_3^j \right) D_y \\ &+ \left(\alpha_1 + b\alpha_3^1 + \sum_{j=2}^r (b - c_j)\alpha_3^j + \alpha_5 \right) D_v \end{aligned} \tag{4.4}$$

Lemma 4.7 *If the elements α_3^j are negative then the divisors from $Diff$ are not of the form (4.4).*

Proof If α_4 was negative, then the coefficient of D_t would be less than or equal to $-r - 1$ and none of the divisors from $Diff$ has got such a coefficient, so α_4 has to be positive. Since α_3 is negative and α_4 is positive, then α_2 has to be negative and α_5 has to be positive. This means that the coefficient of D_y is less then or equal to $-r - 1$. The divisors from $Diff$ are not of this form.

From now on we may assume that α_3 is positive.

Lemma 4.8 *The divisors from $Diff_1$ are not of the form (4.4).*

Proof Suppose that a divisor from $Diff_1$ can be written in a form (4.4). We have:

$$\alpha_4 + \sum_{j=1}^r \alpha_3^j = \alpha_2 - \alpha_5 + \sum_{j=1}^r \alpha_3^j,$$

so $\alpha_4 + \alpha_5 = \alpha_2$. But α_2, α_4 and α_5 cannot be of the same sign, so α_4 and α_5 have to have different signs. As α_3 was positive we see that α_4 is positive, so α_5 and α_1 are negative. Let us notice that:

$$\begin{aligned} & \alpha_1 + b\alpha_3^1 + \left(\sum_{j=2}^r (b - c_j)\alpha_3^j \right) + \alpha_5 \\ & \leq -n + r + b \left(\sum_{j=1}^r \alpha_3^j \right) - 1 \\ & \leq -n + r - 1 + b \left(\alpha_4 + \sum_{j=1}^r \alpha_3^j \right) \end{aligned}$$

This shows precisely that the coefficient of D_v is less than or equal to $-n + r - 1$ plus b times the coefficient of D_t . Let s be the coefficient of D_t . From the definition of $Diff_1$ the coefficient of D_v is at least $-n + r + bs$. This gives us a contradiction.

Lemma 4.9 *The divisors from $Diff_3$ are not of the form (4.4).*

Proof Suppose that a divisor from $Diff_3$ can be written in a form (4.4). We have:

$$\alpha_4 + \sum_{j=1}^r \alpha_3^j = \alpha_2 - \alpha_5 - 1 + \sum_{j=1}^r \alpha_3^j,$$

so $\alpha_4 + \alpha_5 = \alpha_2 - 1$. The rest of the proof is identical to the proof of Lemma 4.8.

Lemma 4.10 *The divisors from $Diff_2$ are not of the form (4.4).*

Proof Suppose that a divisor from $Diff_2$ can be written in a form (4.4). We have:

$$\alpha_4 + \sum_{j=1}^r \alpha_3^j = \alpha_2 - \alpha_5 + 1 + \sum_{j=1}^r \alpha_3^j,$$

so $\alpha_4 + \alpha_5 = \alpha_2 + 1$. But α_2, α_4 and α_5 cannot be of the same sign, so we have two possible cases:

- (1) The coefficients α_4 and α_5 have different signs. In this case the proof is the same as in Lemmas 4.8 and 4.9.
- (2) We have $\alpha_4 = \alpha_5 = 0$ and $\alpha_2 = -1$. In this case α_1 has to be negative, because α_3 was positive. Let $s = \alpha_4 + \sum_{j=1}^r \alpha_3^j$ be the coefficient of D_t . We have:

$$\alpha_1 + b\alpha_3^1 + \sum_{j=2}^r (b - c_j)\alpha_3^j + \alpha_5 \leq -n + r + bs,$$

so the coefficient of D_v is less than or equal to $-n + r + bs$. But from the definition of $Diff_2$ we know that the coefficient of D_v is at least $bs + r - n + 1$ what gives us a contradiction.

Now we only have to order the line bundles corresponding to divisors from Col in such a way that

$$0 = \text{Ext}_{\mathcal{O}_X}^0(\mathcal{O}(D_1), \mathcal{O}(D_2)) = H^0(\mathcal{O}(D_1)^\vee \otimes \mathcal{O}(D_2)) = H^0(\mathcal{O}(D_2 - D_1)).$$

for any divisors $D_1 > D_2$.

Let us define the order by: $L_{s,q} < L'_{s,q} < L_{s,q+1}, L_{s+1,q_1} < L_{s,q_2}$ where

$$L_{s,q} = \mathcal{O}(-sD_t - sD_y + (q - bs - (n - r))D_v)$$

for $s = 0, \dots, r$ and $q = 0, \dots, n - r$ and

$$L'_{s,q} = \mathcal{O}(-sD_t - (s - 1)D_y + (q - bs - (n - r))D_v)$$

for $s = 1, \dots, r - 1$ and $q = 0, \dots, n - r - 1$. It is easy to see that zero cohomology of appropriate differences vanish.

4.3 Generating the derived category

We prove that the strongly exceptional collection from Sect. 4.1 is also full. First we show that it generates all line bundles. Due to [5, Corollary 4.8] the collection generates the derived category. In order to generate all line bundles we need several lemmas:

Lemma 4.11 *Let s and k be any integers. Line bundles $L_q = \mathcal{O}(-sD_t - sD_y + (k + q)D_v)$ for $q = 0, \dots, n - r$ and $L'_q = \mathcal{O}(-sD_t - (s - 1)D_y + (k + q)D_v)$ for $q = 0, \dots, n - r - 1$ generate $\mathcal{O}(-sD_t - (s - 1)D_y + (n - r + k)D_v)$ in the derived category.*

Proof We consider the Koszul complex for $\mathcal{O}(D_y), \mathcal{O}(D_{v_1}), \dots, \mathcal{O}(D_{v_{n-r}})$:

$$0 \rightarrow \mathcal{O}(-D_y - (n - r)D_v) \rightarrow \dots \rightarrow \mathcal{O}(-D_v)^{n-r} \oplus \mathcal{O}(-D_y) \rightarrow \mathcal{O} \rightarrow 0.$$

By tensoring it with $\mathcal{O}(-sD_t - (s - 1)D_y + (k + n - r)D_v)$ we obtain:

$$0 \rightarrow \mathcal{O}(-sD_t - sD_y + kD_v) \rightarrow \dots \rightarrow \mathcal{O}(-sD_t - (s - 1)D_y + (k + n - r - 1)D_v)^{n-1} \oplus \mathcal{O}(-sD_t - sD_y + (k + n - r)D_v) \rightarrow \mathcal{O}(-sD_t - (s - 1)D_y + (k + n - r)D_v) \rightarrow 0.$$

All sheaves that appear in this exact sequence, apart from the last one, are exactly $\mathcal{O}(-sD_t - sD_y + kD_v), \dots, \mathcal{O}(-sD_t - sD_y + (k + n - r)D_v), \mathcal{O}(-sD_t - (s - 1)D_y + kD_v), \dots, \mathcal{O}(-sD_t - (s - 1)D_y + (k + n - r - 1)D_v)$, so indeed we can generate $\mathcal{O}(-sD_t - (s - 1)D_y + (k + n - r)D_v)$.

Lemma 4.12 *Let s and k be any integers. Line bundles $L_q = \mathcal{O}(-sD_t - sD_y + (k + q)D_v)$ for $q = 0, \dots, n - r$ and $L'_q = \mathcal{O}(-sD_t - (s - 1)D_y + (k + q)D_v)$ for $q = 1, \dots, n - r$ generate $\mathcal{O}(-sD_t - (s - 1)D_y + kD_v)$ in the derived category.*

Proof The proof is similar to the last one. We deduce assertion from the same exact sequence of sheaves.

Lemma 4.13 *Let s and k be any integers. Line bundles $L_q = \mathcal{O}(-sD_t - sD_y + (k + q)D_v)$ for $q = 1, \dots, n - r$ and $L'_q = \mathcal{O}(-sD_t - (s - 1)D_y + (k + q)D_v)$ for $q = 0, \dots, n - r$ generate $\mathcal{O}(-sD_t - sD_y + (n - r + k + 1)D_v)$ in the derived category.*

Proof The proof is similar to the first one. We have to consider the Koszul complex for line bundles $\mathcal{O}(D_u), \mathcal{O}(D_{v_1}), \dots, \mathcal{O}(D_{v_{n-r}})$:

$$0 \rightarrow \mathcal{O}(-D_u - (n - r)D_v) \rightarrow \dots \rightarrow \mathcal{O}(-D_v)^{n-r} \oplus \mathcal{O}(-D_u) \rightarrow \mathcal{O} \rightarrow 0$$

we dualize it and we tensor it with $\mathcal{O}(-sD_t - (s - 1)D_y + kD_v)$.

Lemma 4.14 *Let s and k be any integers. Line bundles $L_q = \mathcal{O}(-sD_t - sD_y + (k + q)D_v)$ for $q = 1, \dots, n - r + 1$ and $L'_q = \mathcal{O}(-sD_t - (s - 1)D_y + (k + q)D_v)$ for $q = 1, \dots, n - r$ generate $\mathcal{O}(-sD_t - sD_y + kD_v)$ in the derived category.*

Proof The proof is similar to the last one. We deduce assertion from the same exact sequence of sheaves.

Lemma 4.15 *Let s and k be any integers. Line bundles $L_q = \mathcal{O}(-sD_t - sD_y + (k + q)D_v)$ for $q = 0, \dots, n - r$ and $L'_q = \mathcal{O}(-sD_t - (s - 1)D_y + (k + q)D_v)$ for $q = 0, \dots, n - r - 1$ generate in the derived category line bundles $\mathcal{O}(-sD_t - sD_y + q'D_v)$ and $\mathcal{O}(-sD_t - (s - 1)D_y + q'D_v)$ for an arbitrary integer q' .*

Proof We prove it by induction on $|q'|$. For $q' \geq k + n - r$ we use Lemmas 4.11 and 4.13, for $q' < k$ we use Lemmas 4.12 and 4.14.

Lemma 4.16 *Let k be any integer. Line bundles $L_{s,q} = \mathcal{O}(-sD_t - sD_y + qD_v)$ for $s = k, \dots, k + r$ and arbitrary q and $L'_{s,q} = \mathcal{O}(-sD_t - (s - 1)D_y + qD_v)$ for $s = k, \dots, k + r - 1$ and arbitrary q generate in the derived category line bundles $L'(k + r, q) = \mathcal{O}(-(k + r)D_t - (k + r - 1)D_y + qD_v)$ with arbitrary q .*

Proof Consider the Koszul complex for $\mathcal{O}(D_y), \mathcal{O}(D_{z_1}), \dots, \mathcal{O}(D_{z_r})$:

$$0 \rightarrow \mathcal{O}(-D_{z_1} - (r - 1)D_{z_2} - D_y) \rightarrow \dots \rightarrow \mathcal{O}(-D_{z_1}) \oplus \mathcal{O}(-D_{z_2})^{r-1} \oplus \mathcal{O}(-D_y) \rightarrow \mathcal{O} \rightarrow 0.$$

After tensoring it with $\mathcal{O}(-(k - 1)D_y + q'D_v)$ for appropriate q' we get the assertion.

Lemma 4.17 *Let k be any integer. Line bundles $L_{s,q} = \mathcal{O}(-sD_t - sD_y + qD_v)$ for $s = k, \dots, k+r$ and arbitrary q and $L'_{s,q} = \mathcal{O}(-sD_t - (s-1)D_y + qD_v)$ for $s = k+1, \dots, k+r$ and arbitrary q generate in the derived category line bundles $L'(k, q) = \mathcal{O}(-kD_t - (k-1)D_y + qD_v)$ for arbitrary q .*

Proof The proof is similar to the last one. We deduce assertion from the same exact sequence of sheaves.

Lemma 4.18 *Let k be any integer. Line bundles $L_{s,q} = \mathcal{O}(-sD_t - sD_y + qD_v)$ for $s = k + 1, \dots, k + r$ and arbitrary q and $L'_{s,q} = \mathcal{O}(-sD_t - (s - 1)D_y + qD_v)$ for $s = k + 1, \dots, k + r + 1$ and arbitrary q generate in the derived category line bundles $L(k, q) = \mathcal{O}(-kD_t - kD_y + qD_v)$ for arbitrary q .*

Proof Consider the Koszul complex for $\mathcal{O}(D_{z_1}), \dots, \mathcal{O}(D_{z_r}), \mathcal{O}(D_t)$:

$$0 \rightarrow \mathcal{O}(-D_{z_1} - (r - 1)D_{z_2} - D_t) \rightarrow \dots \rightarrow \mathcal{O}(-D_{z_1}) \oplus \mathcal{O}(-D_{z_2})^{r-1} \oplus \mathcal{O}(-D_t) \rightarrow \mathcal{O} \rightarrow 0.$$

After tensoring it with $\mathcal{O}(-kD_y + q'D_v)$ for appropriate q' we get the assertion.

Lemma 4.19 *Let k be any integer. Line bundles $L_{s,q} = \mathcal{O}(-sD_t - sD_y + qD_v)$ for $s = k, \dots, k+r$ and arbitrary q and $L'_{s,q} = \mathcal{O}(-sD_t - (s-1)D_y + qD_v)$ for $s = k+1, \dots, k+r$ and arbitrary q generate in the derived category line bundles $L'(k + r + 1, q) = \mathcal{O}(-(k + r + 1)D_t - (k + r)D_y + qD_v)$ for arbitrary q .*

Proof The proof is similar to the last one. We deduce assertion from the same exact sequence of sheaves.

Lemma 4.20 *Let k be any integer. Line bundles $L_{s,q} = \mathcal{O}(-sD_t - sD_y + qD_v)$ for $s = k, \dots, k+r$ and arbitrary q and $L'_{s,q} = \mathcal{O}(-sD_t - (s-1)D_y + qD_v)$ for $s = k, \dots, k+r-1$ and arbitrary q generate in the derived category line bundles $L(s, q) = \mathcal{O}(-sD_t - sD_y + qD_v)$ and $L'(s, q) = \mathcal{O}(-sD_t - (s - 1)D_y + qD_v)$ for arbitrary s and q .*

Proof We prove it by induction on $|s|$. For $s \geq k + n - r$ we use Lemmas 4.16 and 4.19, for $r < k$ we use Lemmas 4.17 and 4.18.

Lemma 4.21 *Let k be any integer. Line bundles $\mathcal{O}(-sD_t - (s+k)D_y + qD_v)$ and $\mathcal{O}(-sD_t - (s + k + 1)D_y + qD_v)$ for arbitrary s and q generate in the derived category line bundles $\mathcal{O}(-sD_t - (s + k + 2)D_y + qD_v)$ for arbitrary s and q .*

Proof Consider the Koszul complex for $\mathcal{O}(D_t), \mathcal{O}(D_u)$:

$$0 \rightarrow \mathcal{O}(-D_t - D_u) \rightarrow \mathcal{O}(-D_t) \oplus \mathcal{O}(-D_u) \rightarrow \mathcal{O} \rightarrow 0.$$

After tensoring it with $\mathcal{O}(-k'D_y + q')$ for appropriate k' and q' we get the assertion.

Lemma 4.22 *Let k be any integer. Line bundles $\mathcal{O}(-sD_t - (s+k)D_y + qD_v)$ and $\mathcal{O}(-sD_t - (s+k+1)D_y + qD_v)$ for arbitrary s and q generate in the derived category line bundles $\mathcal{O}(-sD_t - (s+k-1)D_y + qD_v)$ for arbitrary s and q .*

Proof Consider the Koszul complex for $\mathcal{O}(D_t), \mathcal{O}(D_u)$:

$$0 \rightarrow \mathcal{O}(-D_t - D_u) \rightarrow \mathcal{O}(-D_t) \oplus \mathcal{O}(-D_u) \rightarrow \mathcal{O} \rightarrow 0.$$

After tensoring it with $\mathcal{O}(-k'D_y + q')$ for appropriate k' and q' we get the assertion.

Proposition 4.23 *Line bundles*

$$L_{s,q} = \mathcal{O}(-sD_t - sD_y + (q - bs - (n - r))D_v)$$

for $s = 0, \dots, r$ and $q = 0, \dots, n - r$ and

$$L'_{s,q} = \mathcal{O}(-sD_t - (s - 1)D_y + (q - bs - (n - r))D_v)$$

for $s = 0, \dots, r - 1$ and $q = 0, \dots, n - r - 1$ generate in the derived category all line bundles.

Proof We use Lemmas 4.15, 4.20, 4.21 and 4.22.

Summarizing, we have proved:

Theorem 4.24 *Let X be a smooth, complete, n dimensional toric variety with Picard number three and the set of ray generators $X_0 \cup \dots \cup X_4$, where*

$$X_0 = \{v_1, \dots, v_{n-r}\}, \quad X_1 = \{y\}, \quad X_2 = \{z_1, \dots, z_r\}, \quad X_3 = \{t\}, \quad X_4 = \{u\},$$

primitive collections $X_0 \cup X_1, X_1 \cup X_2, \dots, X_4 \cup X_0$ and primitive relations:

$$\begin{aligned} v_1 + \dots + v_{n-r} + y - cz_2 - \dots - cz_r - (b + 1)t &= 0, \\ y + z_1 + \dots + z_r - u &= 0, \\ z_1 + \dots + z_r + t &= 0, \\ t + u - y &= 0, \\ u + v_1 + \dots + v_{n-r} - c_2z_2 - \dots - c_rz_r - bt &= 0, \end{aligned}$$

where b and c are positive integers.

Then the ordered collection of line bundles

$$L_{s,q} = \mathcal{O}(-sD_t - sD_y + (q - bs - (n - r))D_v)$$

for $s = 0, \dots, r$ and $q = 0, \dots, n - r$ and

$$L'_{s,q} = \mathcal{O}(-sD_t - (s - 1)D_y + (q - bs - (n - r))D_v)$$

for $s = 0, \dots, r - 1$ and $q = 0, \dots, n - r - 1$ where the order is defined by $L_{s,q} < L'_{s,q} < L_{s,q+1}, L_{s+1,q_1} < L_{s,q_2}$ is a full, strongly exceptional collection of line bundles.

Proof From Sect. 4.2 we already know that this is a strongly exceptional collection. We have just checked the sufficient condition for fullness in Proposition 4.23.

5 Bondal’s construction not containing a full, strongly exceptional collection

5.1 Example

Let us consider the case when:

$$\begin{aligned} X_0 &= \{v_1\}, & X_1 &= \{y_1, \dots, y_k\}, & X_2 &= \{z_1\}, \\ X_3 &= \{t_1, \dots, t_k\}, & X_4 &= \{u_1, \dots, u_k\} \end{aligned}$$

then we can take

$$v_1, y_2, \dots, y_k, t_1, \dots, t_k, u_2, \dots, u_k$$

to be a basis of the lattice $N = \mathbb{Z}^{3k-1}$. Other vectors are like in 2.2 with all coefficients b_i and c_i equal to zero. We have linear dependencies of divisors:

$$D_{v_1} = D_{u_1} + D_{y_1}, \quad D_{t_i} = D_{z_1} + D_{y_1}, \quad D_{y_i} = D_{y_1}, \quad D_{u_i} = D_{u_1}$$

Let B be the image of the real torus in the Picard group as described in the Sect. 3.1. One can easily see that:

$$\begin{aligned} B &= \left\{ \mathcal{O} \left(\left[\sum_{i=1}^k -\alpha_t^i \right] D_{z_1} + \left[\sum_{i=2}^k -\alpha_u^i - \alpha_v^1 \right] D_{u_1} + [-\alpha_v^1 + \sum_{i=2}^k -\alpha_y^i + \sum_{i=1}^k \alpha_t^i] D_{y_1} \right) : \right. \\ &\quad \left. 0 \leq \alpha_v^i, \alpha_y^i, \alpha_t^i, \alpha_u^i < 1 \right\}. \end{aligned}$$

So B is contained in the set:

$$\begin{aligned} S &:= \{ \mathcal{O}(-aD_{z_1} - bD_{u_1} + (a - c)D_{y_1}) : a, b, c \in \{0, \dots, k\} \} \\ &= \{ \mathcal{O}(-a(D_{z_1} - D_{y_1}) - bD_{u_1} - cD_{y_1}) : a, b, c \in \{0, \dots, k\} \}. \end{aligned}$$

From Corollary 3.23 we know that line bundle is acyclic if and only if it is not isomorphic to any of the following line bundles

$$\begin{aligned} &\mathcal{O}(\alpha_1 D_{v_1} + \alpha_2 D_{y_1} + \alpha_3 D_{z_1} + \alpha_4 D_{t_1} + \alpha_5 D_{u_1}) \\ &= \mathcal{O}((\alpha_3 + \alpha_4)(D_{z_1} - D_{y_1}) + (\alpha_1 + \alpha_2 + \alpha_3)D_{y_1} + (\alpha_1 + \alpha_5)D_{u_1}), \end{aligned}$$

where exactly 2, 3 or 5 consecutive α are negative and if $\alpha_2 < 0$ then $\alpha_2 \leq -k$, if $\alpha_4 < 0$ then $\alpha_4 \leq -k$ and if $\alpha_5 < 0$ then $\alpha_5 \leq -k$. Let us observe that line bundles form the set

$$R = \left\{ \mathcal{O}(a(D_{z_1} - D_{y_1}) + bD_{y_1} + cD_{u_1}) : (a, b, c) \in \left[\frac{k}{2}, k \right] \times \left[-k, -\frac{k}{2} - 1 \right] \times [0, k] \right\}$$

are not acyclic. Indeed fixing $\alpha_1 = -k, \alpha_3 = \frac{k}{2}$ and taking α_4, α_5 nonnegative and α_2 negative we can achieve all of them. Let us define the set of pairs

$$\begin{aligned} P &:= \left\{ -\left(\frac{k}{2} + \frac{a}{2}\right) (D_{z_1} - D_{y_1}) - \left(\frac{k}{2} + \frac{b}{2}\right) D_{y_1} - \left(\frac{k}{2} + \frac{c}{2}\right) D_{u_1}, -\left(\frac{k}{2} - \frac{a}{2}\right) (D_{z_1} - D_{y_1}) \right. \\ &\quad \left. - \left(\frac{k}{2} - \frac{b}{2}\right) D_{y_1} - \left(\frac{k}{2} - \frac{c}{2}\right) D_{u_1} : (a, b, c) \in \left[\frac{k}{2}, k \right] \times \left[-k, -\frac{k}{2} - 1 \right] \times [0, k] \right\}. \end{aligned}$$

It is easy to see that elements of these pairs are distinct and they belong to S . Difference in each pair is an element of R so it is not acyclic line bundle. Hence to have a strongly

exceptional collection C in S we have to exclude at least one element from each pair. To have integer coefficients of divisors in P we should take $a \equiv b \equiv c \equiv k \pmod{2}$, so we have to throw out at least $\frac{k^3}{32}$ elements among $(k + 1)^3$ elements in S . Full, strongly exceptional collection has to have l elements, where l is the rank of the Grothendick group $K^0(X)$ (for toric varieties it is isomorphic to \mathbb{Z}^l , where l is the number of maximal cones). In our case there are at least k^3 maximal cones, since each time we throw out one element from X_2, X_4 and X_5 we get different maximal cone (exact number is $k^3 + 2k^2 + 2k$). So we have proven the following:

Theorem 5.1 *If $(k + 1)^3 - \frac{1}{32}k^3 < k^3 + 2k^2 + 2k$, what is when $k > 32$, then there is no full, strongly exceptional collection among line bundles that come from Bondal’s construction.*

Remark 5.2 Notice that the considered variety is Fano, so is expected to have a full, strongly exceptional collection.

5.2 Our case

Let us consider the case from Sect. 4.1, but with all coefficients c_i equal to $c \leq b$. Let B be the image of the real torus in the Picard group as described in the Sect. 3.1. One can see that:

$$B = \left\{ \mathcal{O} \left(\left[\sum_{i=1}^r -\alpha_z^i \right] D_t + \left[\sum_{i=1}^{n-r} -\alpha_v^i + c \sum_{i=2}^r \alpha_z^i - (b + 1) \sum_{i=1}^r \alpha_z^i \right] D_y + \left[\sum_{i=1}^{n-r} -\alpha_v^i + c \sum_{i=2}^r \alpha_z^i - b \sum_{i=1}^r \alpha_z^i \right] D_u \right) : 0 \leq \alpha_v^i, \alpha_z^i < 1 \right\}.$$

So B is contained in the set:

$$S := \{ \mathcal{O}(-sD_t - sD_y + qD_v), \mathcal{O}(-sD_t - (s - 1)D_y + qD_v) : s \in \{0, \dots, r\}, q \in \{-(n - r) - c - (b - c)s, \dots, (b - c)(-s + 1)\} \}$$

Our collection defined in Sect. 4.1, or its torsion, is contained in the set S unless $cr \leq b$. It can be also shown that if this inequality fails then there is no full strongly exceptional collection among line bundles that come from Bondal’s construction.

Acknowledgments We would like to thank very much Rosa Maria Miró-Roig and Laura Costa for introducing us to this subject. We are also grateful for many useful and interesting talks.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Batyrev, V.V.: On the classification of smooth projective toric varieties. *Tōhoku Math. J.* **43**, 569–585 (1991)
2. Bernardi, A., Tirabassi, S.: Derived Categories of Toric Fano 3-Folds via the Frobenius Morphism. <http://arxiv.org/abs/1002.1666> (preprint)
3. Bondal, A.I.: Derived categories of toric varieties. Oberwolfach Report 5/2006, pp. 284–286
4. Bondal, A.I.: Representation of associative algebras and coherent sheaves. *Math. USSR Izvestiya* **34**(1), 23–42 (1990)

5. Borisov, L., Horja, P.: On the K -theory of smooth toric DM stacks, Snowbird lectures on string geometry. *Contemp. Math.* **401**, 21–42 (2006)
6. Borisov, L., Hua, Z.: On the conjecture of King for smooth toric Deligne-Mumford stacks. *Adv. Math.* **221**, 277–301 (2009)
7. Caldararu, A.: Derived categories of sheaves: a skimming, Snowbird lectures in algebraic geometry. *Contemp. Math.* **388**, 43–75 (2005)
8. Costa, L., Miró-Roig, R.M.: Derived category of toric varieties with small Picard number (preprint)
9. Costa, L., Miró-Roig, R.M.: Tilting sheaves on toric varieties. *Math. Z.* **248**(4), 849–865 (2004)
10. Cox, D., Little, J., Schenck, H.: Toric Varieties. <http://www.cs.amherst.edu/~dac/toric.html> (preprint)
11. Dey, A., Lasoń, M., Michałek, M.: Derived category of toric varieties with Picard number three. *Le Matematiche* **64**(2), 99–116 (2009)
12. Efimov, A.: Maximal lengths of exceptional collections of line bundles. http://arxiv.org/PS_cache/arxiv/pdf/1010/1010.3755v2.pdf (preprint)
13. Fulton, W.: Introduction to Toric Varieties. In: *Annals of Mathematics Studies*. Princeton University Press, Princeton (1993)
14. Gelfand, S., Manin, Y.: *Methods of homological algebra*. Springer Monographs in Mathematics, 2nd edn. Springer, Berlin (2003)
15. Gorodentsev, A.L., Rudakov, A.N.: Exceptional vector bundles on projective spaces. *Duke Math. J.* **54**(1), 115–130 (1987)
16. Gronbaum, B.: *Convex Polytopes*. Wiley, London (1967)
17. Hille, L., Perling, M.: A counterexample to King’s conjecture. *Compos. Math.* **142**, 1507–1521 (2006)
18. Hille, L., Perling, M.: Exceptional Sequences of Invertible Sheaves on Rational Surfaces. http://arxiv.org/PS_cache/arxiv/pdf/0810/0810.1936v2.pdf (preprint)
19. Huybrechts, D.: *Fourier–Mukai Transforms in Algebraic Geometry*. Oxford Mathematical Monographs, Oxford (2006)
20. Kawamata, Y.: Derived categories of toric varieties. *Michigan Math. J.* **54**(3), 517–535 (2006)
21. King, A.: Tilting bundles on some rational surfaces. <http://www.maths.bath.ac.uk/~masadk/papers/> (preprint)
22. Kleinschmidt, P.: A classification of toric varieties with few generators. *Aequationes Math.* **35**, 254–266 (1988)
23. Michałek, M.: Family of counterexamples to King’s conjecture. *Comptes Rendus Math.* **349**(1–2), 67–69 (2011)
24. Mrozek, M., Batko, B.: Coreduction homology algorithm. *Discrete Comput. Geom.* **41**, 96–118 (2009)
25. Oda, T., Park, H.S.: Linear Gale transforms and Gelfand–Kapranov–Zelevinskij decompositions. *Tôhoku Math. J.* **43**, P375–399 (1991)
26. Perling, M.: Examples for exceptional sequences of invertible sheaves on rational surfaces. http://arxiv.org/PS_cache/arxiv/pdf/0904/0904.0529v2.pdf (preprint)
27. Thomsen, J.F.: Frobenius direct images of line bundles on toric varieties. *J. Algebra* **226**, 865–874 (2000)