



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: February 7, 2014

REVISED: March 20, 2014

ACCEPTED: April 1, 2014

PUBLISHED: April 22, 2014

The T_7 flavor symmetry in 3-3-1 model with neutral leptons

V.V. Vien^a and H.N. Long^b

^a*Department of Physics, Tay Nguyen University,
567 Le Duan, Buon Ma Thuot, Vietnam*

^b*Institute of Physics,
VAST, 10 Dao Tan, Ba Dinh, Hanoi, Vietnam*

E-mail: wvienk16@gmail.com, hnlng@iop.vast.ac.vn

ABSTRACT: We construct a 3-3-1 model based on non-Abelian discrete symmetry T_7 responsible for the fermion masses. Neutrinos get masses from only anti-sextets which are in triplets $\underline{3}$ and $\underline{3}^*$ under T_7 . The flavor mixing patterns and mass splitting are obtained without perturbation. The tribimaximal form obtained with the breaking $T_7 \rightarrow Z_3$ in charged lepton sector and both $T_7 \rightarrow Z_3$ and $Z_3 \rightarrow \{\text{Identity}\}$ must be taken place in neutrino sector but only apart in breakings $Z_3 \rightarrow \{\text{Identity}\}$ (without contribution of σ'), and the upper bound on neutrino mass $\sum_{i=1}^3 m_i$ at the level is presented. The Dirac CP violation phase δ is predicted to either $\frac{\pi}{2}$ or $\frac{3\pi}{2}$ which is maximal CP violation. From the Dirac CP violation phase we obtain the relation between Euler's angles which is consistent with the experimental in PDG 2012. On the other hand, the realistic lepton mixing can be obtained if both the direction for breakings $T_7 \rightarrow Z_3$ and $Z_3 \rightarrow \{\text{Identity}\}$ are taken place in neutrino sectors. The CKM matrix is the identity matrix at the tree-level.

KEYWORDS: Beyond Standard Model, Nonperturbative Effects, CP violation, Discrete and Finite Symmetries

ARXIV EPRINT: [1402.1256](https://arxiv.org/abs/1402.1256)

Contents

1	Introduction	1
2	Fermion content	4
3	Charged lepton masses	5
4	Quark masses	6
5	Neutrino mass and mixing	9
5.1	Normal case ($\Delta m_{23}^2 > 0$)	16
5.1.1	The case $\delta = \frac{\pi}{2}$	16
5.1.2	The case $\delta = \frac{3\pi}{2}$	19
5.2	Inverted case ($\Delta m_{23}^2 < 0$)	20
6	Remark on breaking, VEVs and rho parameter	23
7	Conclusions	24
A	T_7 group and Clebsch-Gordan coefficients	25
B	The numbers	26
C	The solutions with $\delta = \frac{\pi}{2}$ in the normal case	26
D	The solutions with $\delta = \frac{3\pi}{2}$ in the normal case	27

1 Introduction

Despite the great success of the Standard Model (SM) of the elementary particle physics, the origin of flavor structure, masses and mixings between generations of matter particles are unknown yet. The neutrino mass and mixing is one of the most important evidence of beyond Standard Model physics. Many experiments show that neutrinos have tiny masses and their mixing is still mysterious [1, 2].

The tri-bimaximal form for explaining the lepton mixing scheme was first proposed by Harrison-Perkins-Scott (HPS), which apart from the phase redefinitions, is given by [3–6]

$$U_{\text{HPS}} = \begin{pmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \end{pmatrix}, \tag{1.1}$$

can be considered as a good approximation for the recent neutrino experimental data.

The most recent data are a clear sign of rather large value θ_{13} [7]. The data in PDG2012 [8–12] imply:

$$\begin{aligned} \sin^2(2\theta_{12}) &= 0.857 \pm 0.024, & \sin^2(2\theta_{13}) &= 0.098 \pm 0.013, & \sin^2(2\theta_{23}) &> 0.95, \\ \Delta m_{21}^2 &= (7.50 \pm 0.20) \times 10^{-5} \text{eV}^2, & \Delta m_{32}^2 &= (2.32_{-0.08}^{+0.12}) \times 10^{-3} \text{eV}^2. \end{aligned} \quad (1.2)$$

These large neutrino mixing angles are completely different from the quark mixing ones defined by the Cabibbo-Kobayashi-Maskawa (CKM) matrix [13, 14]. This has stimulated work on flavor symmetries and non-Abelian discrete symmetries are considered to be the most attractive candidate to formulate dynamical principles that can lead to the flavor mixing patterns for quarks and lepton. There are many recent models based on the non-Abelian discrete symmetries, such as A_4 [15–32, 34], A_5 [35–47], S_3 [48–87], S_4 [88–116], D_4 [117–128], D_5 [129, 130], T' [131–140], T_7 [141–147] and so forth.

Among the possible extensions of SM, a curious choice are the 3-3-1 models which encompass a class of models based on the gauge group $SU(3)_C \otimes SU(3)_L \otimes U(1)_X$ [148–163], that is at first spontaneously broken to the SM group $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$ and then undergoes the spontaneously broken to $SU(3)_C \otimes U(1)_Q$. The extension of the gauge group with respect to SM leads to interesting consequences. The first one is that the requirement of anomaly cancelation together with that of asymptotic freedom of QCD implies that the number of generations must necessarily be equal to the number of colors, hence giving an explanation for the existence of three generations. Furthermore, quark generations should transform differently under the action of $SU(3)_L$. In particular, two quark generations should transform as triplets, one as an antitriplet.

A fundamental relation holds among some of the generators of the group [146, 147]:

$$Q = T_3 + \beta T_8 + X \quad (1.3)$$

where Q indicates the electric charge, T_3 and T_8 are two of the $SU(3)$ generators and X is the generator of $U(1)_X$. β is a key parameter that defines a specific variant of the model.

The model thus provides a partial explanation of the family number, as also required by flavor symmetries such as T_7 for 3-dimensional representations. In addition, due to the anomaly cancelation one family of quarks has to transform under $SU(3)_L$ differently from the two others. T_7 can meet this requirement with three inequivalent representations $\underline{1}, \underline{1}', \underline{1}''$. Note that T_7 has not been considered before in the kind of the 3-3-1 model.

There are two typical variants of the 3-3-1 models as far as lepton sectors are concerned. In the minimal version, three $SU(3)_L$ lepton triplets are (ν_L, l_L, l_R^c) , where l_R are ordinary right-handed charged-leptons [148–152]. In the second version, the third components of lepton triplets are the right-handed neutrinos, (ν_L, l_L, ν_R^c) [152–157]. To have a model with the realistic neutrino mixing matrix, we should consider another variant of the form (ν_L, l_L, N_R^c) where N_R are three new fermion singlets under standard model symmetry with vanishing lepton-numbers [164–167].

In our previous works [164–167], the discrete symmetries have been explored to the 3-3-1 models. In ref. [165] we have studied the 3-3-1 model with neutral fermions based on S_4 group, in which most of the Higgs multiplets are in triplets under S_4 except χ lying in

a singlet,¹ and the exact tribimaximal form [3–6] is obtained, where $\theta_{13} = 0$. As we know, the recent considerations have implied $\theta_{13} \neq 0$ [15–32, 48–116], but small as given in (1.2). This problem has been improved in this type of the model in ref. [166, 167] by adding new $SU(3)_L$ multiplets and one of them is regarded as a small perturbation. The model therefore contains up to eight Higgs multiplets, and the scalar potential of the model is quite complicated [166, 167].

CP violation plays a crucial role in our understanding of the observed baryon asymmetry of the Universe [168]. In the SM, CP symmetry is violated due to a complex phase in the CKM matrix [13, 14]. However, since the extent of CP violation in the SM is not enough for achieving the observed BAU, we need new source of CP violation for successful BAU. On the other hand, CP violations in the lepton sector are imperative if the BAU could be realized through leptogenesis. So, any hint or observation of the leptonic CP violation can strengthen our belief in leptogenesis [168].

The violation of the CP symmetry is a crucial ingredient of any dynamical mechanism which intends to explain both low energy CP violation and the baryon asymmetry. Renormalizable gauge theories are based on the spontaneous symmetry breaking mechanism, and it is natural to have the spontaneous CP violation as an integral part of that mechanism. Determining all possible sources of CP violation is a fundamental challenge for high energy physics. In theoretical and economical viewpoints, the spontaneous CP breaking necessary to generate the baryon asymmetry and leptonic CP violation at low energies brings us to a common source which comes from the phase of the scalar field responsible for the spontaneous CP breaking at a high energy scale [168].

In this paper, we investigate another choice with T_7 , the smallest group with two non-equivalent 3-dimensional irreducible representations, contains two triplet irreducible representations and three singlets which play a crucial role in consistently reproducing fermion masses and mixing. As we will see, T_7 model has some new features since fewer Higgs multiplets are needed in order to allow the fermions to gain masses and to break symmetries and the physics we will see is different from the former. The CP violation is the first time considered under $SU(3)_L \times U(1)_X$ model based on T_7 flavor symmetry in which the T_7 symmetry avoids the mass degeneracy of lepton masses. The light neutrino masses can be generated at tree level, and the vacuum alignment problem which arises in the presence of two T_7 -triplet scalar fields $\underline{3}, \underline{3}^*$ can naturally explain the measured value of θ_{13} and thereby the hierarchy of neutrino masses. The seesaw mechanism can explain the smallness of the measured neutrino masses and the maximal Dirac CP violation.

The rest of this work is organized as follows. In section 2 and 3 we present the necessary elements of the 3-3-1 model with the T_7 symmetry as well as introducing necessary Higgs fields responsible for the charged lepton masses. In section 4, we discuss on quark sector. Section 5 is devoted for the neutrino mass and mixing. We summarize our results and make conclusions in the section 7. Appendix A presents a brief of the T_7 theory. Appendix B provides the lepton number (L) and lepton parity (P_l) of particles in the model.

¹ χ is the unique singlet under S_4 defined in expression (25) in ref. [165].

2 Fermion content

The gauge symmetry is based on $SU(3)_C \otimes SU(3)_L \otimes U(1)_X$, where the electroweak factor $SU(3)_L \otimes U(1)_X$ is extended from those of the Standard Model (SM), and the strong interaction sector is retained. Each lepton family includes a new neutral fermion (N_R) with vanishing lepton number $L(N_R) = 0$ arranged under the $SU(3)_L$ symmetry as a triplet (ν_L, l_L, N_R^c) and a singlet l_R . The residual electric charge operator Q is therefore related to the generators of the gauge symmetry by

$$Q = T_3 - \frac{1}{\sqrt{3}}T_8 + X, \tag{2.1}$$

where T_a ($a = 1, 2, \dots, 8$) are $SU(3)_L$ charges with $\text{Tr}T_a T_b = \frac{1}{2}\delta_{ab}$ and X is the $U(1)_X$ charge. This means that the model under consideration does not contain exotic electric charges in the fundamental fermion, scalar and adjoint gauge boson representations.

Since the particles in the lepton triplet have different lepton number (1 and 0), so the lepton number in the model does not commute with the gauge symmetry unlike the SM. Therefore, it is better to work with a new conserved charge \mathcal{L} commuting with the gauge symmetry and related to the ordinary lepton number by diagonal matrices [164–167, 169]

$$L = \frac{2}{\sqrt{3}}T_8 + \mathcal{L}. \tag{2.2}$$

The lepton charge arranged in this way, i.e. $L(N_R) = 0$, is in order to prevent unwanted interactions due to $U(1)_{\mathcal{L}}$ symmetry and breaking due to the lepton parity to obtain the consistent lepton and quark spectra. By this embedding, exotic quarks U, D as well as new non-Hermitian gauge bosons X^0, Y^\pm possess lepton charges as of the ordinary leptons: $L(D) = -L(U) = L(X^0) = L(Y^-) = 1$.

Under the $[SU(3)_L, U(1)_X, U(1)_{\mathcal{L}}, T_7]$ symmetries as proposed, the fermions of the model transform as follows

$$\begin{aligned} \psi_L \equiv \psi_{1,2,3L} &= (\nu_L \quad l_L \quad N_R^c)^T \sim [3, -1/3, 2/3, \underline{3}], \\ l_{1R} &\sim [1, -1, 1, \underline{1}], \quad l_{2R} \sim [1, -1, 1, \underline{1}'], \quad l_{3R} \sim [1, -1, 1, \underline{1}''], \\ Q_{1L} \equiv (d_{1L} \quad -u_{1L} \quad D_{1L})^T &\sim [3^*, 0, 1/3, \underline{1}'], \\ Q_{2L} \equiv (d_{2L} \quad -u_{2L} \quad D_{2L})^T &\sim [3^*, 0, 1/3, \underline{1}''], \\ Q_{3L} &= (u_{3L} \quad d_{3L} \quad U_L)^T \sim [3, 1/3, -1/3, \underline{1}], \\ u_R &\sim u_{1,2,3R} = [1, 2/3, 0, \underline{3}], \quad d_R \sim [1, -1/3, 0, \underline{3}^*], \\ U_R &\sim [1, 2/3, -1, \underline{1}], \quad D_{1R} \sim [1, -1/3, 1, \underline{1}'], \quad D_{2R} \sim [1, -1/3, 1, \underline{1}']. \end{aligned} \tag{2.3}$$

where the subscript numbers on field indicate to respective families which also in order define components of their T_7 multiplets. U and $D_{1,2}$ are exotic quarks carrying lepton numbers $L(U) = -1$ and $L(D_{1,2}) = 1$, known as leptoquarks. In the following, we consider possibilities of generating the masses for the fermions. The scalar multiplets needed for the purpose are also introduced.

3 Charged lepton masses

The charged lepton masses arise from the couplings of $\bar{\psi}_L l_{1R}$, $\bar{\psi}_L l_{2R}$ and $\bar{\psi}_L l_{3R}$ to scalars, where $\bar{\psi}_L l_{iL}$ ($i = 1, 2, 3$) transforms as 3^* under $SU(3)_L$ and $\underline{3}^*$ under T_7 . To generate masses for charged leptons, we need a $SU(3)_L$ Higgs triplets that lying in $\underline{3}$ under T_7 and transforms as 3 under $SU(3)_L$,

$$\phi_i = \begin{pmatrix} \phi_{i1}^+ \\ \phi_{i2}^0 \\ \phi_{i3}^+ \end{pmatrix} \sim [3, 2/3, -1/3, \underline{3}] \quad (i = 1, 2, 3). \quad (3.1)$$

Following the potential minimization conditions, we have the followings alignments:

- (1) The first alignment: $\langle \phi_1 \rangle = \langle \phi_2 \rangle = \langle \phi_3 \rangle$ then T_7 is broken into Z_3 consisting of the elements $\{e, b, b^2\}$.
- (2) The second alignment: $\langle \phi_1 \rangle \neq \langle \phi_2 \rangle \neq \langle \phi_3 \rangle$ or $\langle \phi_1 \rangle \neq \langle \phi_2 \rangle = \langle \phi_3 \rangle$ or $\langle \phi_2 \rangle \neq \langle \phi_1 \rangle \neq \langle \phi_3 \rangle$ or $\langle \phi_3 \rangle \neq \langle \phi_1 \rangle \neq \langle \phi_2 \rangle$ then T_7 is broken into {Identity}.
- (3) The third alignment: $0 = \langle \phi_1 \rangle \neq \langle \phi_2 \rangle = \langle \phi_3 \rangle \neq 0$ or $0 = \langle \phi_2 \rangle \neq \langle \phi_3 \rangle = \langle \phi_1 \rangle \neq 0$ or $0 = \langle \phi_3 \rangle \neq \langle \phi_1 \rangle = \langle \phi_2 \rangle \neq 0$ then T_7 is broken into {Identity}.
- (4) The fourth alignment: $0 = \langle \phi_1 \rangle \neq \langle \phi_2 \rangle \neq \langle \phi_3 \rangle \neq 0$ or $0 = \langle \phi_2 \rangle \neq \langle \phi_1 \rangle \neq \langle \phi_3 \rangle \neq 0$ or $0 = \langle \phi_3 \rangle \neq \langle \phi_2 \rangle \neq \langle \phi_1 \rangle \neq 0$ then T_7 is broken into {Identity}.
- (5) The fifth alignment: $0 = \langle \phi_1 \rangle = \langle \phi_2 \rangle \neq \langle \phi_3 \rangle \neq 0$ or $0 = \langle \phi_1 \rangle = \langle \phi_3 \rangle \neq \langle \phi_2 \rangle \neq 0$ or $0 = \langle \phi_2 \rangle = \langle \phi_3 \rangle \neq \langle \phi_1 \rangle \neq 0$ then T_7 is broken into {Identity}.

In this work, we argue that only the first alignment of VEV in charged - lepton sector is taken place, i.e, $T_7 \rightarrow Z_3$, and this can be achieved by the Higgs triplet ϕ with the VEV alignment $\langle \phi \rangle = (\langle \phi_1 \rangle, \langle \phi_1 \rangle, \langle \phi_1 \rangle)$ under T_7 , where

$$\langle \phi_1 \rangle = (0 \quad v \quad 0)^T. \quad (3.2)$$

The Yukawa interactions are

$$\begin{aligned} -\mathcal{L}_l &= h_1(\bar{\psi}_L \phi)_{\underline{1}} l_{1R} + h_2(\bar{\psi}_L \phi)_{\underline{1}'} l_{2R} + h_3(\bar{\psi}_L \phi)_{\underline{1}''} l_{3R} + \text{H.c.} \\ &= h_1(\bar{\psi}_{1L} \phi_1 + \bar{\psi}_{2L} \phi_2 + \bar{\psi}_{3L} \phi_3) l_{1R} \\ &\quad + h_2(\bar{\psi}_{1L} \phi_1 + \omega^2 \bar{\psi}_{2L} \phi_2 + \omega \bar{\psi}_{3L} \phi_3) l_{2R} \\ &\quad + h_3(\bar{\psi}_{1L} \phi_1 + \omega \bar{\psi}_{2L} \phi_2 + \omega^2 \bar{\psi}_{3L} \phi_3) l_{3R} + \text{H.c.} \end{aligned} \quad (3.3)$$

The mass Lagrangian for the charged leptons is then given by

$$\begin{aligned} -\mathcal{L}_l^{\text{mass}} &= h_1 v \bar{l}_{1L} l_{1R} + h_2 v \bar{l}_{1L} l_{2R} + h_3 v \bar{l}_{1L} l_{3R} \\ &\quad + h_1 v \bar{l}_{2L} l_{1R} + h_2 v \omega^2 \bar{l}_{2L} l_{2R} + h_3 v \omega \bar{l}_{2L} l_{3R} \\ &\quad + h_1 v \bar{l}_{3L} l_{1R} + h_2 v \omega \bar{l}_{3L} l_{2R} + h_3 v \omega^2 \bar{l}_{3L} l_{3R} + \text{H.c.} \end{aligned} \quad (3.4)$$

The mass Lagrangian for the charged leptons reads

$$-\mathcal{L}_l^{\text{mass}} = (\bar{l}_{1L}, \bar{l}_{2L}, \bar{l}_{3L}) M_l (l_{1R}, l_{2R}, l_{3R})^T + H.c., \quad (3.5)$$

where

$$M_l = \begin{pmatrix} h_1 v & h_2 v & h_3 v \\ h_1 v & h_2 v \omega^2 & h_3 v \omega \\ h_1 v & h_2 v \omega & h_3 v \omega^2 \end{pmatrix}. \quad (3.6)$$

This matrix can be diagonalized as,

$$U_L^\dagger M_l U_R = \begin{pmatrix} \sqrt{3} h_1 v & 0 & 0 \\ 0 & \sqrt{3} h_2 v & 0 \\ 0 & 0 & \sqrt{3} h_3 v \end{pmatrix} \equiv \begin{pmatrix} m_e & 0 & 0 \\ 0 & m_\mu & 0 \\ 0 & 0 & m_\tau \end{pmatrix}, \quad (3.7)$$

where

$$U_L = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{pmatrix}, \quad U_R = 1. \quad (3.8)$$

As will see in section 5, in this case, the exact tribimaximal mixing form is obtained, by choosing the right vev's in the neutrino sector.

The experimental values for masses of the charged leptons at the weak scale are given as [8]:

$$m_e = 0.511 \text{ MeV}, \quad m_\mu = 105.658 \text{ MeV}, \quad m_\tau = 1776.82 \text{ MeV} \quad (3.9)$$

from which it follows that $h_1 \ll h_2 \ll h_3$. On the other hand, if we choose the VEV $v = 100 \text{ GeV}$ then $h_1 \sim 10^{-6}$, $h_2 \sim 10^{-4}$, $h_3 \sim 10^{-2}$.

4 Quark masses

To generate masses for quarks with a minimal Higgs content, we additionally introduce the following Higgs triplets:

$$\eta_i = \begin{pmatrix} \eta_{i1}^0 \\ \eta_{i2}^- \\ \eta_{i3}^0 \end{pmatrix} \sim [3, -1/3, -1/3, \underline{3}] \quad (i = 1, 2, 3), \quad (4.1)$$

$$\chi = \begin{pmatrix} \chi_1^0 \\ \chi_2^- \\ \chi_3^0 \end{pmatrix} \sim [3, -1/3, 2/3, \underline{1}]. \quad (4.2)$$

The Higgs content and Yukawa couplings in the quark sector are summarized in table 1.

The Yukawa interactions are

$$\begin{aligned} -\mathcal{L}_q = & h_3^d \bar{Q}_{3L} (\phi d_R)_1 + h_1^u \bar{Q}_{1L} (\phi^* u_R)_{1''} + h_2^u \bar{Q}_{2L} (\phi^* u_R)_{1'} \\ & + h_3^u \bar{Q}_{3L} (\eta u_R)_1 + h_1^d \bar{Q}_{1L} (\eta^* d_R)_{1''} + h_2^d \bar{Q}_{2L} (\eta^* d_R)_{1'} \\ & + f_3 \bar{Q}_{3L} \chi U_R + f_1 \bar{Q}_{1L} \chi^* D_{1R} + f_2 \bar{Q}_{2L} \chi^* D_{2R} + H.c. \end{aligned}$$

Couplings	Higgs multiplets
$\bar{Q}_{3L}U_R \sim (3^*, \frac{1}{3}, -\frac{2}{3}, \underline{1})$	$\chi \sim (3, -\frac{1}{3}, \frac{2}{3}, \underline{1})$
$\bar{Q}_{2L}D_{2R} \sim (3, -\frac{1}{3}, \frac{2}{3}, \underline{1})$	$\chi^* \sim (3^*, \frac{1}{3}, -\frac{2}{3}, \underline{1})$
$\bar{Q}_{1L}D_{1R} \sim (3, -\frac{1}{3}, \frac{2}{3}, \underline{1})$	$\chi^* \sim (3^*, \frac{1}{3}, -\frac{2}{3}, \underline{1})$
$\bar{Q}_{3L}d_R \sim (3^*, -\frac{2}{3}, \frac{1}{3}, \underline{3}^*)$	$\phi \sim (3, \frac{2}{3}, -\frac{1}{3}, \underline{3})$
$\bar{Q}_{1L}u_R \sim (3, \frac{2}{3}, -\frac{1}{3}, \underline{3})$	$\phi^* \sim (3^*, -\frac{2}{3}, \frac{1}{3}, \underline{3}^*)$
$\bar{Q}_{2L}u_R \sim (3, \frac{2}{3}, -\frac{1}{3}, \underline{3})$	$\phi^* \sim (3^*, -\frac{2}{3}, \frac{1}{3}, \underline{3}^*)$
$\bar{Q}_{3L}u_R \sim (3^*, \frac{1}{3}, \frac{1}{3}, \underline{3})$	$\eta \sim (3, -\frac{1}{3}, -\frac{1}{3}, \underline{3}^*)$
$\bar{Q}_{1L}d_R \sim (3, -\frac{1}{3}, -\frac{1}{3}, \underline{3}^*)$	$\eta^* \sim (3^*, \frac{1}{3}, \frac{1}{3}, \underline{3})$
$\bar{Q}_{2L}d_R \sim (3, -\frac{1}{3}, -\frac{1}{3}, \underline{3}^*)$	$\eta^* \sim (3^*, \frac{1}{3}, \frac{1}{3}, \underline{3})$

Table 1. List of couplings which form a singlet from the invariance under the T_7 .

$$\begin{aligned}
 &= h_3^d \bar{Q}_{3L}(\phi_1 d_{1R} + \phi_2 d_{2R} + \phi_3 d_{3R}) \\
 &\quad + h_1^u \bar{Q}_{1L}(\phi_1^* u_R + \omega^2 \phi_2^* u_{2R} + \omega \phi_3^* u_{3R}) \\
 &\quad + h_2^u \bar{Q}_{2L}(\phi_1^* u_R + \omega \phi_2^* u_{2R} + \omega^2 \phi_3^* u_{3R}) \\
 &\quad + h_3^u \bar{Q}_{3L}(\eta_1 u_{1R} + \eta_2 u_{2R} + \eta_3 u_{3R}) \\
 &\quad + h_1^d \bar{Q}_{1L}(\eta_1^* d_{1R} + \omega^2 \eta_2^* d_{2R} + \omega \eta_3^* d_{3R}) \\
 &\quad + h_2^d \bar{Q}_{2L}(\eta_1^* d_{1R} + \omega \eta_2^* d_{2R} + \omega^2 \eta_3^* d_{3R}) \\
 &\quad + f_3 \bar{Q}_{3L} \chi U_R + f_1 \bar{Q}_{1L} \chi^* D_{1R} + f_2 \bar{Q}_{2L} \chi^* D_{2R} + \text{H.c.} \tag{4.3}
 \end{aligned}$$

We suppose that T_7 is broken into Z_3 like the case of the charged lepton sector, i.e, the VEVs of η and χ are given as $\langle \eta \rangle = (\langle \eta_1 \rangle, \langle \eta_1 \rangle, \langle \eta_1 \rangle)$ with

$$\langle \eta_1 \rangle = (u \quad 0 \quad 0)^T, \tag{4.4}$$

and

$$\langle \chi \rangle = (0 \quad 0 \quad v_\chi)^T. \tag{4.5}$$

The mass Lagrangian for quarks is given by

$$\begin{aligned}
 -\mathcal{L}_q^{\text{mass}} &= -h_1^u v_1 \bar{u}_{1L} u_{1R} - h_1^u v_2 \omega^2 \bar{u}_{1L} u_{2R} - h_1^u v_3 \omega \bar{u}_{1L} u_{3R} \\
 &\quad - h_2^u v_1 \bar{u}_{2L} u_{1R} - h_2^u v_2 \omega \bar{u}_{2L} u_{2R} - h_2^u v_3 \omega^2 \bar{u}_{2L} u_{3R} \\
 &\quad + h_3^u u_1 \bar{u}_{3L} u_{1R} + h_3^u u_2 \bar{u}_{3L} u_{2R} + h_3^u u_3 \bar{u}_{3L} u_{3R} \\
 &\quad + h_1^d u_1 \bar{d}_{1L} d_{1R} + \omega^2 h_1^d u_2 \bar{d}_{1L} d_{2R} + \omega h_1^d u_3 \bar{d}_{1L} d_{3R} \\
 &\quad + h_2^d u_1 \bar{d}_{2L} d_{1R} + \omega h_2^d u_2 \bar{d}_{2L} d_{2R} + \omega^2 h_2^d u_3 \bar{d}_{2L} d_{3R} \\
 &\quad + h_3^d v_1 \bar{d}_{3L} d_{1R} + h_3^d v_2 \bar{d}_{3L} d_{2R} + h_3^d v_3 \bar{d}_{3L} d_{3R} \\
 &\quad + f_3 v_\chi \bar{U}_L U_R + f_1 v_\chi \bar{D}_{1L} D_{1R} + f_2 v_\chi \bar{D}_{2L} D_{2R} + \text{H.c.} \tag{4.6}
 \end{aligned}$$

The exotic quarks get masses

$$m_U = f_3 v_\chi, \quad m_{D_{1,2}} = f_{1,2} v_\chi. \quad (4.7)$$

The mass matrices for ordinary up-quarks and down-quarks are, respectively, obtained as follows:

$$M_u = \begin{pmatrix} -h_1^u v & -h_1^u v \omega^2 & -h_1^u v \omega \\ -h_2^u v & -h_2^u v \omega & -h_2^u v \omega^2 \\ h_3^u u & h_3^u u & h_3^u u \end{pmatrix}, \quad M_d = \begin{pmatrix} h_1^d u & h_1^d u \omega^2 & h_1^d u \omega \\ h_2^d u & h_2^d u \omega & h_2^d u \omega^2 \\ h_3^d v & h_3^d v & h_3^d v \end{pmatrix}. \quad (4.8)$$

The structure of the up- and down-quark mass matrices in (4.8) is similar to those in [33], i.e, in the model under consideration there is no CP violation in the quark mixing matrix. The mass matrices M_u, M_d in (4.8) are diagonalized as follows

$$U_L^{u+} M_u U_R^u = \begin{pmatrix} -\sqrt{3} h_1^u v & 0 & 0 \\ 0 & -\sqrt{3} h_2^u v & 0 \\ 0 & 0 & \sqrt{3} h_3^u u \end{pmatrix} = \begin{pmatrix} m_u & 0 & 0 \\ 0 & m_c & 0 \\ 0 & 0 & m_t \end{pmatrix},$$

$$U_L^{d+} M_d U_R^d = \begin{pmatrix} \sqrt{3} h_1^d u & 0 & 0 \\ 0 & \sqrt{3} h_2^d u & 0 \\ 0 & 0 & \sqrt{3} h_3^d v \end{pmatrix} = \begin{pmatrix} m_d & 0 & 0 \\ 0 & m_s & 0 \\ 0 & 0 & m_b \end{pmatrix}. \quad (4.9)$$

where

$$U_R^u = U_R^d = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ \omega & \omega^2 & 1 \\ \omega^2 & \omega & 1 \end{pmatrix}, \quad U_L^u = U_L^d = 1 \quad (4.10)$$

The unitary matrices, which couple the left-handed up- and down-quarks to those in the mass bases are unit matrices. Therefore we get the Cabibbo-Kobayashi-Maskawa (CKM) matrix

$$U_{\text{CKM}} = U_L^{d\dagger} U_L^u = 1. \quad (4.11)$$

Note that the property in (4.11) is common for some models based on the discrete symmetry groups [164, 165].

The up and down quark masses are

$$\begin{aligned} m_u &= -\sqrt{3} h_1^u v, & m_c &= -\sqrt{3} h_2^u v, & m_t &= \sqrt{3} h_3^u u, \\ m_d &= \sqrt{3} h_1^d u, & m_s &= \sqrt{3} h_2^d u, & m_b &= \sqrt{3} h_3^d v. \end{aligned} \quad (4.12)$$

The current mass values for the quarks are given by [8]:

$$\begin{aligned} m_u &= 2.3_{-0.5}^{+0.7} \text{ MeV}, & m_c &= 1.275 \pm 0.025 \text{ GeV}, & m_t &= 173.5 \pm 0.6 \pm 0.8, \text{ GeV} \\ m_d &= 4.8_{-0.3}^{+0.7} \text{ MeV}, & m_s &= 95 \pm 5 \text{ MeV}, & m_b &= 4.18 \pm 0.03 \text{ GeV}. \end{aligned} \quad (4.13)$$

It is obvious that if $|u| \sim |v|$, the Yukawa coupling hierarchies are $|h_1^u| \ll h_2^u \ll h_3^u$, $h_1^d \ll h_2^d \ll h_3^d$ and the couplings between up-quarks (h_2^u, h_3^u) and Higgs scalar multiplets are slightly heavier than those of down-quarks ($h_2^d \ll h_3^d$), respectively.

5 Neutrino mass and mixing

The neutrino masses arise from the couplings of $\bar{\psi}_L^c \psi_L$ to scalars, where $\bar{\psi}_L^c \psi_L$ transforms as $3^* \oplus 6$ under $SU(3)_L$ and $\underline{3} \oplus \underline{3}^* \oplus \underline{3}^*$ under T_7 . It is worth mentioning that, with the T_7 group, $\underline{3} \times \underline{3} \times \underline{3}$ has two invariants and $\underline{3} \times \underline{3} \times \underline{3}^*$ has one invariant. For the known scalar triplets (ϕ, χ, η) , there is no available interaction because of the \mathcal{L} -symmetry. We will therefore propose new $SU(3)_L$ anti-sextets instead coupling to $\bar{\psi}_L^c \psi_L$ responsible for the neutrino masses. To obtain a realistic neutrino spectrum, the antisextets transform as follows²

$$\sigma_i = \begin{pmatrix} \sigma_{11}^0 & \sigma_{12}^+ & \sigma_{13}^0 \\ \sigma_{12}^+ & \sigma_{22}^{++} & \sigma_{23}^+ \\ \sigma_{13}^0 & \sigma_{23}^+ & \sigma_{33}^0 \end{pmatrix}_i \sim [6^*, 2/3, -4/3, \underline{3}^*] \quad (i = 1, 2, 3), \quad (5.1)$$

Following the potential minimization conditions, we have the followings alignments:

- (1) The first alignment: $\langle \sigma_1 \rangle = \langle \sigma_2 \rangle = \langle \sigma_3 \rangle$ then T_7 is broken into Z_3 consisting of the elements $\{e, b, b^2\}$.
- (2) The second alignment: $\langle \sigma_1 \rangle \neq \langle \sigma_2 \rangle \neq \langle \sigma_3 \rangle$ or $\langle \sigma_1 \rangle \neq \langle \sigma_2 \rangle = \langle \sigma_3 \rangle$ or $\langle \sigma_2 \rangle \neq \langle \sigma_1 \rangle \neq \langle \sigma_3 \rangle$ or $\langle \sigma_3 \rangle \neq \langle \sigma_1 \rangle \neq \langle \sigma_2 \rangle$ then T_7 is broken into $\{\text{Identity}\}$.
- (3) The third alignment: $0 = \langle \sigma_1 \rangle \neq \langle \sigma_2 \rangle = \langle \sigma_3 \rangle \neq 0$ or $0 = \langle \sigma_2 \rangle \neq \langle \sigma_3 \rangle = \langle \sigma_1 \rangle \neq 0$ or $0 = \langle \sigma_3 \rangle \neq \langle \sigma_1 \rangle = \langle \sigma_2 \rangle \neq 0$ then T_7 is broken into $\{\text{Identity}\}$.
- (4) The fourth alignment: $0 = \langle \sigma_1 \rangle \neq \langle \sigma_2 \rangle \neq \langle \sigma_3 \rangle \neq 0$ or $0 = \langle \sigma_2 \rangle \neq \langle \sigma_1 \rangle \neq \langle \sigma_3 \rangle \neq 0$ or $0 = \langle \sigma_3 \rangle \neq \langle \sigma_1 \rangle \neq \langle \sigma_2 \rangle \neq 0$ then T' is broken into $\{\text{Identity}\}$.
- (5) The fifth alignment: $0 = \langle \sigma_1 \rangle = \langle \sigma_2 \rangle \neq \langle \sigma_3 \rangle \neq 0$ or $0 = \langle \sigma_1 \rangle = \langle \sigma_3 \rangle \neq \langle \sigma_2 \rangle \neq 0$ or $0 = \langle \sigma_2 \rangle = \langle \sigma_3 \rangle \neq \langle \sigma_1 \rangle \neq 0$ then T_7 is broken into $\{\text{Identity}\}$.

To obtain a realistic neutrino spectrum, in this work we argue that both the breakings $T_7 \rightarrow Z_3$ and $T_7 \rightarrow \{\text{identity}\}$ (Instead of $Z_3 \rightarrow \{\text{identity}\}$) must be taken place in neutrino sector. However, the VEVs of σ does only one of these tasks. The $T_7 \rightarrow Z_3$ can be achieved by a $SU(3)_L$ anti-sextet σ given in (5.1) with the VEVs is set as $\langle \sigma \rangle = (\langle \sigma_1 \rangle, \langle \sigma_1 \rangle, \langle \sigma_1 \rangle)$ under T_7 , where

$$\langle \sigma_1 \rangle = \begin{pmatrix} \lambda_\sigma & 0 & v_\sigma \\ 0 & 0 & 0 \\ v_\sigma & 0 & \Lambda_\sigma \end{pmatrix}. \quad (5.2)$$

To achieve the second direction of the breakings $T_7 \rightarrow \{\text{Identity}\}$ (equivalently to $Z_3 \rightarrow \{\text{Identity}\}$), we additionally introduce another $SU(3)_L$ anti-sextet Higgs scalar which is either put in $\underline{3}$ or $\underline{3}^*$ under T_7 . This is equivalent to breaking the subgroup Z_3 of the first direction into $\{\text{Identity}\}$, and it can be achieved within each case below.

²Note that in the model under consideration, if the choice is a $SU(3)_L$ triplet [for example $\rho \sim (3, 2/3, -4/3, \underline{3}^*)$] instead of $SU(3)_L$ anti-sextet σ as in (5.1), there will be a contribution from term of $(\bar{\psi}_L^c \psi_L)_{\underline{3}^*} \rho$ added to the elements $(M_D)_{11}, (M_D)_{22}, (M_D)_{33}$ of the matrix M_D which is the same order. The lepton mixing matrix therefore can only reach to U_{HPS} but not $\theta_{13} \neq 0$.

1. A new $SU(3)_L$ anti-sextet s which is put in the $\underline{3}$ under T_7 ,

$$s_i = \begin{pmatrix} s_{11}^0 & s_{12}^+ & s_{13}^0 \\ s_{12}^+ & s_{22}^{++} & s_{23}^+ \\ s_{13}^0 & s_{23}^+ & s_{33}^0 \end{pmatrix}_i \sim [6^*, 2/3, -4/3, \underline{3}], \quad (i = 1, 2, 3) \quad (5.3)$$

with the VEVs given by $\langle s \rangle = (\langle s_1 \rangle, 0, 0)^T$, where

$$\langle s_1 \rangle = \begin{pmatrix} \lambda_s & 0 & v_s \\ 0 & 0 & 0 \\ v_s & 0 & \Lambda_s \end{pmatrix}. \quad (5.4)$$

2. Another $SU(3)_L$ anti-sextet σ' is put in the $\underline{3}^*$ under T_7 , with the VEVs chosen by

$$\sigma'_i = \begin{pmatrix} \sigma_{11}'^0 & \sigma_{12}'^+ & \sigma_{13}'^0 \\ \sigma_{12}'^+ & \sigma_{22}'^{++} & \sigma_{23}'^+ \\ \sigma_{13}'^0 & \sigma_{23}'^+ & \sigma_{33}'^0 \end{pmatrix}_i \sim [6^*, 2/3, -4/3, \underline{3}^*], \quad (i = 1, 2, 3)$$

$$\langle \sigma'_1 \rangle = \begin{pmatrix} \lambda'_\sigma & 0 & v'_\sigma \\ 0 & 0 & 0 \\ v'_\sigma & 0 & \Lambda'_\sigma \end{pmatrix}, \quad \langle \sigma'_2 \rangle = \langle \sigma'_3 \rangle = 0. \quad (5.5)$$

Note that σ' differs from σ only in the VEVs alignment. Combining both cases, after calculation, we obtain the Yukawa interactions:

$$\begin{aligned} -\mathcal{L}_\nu &= \frac{1}{2}x(\bar{\psi}_L^c \sigma)_{\underline{3}^*} \psi_L + y(\bar{\psi}_L^c s)_{\underline{3}^*} \psi_L + \frac{z}{2}(\bar{\psi}_L^c \sigma')_{\underline{3}^*} \psi_L + \text{H.c.} \\ &= \frac{1}{2}x(\bar{\psi}_{1L}^c \sigma_2 \psi_{1L} + \bar{\psi}_{2L}^c \sigma_3 \psi_{2L} + \bar{\psi}_{3L}^c \sigma_1 \psi_{3L}) \\ &\quad + y(\bar{\psi}_{2L}^c s_3 \psi_{1L} + \bar{\psi}_{3L}^c s_1 \psi_{2L} + \bar{\psi}_{1L}^c s_2 \psi_{3L}) \\ &\quad + \frac{z}{2}(\bar{\psi}_{1L}^c \sigma'_2 \psi_{1L} + \bar{\psi}_{2L}^c \sigma'_3 \psi_{2L} + \bar{\psi}_{3L}^c \sigma'_1 \psi_{3L}) + \text{H.c.} \end{aligned} \quad (5.6)$$

The mass Lagrangian for the neutrinos is given by

$$\begin{aligned} -\mathcal{L}_\nu^{\text{mass}} &= \frac{1}{2}x(\lambda_\sigma \bar{\nu}_{1L}^c \nu_{1L} + v_\sigma \bar{\nu}_{1L}^c N_{1R}^c + v_\sigma \bar{N}_{1R} \nu_{1L} + \Lambda_\sigma \bar{N}_{1R} N_{1R}^c) \\ &\quad + \frac{1}{2}x(\lambda_\sigma \bar{\nu}_{2L}^c \nu_{2L} + v_\sigma \bar{\nu}_{2L}^c N_{2R}^c + v_\sigma \bar{N}_{2R} \nu_{2L} + \Lambda_\sigma \bar{N}_{2R} N_{2R}^c) \\ &\quad + \frac{1}{2}x(\lambda_\sigma \bar{\nu}_{3L}^c \nu_{3L} + v_\sigma \bar{\nu}_{3L}^c N_{3R}^c + v_\sigma \bar{N}_{3R} \nu_{3L} + \Lambda_\sigma \bar{N}_{3R} N_{3R}^c) \\ &\quad + y(\lambda_s \bar{\nu}_{3L}^c \nu_{2L} + v_s \bar{\nu}_{3L}^c N_{2R}^c + v_s \bar{N}_{3R} \nu_{2L} + \Lambda_s \bar{N}_{3R} N_{2R}^c) \\ &\quad + \frac{1}{2}z(\lambda'_\sigma \bar{\nu}_{3L}^c \nu_{3L} + v'_\sigma \bar{\nu}_{3L}^c N_{3R}^c + v'_\sigma \bar{N}_{3R} \nu_{3L} + \Lambda'_\sigma \bar{N}_{3R} N_{3R}^c) + \text{H.c.} \end{aligned} \quad (5.7)$$

The neutrino mass Lagrangian can be written in matrix form as follows

$$-\mathcal{L}_\nu^{\text{mass}} = \frac{1}{2} \bar{\chi}_L^c M_\nu \chi_L + \text{h.c.}, \quad (5.8)$$

where

$$\begin{aligned}\chi_L &\equiv (\nu_L \quad N_R^c)^T, & M_\nu &\equiv \begin{pmatrix} M_L & M_D^T \\ M_D & M_R \end{pmatrix}, \\ \nu_L &= (\nu_{1L}, \nu_{2L}, \nu_{3L})^T, & N_R &= (N_{1R}, N_{2R}, N_{3R})^T,\end{aligned}\tag{5.9}$$

and the mass matrices are then obtained by

$$M_{L,R,D} = \begin{pmatrix} a_{L,R,D} & 0 & 0 \\ 0 & a_{L,R,D} & b_{L,R,D} \\ 0 & b_{L,R,D} & a_{L,R,D} + c_{L,R,D} \end{pmatrix},\tag{5.10}$$

with

$$\begin{aligned}a_L &= \lambda_\sigma x, & a_D &= v_\sigma x, & a_R &= \Lambda_\sigma x, \\ b_L &= \lambda_s y, & b_D &= v_s y, & b_R &= \Lambda_s y, \\ c_L &= \lambda'_\sigma z, & c_D &= v'_\sigma z, & c_R &= \Lambda'_\sigma z.\end{aligned}\tag{5.11}$$

Three observed neutrinos gain masses via a combination of type I and type II seesaw mechanisms derived from (5.8) and (5.10) as

$$M_{\text{eff}} = M_L - M_D^T M_R^{-1} M_D = \begin{pmatrix} A & 0 & 0 \\ 0 & B_1 & C \\ 0 & C & B_2 \end{pmatrix},\tag{5.12}$$

where

$$\begin{aligned}A &= a_L - \frac{a_D^2}{a_R}, \\ B_1 &= a_L - \frac{a_R b_D^2 - 2a_D b_D b_R + a_D^2 (a_R + d_R)}{a_R^2 - b_R^2 + a_R d_R}, \\ B_2 &= B_1 + d_L + \frac{2(b_D b_R - a_D a_R) d_D + (a_D^2 - b_D^2) d_R - a_R d_D^2}{a_R^2 - b_R^2 + a_R d_R}, \\ C &= b_L - \frac{(a_D^2 + b_D^2) b_R - (2a_D a_R + a_D d_R) b_D + (a_D b_R - a_R b_D) d_D}{a_R^2 - b_R^2 + a_R d_R}.\end{aligned}\tag{5.13}$$

We can diagonalize the mass matrix (5.12) as follows $U_\nu^T M_{\text{eff}} U_\nu = \text{diag}(m_1, m_2, m_3)$, with

$$\begin{aligned}m_1 &= \frac{1}{2} \left(B_1 + B_2 + \sqrt{(B_1 + B_2)^2 + 4C^2} \right), \\ m_2 &= A, \\ m_3 &= \frac{1}{2} \left(B_1 + B_2 - \sqrt{(B_1 + B_2)^2 + 4C^2} \right),\end{aligned}\tag{5.14}$$

and the corresponding neutrino mixing matrix:

$$U_\nu = \begin{pmatrix} 0 & 1 & 0 \\ \frac{1}{\sqrt{K^2+1}} & 0 & \frac{K}{\sqrt{K^2+1}} \\ -\frac{K}{\sqrt{K^2+1}} & 0 & \frac{1}{\sqrt{K^2+1}} \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & i \end{pmatrix},\tag{5.15}$$

where

$$K = \frac{B_1 - B_2 - \sqrt{4C^2 + (B_1 - B_2)^2}}{2C}. \quad (5.16)$$

Combining (3.8) and (5.15), we get the lepton mixing matrix:

$$U_L^\dagger U_\nu = \frac{1}{\sqrt{3}} \begin{pmatrix} \frac{1-K}{\sqrt{K^2+1}} & 1 & \frac{1+K}{\sqrt{K^2+1}} \\ \frac{\omega(1-K\omega)}{\sqrt{K^2+1}} & 1 & \frac{\omega(\omega+K)}{\sqrt{K^2+1}} \\ \frac{\omega(\omega-K)}{\sqrt{K^2+1}} & 1 & \frac{\omega(K\omega+1)}{\sqrt{K^2+1}} \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & i \end{pmatrix}. \quad (5.17)$$

It is worth noting that in our model, K given in (5.16) is an arbitrary number. Hence in general the lepton mixing matrix given in (5.17) is different to U_{HPS} in (1.1), but similar to the original version of trimaximal mixing considered in ref. [170] which is based on the $\Delta(27)$ group extension of the Standard Model. Although there are some phenomenological predictions of the model are similar to those of in ref. [170] but the fundamental difference between our model with the well known one is the prediction of CP violation (our model predicts maximal CP violation $\delta = \pi/2, 3\pi/2$ with $\theta_{23} \neq \pi/4$, but in ref. [170], the maximal CP violation $\delta = \pi/2, 3\pi/2$ achieved with $\theta_{23} = \pi/4$). In the case where T_7 is broken into Identity (Instead of $Z_3 \rightarrow$ Identity) only by s , i.e, without contribution of σ' (or $\lambda'_\sigma = \nu'_\sigma = \Lambda'_\sigma = 0$), the lepton mixing matrix (5.17) being equal to U_{HPS} as given in (1.1). This is a good features of T_7 with tensor product $\mathfrak{3} \otimes \mathfrak{3}$ given in (A.3).

In the standard Particle Data Group (PDG) parametrization, the lepton mixing matrix (U_{PMNS}) can be parametrized as

$$U_{PMNS} = \begin{pmatrix} c_{12}c_{13} & -s_{12}c_{13} & -s_{13}e^{-i\delta} \\ s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & -s_{23}c_{13} \\ s_{12}s_{23} + c_{12}c_{23}s_{13}e^{i\delta} & c_{12}s_{23} + s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \times P. \quad (5.18)$$

where $P = \text{diag}(1, e^{i\alpha}, e^{i\beta})$, and $c_{ij} = \cos\theta_{ij}$, $s_{ij} = \sin\theta_{ij}$ with θ_{12} , θ_{23} and θ_{13} being the solar angle, atmospheric angle and the reactor angle respectively. δ is the Dirac CP violating phase while α and β are the two Majorana CP violating phases.

From the (5.17) and (5.18) we rule out $\alpha = 0, \beta = \frac{\pi}{2}$, and the lepton mixing matrix in (5.17) can be parameterized in three Euler's angles θ_{ij} as follows:

$$s_{13}e^{-i\delta} = \frac{-1 - K}{\sqrt{3}\sqrt{K^2 + 1}}, \quad (5.19)$$

$$t_{12} = \frac{\sqrt{K^2 + 1}}{K - 1}, \quad (5.20)$$

$$t_{23} = -\frac{\omega + K}{1 + K\omega}. \quad (5.21)$$

Substituting $\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$ into (5.21) yields:

$$K = k_1 + ik_2, \quad (5.22)$$

$$k_1 = \frac{t_{23}^2 - 4t_{23} + 1}{2(t_{23}^2 - t_{23} + 1)}, \quad k_2 = \frac{\sqrt{3}(t_{23}^2 - 1)}{2(t_{23}^2 - t_{23} + 1)}.$$

The expression (5.22) tells us that $k_1^2 + k_2^2 \equiv |K|^2 = 1$. Combining (5.19) and (5.20) yields:

$$\begin{aligned}
 e^{-i\delta} &= \frac{1}{\sqrt{3}s_{13}t_{12}} \frac{1+K}{1-K} = \frac{1}{\sqrt{3}s_{13}t_{12}} \left[\frac{1-k_1^2-k_2^2}{[(1-k_1)^2+k_2^2]} + \frac{2k_2}{[(1-k_1)^2+k_2^2]}i \right] \\
 &= -\frac{i(1-t_{23})}{s_{13}t_{12}(1+t_{23})} \equiv \cos \delta - i \sin \delta
 \end{aligned}$$

or

$$\cos \delta = 0, \quad \sin \delta = \frac{1-t_{23}}{s_{13}t_{12}(t_{23}+1)}. \tag{5.23}$$

Since $\cos \delta = 0$ so that $\sin \delta$ must be equal to ± 1 , it is then $\delta = \frac{\pi}{2}$ or $\delta = \frac{3\pi}{2}$. Thus, our model predicts the maximal Dirac CP violating phase which is the same as in refs. [170, 171] the maximal CP violation $\delta = \frac{\pi}{2}, \frac{3\pi}{2}$ is achieved with $\theta_{23} = \pi/4$,³ and this is one of the most striking prediction of the model under consideration.

Up to now the precise evaluation of θ_{23} is still an open problem while θ_{12} and θ_{13} are now very constrained [8]. From (5.23), our model can provide constraints on θ_{23} from θ_{12} and θ_{13} which satisfy [8] as follows.

- (i) In the case $\delta = \frac{\pi}{2}$, from (5.23) we have the relation among three Euler's angles as follows:

$$\begin{aligned}
 t_{23} &= \frac{1-s_{13}t_{12}}{1+s_{13}t_{12}} \\
 &= \frac{2-\sqrt{2}\sqrt{\frac{2-2\sqrt{1-\sin^2(2\theta_{12})+\sin^2(2\theta_{12})}}{\sin^2(2\theta_{12})}}\sqrt{1-\sqrt{1-\sin^2(2\theta_{13})}}}{2+\sqrt{2}\sqrt{\frac{2-2\sqrt{1-\sin^2(2\theta_{12})+\sin^2(2\theta_{12})}}{\sin^2(2\theta_{12})}}\sqrt{1-\sqrt{1-\sin^2(2\theta_{13})}}}. \tag{5.25}
 \end{aligned}$$

In figure 1, we have plotted the values of t_{23} as functions of $\sin^2(2\theta_{13})$ and $\sin^2(2\theta_{12})$ with $\sin^2(2\theta_{12}) \in (0.833, 0.881)$, $\sin^2(2\theta_{13}) \in (0.085, 0.111)$ given in (1.2). If $\sin^2(2\theta_{13}) = 0.098$ ($\theta_{13} = 9.11^\circ$) we have the relation between t_{23} and $\sin^2(2\theta_{12})$ as shown in figure 2.

For the best fit values of θ_{12} and θ_{13} given in [8], $\sin^2(2\theta_{12}) = 0.857$, $\sin^2(2\theta_{13}) = 0.098$ we obtain $t_{23} = 0.8075$ ($\theta_{23} = 38.92^\circ$), and

$$K = -0.930528 - 0.366221i, \quad (|K| = 1). \tag{5.26}$$

The lepton mixing matrix in (5.17) then takes the form:

$$U \simeq \begin{pmatrix} 0.831597 & 0.57735 & 0.157754 \\ -0.552417 & 0.57735 & -0.799061 \\ -0.27918 & 0.57735 & 0.641307 \end{pmatrix}. \tag{5.27}$$

³It would be interesting to note that the above conclusion is superficial. From eq. (16) in [170], it follows

$$\cos \delta \frac{2 \tan \theta_{23}}{1 - \tan^2 \theta_{23}} = \frac{2 \cos_{13} \cot 2\theta_{13}}{\sqrt{2 - 3 \sin^2 \theta_{13}}}. \tag{5.24}$$

If $\delta = \frac{\pi}{2}, \frac{3\pi}{2}$ and $\theta_{13} \neq 0$ then $\cos \delta = 0$, $\sin \theta_{13} \neq 0$, with $\theta_{23} = \frac{\pi}{4}$ (maximal), denominator in the left-hand side (5.24) vanishes. Thus, the left-hand side of (5.24) gets the form $\frac{0}{0}$, which does not have meaning. (as will see bellow in our model $\theta_{23} \neq \pi/4$).




Figure 1. t_{23} as a function of $\sin^2(2\theta_{12})$ and $\sin^2(2\theta_{13})$ with $\sin^2(2\theta_{12}) \in (0.833, 0.881)$ and $\sin^2(2\theta_{13}) \in (0.085, 0.111)$ in the case $\delta = \frac{\pi}{2}$.




Figure 2. t_{23} as a function of $\sin^2(2\theta_{12})$ with $\sin^2(2\theta_{12}) \in (0.833, 0.881)$ and $\sin^2(2\theta_{13}) = 0.098$ in the case $\delta = \frac{\pi}{2}$.

These results also implies that in the model under consideration, the value of the Jarlskog invariant J_{CP} which determines the magnitude of CP violation in neutrino oscillations is determined [172]:

$$J_{CP} = \frac{1}{8} \cos \theta_{13} \sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \sin \delta = 0.03527. \quad (5.28)$$

(ii) In the case $\delta = \frac{3\pi}{2}$, we have the relation among three Euler's angles as follows:

$$t_{23} = \frac{1 + s_{13}t_{12}}{1 - s_{13}t_{12}} = \frac{2 + \sqrt{2} \sqrt{\frac{2 - 2\sqrt{1 - \sin^2(2\theta_{12}) + \sin^2(2\theta_{12})}}{\sin^2(2\theta_{12})}} \sqrt{1 - \sqrt{1 - \sin^2(2\theta_{13})}}}{2 - \sqrt{2} \sqrt{\frac{2 - 2\sqrt{1 - \sin^2(2\theta_{12}) + \sin^2(2\theta_{12})}}{\sin^2(2\theta_{12})}} \sqrt{1 - \sqrt{1 - \sin^2(2\theta_{13})}}}. \quad (5.29)$$

In figure 3, we have plotted the values of t_{23} as a function of $\sin^2(2\theta_{12})$ and $\sin^2(2\theta_{13})$ with $\sin^2(2\theta_{12}) \in (0.833, 0.881)$ and $\sin^2(2\theta_{13}) \in (0.085, 0.111)$ in the case $\delta = \frac{3\pi}{2}$.




Figure 3. t_{23} as a function of $\sin^2(2\theta_{12})$ and $\sin^2(2\theta_{13})$ with $\sin^2(2\theta_{12}) \in (0.833, 0.881)$ and $\sin^2(2\theta_{13}) \in (0.085, 0.111)$ in the case $\delta = \frac{3\pi}{2}$.

For the best fit values of θ_{12} and θ_{13} given in [8], $\sin^2(2\theta_{12}) = 0.857$, $\sin^2(2\theta_{13}) = 0.098$ we obtain $t_{23} = 1.2384$ ($\theta_{23} = 51.079^\circ$), and

$$K = -0.930527 + 0.366223i, \quad (|K| = 1). \quad (5.30)$$

The lepton mixing matrix in (5.17) in this case takes the form:

$$U \simeq \begin{pmatrix} 0.831597 & 0.57735 & -0.157755 \\ -0.279179 & 0.57735 & -0.641306 \\ -0.552418 & 0.57735 & 0.799061 \end{pmatrix}, \quad (5.31)$$

and the value of the Jarlskog invariant J_{CP} is determined [172]:

$$J_{CP} = \frac{1}{8} \cos \theta_{13} \sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \sin \delta \simeq -0.03527. \quad (5.32)$$

Until now values of neutrino masses (or the absolute neutrino masses) as well as the mass ordering of neutrinos is unknown. The tritium experiment [173, 174] provides an upper bound on the absolute value of neutrino mass

$$m_i \leq 2.2 \text{ eV} \quad (5.33)$$

A more stringent bound was found from the analysis of the latest cosmological data [175]

$$m_i \leq 0.6 \text{ eV}, \quad (5.34)$$

while arguments from the growth of large-scale structure in the early Universe yield the upper bound [176]

$$\sum_{i=1}^3 m_i \leq 0.5 \text{ eV}. \quad (5.35)$$

The neutrino mass spectrum can be the normal mass hierarchy ($|m_1| \simeq |m_2| < |m_3|$), the inverted hierarchy ($|m_3| < |m_1| \simeq |m_2|$) or nearly degenerate ($|m_1| \simeq |m_2| \simeq |m_3|$). The mass ordering of neutrino depends on the sign of Δm_{23}^2 which is currently unknown. In the case of 3-neutrino mixing, in the model under consideration, the two possible signs of Δm_{23}^2 correspond to two types of neutrino mass spectrum can be provided. Combining (5.14) and the two experimental constraints on squared mass differences of neutrinos as shown in (1.2) and the values of K in (5.26) or in (5.30), we have the solutions as shown bellows.

5.1 Normal case ($\Delta m_{23}^2 > 0$)

5.1.1 The case $\delta = \frac{\pi}{2}$

In this case, combining (5.16) and the values of K in (5.26), we obtain

$$B_1 = B_2 - (1.67146 \times 10^{-7} + 0.732442i)C. \quad (5.36)$$

Substituting B_1 from (5.36) into (5.14) and combining with the two experimental constraints on squared mass differences of neutrinos as shown in (1.2), we get the solutions (in [eV]) given in appendix C.

From (5.36), (C.1) and (C.5) we see that A must be a real number in order to make the light neutrino masses $m_{1,2,3}$ to be real. In general, $B_{1,2}, C$ are complex numbers, α is also a complex number but $\text{Im}(\alpha) \ll \text{Re}(\alpha)$; $m_{1,2}$ are real numbers, m_3 is a complex number with $\text{Im}(m_3) \ll \text{Re}(m_3)$ but as will see below in the regions of the model parameters m_3 is real number, too.

The solutions in equations from (C.1) to (C.4) have the same absolute values of $m_{1,2,3}$, the unique difference is the sign of $m_{1,3}$. So, here we only consider in detail the case in (C.1)⁴ Using the upper bound on the absolute value of neutrino mass (5.34) we can restrict the values of A : $A \leq 0.6$ eV. However, in the case in (C.1), $A \in (0.0087, 0.01)$ eV or $A \in (-0.01, -0.0087)$ eV are good regions of A that can reach the realistic neutrino mass hierarchy. In this region of A , $B_{1,2}$ and C are complex numbers. The real part and the imaginary part of $B_{1,2}$ and C as functions of A (or m_2) are plotted in figures 4a and 4b, respectively. In figure 5, we have plotted the absolute value $|m_{1,3}|$ as functions of m_2 with $m_2 \in (0.00867, 0.05)$ eV. This figure shows that there exist allowed regions for values m_2 (or A) where either normal or quasi-degenerate neutrino masses spectrum is achieved. The quasi-degenerate mass hierarchy is obtained when A lies in a region $[0.05 \text{ eV}, +\infty]$ (A increases but must be small enough because of the scale of $m_{1,2,3}$). The normal mass hierarchy will be obtained if A takes the values around $(0.0087, 0.01)$ eV or $(-0.01, -0.0087)$ eV. The figures 6a and 6b give the sum $\sum_{i=1}^3 m_i$ and $\sum_{i=1}^3 |m_i|$ with $m_2 \in (0.0087, 0.05)$ eV, respectively.

⁴The expressions from (C.1) to (C.4) show that m_i ($i = 1, 2, 3$) depends only on a parameter $A = m_2$ so we consider $m_{1,3}$ as functions of m_2 . However, to have an explicit hierarchy on neutrino masses, in the following figures, m_2 should be included.




Figure 4. (a) The real part of $B_{1,2}$ and C as functions of m_2 , (b) The imaginary part of $B_{1,2}$ and C as functions of m_2 in the case of $\Delta m_{23}^2 > 0$.




Figure 5. $|m_{1,3}|$ as functions of m_2 with $m_2 \in (0.00867, 0.05)$ eV in the case of $\Delta m_{23}^2 > 0$.




Figure 6. (a) The sum $\sum_{i=1}^3 m_i$ as a function of A with $A \in (0.00867, 0.05)$ eV; (b) The sum $\sum_{i=1}^3 |m_i|$ as a function of A with $A \in (0.00867, 0.05)$ eV in the case of $\Delta m_{23}^2 > 0$.

From the expressions (5.17) , (5.30) and (C.1), it is easily to obtain the effective mass $\langle m_{ee} \rangle$ governing neutrinoless double beta decay [177–182],

$$\begin{aligned} \langle m_{ee} \rangle &= \left| \sum_{i=1}^3 U_{ei}^2 m_i \right| \\ &= \frac{A}{3} - (0.333333 - 8.90616 \times 10^{-18}i) \sqrt{4A^2 - 0.0003} \\ &\quad + (0.0248863 - 1.26619 \times 10^{-8}i) \sqrt{\Gamma}, \end{aligned} \quad (5.37)$$

$$\Gamma = 0.002245 + (2 - 2.6469 \times 10^{-23}i)A^2 - (1.73176 + 1.22425 \times 10^{-7}i)\sqrt{\gamma}, \quad (5.38)$$

$$\begin{aligned} \gamma &= (1.88579 \times 10^{-7} + 1.33377i)(2.44973 \times 10^{-19} + 0.00866025i - Ai) \\ &\quad \times (4.79026 \times 10^{-19} + 0.0481664i + A)(1.84151 \times 10^{-18} - 0.0481664i + A) \\ &\quad \times (0.00866025 - 6.69766 \times 10^{-26}i + A). \end{aligned} \quad (5.39)$$

$$m_\beta = \sqrt{\sum_{i=1}^3 |U_{ei}|^2 m_i^2}, \quad (5.40)$$

where

$$\begin{aligned} \sum_{i=1}^3 |U_{ei}|^2 m_i^2 &= 2.13672 \times 10^{-6} + 1.09955A^2 \\ &\quad - (0.0430971 + 3.0467 \times 10^{-9}i)\sqrt{\gamma'} + 0.0248863\sqrt{4A^2 - 0.0003}\sqrt{\Gamma'}, \end{aligned} \quad (5.41)$$

with

$$\begin{aligned} \gamma' &= -2.32077 \times 10^{-7} + 3.28127 \times 10^{-14}i + (0.00299432 - 4.23359 \times 10^{-10}i)A^2 \\ &\quad + (1.33377 - 1.88579 \times 10^{-7}i)A^4, \\ \Gamma' &= 0.002245 + (2 - 2.64698 \times 10^{-23}i)A^2 - (1.73176 + 1.22425 \times 10^{-7}i\sqrt{\gamma'}). \end{aligned} \quad (5.42)$$

We also note that in the normal spectrum, $|m_1| \approx |m_2| < |m_3|$, so m_1 given in (C.1) is the lightest neutrino mass. Hence, it is denoted as $m_1 \equiv m_{\text{light}}$. In figures 7a and 7b, we have plotted the value $|m_{ee}|$, $|m_\beta|$ and $|m_{\text{light}}|$ as functions of m_2 with $m_2 \in (-0.05, -0.0087)$ eV and $m_2 \in (0.0087, 0.05)$ eV, respectively.

To get explicit values of the model parameters, we assume $m_2 = 10^{-2}$ eV, which is safely small. Then the other neutrino masses are explicitly given as $m_1 \simeq -5.298 \times 10^{-3}$ eV, $m_2 \simeq 10^{-2}$ eV, $m_3 \simeq -4.95 \times 10^{-2}$ eV and $|m_{ee}| \simeq 1.09 \times 10^{-3}$ eV, $|m_\beta| \simeq 1.178 \times 10^{-2}$ eV. This solution means a normal neutrino mass spectrum as mentioned above and consistent with the recent experimental data [8, 183, 184]. It follows that

$$\begin{aligned} C &\simeq 0.0237465 - 8.39362 \times 10^{-10}i \simeq 0.0237465 \text{ eV}, \\ B_1 &= -0.0270968 - 0.00869645i, \quad B_2 = -0.0276928 + 0.0232392i. \end{aligned} \quad (5.43)$$




Figure 7. The $|m_{ee}|$, $|m_\beta|$ and $|m_{light}|$ as functions of m_2 from (C.1) in the case of $\Delta m_{23}^2 > 0$. (a) $m_2 \in (0.00867, 0.05)$ eV, (b) $m_2 \in (-0.05, -0.00867)$ eV.

Furthermore, by assuming that

$$\lambda_s = -\lambda_\sigma = -\lambda_{\sigma'} = -1 \text{ eV}, \quad v_s = v_\sigma = -v'_{\sigma'}, \quad \Lambda_s = -\Lambda_\sigma = -\Lambda'_{\sigma'} = -v^2, \quad (5.44)$$

we obtain

$$\begin{aligned} A &= 2x, & C &= y \left(-2 + \frac{4x^2}{x^2 - y^2 + xz} \right), \\ B_1 &= x \left(2 + \frac{4y^2}{x^2 - y^2 + xz} \right), & B_2 &= 2(z - x) + \frac{4x^3}{x^2 - y^2 + xz}. \end{aligned} \quad (5.45)$$

Combining (5.43) and (5.45) yields: $x \simeq 5 \times 10^{-3}$, $y \simeq (-4.52717 - 7.71265i) \times 10^{-3}$, $z \simeq (-10.4861 + 5.89481i) \times 10^{-3}$.

5.1.2 The case $\delta = \frac{3\pi}{2}$

In this case, combining (5.16) and the values of K in (5.30), we obtain

$$B_1 = B_2 + (2.01498 \times 10^{-7} + 0.732446i)C. \quad (5.46)$$

Substituting B_1 from (5.46) into (5.14) and combining with the two experimental constraints on squared mass differences of neutrinos as shown in (1.2), we obtain four solutions (in [eV]) given in appendix D.

Similar to the case $\delta = \frac{\pi}{2}$ in subsection 5.1.1, in this case we also have four solutions in which $m_{1,3}$ have the same absolute values, the unique difference is the sign of $m_{1,3}$. So, we only consider one solution in the case (D.1). The values $|m_{1,3}|$ as functions of m_2 are shown in figures 8a, 8b. Furthermore, if we assume $m_2 = 10^{-2}$ eV, which is safely small. Then the other neutrino masses are explicitly given as $m_1 \simeq 8.997 \times 10^{-3}$ eV, $m_3 \simeq 5.00 \times 10^{-3}$ eV. It follows that $C \simeq 2.37465 - 1.01188 \times 10^{-7}i$ eV, $B_1 \simeq -2.70967 + 0.869651i$, $B_2 \simeq -2.76928 - 2.32392i$. Furthermore, with the assumption (5.44), we obtain $x \simeq 5 \times 10^{-3}$, $y \simeq (-4.52717 + 7.71265i) \times 10^{-3}$, $z \simeq (-10.4861 - 5.8948i) \times 10^{-2}$.




Figure 8. $|m_{1,3}|$ as functions of m_2 from (D.1) in the case of $\Delta m_{23}^2 > 0$. (a) The $|m_{1,3}|$ as functions of m_2 with $m_2 \in (0.00867, 0.05)$ eV, (b) The $|m_{1,3}|$ as functions of m_2 with $m_2 \in (-0.05, -0.00867)$ eV.

5.2 Inverted case ($\Delta m_{23}^2 < 0$)

Similar to the normal case, in this case we also have four solutions in which $m_{1,3}$ have the same absolute values, the unique difference is the sign of $m_{1,3}$. So, in the case $K = -0.930528 - 0.366221i$, we only consider one solution in the form:

$$\begin{aligned}
 B_1 &= B_2 - (1.67146 \times 10^{-7} + 0.732442i)C, \\
 C &= 0.5\sqrt{\alpha_1 - 2\sqrt{\beta_1}}, \\
 B_2 &= 0.5\sqrt{4A^2 - 0.0003} + (8.3573 \times 10^{-8} + 0.366221i)C \\
 &\quad - 0.5\sqrt{(3.46353 + 2.4485 \times 10^{-7})C^2}, \\
 m_1 &= 0.5\sqrt{4A^2 - 0.0003}, \quad m_2 = A, \\
 m_3 &= 0.5\sqrt{4A^2 - 0.0003} - \sqrt{\Gamma}.
 \end{aligned} \tag{5.47}$$

where

$$\begin{aligned}
 \Gamma &= -0.002395 + 2.58494 \times 10^{-26}i + (2 - 2.64698 \times 10^{-23}i)A^2 \\
 &\quad - (1.73176 + 1.22425 \times 10^{-7}i)\sqrt{\beta_1}, \\
 \alpha_1 &= -0.00276597 + 1.95536 \times 10^{-10}i + (2.30978 - 1.63287 \times 10^{-7}i)A^2, \\
 \beta_1 &= (2.32077 \times 10^{-7} - 3.28127 \times 10^{-14}i) - (0.00319439 - 4.51646 \times 10^{-10}i)A^2 \\
 &\quad + (1.33377 - 1.88579 \times 10^{-7}i)A^4.
 \end{aligned} \tag{5.48}$$

In figures 9a and 9b, we have plotted the real and the imaginary part of $B_{1,2}$ and C in (5.47) as functions of m_2 with $m_2 \in (0.0482, 0.05)$ eV, respectively, in the case of $\Delta m_{23}^2 < 0$ and $\delta = \frac{\pi}{2}$. The absolute value $|m_{1,3}|$ as functions of m_2 with $m_2 \in (-0.1, -0.0482)$ eV and $m_2 \in (0.0482, 0.1)$ eV are plotted in figures 10a and 10b, respectively. These figures show that there exist allowed regions for value of m_2 (or A) where either inverted or quasi-degenerate neutrino mass hierarchy achieved. The quasi-degenerate mass hierarchy obtained when A lies in a region $[0.1 \text{ eV}, +\infty]$ or $[-\infty, -0.1 \text{ eV}]$ ($|A|$ increases but must be




Figure 9. (a) The real part of $B_{1,2}$ and C as functions of m_2 , (b) The imaginary part of $B_{1,2}$ and C as functions of m_2 in the case of $\Delta m_{23}^2 < 0$ and $\delta = \frac{\pi}{2}$.




Figure 10. (a) The $|m_{1,3}|$ as functions of m_2 in the case of $\Delta m_{23}^2 < 0$ and $\delta = \frac{\pi}{2}$ with (a) $m_2 \in (-0.0482, -0.1)$ eV, (b) $m_2 \in (0.0482, 0.1)$ eV.

small enough because of the scale of $m_{1,2,3}$). The inverted mass hierarchy will be obtained if A takes the values around $(-0.1, -0.0482)$ eV or $(0.0482, 0.1)$ eV. Figures 11a and 11b give the sum $\sum_{i=1}^3 |m_i|$ with $m_2 \in (-0.1, -0.0482)$ eV and $m_2 \in (0.0482, 0.1)$ eV, respectively.

In the inverted spectrum, $|m_3| \approx |m_1| \simeq |m_2|$, and $m_3 \equiv m_{\text{light}}$ given in (5.47) is the lightest neutrino mass. $|m_{ee}|$, $|m_\beta|$ and $|m_{\text{light}}|$ as functions of m_2 with $m_2 \in (-0.1, -0.0482)$ eV and $m_2 \in (0.0482, 0.1)$ eV are plotted in figures 12a and 12b, respectively.

In similarity to the normal case, to get explicit values of the model parameters, we assume $m_2 = 5 \times 10^{-2}$ eV, which is safely small. Then the other neutrino masses are explicitly given as $m_1 \simeq 4.925 \times 10^{-2}$ eV and $m_3 \simeq 1.342 \times 10^{-2}$ eV. It follows that $C \simeq 0.0192514 - 6.80475 \times 10^{-10}i$ eV, $B_1 \simeq 0.0313303 - 0.00705026i$ eV, $B_2 \simeq 0.0313303 + 0.00705026i$ eV. Furthermore, with the assumption (5.44), we obtain $x \simeq 2.50 \times 10^{-2}$, $y \simeq (-0.778994 + 1.4801i) \times 10^{-2}$, $z \simeq (0.0146926 + 0.0228004i) \times 10^{-2}$.

The relations between m_{ee} , m_β and m_{light} in both normal and inverted hierarchy are shown in figures 13, 14 and 15, respectively.




Figure 11. (a) The sum $\sum_{i=1}^3 m_i$ as a function of m_2 with $m_2 \in (0.0482, 0.05)$ eV in the case of $\Delta m_{23}^2 < 0$ and $K = -0.930528 - 0.366221i$, (b) The sum $\sum_{i=1}^3 |m_i|$ as a function of m_2 with $m_2 \in (0.0482, 0.05)$ eV in the case of $\Delta m_{23}^2 < 0$ and $K = -0.930528 - 0.366221i$.




Figure 12. The $|m_{ee}|$, $|m_\beta|$ and $|m_{light}|$ as functions of m_2 from (5.47) in the case of $\Delta m_{23}^2 < 0$ and $\delta = \frac{\pi}{2}$. (a) $m_2 \in (-0.1, -0.0482)$ eV, (b) $m_2 \in (0.0482, 0.1)$ eV.




Figure 13. $|m_{ee}|$ as functions of m_2 in normal and inverted hierarchy with $\delta = \frac{\pi}{2}$.




Figure 14. $|m_\beta|$ as functions of m_2 in normal and inverted hierarchy with $\delta = \frac{\pi}{2}$.




Figure 15. $|m_{light}|$ as functions of m_2 in normal and inverted hierarchy with $\delta = \frac{\pi}{2}$.

6 Remark on breaking, VEVs and rho parameter

To obtain a realistic neutrino spectrum, in this work we argue that both the breakings $T_7 \rightarrow Z_3$ and $T_7 \rightarrow \{\text{identity}\}$ must be taken place in neutrino sector while only the breaking $T_7 \rightarrow Z_3$ is taken place in charged lepton and quark sectors. The quark masses at the tree-level can be fitted but then the CKM matrix is diagonal.

A breaking of the lepton parity due to the odd VEVs $\langle \eta_3^0 \rangle$, $\langle \chi_1^0 \rangle$, or a violation of \mathcal{L} and/nor S_3 symmetry in terms of Yukawa interactions will disturb the tree level matrix resulting in mixing between SM and exotic quarks and/or possibly providing the desirable quark mixing pattern [165–167]. To get a realistic pattern of SM quarks mixing, we should add radiative correction or use the effective six dimensional operators (for details, see ref. [187]). However, detailed study on this problem is out of the scope of this work.

Note that $\Lambda_\sigma, \Lambda_s, \Lambda_{\sigma'}$ are needed to the same order and not to be so large that can naturally be taken at TeV scale as the VEV v_χ of χ . This is because v_σ, v_s and $v_{\sigma'}$ carry lepton number, simultaneously breaking the lepton parity which is naturally constrained to be much smaller than the electroweak scale [160, 161, 164–166]. This is also behind a theoretical fact that v_χ, Λ_σ are scales for the gauge symmetry breaking in the first stage

from $SU(3)_L \otimes U(1)_X \rightarrow SU(2)_L \otimes U(1)_Y$ in the original form of 3-3-1 models [160, 161, 169]. They will provide masses for the new gauge bosons: Z' , X and Y . Also, the exotic quarks gain masses from v_χ while the neutral fermions masses arise from $\Lambda_\sigma, \Lambda_s, \Lambda_{\sigma'}$. The second stage of the gauge symmetry breaking from $SU(2)_L \otimes U(1)_Y \rightarrow U(1)_Q$ is achieved by the electroweak scale VEVs such as u, v responsible for ordinary particle masses. In combination with those of type II seesaw as determined, in our model, the following limit is often taken into account [160, 161, 164–166]:

$$(\text{eV})^2 \sim \lambda_\sigma^2, \lambda_s^2, \lambda_{\sigma'}^2 \ll v_\sigma^2, v_s^2, v_{\sigma'}^2 \ll u^2, v^2 \ll v_\chi^2 \sim \Lambda_\sigma^2 \sim \Lambda_s^2 \sim \Lambda_{\sigma'}^2 \sim (\text{TeV})^2. \quad (6.1)$$

Our model contains a lot of $SU(3)_L$ scalar triplets that may modify the precision electroweak parameter such as S, T, U [188, 189] and ρ parameters. The most serious one can result from the tree-level contributions to the ρ parameter. To see this let us approximate W, Z mass and ρ parameter:⁵

$$\begin{aligned} M_W^2 &\simeq \frac{g^2}{2} v_W^2, & M_Z^2 &\simeq \frac{g^2 v_W^2}{2c_W^2}, \\ M_Y^2 &\simeq \frac{g^2}{2} (6\Lambda_\sigma^2 + 2\Lambda_s^2 + 2\Lambda_{\sigma'}^2 + v_\chi^2), \end{aligned} \quad (6.2)$$

and

$$\rho = \frac{M_W^2}{c_W^2 M_Z^2} \simeq 1 + \frac{\lambda_s^2}{v_W^2}, \quad (6.3)$$

where $v_W^2 \simeq (3u^2 + 3v^2) = (246 \text{ GeV})^2$ is naturally given according to (6.1) with $u \sim v \sim 100 \text{ GeV}$. Since $\lambda_s = 6v_\sigma^2 + 2v_s^2 + 2v_{\sigma'}^2$ is in eV scale responsible for the observed neutrino masses, the ρ in (6.3) is absolutely close to the unity and in agreement with the data [8].

The mixings between the charged gauge bosons $W - Y$ and the neutral ones $Z' - W_4$ are in the same order since they are proportional to $\frac{v_\sigma}{\Lambda_\sigma}$, and in the limit $\lambda_s, \lambda_\sigma, v_s, v_\sigma \rightarrow 0$ these mixing angles tend to zero. In addition, from (6.1) and (6.2), it follows that M_W^2 is much smaller than M_Y^2 .

7 Conclusions

In this paper, we have constructed the T_7 model based on $SU(3)_C \otimes SU(3)_L \otimes U(1)_X$ gauge symmetry responsible for fermion masses and mixing. Neutrinos get masses from only anti-sextets which are in triplets $\underline{3}$ and $\underline{3}^*$ under T_7 . The flavor mixing patterns and mass splitting are obtained without perturbation. The number of Higgs multiplets needed in order to allow the fermions to gain masses are less than those of S_3, S_4 and D_4 [165–167]. The tribimaximal form obtained with the breaking $T_7 \rightarrow Z_3$ in charged lepton sector and both $T_7 \rightarrow Z_3$ and $Z_3 \rightarrow \{\text{Identity}\}$ must be taken place in neutrino sector but only apart in

⁵We have used the notation $s_W = \sin \theta_W, c_W = \cos \theta_W, t_W = \tan \theta_W$, and the continuation of the gauge coupling constant g of the $SU(3)_L$ at the spontaneous symmetry breaking point [167, 185, 186], $t = \frac{3\sqrt{2}s_W}{\sqrt{3-4s_W^2}}$ was used.

class	n	h	χ_1	$\chi_{1'}$	$\chi_{1''}$	χ_3	χ_{3^*}
C_1	1	1	1	1	1	3	3
C_2	7	3	1	ω	ω^2	0	0
C_3	7	3	1	ω^2	ω	0	0
C_4	3	7	1	1	1	ξ	ξ^*
C_5	3	7	1	1	1	ξ^*	ξ

Table 2. Character table of T_7 group.

breakings $Z_3 \rightarrow \{\text{Identity}\}$ (without contribution of σ'), and the upper bound on neutrino mass $\sum_{i=1}^3 m_i$ at the level is presented. From the Dirac CP violation phase we obtain the relation between Euler's angles which is consistent with the experimental in PDG 2012. On the other hand, the realistic lepton mixing can be obtained if both the direction for breakings $T_7 \rightarrow Z_3$ and $Z_3 \rightarrow \{\text{Identity}\}$ are taken place in neutrino sectors. The CKM matrix is the identity matrix at the tree-level. The Dirac CP violation phase δ is predicted to either $\frac{\pi}{2}$ or $\frac{3\pi}{2}$ which is maximal CP violation.

Acknowledgments

This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 103.01-2011.63.

A T_7 group and Clebsch-Gordan coefficients

The tetrahedral group A_4 has 12 elements and four equivalence classes with three inequivalent one-dimensional representations and one three-dimensional one, which is the smallest group with only a real $\underline{3}$ representation. The Frobenius group T_7 has 21 elements and five equivalence classes with three inequivalent one-dimensional representations and two three-dimensional ones, which is the smallest group with a pair of complex $\underline{3}$ and $\underline{3}^*$ representations. It is generated by

$$a = \begin{pmatrix} \rho & 0 & 0 \\ 0 & \rho^2 & 0 \\ 0 & 0 & \rho^4 \end{pmatrix}, \quad b = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \tag{A.1}$$

where $\rho = \exp(2\pi i/7)$, so that $a^7 = 1$, $b^3 = 1$, and $ab = ba^4$. The character table of T_7 is given in table 2, where $\xi = -1/2 + i\sqrt{7}/2$.

The conjugacy classes generated from a and b are

$$\begin{aligned} C_1 &: \{e\}, \\ C_2 &: \{b, ba, ba^2, ba^3, ba^4, ba^5, ba^6\}, \\ C_3 &: \{b^2, b^2a, b^2a^2, b^2a^3, b^2a^4, b^2a^5, b^2a^6\}, \end{aligned}$$

$$\begin{aligned}
 C_4 &: \{a, a^2, a^4\}, \\
 C_5 &: \{a^3, a^5, a^6\}.
 \end{aligned}
 \tag{A.2}$$

Let us put $\underline{3}(1, 2, 3)$ which means some $\underline{3}$ multiplet such as $x = (x_1, x_2, x_3) \sim \underline{3}$ or $y = (y_1, y_2, y_3) \sim \underline{3}$ and so on, and similarly for the other representations. Moreover, the numbered multiplets such as (\dots, ij, \dots) mean $(\dots, x_i y_j, \dots)$ where x_i and y_j are the multiplet components of different representations x and y , respectively. In the following the components of representations in l.h.s. will be omitted and should be understood, but they always exist in order in the components of decompositions in r.h.s. All the group multiplication rules of T_7 as given below.

$$\begin{aligned}
 \underline{1} \otimes \underline{1} &= \underline{1}(11), & \underline{1} \otimes \underline{1}' &= \underline{1}'(11), & \underline{1} \otimes \underline{1}'' &= \underline{1}''(11), \\
 \underline{1}' \otimes \underline{1}'' &= \underline{1}(11), & \underline{1}' \otimes \underline{1}' &= \underline{1}''(11), & \underline{1}'' \otimes \underline{1}'' &= \underline{1}'(11), \\
 \underline{1} \otimes \underline{3} &= \underline{3}(11, 12, 13), & \underline{1}' \otimes \underline{3} &= \underline{3}(11, \omega 12, \omega^2 13), & \underline{1}'' \otimes \underline{3} &= \underline{3}(11, \omega^2 12, \omega 13), \\
 \underline{1} \otimes \underline{3}^* &= \underline{3}^*(11, 12, 13), & \underline{1}' \otimes \underline{3}^* &= \underline{3}^*(11, \omega 12, \omega^2 13), & \underline{1}'' \otimes \underline{3}^* &= \underline{3}^*(11, \omega^2 12, \omega 13), \\
 \underline{3} \otimes \underline{3} &= \underline{3}(33, 11, 22) \oplus \underline{3}^*(23, 31, 12) \oplus \underline{3}^*(32, 13, 21), \\
 \underline{3}^* \otimes \underline{3}^* &= \underline{3}^*(33, 11, 22) \oplus \underline{3}(23, 31, 12) \oplus \underline{3}(32, 13, 21), \\
 \underline{3} \otimes \underline{3}^* &= \underline{1}(11 + 22 + 33) \oplus \underline{1}'(11 + \omega 22 + \omega^2 33) \\
 &\quad \oplus \underline{1}''(11 + \omega^2 22 + \omega 33) \oplus \underline{3}(21, 32, 13) \oplus \underline{3}^*(12, 23, 31).
 \end{aligned}
 \tag{A.3}$$

Note that $\underline{3} \times \underline{3} \times \underline{3}$ has two invariants and $\underline{3} \times \underline{3} \times \underline{3}^*$ has one invariant.

B The numbers

In the following we will explicitly point out the lepton number (L) and lepton parity (P_l) of the model particles (notice that the family indices are suppressed):

Particles	L	P_l
$N_R, u, d, \phi_1^+, \phi_1'^+, \phi_2^0, \phi_2'^0, \eta_1^0, \eta_1'^0, \eta_2^-, \eta_2'^-, \chi_3^0, \sigma_{33}^0, s_{33}^0$	0	1
$\nu_L, l, U, D^*, \phi_3^+, \phi_3'^+, \eta_3^0, \eta_3'^0, \chi_1^{0*}, \chi_2^+, \sigma_{13}^0, \sigma_{23}^+, s_{13}^0, s_{23}^+$	-1	-1
$\sigma_{11}^0, \sigma_{12}^+, \sigma_{22}^{++}, s_{11}^0, s_{12}^+, s_{22}^{++}$	-2	1

C The solutions with $\delta = \frac{\pi}{2}$ in the normal case

- The first case:

$$\begin{aligned}
 C &= 0.5\sqrt{\alpha - 2\sqrt{\beta}}, \\
 B_2 &= -0.5\sqrt{4A^2 - 0.0003} + (8.3573 \times 10^{-8} + 0.366221i)C \\
 &\quad - 0.5\sqrt{(3.46353 + 2.4485 \times 10^{-7})C^2}, \\
 m_1 &= -0.5\sqrt{4A^2 - 0.0003}, \quad m_2 = A, \\
 m_3 &= -0.5\sqrt{4A^2 - 0.0003} \\
 &\quad - \sqrt{0.002245 + (2 - 2.64698 \times 10^{-23}i)A^2 - (1.73176 + 1.22425 \times 10^{-7}i)\sqrt{\beta}}.
 \end{aligned}
 \tag{C.1}$$

- The second case:

$$\begin{aligned}
 C &= 0.5\sqrt{\alpha + 2\sqrt{\beta}}, \\
 B_2 &= -0.5\sqrt{4A^2 - 0.0003} + (8.3573 \times 10^{-8} + 0.366221i)C \\
 &\quad - 0.5\sqrt{(3.46353 + 2.4485 \times 10^{-7})C^2}, \\
 m_1 &= -0.5\sqrt{4A^2 - 0.0003}, \quad m_2 = A, \\
 m_3 &= -0.5\sqrt{4A^2 - 0.0003} \\
 &\quad - \sqrt{0.002245 + (2 - 2.64698 \times 10^{-23}i)A^2 + (1.73176 + 1.22425 \times 10^{-7}i)\sqrt{\beta}},
 \end{aligned} \tag{C.2}$$

- The third case:

$$\begin{aligned}
 C &= -0.5\sqrt{\alpha - 2\sqrt{\beta}}, \\
 B_2 &= 0.5\sqrt{4A^2 - 0.0003} + (8.3573 \times 10^{-8} + 0.366221i)C \\
 &\quad - 0.5\sqrt{(3.46353 + 2.4485 \times 10^{-7})C^2}, \\
 m_1 &= 0.5\sqrt{4A^2 - 0.0003}, \quad m_2 = A, \\
 m_3 &= 0.5\sqrt{4A^2 - 0.0003} \\
 &\quad - \sqrt{0.002245 + (2 - 2.64698 \times 10^{-23}i)A^2 + (1.73176 + 1.22425 \times 10^{-7}i)\sqrt{\beta}},
 \end{aligned} \tag{C.3}$$

- The fourth case:

$$\begin{aligned}
 C &= 0.5\sqrt{\alpha + 2\sqrt{\beta}}, \\
 B_2 &= 0.5\sqrt{4A^2 - 0.0003} + (8.3573 \times 10^{-8} + 0.366221i)C \\
 &\quad - 0.5\sqrt{(3.46353 + 2.4485 \times 10^{-7})C^2}, \\
 m_1 &= 0.5\sqrt{4A^2 - 0.0003}, \quad m_2 = A, \\
 m_3 &= 0.5\sqrt{4A^2 - 0.0003} \\
 &\quad - \sqrt{0.002245 + (2 - 2.64698 \times 10^{-23}i)A^2 + (1.73176 + 1.22425 \times 10^{-7}i)\sqrt{\beta}},
 \end{aligned} \tag{C.4}$$

where

$$\begin{aligned}
 \alpha &= (0.00259273 - 1.8329 \times 10^{-10}i) + (2.30978 - 1.63287 \times 10^{-7}i)A^2, \\
 \beta &= -2.32077 \times 10^{-7} + 3.28127 \times 10^{-14}i \\
 &\quad + (0.00299432 - 4.23359 \times 10^{-10}i)A^2 + (1.33377 - 1.88579 \times 10^{-7}i)A^4.
 \end{aligned} \tag{C.5}$$

D The solutions with $\delta = \frac{3\pi}{2}$ in the normal case

- The first case:

$$\begin{aligned}
 C &= 0.5\sqrt{\alpha' - 2\sqrt{\beta'}}, \\
 B_2 &= -0.5\sqrt{4A^2 - 0.0003} + (1.00749 \times 10^{-7} + 0.366223i)C \\
 &\quad - 0.5\sqrt{(3.46352 + 2.95173 \times 10^{-7})C^2},
 \end{aligned}$$

$$\begin{aligned}
 m_1 &= -0.5\sqrt{4A^2 - 0.0003}, & m_2 &= A, \\
 m_3 &= -0.5\sqrt{4A^2 - 0.0003} \\
 &\quad - \sqrt{0.002245 + 2.58494 \times 10^{-26}i + 2A^2 - (1.73176 + 1.47587 \times 10^{-7}i)\sqrt{\beta'}}.
 \end{aligned} \tag{D.1}$$

- The second case:

$$\begin{aligned}
 C &= 0.5\sqrt{\alpha + 2\sqrt{\beta}}, \\
 B_2 &= -0.5\sqrt{4A^2 - 0.0003} + (1.00749 \times 10^{-7} + 0.366223i)C \\
 &\quad - 0.5\sqrt{(3.46352 + 2.95173 \times 10^{-7})C^2}, \\
 m_1 &= -0.5\sqrt{4A^2 - 0.0003}, & m_2 &= A, \\
 m_3 &= -0.5\sqrt{4A^2 - 0.0003} \\
 &\quad - \sqrt{0.002245 + 2.58494 \times 10^{-26}i + 2A^2 + (1.73176 + 1.47587 \times 10^{-7}i)\sqrt{\beta'}}.
 \end{aligned} \tag{D.2}$$

- The third case:

$$\begin{aligned}
 C &= 0.5\sqrt{\alpha' - 2\sqrt{\beta'}}, \\
 B_2 &= 0.5\sqrt{4A^2 - 0.0003} - (1.00749 \times 10^{-7} + 0.366223i)C \\
 &\quad - 0.5\sqrt{(3.46352 + 2.95173 \times 10^{-7})C^2}, \\
 m_1 &= 0.5\sqrt{4A^2 - 0.0003}, & m_2 &= A, \\
 m_3 &= 0.5\sqrt{4A^2 - 0.0003} \\
 &\quad - \sqrt{0.002245 + 2.58494 \times 10^{-26}i + 2A^2 - (1.73176 + 1.47587 \times 10^{-7}i)\sqrt{\beta'}}.
 \end{aligned} \tag{D.3}$$

- The fourth case:

$$\begin{aligned}
 C &= 0.5\sqrt{\alpha' + 2\sqrt{\beta'}}, \\
 B_2 &= 0.5\sqrt{4A^2 - 0.0003} - (1.00749 \times 10^{-7} + 0.366223i)C \\
 &\quad - 0.5\sqrt{(3.46352 + 2.95173 \times 10^{-7})C^2}, \\
 m_1 &= 0.5\sqrt{4A^2 - 0.0003}, & m_2 &= A, \\
 m_3 &= 0.5\sqrt{4A^2 - 0.0003} \\
 &\quad - \sqrt{0.002245 + (2 - 2.64698 \times 10^{-23}i)A^2 + (1.73176 + 1.47587 \times 10^{-7}i)\sqrt{\beta'}},
 \end{aligned} \tag{D.4}$$

where

$$\begin{aligned}
 \alpha' &= (0.00259274 - 2.20962 \times 10^{-10}i) + (2.30979 - 1.96848 \times 10^{-7}i)A^2, \\
 \beta' &= -2.32078 \times 10^{-7} + 3.95569 \times 10^{-14}i \\
 &\quad + (0.00299433 - 5.10375 \times 10^{-10}i)A^2 + (1.33378 - 2.27338 \times 10^{-7}i)A^4.
 \end{aligned} \tag{D.5}$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] G. Altarelli, *An overview of neutrino mixing*, *Nucl. Phys. Proc. Suppl.* **241-242** (2013) 77 [[arXiv:1210.3467](#)] [[INSPIRE](#)].
- [2] G. Altarelli, *Neutrino mixing: theoretical overview*, [arXiv:1304.5047](#) [[INSPIRE](#)].
- [3] P. Harrison, D. Perkins and W.G. Scott, *Tri-bimaximal mixing and the neutrino oscillation data*, *Phys. Lett. B* **530** (2002) 167 [[hep-ph/0202074](#)] [[INSPIRE](#)].
- [4] Z.-z. Xing, *Nearly tri bimaximal neutrino mixing and CP-violation*, *Phys. Lett. B* **533** (2002) 85 [[hep-ph/0204049](#)] [[INSPIRE](#)].
- [5] X.G. He and A. Zee, *Some simple mixing and mass matrices for neutrinos*, *Phys. Lett. B* **560** (2003) 87 [[hep-ph/0301092](#)] [[INSPIRE](#)].
- [6] X.-G. He and A. Zee, *Neutrino masses with 'zero sum' condition: $m(\nu_1) + m(\nu_2) + m(\nu_3) = 0$* , *Phys. Rev. D* **68** (2003) 037302 [[hep-ph/0302201](#)] [[INSPIRE](#)].
- [7] A.Y. Smirnov, *Neutrino 2012: outlook — Theory*, *Nucl. Phys. Proc. Suppl.* **235-236** (2013) 431 [[arXiv:1210.4061](#)] [[INSPIRE](#)].
- [8] PARTICLE DATA GROUP collaboration, J. Beringer et al., *Review of particle physics*, *Phys. Rev. D* **86** (2012) 010001 [[INSPIRE](#)].
- [9] T. Schwetz, M. Tortola and J. Valle, *Where we are on θ_{13} : addendum to 'Global neutrino data and recent reactor fluxes: status of three-flavour oscillation parameters'*, *New J. Phys.* **13** (2011) 109401 [[arXiv:1108.1376](#)] [[INSPIRE](#)].
- [10] T2K collaboration, K. Abe et al., *Indication of electron neutrino appearance from an accelerator-produced off-axis muon neutrino beam*, *Phys. Rev. Lett.* **107** (2011) 041801 [[arXiv:1106.2822](#)] [[INSPIRE](#)].
- [11] MINOS collaboration, P. Adamson et al., *Improved search for muon-neutrino to electron-neutrino oscillations in MINOS*, *Phys. Rev. Lett.* **107** (2011) 181802 [[arXiv:1108.0015](#)] [[INSPIRE](#)].
- [12] G. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. Rotunno, *Evidence of $\theta_{13} > 0$ from global neutrino data analysis*, *Phys. Rev. D* **84** (2011) 053007 [[arXiv:1106.6028](#)] [[INSPIRE](#)].
- [13] N. Cabibbo, *Unitary symmetry and leptonic decays*, *Phys. Rev. Lett.* **10** (1963) 531 [[INSPIRE](#)].
- [14] M. Kobayashi and T. Maskawa, *CP violation in the renormalizable theory of weak interaction*, *Prog. Theor. Phys.* **49** (1973) 652 [[INSPIRE](#)].
- [15] E. Ma and G. Rajasekaran, *Softly broken A_4 symmetry for nearly degenerate neutrino masses*, *Phys. Rev. D* **64** (2001) 113012 [[hep-ph/0106291](#)] [[INSPIRE](#)].
- [16] K. Babu, E. Ma and J. Valle, *Underlying A_4 symmetry for the neutrino mass matrix and the quark mixing matrix*, *Phys. Lett. B* **552** (2003) 207 [[hep-ph/0206292](#)] [[INSPIRE](#)].
- [17] G. Altarelli and F. Feruglio, *Tri-bimaximal neutrino mixing from discrete symmetry in extra dimensions*, *Nucl. Phys. B* **720** (2005) 64 [[hep-ph/0504165](#)] [[INSPIRE](#)].
- [18] E. Ma, *Tribimaximal neutrino mixing from a supersymmetric model with A_4 family symmetry*, *Phys. Rev. D* **73** (2006) 057304 [[hep-ph/0511133](#)] [[INSPIRE](#)].

- [19] X.-G. He, Y.-Y. Keum and R.R. Volkas, *A_4 flavor symmetry breaking scheme for understanding quark and neutrino mixing angles*, *JHEP* **04** (2006) 039 [[hep-ph/0601001](#)] [[INSPIRE](#)].
- [20] S. Morisi, M. Picariello and E. Torrente-Lujan, *Model for fermion masses and lepton mixing in $SO(10) \times A_4$* , *Phys. Rev. D* **75** (2007) 075015 [[hep-ph/0702034](#)] [[INSPIRE](#)].
- [21] C. Lam, *Symmetry of lepton mixing*, *Phys. Lett. B* **656** (2007) 193 [[arXiv:0708.3665](#)] [[INSPIRE](#)].
- [22] F. Bazzocchi, S. Kaneko and S. Morisi, *A SUSY A_4 model for fermion masses and mixings*, *JHEP* **03** (2008) 063 [[arXiv:0707.3032](#)] [[INSPIRE](#)].
- [23] A. Blum, C. Hagedorn and M. Lindner, *Fermion masses and mixings from dihedral flavor symmetries with preserved subgroups*, *Phys. Rev. D* **77** (2008) 076004 [[arXiv:0709.3450](#)] [[INSPIRE](#)].
- [24] F. Bazzocchi, M. Frigerio and S. Morisi, *Fermion masses and mixing in models with $SO(10) \times A_4$ symmetry*, *Phys. Rev. D* **78** (2008) 116018 [[arXiv:0809.3573](#)] [[INSPIRE](#)].
- [25] G. Altarelli, F. Feruglio and C. Hagedorn, *A SUSY $SU(5)$ grand unified model of tri-bimaximal mixing from A_4* , *JHEP* **03** (2008) 052 [[arXiv:0802.0090](#)] [[INSPIRE](#)].
- [26] M. Hirsch, S. Morisi and J. Valle, *Tri-bimaximal neutrino mixing and neutrinoless double β decay*, *Phys. Rev. D* **78** (2008) 093007 [[arXiv:0804.1521](#)] [[INSPIRE](#)].
- [27] E. Ma, *Dark scalar doublets and neutrino tribimaximal mixing from A_4 symmetry*, *Phys. Lett. B* **671** (2009) 366 [[arXiv:0808.1729](#)] [[INSPIRE](#)].
- [28] G. Altarelli and D. Meloni, *A simplest A_4 model for tri-bimaximal neutrino mixing*, *J. Phys. G* **36** (2009) 085005 [[arXiv:0905.0620](#)] [[INSPIRE](#)].
- [29] Y. Lin, *A predictive A_4 model, charged lepton hierarchy and tri-bimaximal sum rule*, *Nucl. Phys. B* **813** (2009) 91 [[arXiv:0804.2867](#)] [[INSPIRE](#)].
- [30] Y. Ahn and C.-S. Chen, *Non-zero U_{e3} and TeV-leptogenesis through A_4 symmetry breaking*, *Phys. Rev. D* **81** (2010) 105013 [[arXiv:1001.2869](#)] [[INSPIRE](#)].
- [31] J. Barry and W. Rodejohann, *Deviations from tribimaximal mixing due to the vacuum expectation value misalignment in A_4 models*, *Phys. Rev. D* **81** (2010) 093002 [Erratum *ibid.* **D 81** (2010) 119901] [[arXiv:1003.2385](#)] [[INSPIRE](#)].
- [32] G.-J. Ding and D. Meloni, *A model for tri-bimaximal mixing from a completely broken A_4* , *Nucl. Phys. B* **855** (2012) 21 [[arXiv:1108.2733](#)] [[INSPIRE](#)].
- [33] L. Lavoura and H. Kuhbock, *A_4 model for the quark mass matrices*, *Eur. Phys. J. C* **55** (2008) 303 [[arXiv:0711.0670](#)] [[INSPIRE](#)].
- [34] P.V. Dong, L.T. Hue, H.N. Long and D.V. Soa, *The 3-3-1 model with A_4 flavor symmetry*, *Phys. Rev. D* **81** (2010) 053004 [[arXiv:1001.4625](#)] [[INSPIRE](#)].
- [35] A. Datta, F.S. Ling and P. Ramond, *Correlated hierarchy, Dirac masses and large mixing angles*, *Nucl. Phys. B* **671** (2003) 383 [[hep-ph/0306002](#)] [[INSPIRE](#)].
- [36] C. Luhn, S. Nasri and P. Ramond, *Simple finite non-abelian flavor groups*, *J. Math. Phys.* **48** (2007) 123519 [[arXiv:0709.1447](#)] [[INSPIRE](#)].
- [37] Y. Kajiyama, M. Raidal and A. Strumia, *The Golden ratio prediction for the solar neutrino mixing*, *Phys. Rev. D* **76** (2007) 117301 [[arXiv:0705.4559](#)] [[INSPIRE](#)].

- [38] C. Luhn and P. Ramond, *Quintics with finite simple symmetries*, *J. Math. Phys.* **49** (2008) 053525 [[arXiv:0803.0526](#)] [[INSPIRE](#)].
- [39] L.L. Everett and A.J. Stuart, *Icosahedral (A_5) family symmetry and the golden ratio prediction for solar neutrino mixing*, *Phys. Rev. D* **79** (2009) 085005 [[arXiv:0812.1057](#)] [[INSPIRE](#)].
- [40] A. Adulpravitchai, A. Blum and W. Rodejohann, *Golden ratio prediction for solar neutrino mixing*, *New J. Phys.* **11** (2009) 063026 [[arXiv:0903.0531](#)] [[INSPIRE](#)].
- [41] C.-S. Chen, T.W. Kephart and T.-C. Yuan, *An A_5 model of four lepton generations*, *JHEP* **04** (2011) 015 [[arXiv:1011.3199](#)] [[INSPIRE](#)].
- [42] I.K. Cooper, S.F. King and A.J. Stuart, *A golden A_5 model of leptons with a minimal NLO correction*, *Nucl. Phys. B* **875** (2013) 650 [[arXiv:1212.1066](#)] [[INSPIRE](#)].
- [43] C.-S. Chen, T.W. Kephart and T.-C. Yuan, *Binary icosahedral flavor symmetry for four generations of quarks and leptons*, *PTEP* **2013** (2013) 103B01 [[arXiv:1110.6233](#)] [[INSPIRE](#)].
- [44] K. Hashimoto and H. Okada, *Lepton flavor model and decaying dark matter in the binary icosahedral group symmetry*, [arXiv:1110.3640](#) [[INSPIRE](#)].
- [45] G.-J. Ding, L.L. Everett and A.J. Stuart, *Golden ratio neutrino mixing and A_5 flavor symmetry*, *Nucl. Phys. B* **857** (2012) 219 [[arXiv:1110.1688](#)] [[INSPIRE](#)].
- [46] F. Feruglio and A. Paris, *The golden ratio prediction for the solar angle from a natural model with A_5 flavour symmetry*, *JHEP* **03** (2011) 101 [[arXiv:1101.0393](#)] [[INSPIRE](#)].
- [47] L.L. Everett and A.J. Stuart, *The double cover of the icosahedral symmetry group and quark mass textures*, *Phys. Lett. B* **698** (2011) 131 [[arXiv:1011.4928](#)] [[INSPIRE](#)].
- [48] L. Wolfenstein, *Oscillations among three neutrino types and CP-violation*, *Phys. Rev. D* **18** (1978) 958 [[INSPIRE](#)].
- [49] S. Pakvasa and H. Sugawara, *Discrete symmetry and Cabibbo angle*, *Phys. Lett. B* **73** (1978) 61 [[INSPIRE](#)].
- [50] S. Pakvasa and H. Sugawara, *Mass of the t quark in $SU(2) \times U(1)$* , *Phys. Lett. B* **82** (1979) 105 [[INSPIRE](#)].
- [51] E. Derman and H.-S. Tsao, *$SU(2) \times U(1) \times S(n)$ flavor dynamics and a bound on the number of flavors*, *Phys. Rev. D* **20** (1979) 1207 [[INSPIRE](#)].
- [52] Y. Yamanaka, H. Sugawara and S. Pakvasa, *Permutation symmetries and the fermion mass matrix*, *Phys. Rev. D* **25** (1982) 1895 [*Erratum ibid.* **D 29** (1984) 2135] [[INSPIRE](#)].
- [53] K. Kang, J.E. Kim and P. Ko, *A simple modification of the maximal mixing scenario for three light neutrinos*, *Z. Phys. C* **72** (1996) 671 [[hep-ph/9503436](#)] [[INSPIRE](#)].
- [54] H. Fritzsch and Z.-Z. Xing, *Lepton mass hierarchy and neutrino oscillations*, *Phys. Lett. B* **372** (1996) 265 [[hep-ph/9509389](#)] [[INSPIRE](#)].
- [55] K. Kang, S.K. Kang, J.E. Kim and P.-w. Ko, *Almost maximally broken permutation symmetry for neutrino mass matrix*, *Mod. Phys. Lett. A* **12** (1997) 1175 [[hep-ph/9611369](#)] [[INSPIRE](#)].
- [56] M. Fukugita, M. Tanimoto and T. Yanagida, *Atmospheric neutrino oscillation and a phenomenological lepton mass matrix*, *Phys. Rev. D* **57** (1998) 4429 [[hep-ph/9709388](#)] [[INSPIRE](#)].

- [57] H. Fritzsch and Z.-z. Xing, *Large leptonic flavor mixing and the mass spectrum of leptons*, *Phys. Lett. B* **440** (1998) 313 [[hep-ph/9808272](#)] [[INSPIRE](#)].
- [58] Y. Koide, *Universal seesaw mass matrix model with an S_3 symmetry*, *Phys. Rev. D* **60** (1999) 077301 [[hep-ph/9905416](#)] [[INSPIRE](#)].
- [59] H. Fritzsch and Z.-z. Xing, *Maximal neutrino mixing and maximal CP-violation*, *Phys. Rev. D* **61** (2000) 073016 [[hep-ph/9909304](#)] [[INSPIRE](#)].
- [60] M. Tanimoto, *Large mixing angle MSW solution in S_3 flavor symmetry*, *Phys. Lett. B* **483** (2000) 417 [[hep-ph/0001306](#)] [[INSPIRE](#)].
- [61] G. Branco and J. Silva-Marcos, *The symmetry behind extended flavor democracy and large leptonic mixing*, *Phys. Lett. B* **526** (2002) 104 [[hep-ph/0106125](#)] [[INSPIRE](#)].
- [62] M. Fujii, K. Hamaguchi and T. Yanagida, *Leptogenesis with almost degenerate Majorana neutrinos*, *Phys. Rev. D* **65** (2002) 115012 [[hep-ph/0202210](#)] [[INSPIRE](#)].
- [63] J. Kubo, A. Mondragon, M. Mondragon and E. Rodriguez-Jauregui, *The flavor symmetry*, *Prog. Theor. Phys.* **109** (2003) 795 [*Erratum ibid.* **114** (2005) 287] [[hep-ph/0302196](#)] [[INSPIRE](#)].
- [64] P. Harrison and W. Scott, *Permutation symmetry, tri-bimaximal neutrino mixing and the S^3 group characters*, *Phys. Lett. B* **557** (2003) 76 [[hep-ph/0302025](#)] [[INSPIRE](#)].
- [65] S.-L. Chen, M. Frigerio and E. Ma, *Large neutrino mixing and normal mass hierarchy: a discrete understanding*, *Phys. Rev. D* **70** (2004) 073008 [*Erratum ibid.* **D 70** (2004) 079905] [[hep-ph/0404084](#)] [[INSPIRE](#)].
- [66] H. Fritzsch and Z.-z. Xing, *Democratic neutrino mixing reexamined*, *Phys. Lett. B* **598** (2004) 237 [[hep-ph/0406206](#)] [[INSPIRE](#)].
- [67] F. Caravaglios and S. Morisi, *Neutrino masses and mixings with an S_3 family permutation symmetry*, [hep-ph/0503234](#) [[INSPIRE](#)].
- [68] W. Grimus and L. Lavoura, *$S_3 \times Z_2$ model for neutrino mass matrices*, *JHEP* **08** (2005) 013 [[hep-ph/0504153](#)] [[INSPIRE](#)].
- [69] R. Mohapatra, S. Nasri and H.-B. Yu, *S_3 symmetry and tri-bimaximal mixing*, *Phys. Lett. B* **639** (2006) 318 [[hep-ph/0605020](#)] [[INSPIRE](#)].
- [70] R. Jora, S. Nasri and J. Schechter, *An approach to permutation symmetry for the electroweak theory*, *Int. J. Mod. Phys. A* **21** (2006) 5875 [[hep-ph/0605069](#)] [[INSPIRE](#)].
- [71] J.E. Kim and J.-C. Park, *Quantum numbers of heavy neutrinos, tri-bi-maximal mixing through double seesaw with permutation symmetry and comment on $\theta_{\text{sol}} + \theta_c \simeq \pi/4$* , *JHEP* **05** (2006) 017 [[hep-ph/0512130](#)] [[INSPIRE](#)].
- [72] Y. Koide, *S_3 symmetry and neutrino masses and mixings*, *Eur. Phys. J. C* **50** (2007) 809 [[hep-ph/0612058](#)] [[INSPIRE](#)].
- [73] A. Mondragon, M. Mondragon and E. Peinado, *Lepton masses, mixings and FCNC in a minimal $S(3)$ -invariant extension of the standard model*, *Phys. Rev. D* **76** (2007) 076003 [[arXiv:0706.0354](#)] [[INSPIRE](#)].
- [74] A. Mondragon, M. Mondragon and E. Peinado, *Nearly tri-bimaximal mixing in the S_3 flavour symmetry*, *AIP Conf. Proc.* **1026** (2008) 164 [[arXiv:0712.2488](#)] [[INSPIRE](#)].
- [75] M. Picariello, *Neutrino CP-violating parameters from nontrivial quark-lepton correlation: a $S_3 \times GUT$ model*, *Int. J. Mod. Phys. A* **23** (2008) 4435 [[hep-ph/0611189](#)] [[INSPIRE](#)].

- [76] C.-Y. Chen and L. Wolfenstein, *Consequences of approximate S_3 symmetry of the neutrino mass matrix*, *Phys. Rev. D* **77** (2008) 093009 [[arXiv:0709.3767](#)] [[INSPIRE](#)].
- [77] R. Jora, J. Schechter and M. Naeem Shahid, *Perturbed S_3 neutrinos*, *Phys. Rev. D* **80** (2009) 093007 [*Erratum ibid.* **D 82** (2010) 079902] [[arXiv:0909.4414](#)] [[INSPIRE](#)].
- [78] D.A. Dicus, S.-F. Ge and W.W. Repko, *Neutrino mixing with broken S_3 symmetry*, *Phys. Rev. D* **82** (2010) 033005 [[arXiv:1004.3266](#)] [[INSPIRE](#)].
- [79] Z.-z. Xing, D. Yang and S. Zhou, *Broken S_3 flavor symmetry of leptons and quarks: mass spectra and flavor mixing patterns*, *Phys. Lett. B* **690** (2010) 304 [[arXiv:1004.4234](#)] [[INSPIRE](#)].
- [80] R. Jora, J. Schechter and M.N. Shahid, *Doubly perturbed S_3 neutrinos and the s_{13} mixing parameter*, *Phys. Rev. D* **82** (2010) 053006 [[arXiv:1006.3307](#)] [[INSPIRE](#)].
- [81] S. Dev, S. Gupta and R.R. Gautam, *Broken S_3 symmetry in the neutrino mass matrix*, *Phys. Lett. B* **702** (2011) 28 [[arXiv:1106.3873](#)] [[INSPIRE](#)].
- [82] D. Meloni, S. Morisi and E. Peinado, *Fritzsch neutrino mass matrix from S_3 symmetry*, *J. Phys. G* **38** (2011) 015003 [[arXiv:1005.3482](#)] [[INSPIRE](#)].
- [83] G. Bhattacharyya, P. Leser and H. Pas, *Exotic Higgs boson decay modes as a harbinger of S_3 flavor symmetry*, *Phys. Rev. D* **83** (2011) 011701 [[arXiv:1006.5597](#)] [[INSPIRE](#)].
- [84] T. Kaneko and H. Sugawara, *Broken S_3 symmetry in flavor physics*, *Phys. Lett. B* **697** (2011) 329 [[arXiv:1011.5748](#)] [[INSPIRE](#)].
- [85] S. Zhou, *Relatively large θ_{13} and nearly maximal θ_{23} from the approximate S_3 symmetry of lepton mass matrices*, *Phys. Lett. B* **704** (2011) 291 [[arXiv:1106.4808](#)] [[INSPIRE](#)].
- [86] F. González Canales, A. Mondragón, M. Mondragón, U.J. Saldaña Salazar and L. Velasco-Sevilla, *Quark sector of S_3 models: classification and comparison with experimental data*, *Phys. Rev. D* **88** (2013) 096004 [[arXiv:1304.6644](#)] [[INSPIRE](#)].
- [87] E. Ma and B. Melic, *Updated S_3 model of quarks*, *Phys. Lett. B* **725** (2013) 402 [[arXiv:1303.6928](#)] [[INSPIRE](#)].
- [88] R. Mohapatra, M. Parida and G. Rajasekaran, *High scale mixing unification and large neutrino mixing angles*, *Phys. Rev. D* **69** (2004) 053007 [[hep-ph/0301234](#)] [[INSPIRE](#)].
- [89] C. Hagedorn, M. Lindner and R. Mohapatra, *S_4 flavor symmetry and fermion masses: towards a grand unified theory of flavor*, *JHEP* **06** (2006) 042 [[hep-ph/0602244](#)] [[INSPIRE](#)].
- [90] E. Ma, *Neutrino mass matrix from S_4 symmetry*, *Phys. Lett. B* **632** (2006) 352 [[hep-ph/0508231](#)] [[INSPIRE](#)].
- [91] H. Zhang, *Flavor $S_4 \times Z_2$ symmetry and neutrino mixing*, *Phys. Lett. B* **655** (2007) 132 [[hep-ph/0612214](#)] [[INSPIRE](#)].
- [92] Y. Koide, *S_4 flavor symmetry embedded into SU(3) and lepton masses and mixing*, *JHEP* **08** (2007) 086 [[arXiv:0705.2275](#)] [[INSPIRE](#)].
- [93] H. Ishimori, Y. Shimizu and M. Tanimoto, *S_4 flavor symmetry of quarks and leptons in SU(5) GUT*, *Prog. Theor. Phys.* **121** (2009) 769 [[arXiv:0812.5031](#)] [[INSPIRE](#)].
- [94] F. Bazzocchi, L. Merlo and S. Morisi, *Fermion masses and mixings in a S_4 -based model*, *Nucl. Phys. B* **816** (2009) 204 [[arXiv:0901.2086](#)] [[INSPIRE](#)].

- [95] F. Bazzocchi and S. Morisi, S_4 as a natural flavor symmetry for lepton mixing, *Phys. Rev. D* **80** (2009) 096005 [[arXiv:0811.0345](#)] [[INSPIRE](#)].
- [96] G. Altarelli and F. Feruglio, Discrete flavor symmetries and models of neutrino mixing, *Rev. Mod. Phys.* **82** (2010) 2701 [[arXiv:1002.0211](#)] [[INSPIRE](#)].
- [97] G.-J. Ding, Fermion masses and flavor mixings in a model with S_4 flavor symmetry, *Nucl. Phys. B* **827** (2010) 82 [[arXiv:0909.2210](#)] [[INSPIRE](#)].
- [98] Y. Ahn, S.K. Kang, C. Kim and T.P. Nguyen, A direct link between neutrinoless double beta decay and leptogenesis in a seesaw model with S_4 symmetry, *Phys. Rev. D* **82** (2010) 093005 [[arXiv:1004.3469](#)] [[INSPIRE](#)].
- [99] Y. Daikoku and H. Okada, Cancellation mechanism of FCNCs in $S_4 \times Z_2$ flavor symmetric extra U(1) model, [arXiv:1008.0914](#) [[INSPIRE](#)].
- [100] H. Ishimori, Y. Shimizu, M. Tanimoto and A. Watanabe, Neutrino masses and mixing from S_4 flavor twisting, *Phys. Rev. D* **83** (2011) 033004 [[arXiv:1010.3805](#)] [[INSPIRE](#)].
- [101] H. Ishimori and M. Tanimoto, Slepton mass matrices, $\mu \rightarrow e\gamma$ decay and EDM in SUSY S_4 flavor model, *Prog. Theor. Phys.* **125** (2011) 653 [[arXiv:1012.2232](#)] [[INSPIRE](#)].
- [102] R.-Z. Yang and H. Zhang, Minimal seesaw model with S_4 flavor symmetry, *Phys. Lett. B* **700** (2011) 316 [[arXiv:1104.0380](#)] [[INSPIRE](#)].
- [103] S. Morisi and E. Peinado, Admixture of quasi-Dirac and Majorana neutrinos with tri-bimaximal mixing, *Phys. Lett. B* **701** (2011) 451 [[arXiv:1104.4961](#)] [[INSPIRE](#)].
- [104] S. Morisi, K.M. Patel and E. Peinado, Model for $T2K$ indication with maximal atmospheric angle and tri-maximal solar angle, *Phys. Rev. D* **84** (2011) 053002 [[arXiv:1107.0696](#)] [[INSPIRE](#)].
- [105] L. Dorame, S. Morisi, E. Peinado, J. Valle and A.D. Rojas, A new neutrino mass sum rule from inverse seesaw, *Phys. Rev. D* **86** (2012) 056001 [[arXiv:1203.0155](#)] [[INSPIRE](#)].
- [106] D. Hernandez and A.Y. Smirnov, Lepton mixing and discrete symmetries, *Phys. Rev. D* **86** (2012) 053014 [[arXiv:1204.0445](#)] [[INSPIRE](#)].
- [107] Z.-h. Zhao, Understanding for flavor physics in the lepton sector, *Phys. Rev. D* **86** (2012) 096010 [[arXiv:1207.2545](#)] [[INSPIRE](#)].
- [108] R. Krishnan, P. Harrison and W. Scott, Simplest neutrino mixing from S_4 symmetry, *JHEP* **04** (2013) 087 [[arXiv:1211.2000](#)] [[INSPIRE](#)].
- [109] R. Krishnan, A model for large θ_{13} constructed using the eigenvectors of the S_4 rotation matrices, *J. Phys. Conf. Ser.* **447** (2013) 012043 [[arXiv:1211.3364](#)] [[INSPIRE](#)].
- [110] I. de Medeiros Varzielas and L. Lavoura, Flavour models for TM_1 lepton mixing, *J. Phys. G* **40** (2013) 085002 [[arXiv:1212.3247](#)] [[INSPIRE](#)].
- [111] W. Grimus, Discrete symmetries, roots of unity and lepton mixing, *J. Phys. G* **40** (2013) 075008 [[arXiv:1301.0495](#)] [[INSPIRE](#)].
- [112] S.F. King and C. Luhn, Neutrino mass and mixing with discrete symmetry, *Rept. Prog. Phys.* **76** (2013) 056201 [[arXiv:1301.1340](#)] [[INSPIRE](#)].
- [113] R. González Felipe, H. Serôdio and J.P. Silva, Models with three Higgs doublets in the triplet representations of A_4 or S_4 , *Phys. Rev. D* **87** (2013) 055010 [[arXiv:1302.0861](#)] [[INSPIRE](#)].

- [114] Y. Daikoku and H. Okada, *Phenomenology of S_4 flavor symmetric extra U(1) model*, *Phys. Rev. D* **88** (2013) 015034 [[arXiv:1303.7056](#)] [[INSPIRE](#)].
- [115] F. Feruglio, C. Hagedorn and R. Ziegler, *A realistic pattern of lepton mixing and masses from S_4 and CP*, *Eur. Phys. J. C* **74** (2014) 2753 [[arXiv:1303.7178](#)] [[INSPIRE](#)].
- [116] C. Luhn, *Trimaximal TM_1 neutrino mixing in S_4 with spontaneous CP-violation*, *Nucl. Phys. B* **875** (2013) 80 [[arXiv:1306.2358](#)] [[INSPIRE](#)].
- [117] P. Frampton and T. Kephart, *Simple non-Abelian finite flavor groups and fermion masses*, *Int. J. Mod. Phys. A* **10** (1995) 4689 [[hep-ph/9409330](#)] [[INSPIRE](#)].
- [118] P.H. Frampton and T.W. Kephart, *Classification of conformality models based on non-Abelian orbifolds*, *Phys. Rev. D* **64** (2001) 086007 [[hep-th/0011186](#)] [[INSPIRE](#)].
- [119] W. Grimus and L. Lavoura, *A discrete symmetry group for maximal atmospheric neutrino mixing*, *Phys. Lett. B* **572** (2003) 189 [[hep-ph/0305046](#)] [[INSPIRE](#)].
- [120] W. Grimus and L. Lavoura, *Models of maximal atmospheric neutrino mixing and leptogenesis*, [hep-ph/0405261](#) [[INSPIRE](#)].
- [121] W. Grimus, A.S. Joshipura, S. Kaneko, L. Lavoura and M. Tanimoto, *Lepton mixing angle $\theta_{13} = 0$ with a horizontal symmetry D_4* , *JHEP* **07** (2004) 078 [[hep-ph/0407112](#)] [[INSPIRE](#)].
- [122] M. Frigerio, S. Kaneko, E. Ma and M. Tanimoto, *Quaternion family symmetry of quarks and leptons*, *Phys. Rev. D* **71** (2005) 011901 [[hep-ph/0409187](#)] [[INSPIRE](#)].
- [123] K.S. Babu and J. Kubo, *Dihedral families of quarks, leptons and Higgses*, *Phys. Rev. D* **71** (2005) 056006 [[hep-ph/0411226](#)] [[INSPIRE](#)].
- [124] M. Honda, R. Takahashi and M. Tanimoto, *Embedding the texture of the neutrino mass matrix into the MaVaNs scenario*, *JHEP* **01** (2006) 042 [[hep-ph/0510018](#)] [[INSPIRE](#)].
- [125] H. Ishimori et al., *$D(4)$ flavor symmetry for neutrino masses and mixing*, *Phys. Lett. B* **662** (2008) 178 [[arXiv:0802.2310](#)] [[INSPIRE](#)].
- [126] H. Abe, K.-S. Choi, T. Kobayashi and H. Ohki, *Non-abelian discrete flavor symmetries from magnetized/intersecting brane models*, *Nucl. Phys. B* **820** (2009) 317 [[arXiv:0904.2631](#)] [[INSPIRE](#)].
- [127] T. Araki, T. Kobayashi, J. Kubo, S. Ramos-Sanchez, M. Ratz et al., *(Non-)abelian discrete anomalies*, *Nucl. Phys. B* **805** (2008) 124 [[arXiv:0805.0207](#)] [[INSPIRE](#)].
- [128] A. Adulpravitchai, A. Blum and C. Hagedorn, *A supersymmetric D_4 model for μ - τ symmetry*, *JHEP* **03** (2009) 046 [[arXiv:0812.3799](#)] [[INSPIRE](#)].
- [129] C. Hagedorn, M. Lindner and F. Plentinger, *The discrete flavor symmetry D_5* , *Phys. Rev. D* **74** (2006) 025007 [[hep-ph/0604265](#)] [[INSPIRE](#)].
- [130] T. Kobayashi, H.P. Nilles, F. Plöger, S. Raby and M. Ratz, *Stringy origin of non-Abelian discrete flavor symmetries*, *Nucl. Phys. B* **768** (2007) 135 [[hep-ph/0611020](#)] [[INSPIRE](#)].
- [131] F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, *Tri-bimaximal neutrino mixing and quark masses from a discrete flavour symmetry*, *Nucl. Phys. B* **775** (2007) 120 [Erratum *ibid.* **836** (2010) 127-128] [[hep-ph/0702194](#)] [[INSPIRE](#)].
- [132] M.-C. Chen and K. Mahanthappa, *CKM and tri-bimaximal MNS matrices in a $SU(5) \times {}^{(d)}T$ model*, *Phys. Lett. B* **652** (2007) 34 [[arXiv:0705.0714](#)] [[INSPIRE](#)].

- [133] P.H. Frampton and T.W. Kephart, *Flavor symmetry for quarks and leptons*, *JHEP* **09** (2007) 110 [[arXiv:0706.1186](#)] [[INSPIRE](#)].
- [134] G.-J. Ding, *Fermion mass hierarchies and flavor mixing from T' symmetry*, *Phys. Rev. D* **78** (2008) 036011 [[arXiv:0803.2278](#)] [[INSPIRE](#)].
- [135] P.H. Frampton and S. Matsuzaki, *T' predictions of PMNS and CKM angles*, *Phys. Lett. B* **679** (2009) 347 [[arXiv:0902.1140](#)] [[INSPIRE](#)].
- [136] D.A. Eby, P.H. Frampton and S. Matsuzaki, *Predictions of neutrino mixing angles in a T' model*, *Phys. Lett. B* **671** (2009) 386 [[arXiv:0810.4899](#)] [[INSPIRE](#)].
- [137] C.M. Ho and T.W. Kephart, *Electron and muon $g - 2$ contributions from the T' Higgs sector*, *Phys. Lett. B* **687** (2010) 201 [[arXiv:1001.3696](#)] [[INSPIRE](#)].
- [138] P.H. Frampton, C.M. Ho, T.W. Kephart and S. Matsuzaki, *LHC Higgs production and decay in the T' model*, *Phys. Rev. D* **82** (2010) 113007 [[arXiv:1009.0307](#)] [[INSPIRE](#)].
- [139] D.A. Eby, P.H. Frampton, X.-G. He and T.W. Kephart, *Quartification with T' flavor*, *Phys. Rev. D* **84** (2011) 037302 [[arXiv:1103.5737](#)] [[INSPIRE](#)].
- [140] P.H. Frampton, C.M. Ho and T.W. Kephart, *“Heterotic” discrete flavor model*, *Phys. Rev. D* **89** (2014) 027701 [[arXiv:1305.4402](#)] [[INSPIRE](#)].
- [141] C. Luhn, S. Nasri and P. Ramond, *Tri-bimaximal neutrino mixing and the family symmetry semidirect product of Z_7 and Z_3* , *Phys. Lett. B* **652** (2007) 27 [[arXiv:0706.2341](#)] [[INSPIRE](#)].
- [142] C. Hagedorn, M.A. Schmidt and A.Y. Smirnov, *Lepton mixing and cancellation of the Dirac mass hierarchy in SO(10) GUTs with flavor symmetries T^7 and Σ_{81}* , *Phys. Rev. D* **79** (2009) 036002 [[arXiv:0811.2955](#)] [[INSPIRE](#)].
- [143] Q.-H. Cao, S. Khalil, E. Ma and H. Okada, *Observable T_7 lepton flavor symmetry at the Large Hadron Collider*, *Phys. Rev. Lett.* **106** (2011) 131801 [[arXiv:1009.5415](#)] [[INSPIRE](#)].
- [144] Q.-H. Cao, S. Khalil, E. Ma and H. Okada, *Nonzero θ_{13} for neutrino mixing in a supersymmetric B-L gauge model with T_7 lepton flavor symmetry*, *Phys. Rev. D* **84** (2011) 071302 [[arXiv:1108.0570](#)] [[INSPIRE](#)].
- [145] H. Ishimori, S. Khalil and E. Ma, *CP phases of neutrino mixing in a supersymmetric B-L gauge model with T_7 lepton flavor symmetry*, *Phys. Rev. D* **86** (2012) 013008 [[arXiv:1204.2705](#)] [[INSPIRE](#)].
- [146] A.J. Buras, F. De Fazio and J. Girrbach, *331 models facing new $b \rightarrow s\mu^+\mu^-$ data*, *JHEP* **02** (2014) 112 [[arXiv:1311.6729](#)] [[INSPIRE](#)].
- [147] F. De Fazio, *Quark flavour observables in 331 models in the flavour precision era*, [arXiv:1310.4614](#) [[INSPIRE](#)].
- [148] J. Valle and M. Singer, *Lepton number violation with quasi Dirac neutrinos*, *Phys. Rev. D* **28** (1983) 540 [[INSPIRE](#)].
- [149] F. Pisano and V. Pleitez, *An $SU(3) \times U(1)$ model for electroweak interactions*, *Phys. Rev. D* **46** (1992) 410 [[hep-ph/9206242](#)] [[INSPIRE](#)].
- [150] P. Frampton, *Chiral dilepton model and the flavor question*, *Phys. Rev. Lett.* **69** (1992) 2889 [[INSPIRE](#)].
- [151] R. Foot, O.F. Hernandez, F. Pisano and V. Pleitez, *Lepton masses in an $SU(3)_L \times U(1)_N$ gauge model*, *Phys. Rev. D* **47** (1993) 4158 [[hep-ph/9207264](#)] [[INSPIRE](#)].

- [152] J. Montero, F. Pisano and V. Pleitez, *Neutral currents and GIM mechanism in $SU(3)_L \times U(1)_N$ models for electroweak interactions*, *Phys. Rev. D* **47** (1993) 2918 [[hep-ph/9212271](#)] [[INSPIRE](#)].
- [153] M. Singer, J. Valle and J. Schechter, *Canonical neutral current predictions from the weak electromagnetic gauge group $SU(3) \times U(1)$* , *Phys. Rev. D* **22** (1980) 738 [[INSPIRE](#)].
- [154] R. Foot, H.N. Long and T.A. Tran, *$SU(3)_L \times U(1)_N$ and $SU(4)_L \times U(1)_N$ gauge models with right-handed neutrinos*, *Phys. Rev. D* **50** (1994) R34 [[hep-ph/9402243](#)] [[INSPIRE](#)].
- [155] H.N. Long, *$SU(3)_L \times U(1)_N$ model for right-handed neutrino neutral currents*, *Phys. Rev. D* **54** (1996) 4691 [[hep-ph/9607439](#)] [[INSPIRE](#)].
- [156] H.N. Long, *The 331 model with right handed neutrinos*, *Phys. Rev. D* **53** (1996) 437 [[hep-ph/9504274](#)] [[INSPIRE](#)].
- [157] H.N. Long, *Scalar sector of the 331 model with three Higgs triplets*, *Mod. Phys. Lett. A* **13** (1998) 1865 [[hep-ph/9711204](#)] [[INSPIRE](#)].
- [158] W.A. Ponce, Y. Giraldo and L.A. Sanchez, *Minimal scalar sector of 3-3-1 models without exotic electric charges*, *Phys. Rev. D* **67** (2003) 075001 [[hep-ph/0210026](#)] [[INSPIRE](#)].
- [159] P. Dong, H.N. Long, D. Nhung and D. Soa, *$SU(3)_C \times SU(3)_L \times U(1)_X$ model with two Higgs triplets*, *Phys. Rev. D* **73** (2006) 035004 [[hep-ph/0601046](#)] [[INSPIRE](#)].
- [160] P. Dong, D. Huong, T. Huong and H.N. Long, *Fermion masses in the economical 3-3-1 model*, *Phys. Rev. D* **74** (2006) 053003 [[hep-ph/0607291](#)] [[INSPIRE](#)].
- [161] P. Dong, H.N. Long and D. Soa, *Neutrino masses in the economical 3-3-1 model*, *Phys. Rev. D* **75** (2007) 073006 [[hep-ph/0610381](#)] [[INSPIRE](#)].
- [162] P. Dong, H. Long and H. Hung, *Question of Peccei-Quinn symmetry and quark masses in the economical 3-3-1 model*, *Phys. Rev. D* **86** (2012) 033002 [[arXiv:1205.5648](#)] [[INSPIRE](#)].
- [163] P.V. Dong and H.N. Long, *The economical $SU(3)_C \times SU(3)_L \times U(1)_X$ model*, *Adv. High Energy Phys.* **2008** (2008) 739492 [[arXiv:0804.3239](#)] [[INSPIRE](#)].
- [164] P.V. Dong, L.T. Hue, H. Long and D.V. Soa, *The 3-3-1 model with A_4 flavor symmetry*, *Phys. Rev. D* **81** (2010) 053004 [[arXiv:1001.4625](#)] [[INSPIRE](#)].
- [165] P.V. Dong, H.N. Long, D.V. Soa and V.V. Vien, *The 3-3-1 model with S_4 flavor symmetry*, *Eur. Phys. J. C* **71** (2011) 1544 [[arXiv:1009.2328](#)] [[INSPIRE](#)].
- [166] P.V. Dong, H.N. Long, C.H. Nam and V.V. Vien, *The S_3 flavor symmetry in 3-3-1 models*, *Phys. Rev. D* **85** (2012) 053001 [[arXiv:1111.6360](#)] [[INSPIRE](#)].
- [167] V.V. Vien and H.N. Long, *The D_4 flavor symmetry in 3-3-1 model with neutral leptons*, *Int. J. Mod. Phys. A* **28** (2013) 1350159 [[arXiv:1312.5034](#)] [[INSPIRE](#)].
- [168] Y.H. Ahn, S.K. Kang and C.S. Kim, *Spontaneous CP-violation in A_4 flavor symmetry and leptogenesis*, *Phys. Rev. D* **87** (2013) 113012 [[arXiv:1304.0921](#)] [[INSPIRE](#)].
- [169] D. Chang and H.N. Long, *Interesting radiative patterns of neutrino mass in an $SU(3)_C \times SU(3)_L \times U(1)_X$ model with right-handed neutrinos*, *Phys. Rev. D* **73** (2006) 053006 [[hep-ph/0603098](#)] [[INSPIRE](#)].
- [170] C.H. Albright and W. Rodejohann, *Comparing trimaximal mixing and its variants with deviations from tri-bimaximal mixing*, *Eur. Phys. J. C* **62** (2009) 599 [[arXiv:0812.0436](#)] [[INSPIRE](#)].

- [171] D. Marzocca, S. Petcov, A. Romanino and M. Sevilla, *Nonzero $|U_{e3}|$ from charged lepton corrections and the atmospheric neutrino mixing angle*, *JHEP* **05** (2013) 073 [[arXiv:1302.0423](#)] [[INSPIRE](#)].
- [172] X. Zhang and B.-Q. Ma, *A prediction of neutrino mixing matrix with CP-violating phase*, *Phys. Lett. B* **713** (2012) 202 [[arXiv:1203.2906](#)] [[INSPIRE](#)].
- [173] ICECUBE collaboration, J. Ahrens et al., *IceCube — The next generation neutrino telescope at the south pole*, *Nucl. Phys. Proc. Suppl.* **118** (2003) 388 [[astro-ph/0209556](#)] [[INSPIRE](#)].
- [174] V. Lobashev et al., *Direct search for neutrino mass and anomaly in the tritium β -spectrum: status of ‘Troitsk neutrino mass’ experiment*, *Nucl. Phys. Proc. Suppl.* **91** (2001) 280.
- [175] SDSS collaboration, M. Tegmark et al., *Cosmological parameters from SDSS and WMAP*, *Phys. Rev. D* **69** (2004) 103501 [[astro-ph/0310723](#)] [[INSPIRE](#)].
- [176] T.J. Weiler, *Oscillation and mixing among the three neutrino flavors*, [arXiv:1308.1715](#) [[INSPIRE](#)].
- [177] W. Rodejohann, *Neutrino-less double beta decay and particle physics*, *Int. J. Mod. Phys. E* **20** (2011) 1833 [[arXiv:1106.1334](#)] [[INSPIRE](#)].
- [178] M. Mitra, G. Senjanović and F. Vissani, *Neutrinoless double β decay and heavy sterile neutrinos*, *Nucl. Phys. B* **856** (2012) 26 [[arXiv:1108.0004](#)] [[INSPIRE](#)].
- [179] S. Bilenky and C. Giunti, *Neutrinoless double- β decay: a brief review*, *Mod. Phys. Lett. A* **27** (2012) 1230015 [[arXiv:1203.5250](#)] [[INSPIRE](#)].
- [180] W. Rodejohann, *Neutrinoless double β decay and neutrino physics*, *J. Phys. G* **39** (2012) 124008 [[arXiv:1206.2560](#)] [[INSPIRE](#)].
- [181] A. Merle, *keV neutrino model building*, *Int. J. Mod. Phys. D* **22** (2013) 1330020 [[arXiv:1302.2625](#)] [[INSPIRE](#)].
- [182] J. Vergados, H. Ejiri and F. Simkovic, *Theory of neutrinoless double β decay*, *Rept. Prog. Phys.* **75** (2012) 106301 [[arXiv:1205.0649](#)] [[INSPIRE](#)].
- [183] C. Kraus et al., *Final results from phase II of the Mainz Neutrino Mass Search in tritium β decay*, *Eur. Phys. J. C* **40** (2005) 447 [[hep-ex/0412056](#)] [[INSPIRE](#)].
- [184] TROITSK collaboration, V. Aseev et al., *An upper limit on electron antineutrino mass from Troitsk experiment*, *Phys. Rev. D* **84** (2011) 112003 [[arXiv:1108.5034](#)] [[INSPIRE](#)].
- [185] P.V. Dong and H.N. Long, *$U(1)_Q$ invariance and $SU(3)_C \times SU(3)_L \times U(1)_X$ models with β arbitrary*, *Eur. Phys. J. C* **42** (2005) 325 [[hep-ph/0506022](#)] [[INSPIRE](#)].
- [186] P.V. Dong, V.T.N. Huyen, H.N. Long and H.V. Thuy, *Gauge boson mixing in the 3-3-1 models with discrete symmetries*, *Adv. High Energy Phys.* **2012** (2012) 715038
- [187] A.E.C. Hernández, R. Martínez and J. Nisperuza, *S_3 flavour symmetry breaking scheme for understanding the quark mass and mixing pattern in $SU(3)_C \otimes SU(3)_L \otimes U(1)_X$ models*, [arXiv:1401.0937](#) [[INSPIRE](#)].
- [188] P.H. Frampton and M. Harada, *Constraints from precision electroweak data on leptoquarks and bileptons*, *Phys. Rev. D* **58** (1998) 095013 [[hep-ph/9711448](#)] [[INSPIRE](#)].
- [189] H.N. Long and T. Inami, *S , T , U parameters in $SU(3)_C \times SU(3)_L \times U(1)$ model with right-handed neutrinos*, *Phys. Rev. D* **61** (2000) 075002 [[hep-ph/9902475](#)] [[INSPIRE](#)].