



ORAL PRESENTATION

Open Access

Antibody-mediated inhibition of HIV-1 elicited by HIV-I DNA priming and boosting with heterologous HIV-1 recombinant MVA in healthy Tanzanian adults

A Joachim^{1*}, C Nilsson², S Aboud³, EF Lyamuya³, M Robb⁴, M Marovich⁵, C Ochsenauber⁶, B Wahren², E Sandström⁷, G Biberfeld², G Ferrari⁸, VR Polonis⁵

From AIDS Vaccine 2012
Boston, MA, USA. 9-12 September 2012

Background

We evaluated HIV antibody (Ab) responses elicited by immunization, in a phase I/II placebo-controlled double blind trial using multiclade, multigene HIV-1-DNA prime boosted with HIV-MVA conducted among healthy volunteers in Tanzania (HIVIS03).

Methods

Sixty HIV-uninfected volunteers, randomized into groups of 20 received placebo or 1 mg HIV-DNA intradermally (id) or 3.8 mg intramuscularly (im). DNA plasmids containing HIV-1 gp160 subtypes A, B, C; rev B; p17/p24 gag A, B and RTmut B were given at months 0, 1 and 3 using a needle-free Biojector device. HIV-MVA expressing CRF01_AE HIV-1 env, gag, pol was administered im by needle at months 9 and 21. Sera were tested at baseline, two months post-first and four weeks post-second HIV-MVA boosting. HIV Ab responses were tested using pseudoviruses and TZM-bl cells as well as luciferase-expressing infectious molecular clones (IMC-LucR) in PBMC-based assays. ADCC responses were tested using the flow cytometry GranToxiLux-based assay.

Results

Neutralizing Ab activity was demonstrated only in the PBMC assay, and after the second MVA boost in 24 (83%) of 29 vaccinees against the clade CRF01_AE CM235 IMC and in 21 (72%) of 29 vaccinees against

clade B SF162-IMC. NK cell depletion from PBMC targets resulted in a significant loss of HIV inhibition by vaccinee sera, indicating a role of Ab-mediated Fcγ-receptor function. Vaccine-induced ADCC responses were detected in 21 (75%) of 28 vaccinees after the second HIV-MVA boost. ADCC Ab titers did not differ significantly between id- (median 840, range 300-5400) and im-primed (median 880, range 400-3600) vaccinees (p=0.45).

Conclusion

HIV-DNA priming followed by two HIV-MVA boosts elicited HIV-specific inhibitory and/or ADCC-mediating antibody responses in a high proportion of Tanzanian adults.

Author details

¹Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania, United Republic of. ²Karolinska Institutet / Institute for Infectious Disease Control (SMI), Stockholm, Sweden. ³Muhimbili University of Health and Allied Sciences (MUHAS), Dar es Salaam, Tanzania, United Republic of. ⁴The Henry M. Jackson Foundation, Rockville, MD, USA. ⁵Walter Reed Army Institute of Research (WRAIR), Rockville, MD, USA. ⁶Department of Medicine, University of Alabama at Birmingham, Birmingham, AL, USA. ⁷Venhälsan, Karolinska Institutet (KI) at Södersjukhuset, Stockholm, Sweden. ⁸Department of Surgery, Duke University Medical Center, Durham, NC, USA.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-O53

Cite this article as: Joachim et al.: Antibody-mediated inhibition of HIV-1 elicited by HIV-I DNA priming and boosting with heterologous HIV-1 recombinant MVA in healthy Tanzanian adults. *Retrovirology* 2012 **9** (Suppl 2):O53.

¹Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania, United Republic of

Full list of author information is available at the end of the article