

## RESEARCH ARTICLE

## Open Access

# Genetic map of *Triticum turgidum* based on a hexaploid wheat population without genetic recombination for D genome

Li Zhang<sup>1†</sup>, Jiang-Tao Luo<sup>1†</sup>, Ming Hao<sup>1†</sup>, Lian-Quan Zhang<sup>1</sup>, Zhong-Wei Yuan<sup>1</sup>, Ze-Hong Yan<sup>1</sup>, Ya-Xi Liu<sup>1</sup>, Bo Zhang<sup>2</sup>, Bao-Long Liu<sup>2</sup>, Chun-Ji Liu<sup>3</sup>, Huai-Gang Zhang<sup>2</sup>, You-Liang Zheng<sup>1</sup> and Deng-Cai Liu<sup>1,2\*</sup>

## Abstract

**Background:** A synthetic doubled-haploid hexaploid wheat population, SynDH1, derived from the spontaneous chromosome doubling of triploid F<sub>1</sub> hybrid plants obtained from the cross of hybrids *Triticum turgidum* ssp. *durum* line Langdon (LDN) and ssp. *turgidum* line AS313, with *Aegilops tauschii* ssp. *tauschii* accession AS60, was previously constructed. SynDH1 is a tetraploidization-hexaploid doubled haploid (DH) population because it contains recombinant A and B chromosomes from two different *T. turgidum* genotypes, while all the D chromosomes from *Ae. tauschii* are homogenous across the whole population. This paper reports the construction of a genetic map using this population.

**Results:** Of the 606 markers used to assemble the genetic map, 588 (97%) were assigned to linkage groups. These included 513 Diversity Arrays Technology (DART) markers, 72 simple sequence repeat (SSR), one insertion site-based polymorphism (ISBP), and two high-molecular-weight glutenin subunit (HMW-GS) markers. These markers were assigned to the 14 chromosomes, covering 2048.79 cM, with a mean distance of 3.48 cM between adjacent markers. This map showed good coverage of the A and B genome chromosomes, apart from 3A, 5A, 6A, and 4B. Compared with previously reported maps, most shared markers showed highly consistent orders. This map was successfully used to identify five quantitative trait loci (QTL), including two for spikelet number on chromosomes 7A and 5B, two for spike length on 7A and 3B, and one for 1000-grain weight on 4B. However, differences in crossability QTL between the two *T. turgidum* parents may explain the segregation distortion regions on chromosomes 1A, 3B, and 6B.

**Conclusions:** A genetic map of *T. turgidum* including 588 markers was constructed using a synthetic doubled haploid (SynDH) hexaploid wheat population. Five QTLs for three agronomic traits were identified from this population. However, more markers are needed to increase the density and resolution of this map in the future study.

**Keywords:** Allopolyploid, Crossability, Doubled haploid, Segregation distortion

\* Correspondence: [dcliu7@yahoo.com](mailto:dcliu7@yahoo.com)

†Equal contributors

<sup>1</sup>Triticaceae Research Institute, Sichuan Agricultural University, Wenjiang, Chengdu, Sichuan 611130, P.R. China

<sup>2</sup>Key Laboratory of Adaptation and Evolution of Plateau Biota, Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining 810001, P.R. China

Full list of author information is available at the end of the article

## Background

Many important crops such as bread wheat, durum wheat, cotton, oat, coffee, and tobacco are allopolyploids. They originated from the merger of two or more distinct but related genomes, by interspecific hybridization and then genome doubling. Because the majority of accessions of ancestral species were not involved in speciation, many of their unique genes may not present in crops [1,2]. Thus unraveling the genes coding for agronomically-important traits in ancestral species is important for crop improvement. The differences in polyploid levels between a crop and its ancestral species may cause differences in gene expression, especially for a quantitative trait locus (QTL). Therefore, genetic populations with the same ploidy level as a given crop are needed to analyze ancestral genes for crop improvement; however their construction is labor-intensive and time-consuming.

We recently developed a method for synthesizing doubled haploid (SynDH) populations specific for allopolyploid species [3]. *Triticum turgidum* L. ( $2n = 4x = 28$ , AABB) and *Aegilops tauschii* Coss. ( $2n = 2x = 14$ , DD), the two ancestral species of common wheat (*Triticum aestivum* L.,  $2n = 6x = 42$ , AABBDD), were used to demonstrate the method. Unlike existing methods for producing DH populations, this method eliminates the need for *in vitro* culture for extracting haploids and/or chemical treatment for chromosome doubling. Moreover, genetic recombination at the hexaploid level can be restricted to the D genome (thus called diploidization-hexaploid SynDH) or A and B genomes (tetraploidization-hexaploid SynDH), making genetic analysis simpler [3]. However, the effects of applying such a strategy to the construction of genetic maps are not clear.

The objectives of this study were to (i) develop a genetic map of the A and B genomes based on DArT (Diversity Arrays Technology), SSR (simple sequence repeat), ISBP (insertion site-based polymorphism), and high-molecular-weight glutenin subunit (HMW-GS) markers using a tetraploidization-hexaploid SynDH population, (ii) assess the extent of segregation distortion, (iii) evaluate the quality of linkage maps constructed using this strategy by comparing with previously reported maps, and (iv) identify QTLs in this population.

## Results

### Construction of genetic map and QTL analysis

Although this SynDH population was derived from the cross of three parents (LDN/AS313//AS60), all of the D chromosomes came from the single genotype AS60. Thus, segregation of this population was restricted to the A or B chromosomes, which came from the two different tetraploid parents, *T. turgidum* AS313 and LDN. Therefore, polymorphic markers between the two *T.*

*turgidum* parents can be used in genetic mapping. A total of 606 polymorphic markers between *T. turgidum* AS313 and LDN were identified, including 521 DArT, 81 SSR, two ISBP, and two HMW-GS markers. These markers were used to genotype the SynDH population.

Of the 606 markers, 588 (97%) were included in the genetic map. They included 513 DArT (98%), 72 SSR (89%), one ISBP, and two HMW-GS markers (Table 1). These markers were assigned to 24 linkage groups on 14 chromosomes, giving a total map length of 2048.79 cM (Table 1). This map showed good coverage of the A and B genome chromosomes, apart from 3A, 5A, 6A, and 4B (Figure 1, 2). Map length of the A genome (796.18 cM) was shorter than that of the B genome (1252.61 cM). Chromosomes 1A, 3A, 4A, 1B, 2B, 5B, 6B, and 7B were each represented by a single linkage group, while 2A, 5A, 6A, 7A, 3B, and 4B were each represented by two or more linkage groups (Figures 1, 2). The average length of these linkage groups was 146.34 cM, with a maximum of 263.48 cM, for 2B, and a minimum of 32.90 cM, for 4B. The average distance between markers on these linkage groups ranged from 1.12 cM for 6A to 5.95 cM for 7B. The overall average was 3.48 cM between any two markers across the linkage map (Table 1).

A total of five QTLs were identified in this population. Two QTLs, located on chromosomes 7A and 5B, were for spikelet number (Table 2). These two QTLs explained 19.2% and 10.2% of the phenotypic variance, respectively. Two QTLs, on 7A and 3B, for spike length explained 19.3% and 15.1% of the phenotypic variance, respectively. The final QTL, on 4B, for 1000-grain weight explained 14.6% of the variance.

### Segregation distortion


Of the genotypic data collected in this study, 49.8% of the alleles in the SynDH population were derived from AS313, 46.1% were from LDN, and the remaining 4.1% was missing data. The overall allele ratio between those from AS313 and LDN was not significantly different at a significance level of 5%. However, significant deviation was found for 70 (11.90%) out of the 588 markers. Of these, 28 (4.7%) were in favor of LDN and 42 (7.1%) were in favor of AS313 (Table 1; Figure 1), consisting of seven (9.72%) SSR and 63 (12.9%) DArT markers. These distorted markers were distributed on 11 of the 14 chromosomes (Figure 1, 2; Table 1). Among them, 13 were on chromosome 1A, 16 were on 3B, and 28 were on 6B. All of the distorted markers on 3B favored LDN alleles, while all of the 28 distorted markers on 6B favored AS313 alleles. Of the 13 distorted markers on 1A, ten favored AS313 alleles and the other three (*wPt-6654*, *wPt-5577*, and *wPt-8245*) favored LDN alleles (Figure 1).

**Table 1 Marker distribution on A- and B-genome chromosomes**

Chromosome	Length (cM)	Number of target markers						Average length cM/marker	Mean density**	
		SSR	ISBP	DARt			Glutelin			Total
				wPt	tPt	rPt				
1A	183.61	5	-	48	4	0	<i>Glu-A1</i>	58 (13)*	3.17	7.06
2A	141.23	4	-	28	2	0	-	34 (0)	4.15	7.06
3A	39.21	0	-	10	0	0	-	10 (3)	3.92	9.80
4A	181.61	4	-	40	0	3	-	47 (1)	3.86	5.86
5A	56.01	0	-	11	2	0	-	13 (0)	4.31	7.00
6A	74.17	4	-	57	4	1	-	66 (0)	1.12	3.53
7A	120.34	4	-	20	0	0	-	24 (1)	5.01	6.69
A genome	796.18	21	-	214	12	4	1	252 (18)	3.16	5.94
1B	187.43	8	-	42	1	0	<i>Glu-B1</i>	52 (2)	3.60	6.25
2B	263.48	4	-	54	2	2	-	62 (3)	4.25	7.53
3B	241.31	16	1	71	3	1	-	92 (16)	2.62	4.64
4B	32.9	0	-	9	0	0	-	9 (1)	3.66	8.23
5B	185.53	10	-	26	1	1	-	38 (1)	4.88	7.42
6B	181.18	9	-	43	3	1	-	56 (28)	3.24	6.47
7B	160.78	4	-	22	0	1	-	27 (1)	5.95	8.46
B genome	1252.61	52	1	267	10	6	1	336 (52)	3.73	6.29
Total	2048.79	72	1	481	22	10	2	588 (70)	3.48	6.13


\*Number of markers with segregation distortion is shown in parentheses.

\*\*the mean density equal to  $L/(n-1)$  where  $n$  is the number of unique markers per chromosome length  $L$ .



**Figure 1 Genetic linkage map of A genomes obtained using 113 doubled-haploid lines from the AS313/LDN//AS60 population.**

Numbers in parentheses indicate the number of markers that are not present at each locus. Segregation distortion of markers are indicated by asterisks at significance levels of 0.05 (\*), 0.01 (\*\*), 0.005 (\*\*\*), 0.001 (\*\*\*\*), 0.0005 (\*\*\*\*\*), and 0.0001 (\*\*\*\*\*). The black arrow indicates the point of separation between two different linkage groups. The scales on the left indicate distances in centiMorgans (Kosambi).



**Map comparisons**

To evaluate the quality of the genetic map developed in this study (SynDH1), locations of the shared markers were compared with those in the CIMMYT integrated map (CIMMYT) [4], durum wheat integrated map (C-L) [5], and triticale genetic map (S-M) [6]. Discrepant marker orders were observed in some regions, mostly on chromosomes 1A and 2A. The order of the majority of shared markers was highly consistent among these four linkage maps (see Additional file 1). The comparison of chromosome 3B with the four linkage maps is shown in Figure 3. Furthermore, the genetic map of chromosome

3B from this study was also highly consistent with its physical map (Figure 3) [4-7].

**Discussion**

The SynDH1 population used in this study was derived from the interspecific hybridization of *T. turgidum* with *Ae. tauschii*, followed by spontaneous chromosome doubling [3]. We demonstrated in this study that such a population can be effectively used to generate genetic maps. This map has been successfully used to locate high-molecular-weight glutenin subunits of loci *Glu-A1* and *Glu-B1*, and to identify five QTLs of three


**Table 2 Putative QTLs detected in the SynDH population**

Trait	QTL	Marker interval	LOD score	R <sup>2</sup> (%) <sup>a</sup>	Estimated additive effect <sup>b</sup>
Spikelet number	<i>QSpn.scau-7A</i>	<i>wPt-4731- Xgwm130</i>	6.08	19.22	-0.50
	<i>QSpn.scau-5B</i>	<i>wPt-730009- Xwmc28</i>	3.32	10.18	0.36
Spike length	<i>QSl.scau-7A</i>	<i>wPt-4731- Xgwm130</i>	5.53	19.26	-0.49
	<i>QSl.scau-3B</i>	<i>Xgwm389- wPt-7225</i>	4.48	15.13	-0.44
1000-grain weight	<i>QTgw.scau-4B</i>	<i>wPt-7233- wPt-5559</i>	3.80	14.58	1.72

<sup>a</sup> Percentage of phenotypic variation explained by each QTL.

<sup>b</sup> indicates a positive or negative effect from the AS313 allele.





**Figure 3 Comparison of chromosome 3B maps.** The SynDH1 map (SynDH1-3B) obtained in this study was compared with the physical map (physical-3B) [7], the CIMMYT-integrated map (CIMMYT-3B) [4], the durum wheat integrated map (C-L-3B) [5], and triticale genetic map (S-M-3B) [6]. The scales on the left side indicate distances in cM (Kosambi). The black arrow indicates the point of separation between two different linkages. To reduce complexity, only markers shared between these maps are shown. Map comparison was performed using the JoinMap 4.0 program [44].

agronomic traits. However, this study identified as many as 24 linkage groups, which is significantly larger than the number of the 14 haploid chromosomes of the A and B genomes. To reduce the number of the linkage groups, additional markers are needed for a better covered, high-density map.

Low map coverage, duplicated marker loci, or segregation distortion can cause inconsistent marker order in genetic maps [8]. Chromosome rearrangements (such as small translocations, deletions, and inversions) can also result in marker inconsistency [8]. Compared to previously reported CIMMYT integrated map (CIMMYT) [4], durum wheat integrated map (C-L) [5], and triticale genetic map (S-M) [6], discrepant marker orders were observed in some regions, mostly on chromosomes 1A and 2A. According to the marker order in the genetic maps of these two chromosomes (see Additional file 1), these discrepancies appear to have been caused by chromosome inversions. In a recently reported genetic map for triticale, deletion of a fragment of chromosome 1A and translocation of 2A caused discrepancies in marker positions and order across populations [6].

Segregation distortion is a common phenomenon that can be influenced by factors affecting fertility of either

gametes or zygotes [9]. Environmental effects also affect segregation distortion and are assumed to influence gametophyte selection [10]. Compared with  $F_2$  and DH, RILs (recombinant inbred lines) are more prone to segregation distortion because of repeated selective forces [11]. Segregation distortion markers have been reported in common wheat [8,12-15] and its two ancestral species of *T. turgidum* [16-20] and *Ae. tauschii* [21].

If a biological segregation distortion locus exists, the concerned locus and those flanking region would all deviate from the expected Mendelian segregation ratio [22]. Therefore, biological segregation distortion affects a cluster of loci that form a segregation distorted region (SDR). Using the criterion of Paillard *et al.* [13], a SDR should contain at least three closely adjacent loci. Based on this criterion, SDRs were only found on chromosomes 1A, 3B, and 6B in this study (Figure 1, 2). Other distortion loci scattered along the chromosomes 3A, 4A, 7A, 1B, 2B, 4B, 5B, and 7B were likely a result of non-biological factors. The limited number of the SynDH lines used could be one of these factors.

Because all the  $F_1$  haploid hybrid plants from LDN/AS313//AS60 produced  $F_2$  doubled haploid plants [3], segregation distortion in the present study was most

likely generated in the production process of the haploid hybrids by wide hybridization between *T. turgidum* and *Ae. tauschii*. Because no *in vitro* culture was applied during this interspecific hybridization procedure [3], its involvement in the segregation distortion can be ruled out. The production of conventional haploids via *in vitro* culture may lead to segregation distortion, because the ability of *in vitro* regeneration is genotype-dependent. When crossing *T. turgidum* with *Ae. tauschii*, different alleles of crossability QTL between LDN and AS313 may affect segregation differently. It is known that crossability genes control the ability of crossing between different species, and affect the seed-setting of interspecific crosses [23,24]. Crossability can be promoted by recessive alleles and inhibited by dominant alleles. The inhibiting effect of chromosome 6B [25] and the promoting effect of 3B [26] on crossability in LDN may cause marker segregation favoring AS313 on 6B and LDN on 3B (Figure 2). Because *Xgwm626* and *Xbarc79* on chromosome 6B gave the most severe segregation distortions, and their flanking markers were all less skewed, the crossability QTL may be situated between these two markers. Likewise, a QTL affecting segregation may also exist near *wPt-6945* on 3B. The distorted segregation favoring AS313 near *wPt-3836* on 1A suggests that this genotype may have a recessive crossability QTL in this region [27]. These results suggest that the distorted segregation in the SynDH population depends on variations in crossability alleles between the two *T. turgidum* parents. In this regard, genotypic effects on crossability need to be considered when using SynDH populations for genetic mapping.

## Conclusions

A genetic map of *T. turgidum* including 588 markers was constructed using a synthetic doubled haploid (SynDH) hexaploid wheat population, derived from the interspecific hybridization of *T. turgidum* with *Ae. tauschii*, followed by spontaneous chromosome doubling. This map has been successfully used to locate high-molecular-weight glutenin subunits on loci *Glu-A1* and *Glu-B1*, and to identify five QTLs of three agronomic traits. However, more markers are needed to increase the density and resolution of this map in the future study. Moreover, the distorted segregation in the SynDH population may be caused by variations in crossability alleles between the two *T. turgidum* parents. In this regard, genotypic effects on crossability need to be considered when using SynDH populations for genetic mapping.

## Methods

### Plant material

Plant materials used in this study included 113 synthesized doubled haploid (SynDH) lines and their parents,

*T. turgidum* ssp. *turgidum* line AS313 and ssp. *durum* line Langdon, and *Ae. tauschii* ssp. *tauschii* accession AS60. The procedure of SynDH was described previously by Zhang *et al.* [3]. Briefly, the F<sub>1</sub> hybrids between AS313 and LDN were pollinated by AS60. No embryo rescue technique or hormone treatment was applied during this interspecific hybridization procedure. The haploid interspecific hybrids were self-pollinated and SynDHs were produced by spontaneous chromosome doubling with the help of unreduced gametes [3]. This SynDH population has recombinant A and B chromosomes between *T. turgidum* lines AS313 and LDN in a background of non-recombinant D chromosomes from *Ae. tauschii* AS60.

### Field trials and trait evaluation

Field trials were conducted in the crop season of 2010–2011 at the Triticeae Research Institute experimental station in Wenjiang, Sichuan Province, China. Individual plants of the 113 SynDH lines were spaced 10 cm apart within 2 m long rows, and the row spacing was 30 cm. This experiment consisted of two replicates. Five agronomic traits (plant height, tiller number per plant, spike length, spikelet number, and 1000-grain weight) were evaluated from 10 random selected plants from each plot at maturity. Plant height (HT) was calculated as the average height in centimeters measured from the soil surface to the tip of the spike (awns excluded). Spike length (SI), spikelet number (Spn), and 1000-grain weight were measured on 10 selected spikes from one line (usually the main spike per plant was chosen). Kernels from a sample of 10 spikes from each plot were threshed, counted, and weighed to calculate 1000-grain weight.

### HMW-GS markers

Two high-molecular-weight glutenin subunit (HMW-GS) loci, *Glu-A1* on 1AL and *Glu-B1* on 1BL, were used for mapping [3].

### SSR, ISBP, and DaRT genotyping

A bulk leaf sample from five plants for each of the SynDH lines and parents was used for DNA isolation using the 2×CTAB method [28]. Fifty-nine ISBP (insertion site-based polymorphism) markers [7], and 539 SSR markers including 195 *wmc* [29,30], 159 *gwm* [31], 113 *barc* [32], 49 *gpw* [33], 12 *gdm* [34], 8 *cfb* [35], 2 *cfa* [33], and 1 *psp* [36] were selected to screen the parents. PCR amplifications were performed as described by Zhang *et al.* [3]. The amplified fragments were separated by electrophoresis on 6% polyacrylamide denaturing gels and visualized using the silver-staining method [37].

Genomic DNA from SynDH lines and their parents was sent to Triticarte (Yarralumla, Australia) for profiling using Diversity Arrays Technology (DArT) [38,39]. The common wheat *PstI* (*TaqI*) v3.0 DArT array, which comprises 19,000 clones, was used. Of these clones, about 7000 are known to be polymorphic in a wide range of wheat cultivars, and about 3500 markers have been assigned to chromosomes using nulli-tetrasomic lines derived from Chinese Spring. Hybridization of genomic DNA to the DArT array, image analysis, and polymorphism scoring were carried out as previously described by Akbari *et al.* [40]. DArTs are biallelic dominant markers. Each marker is scored for each sample: 0, 1, or -, where "-" indicates that the marker could not be reliably scored for that sample. DArT calls were converted into "A" (AS313), "B" (LDN), and "-" (missing data) by comparison against parental scores.

### Map construction

Segregation data were analyzed with QTL IciMapping v3.1 [41], and markers were grouped using a logarithm of odds (LOD) >5. The markers within each group were then ordered with RECORD [42] and rippled with SARF (sum of adjacent recombination frequencies). The marker order was used to sort markers within linkage groups and graphical genotypes were examined in Excel 2003 (see Additional file 2). At this step, singletons (a single locus in one progeny line that appears to have recombined with both of its directly-neighboring loci) were replaced by missing values in the dataset and calculations were repeated until no singletons were found. Ungrouped (not corrected data) markers were anchored to previous linkages using a LOD >4.5. All calculations were repeated for new linkage groups. Independent linkages on the same chromosomes with Kosambi distances [43] between subsequent markers below 50 cM were integrated as one linkage group. The  $\chi^2$  analysis, map drawing, and map comparison were performed using the JoinMap 4.0 program [44]. A  $\chi^2$  goodness of fit analysis for each marker was performed to test for deviation from the 1:1 expected segregation ratio in doubled haploids at a 5% level of significance.

### QTL analysis

QTL IciMapping v3.1, based on the inclusive composite interval mapping (ICIM) model [41], was used for QTL analysis. Threshold values were calculated using 1,000 permutations and QTLs were considered real when ICIM showed the presence of a significant peak at a level of 0.05. Estimates of the positions of QTL corresponded to the peak of the ICIM scans. The percentage of phenotypic variation explained by each QTL was calculated with a single factor regression ( $R^2$ ).

## Additional files

**Additional file 1: Map comparison of A and B genome chromosomes between SynDH1 (this study) and the CIMMYT-integrated map (CIMMYT) [4], the durum wheat integrated map (C-L) [5], and the triticale genetic map (S-M) [6].** The scales on the left indicate distances in cM (Kosambi). To reduce complexity, only markers shared between these maps are shown. Map comparison was performed using the JoinMap 4.0 program [44].

**Additional file 2: Marker information used for linkage group construction.**

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

LZ, JTL, and MH performed most of the research and drafted the manuscript. LQZ, ZWY, ZHY, BZ, BLL, and DCL assisted in the collection of genotypic and phenotypic data. YXL assisted in DArT analysis. CJL revised the manuscript. HGZ, YLZ, and DCL supervised and coordinated the study. All authors read and approved the final manuscript.

### Acknowledgments

This research was supported by the National Natural Science Foundation of China (31071420, 31101140), the Science and Technology Commission of Sichuan Province (2011JQ0016), and by the 100-Talent Program of CAS.

### Author details

<sup>1</sup>Triticaceae Research Institute, Sichuan Agricultural University, Wenjiang, Chengdu, Sichuan 611130, P.R. China. <sup>2</sup>Key Laboratory of Adaptation and Evolution of Plateau Biota, Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining 810001, P.R. China. <sup>3</sup>CSIRO Plant Industry, 306 Carmody Road, St Lucia, QLD 4067, Australia.

Received: 20 February 2012 Accepted: 8 August 2012

Published: 13 August 2012

### References

1. Reif JC, Zhang P, Dreisigacker S, Warburton ML, van Ginkel M, Hoisington D, Bohn M, Melchinger AE: **Wheat genetic diversity trends during domestication and breeding.** *Theor Appl Genet* 2005, **110**:859–864.
2. Warburton ML, Crossa J, Franco J, Kazi M, Trethowan R, Rajaram S, Pfeiffer W, Zhang P, Dreisigacker S, van Ginkel M: **Bringing wild relatives back into the family: recovering genetic diversity in CIMMYT improved wheat germplasm.** *Euphytica* 2006, **149**:289–301.
3. Zhang L, Zhang L, Luo J, Chen W, Hao M, Liu B, Yan Z, Zhang B, Zhang H, Zheng Y, Liu D, Yen Y: **Synthesizing double haploid hexaploid wheat populations based on a spontaneous allopoloidization process.** *J Genet Genomics* 2011, **38**:89–94.
4. Crossa J, Burguño J, Dreisigacker S, Vargas M, Herrera-Foessel SA, Lillemo M, Singh RP, Trethowan R, Warburton M, Franco J, Reynolds M, Crouch JH, Ortiz R: **Association analysis of historical bread wheat germplasm using additive genetic covariance of relatives and population structure.** *Genetics* 2007, **177**:1889–1913.
5. Mantovani P, Maccaferri M, Sanguineti M, Tuberosa R, Catizone I, Wenzl P, Thomson B, Carling J, Huttner E, DeAmbrogio E, Kilian A: **An integrated DArT-SSR linkage map of durum wheat.** *Mol Breed* 2008, **22**:629–648.
6. Tyrka M, Bednarek PT, Kilian A, Wedzony M, Hura T, Bauer E: **Genetic map of triticale compiling DArT, SSR, and AFLP markers.** *Genome* 2011, **54**:391–401.
7. Paux E, Sourdille P, Salse J, Sautenac C, Choulet F, Leroy P, Korol A, Michalak M, Kianian S, Spielmeier W, Lagudah E, Somers D, Kilian A, Alaux M, Vautrin S, Berges H, Eversole K, Appels R, Safar J, Simkova H, Dolezel J, Bernard M, Feuillet C: **A physical map of the 1-gigabase bread wheat chromosome 3B.** *Science* 2008, **322**:101–104.
8. Francki MG, Walker E, Crawford AC, Broughton S, Ohm HW, Barclay I, Wilson RE, McLean R: **Comparison of genetic and cytogenetic maps of hexaploid wheat (*Triticum aestivum* L.) using SSR and DArT markers.** *Mol Genet Genomics* 2009, **281**:181–191.
9. Lyttle TW: **Segregation distorters.** *Annu Rev Genet* 1991, **25**:511–557.



10. Xu Y, Zhu L, Xiao J, Huang N, McCouch SR: **Chromosomal regions associated with segregation distortion of molecular markers in  $F_2$ , backcross, doubled haploid, and recombinant inbred populations in rice (*Oryza sativa*L.).** *Mol Gen Genet* 1997, **253**:535–545.
11. Singh K, Ghai M, Garg M, Chhuneja P, Kaur P, Schnurbusch T, Keller B, Dhaliwal HS: **An integrated molecular linkage map of diploid wheat based on a *Triticum boeoticum* × *T. monococcum* RIL population.** *Theor Appl Genet* 2007, **115**:301–312.
12. Messmer MM, Keller M, Zanetti S, Keller B: **Genetic linkage map of a wheat × spelt cross.** *Theor Appl Genet* 1999, **98**:1163–1170.
13. Paillard S, Schnurbusch T, Winzeler M, Messmer M, Sourdille P, Abderhalden O, Keller B, Schachermayr G: **An integrative genetic linkage map of winter wheat (*Triticum aestivum* L.).** *Theor Appl Genet* 2003, **107**:1235–1242.
14. Semagn K, Bjørnstad Å, Skinnen H, Marøy AG, Tarkegne Y, William M: **Distribution of DArT, AFLP, and SSR markers in a genetic linkage map of a doubled-haploid hexaploid wheat population.** *Genome* 2006, **49**:545–555.
15. Ma J, Li HB, Zhang CY, Yang XM, Liu YX, Yan GJ, Liu CJ: **Identification and validation of a major QTL conferring crown rot resistance in hexaploid wheat.** *Theor Appl Genet* 2010, **120**:1119–1128.
16. Blanco A, Bellomo MP, Cenci A, De Giovanni C, D'Ovidio R, Iacono E, Laddomada B, Pagnotta MA, Porceddu E, Sciancalepore A, Simeone R, Tanzarella OA: **A genetic linkage map of durum wheat.** *Theor Appl Genet* 1998, **97**:721–728.
17. Peng J, Korol AB, Fahima T, Röder MS, Ronin YI, Li YC, Nevo E: **Molecular genetic maps in wild emmer wheat, *Triticum dicoccoides*: genome-wide coverage, massive negative interference, and putative quasi-linkage.** *Genome Res* 2000, **10**:1509–1531.
18. Kumar S, Gill BS, Faris JD: **Identification and characterization of segregation distortion loci along chromosome 5B in tetraploid wheat.** *Mol Genet Genomics* 2007, **278**:187–196.
19. Elouafi I, Nachit MM: **A genetic linkage map of the Durum × *Triticum dicoccoides* backcross population based on SSRs and AFLP markers, and QTL analysis for milling traits.** *Theor Appl Genet* 2004, **108**:401–413.
20. Peleg Z, Saranga Y, Suprunova T, Ronin Y, Röder MS, Kilian A, Korol AB, Fahima T: **High-density genetic map of durum wheat × wild emmer wheat based on SSR and DArT markers.** *Theor Appl Genet* 2008, **117**:103–115.
21. Faris JD, Laddomada B, Gill BS: **Molecular mapping of segregation distortion loci in *Aegilops tauschii*.** *Genetics* 1998, **149**:319–327.
22. Alheit KV, Reif JC, Maurer HP, Hahn V, Weissmann EA, Miedaner T, Würschum T: **Detection of segregation distortion loci in triticale (× *Triticosecale* Wittmack) based on a high-density DArT marker consensus genetic linkage map.** *BMC Genomics* 2011, **12**:380.
23. Riley R, Chapman V: **The inheritance in wheat of crossability with rye.** *Genetical Res* 1967, **9**:259–267.
24. Tixier MH, Sourdille P, Charmet G, Gay G, Jaby C, Cadalen T, Bernard S, Nicolas P, Bernard M: **Detection of QTLs for crossability in wheat using a doubled-haploid population.** *Theor Appl Genet* 1998, **97**:1076–1082.
25. Genc I, Ozkan H, Yagbasanlar T: **Crossability of D-genome chromosome substitution lines of durum wheat (*Triticum turgidum* ssp. *turgidum* conv. *durum*) with *Secale cereale* and *Aegilops squarrosa*.** *Wheat Information Service* 1996, **83**:1–6.
26. Liu DC, Yen C, Yang JL, Lan XJ, Zheng YL: **The chromosome distribution of crossability gene in durum wheat cv. Langdon.** *Wheat Information Service* 1998, **87**:1–4.
27. Liu DC, Yen C, Yang JL, Zheng YL, Lan XJ: **The chromosomal locations of high crossability genes in tetraploid wheat *Triticum turgidum* L. cv. Ailanmai native to Sichuan, China.** *Euphytica* 1999, **108**:79–82.
28. Saghai-Marouf MA, Soliman KM, Jorgensen RA, Allard RW: **Ribosomal DNA spacer length polymorphism in barley: Mendelian inheritance, chromosomal location and population dynamics.** *Proc Natl Acad Sci USA* 1984, **81**:8014–8018.
29. Gupta P, Balyan H, Edwards K, Isaac P, Korzun V, Röder M, Gautier MF, Joudrier P, Schlatter A, Dubcovsky J, De la Pena R, Khairallah M, Penner G, Hayden M, Sharp P, Keller B, Wang R, Hardouin J, Jack P, Leroy P: **Genetic mapping of 66 new microsatellite (SSR) loci in bread wheat.** *Theor Appl Genet* 2002, **105**:413–422.
30. Somers DJ, Isaac P, Edwards K: **A high-density microsatellite consensus map for bread wheat (*Triticum aestivum* L.).** *Theor Appl Genet* 2004, **109**:1105–1114.
31. Röder MS, Korzun V, Wendehake K, Plaschke J, Tixier MH, Leroy P, Ganal MW: **A microsatellite map of wheat.** *Genetics* 1998, **149**:2007–2023.
32. Song QJ, Shi JR, Singh S, Fickus EW, Costa JM, Lewis J, Gill BS, Ward R, Cregan PB: **Development and mapping of microsatellite (SSR) markers in wheat.** *Theor Appl Genet* 2005, **110**:550–560.
33. Sourdille P, Singh S, Cadalen T, Brown-Guedira GL, Gay G, Qi L, Gill BS, Dufour P, Murigneux A, Bernard M: **Microsatellite-based deletion bin system for the establishment of genetic-physical map relationships in wheat (*Triticum aestivum* L.).** *Funct Integr Genomics* 2004, **4**:12–25.
34. Pestsova E, Ganal MW, Röder MS: **Isolation and mapping of microsatellite markers specific for the D genome of bread wheat.** *Genome* 2000, **43**:689–697.
35. Guyomarç'h H, Sourdille P, Charmet G, Edwards K, Bernard M: **Characterisation of polymorphic microsatellite markers from *Aegilops tauschii* and transferability to the D-genome of bread wheat.** *Theor Appl Genet* 2002, **104**:1164–1172.
36. Stephenson P, Bryan G, Kirby J, Collins A, Devos K, Busso C, Gale M: **Fifty new microsatellite loci for the wheat genetic map.** *Theor Appl Genet* 1998, **97**:946–949.
37. Zhang L, Liu D, Yan Z, Lan X, Zheng Y, Zhou Y: **Rapid changes of microsatellite flanking sequence in the allopolyploidization of new synthesized hexaploid wheat.** *Sci China C Life Sci* 2004, **47**:553–561.
38. Jaccoud D, Peng K, Feinstein D, Kilian A: **Diversity arrays: a solid state technology for sequence information independent genotyping.** *Nucleic Acids Res* 2001, **29**:E25.
39. Wenzl P, Carling J, Kudrna D, Jaccoud D, Huttner E, Kleinohs A, Kilian A: **Diversity Arrays Technology (DArT) for whole-genome profiling of barley.** *Proc Natl Acad Sci USA* 2004, **101**:9915–9920.
40. Akbari M, Wenzl P, Caig V, Carling J, Xia L, Yang S, Uszynski G, Mohler V, Lehmensiek A, Kuchel H, Hayden MJ, Howes N, Sharp P, Vaughan P, Rathmell B, Huttner E, Kilian A: **Diversity arrays technology (DArT) for high-throughput profiling of the hexaploid wheat genome.** *Theor Appl Genet* 2006, **113**:1409–1420.
41. Li H, Ye G, Wang J: **A modified algorithm for the improvement of composite interval mapping.** *Genetics* 2007, **175**:361–374.
42. Van Os H, Stam P, Visser RG, Van Eck HJ: **RECORD: a novel method for ordering loci on a genetic linkage map.** *Theor Appl Genet* 2005, **112**:30–40.
43. Kosambi DD: **The estimation of map distances from recombination values.** *Ann Hum Genet* 1943, **12**:172–175.
44. Van Ooijen JW: **JoinMap<sup>®</sup> 4 Software for calculation of genetic linkage map in experimental populations.** Netherlands: Kyzacma B. V. Wageningen; 2006.

doi:10.1186/1471-2156-13-69

**Cite this article as:** Zhang et al.: Genetic map of *Triticum turgidum* based on a hexaploid wheat population without genetic recombination for D genome. *BMC Genetics* 2012 **13**:69.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

