



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: April 17, 2010

ACCEPTED: July 29, 2010

PUBLISHED: August 12, 2010

Non-perturbative renormalization of quark bilinear operators with $N_f = 2$ (tmQCD) Wilson fermions and the tree-level improved gauge action

European Twisted Mass collaboration



M. Constantinou,^a P. Dimopoulos,^b R. Frezzotti,^{c,d} G. Herdoiza,^e K. Jansen,^e
V. Lubicz,^{f,g,1} H. Panagopoulos,^a G.C. Rossi,^{c,d} S. Simula,^g F. Stylianou^a and
A. Vladikas^d

^aDepartment of Physics, University of Cyprus, P.O.Box 20537, Nicosia CY-1678, Cyprus

^bDip. di Fisica, Università di Roma “La Sapienza” and INFN, Sezione di Roma,
P.le A. Moro 2, I-00185 Rome, Italy

^cDip. di Fisica, Università di Roma “Tor Vergata”,
Via della Ricerca Scientifica 1, I-00133 Rome, Italy

^dINFN, Sez. di Roma “Tor Vergata”, c/o Dip. di Fisica, Università di Roma “Tor Vergata”,
Via della Ricerca Scientifica 1, I-00133 Rome, Italy

^eNIC, DESY, Platanenallee 6, D-15738 Zeuthen, Germany

^fDip. di Fisica, Università di Roma Tre, Via della Vasca Navale 84, I-00146 Rome, Italy

^gINFN, Sez. di Roma III, Via della Vasca Navale 84, I-00146 Rome, Italy

E-mail: lubicz@fis.uniroma3.it

¹Corresponding author.

ABSTRACT: We present results for the renormalization constants of bilinear quark operators obtained by using the tree-level Symanzik improved gauge action and the $N_f = 2$ twisted mass fermion action at maximal twist, which guarantees automatic $\mathcal{O}(a)$ -improvement. Our results are also relevant for the corresponding standard (un-twisted) Wilson fermionic action since the two actions only differ, in the massless limit, by a chiral rotation of the quark fields. The scale-independent renormalization constants Z_V , Z_A and the ratio Z_P/Z_S have been computed using the RI-MOM approach, as well as other alternative methods. For Z_A and Z_P/Z_S , the latter are based on both standard twisted mass and Osterwalder-Seiler fermions, while for Z_V a Ward Identity has been used. The quark field renormalization constant Z_q and the scale dependent renormalization constants Z_S , Z_P and Z_T are determined in the RI-MOM scheme. Leading discretization effects of $\mathcal{O}(g^2 a^2)$, evaluated in one-loop perturbation theory, are explicitly subtracted from the RI-MOM estimates.

KEYWORDS: Lattice QCD, Renormalization Regularization and Renormalons

ARXIV EPRINT: [1004.1115](https://arxiv.org/abs/1004.1115)

Contents

1	Introduction	1
2	A novel approach to the calculation of the scale independent RCs	2
2.1	Calculation of Z_P/Z_S	4
2.2	Calculation of Z_A	4
2.3	Calculation of Z_V	5
2.4	Results	5
3	Renormalization constants in the RI-MOM scheme	9
3.1	Analysis of the twisted mass quark propagator	11
3.2	Renormalization constants	13
3.2.1	Chiral extrapolations	13
3.2.2	Renormalization scale dependence and subtraction of the $\mathcal{O}(g^2 a^2)$ effects	14
3.3	Analysis of systematic errors	22
3.4	Final results and comparison with perturbation theory	25
4	Conclusions	28
A	$\mathcal{O}(a)$-improvement of RI-MOM renormalization constants	28
A.1	Improvement at maximal twist	28
A.1.1	The proof	30
A.2	Improvement at a generic value of the twist angle	30
A.2.1	The proof	30
A.3	A final comment	31

1 Introduction

Non-perturbative renormalization is an essential ingredient of lattice QCD calculations which aim at a percent level accuracy. In this paper, we present calculation methods and results for the renormalization constants (RCs) of bilinear quark operators for the lattice action used by the ETM Collaboration. The simulations with $N_f = 2$ dynamical flavours employ the tree-level Symanzik improved gauge action and the twisted mass fermionic action at maximal twist. In the chiral limit, the latter is related to the standard Wilson fermionic action by a chiral transformation, under which quark composite operators behave in a definite and simple way. Therefore the RCs we compute here can be also employed to renormalize bilinear quark operator matrix elements computed with the same glue action but standard (un-twisted) Wilson quarks.

For the computation of the bilinear operator RCs we have used the RI-MOM method [1]. For the calculation of the scale independent RCs, Z_V , Z_A and the ratio Z_P/Z_S , a set of alternative methods have also been used. In particular, we have determined Z_V by imposing the validity of the axial Ward identity whereas for Z_A and Z_P/Z_S we have employed a new method which makes a combined use of standard twisted mass (tm) and Osterwalder-Seiler (OS) formalism. A comparison of the results obtained with different approaches provides an estimate of the size of the systematic errors affecting the calculation of the RCs at fixed lattice bare coupling. Needless to say, in renormalized quantities the discretization errors coming from both RCs and bare quantities will be removed by continuum extrapolation, as illustrated with an example in section 3.3.

Throughout the paper we follow the standard practice of labelling the RCs according to the notation adopted in the un-twisted Wilson case. Thus, for instance, Z_V and Z_A denote the RCs of the local vector and axial-vector currents of the standard Wilson action even though, in the maximal twist case, they renormalize in the physical basis the local (charged) axial-vector and polar-vector currents respectively. We refer to table 1 for the explicit presentation of the renormalization pattern.

We have computed the RCs at three values of the gauge coupling, namely $\beta = 3.80$, 3.90 and 4.05, which correspond to inverse lattice spacing $a^{-1} \simeq 2.0$, 2.3 and 2.9 GeV. Preliminary results of this work have been presented in ref. [2]. Further details concerning the ETMC simulations can be found in refs. [3]–[6].

The paper’s contents are the following. In section 2 we present the novel approach of computing the scale independent RCs Z_A and Z_P/Z_S and the Ward identity determination of Z_V . In section 3 we discuss the RI-MOM approach applied to tm quarks and provide the details of the numerical analysis. A collection of our final results for the RCs can be found in tables 4, 5 and 6. In tables 4 and 5 the predictions of one loop boosted perturbation theory [7, 8] for the various RCs are also given. Finally in the appendix we present the proof of the automatic $\mathcal{O}(a)$ -improvement of the RI-MOM RCs calculated with maximally twisted quarks.

An RI-MOM calculation of the RCs of bilinear quark operators similar to the one described in the present paper but for the standard Wilson’s plaquette gauge action and non-perturbatively improved Wilson (clover) fermions has been recently presented in ref. [9].

2 A novel approach to the calculation of the scale independent RCs

In this section we present a calculation of the scale independent RCs, namely Z_V , Z_A and Z_P/Z_S . The evaluation of Z_V is based on the PCAC Ward identity, which in the quenched approximation has led to very precise results (see e.g. refs. [10, 11]). On the other hand, Z_A and Z_P/Z_S are computed from two-point correlators only with a new method, which is based on the simultaneous use of two regularizations of the valence quark action. One is the standard tm action at maximal twist, while the other is the OS variant [12, 13]. In the so called physical basis these actions can be written in the form:

$$S_{\text{val}} = a^4 \sum_x \sum_{q=u,d} \bar{q}(x) \left(\gamma \tilde{\nabla} - i\gamma_5 r_q W_{\text{cr}} + \mu_q \right) q(x) , \tag{2.1}$$

OS case	tm case
$(A_R)_{\mu,ud} = Z_A A_{\mu,ud} = Z_A A'_{\mu,ud}$	$(A_R)_{\mu,ud} = Z_V A_{\mu,ud} = -i Z_V V'_{\mu,ud}$
$(V_R)_{\mu,ud} = Z_V V_{\mu,ud} = Z_V V'_{\mu,ud}$	$(V_R)_{\mu,ud} = Z_A V_{\mu,ud} = -i Z_A A'_{\mu,ud}$
$(P_R)_{ud} = Z_S P_{ud} = i Z_S S'_{ud}$	$(P_R)_{ud} = Z_P P_{ud} = Z_P P'_{ud}$
$(T_R)_{\mu\nu,ud} = Z_T T_{\mu\nu,ud} = i Z_T \tilde{T}'_{\mu\nu,ud}$	$(T_R)_{\mu\nu,ud} = Z_T T_{\mu\nu,ud} = Z_T T'_{\mu\nu,ud}$

Table 1. Renormalization pattern of the bilinear quark operators for the OS and tm case at maximal twist. The unprimed operators refer to the physical basis while the primed ones to the twisted basis. The symbols P_{ud} and $\tilde{T}_{\mu\nu,ud}$ indicate the operators $\bar{u}\gamma_5 d$ and $\bar{u}\sigma_{\mu\nu}\gamma_5 d$.

with $W_{cr} = -\frac{a}{2} \sum_{\mu} \nabla_{\mu}^* \nabla_{\mu} + M_{cr}$. The Wilson parameters are $r_u = -r_d = 1$ for the standard tm case and $r_u = r_d = 1$ for OS quarks. In the sea sector we have two degenerate quarks, regularized in the standard tm framework.

Let us illustrate the general idea of the calculation of the scale independent RCs, Z_A and Z_P/Z_S . It is based on the observation that the axial transformations of the quark fields from the physical basis to the so called twisted basis (primed fields),

$$q = \exp(i\gamma_5 r_q \pi/4) q' \quad \text{and} \quad \bar{q} = \bar{q}' \exp(i\gamma_5 r_q \pi/4), \quad (2.2)$$

transform the tm and OS actions of eq. (2.1) into an action with the Wilson term in the standard form (i.e. having no γ_5 and $r_u = r_d = 1$) and a mass term which takes the form $(i\mu_u \bar{u}' \gamma_5 u' - i\mu_d \bar{d}' \gamma_5 d')$ in the tm case and $(i\mu_u \bar{u}' \gamma_5 u' + i\mu_d \bar{d}' \gamma_5 d')$ in the OS case. This also implies, in turn, that the RCs for the operators in the twisted basis, being defined in the chiral limit, are the same for the Wilson, tm and OS cases. Consider, now, a non-singlet quark bilinear operator, $O_{\Gamma} = \bar{u}\Gamma d$, defined in terms of the fields of the physical basis. Under the axial transformations of eq. (2.2) this operator transforms into an operator in the twisted basis, denoted as $O_{\tilde{\Gamma}}$ and $O_{\hat{\Gamma}}$ for the tm and OS case respectively. In general, the two operators are not of the same form. However their renormalized matrix elements between given physical states, say $Z_{O_{\tilde{\Gamma}}} \langle \alpha | O_{\tilde{\Gamma}} | \beta \rangle^{\text{tm}}$ and $Z_{O_{\hat{\Gamma}}} \langle \alpha | O_{\hat{\Gamma}} | \beta \rangle^{\text{OS}}$ are estimates of the same physical matrix element $\langle \alpha | (O_{\Gamma})_R | \beta \rangle$ up to $\mathcal{O}(a^2)$ errors:

$$\langle \alpha | (O_{\Gamma})_R | \beta \rangle = Z_{O_{\tilde{\Gamma}}} \langle \alpha | O_{\tilde{\Gamma}} | \beta \rangle^{\text{tm}} + \mathcal{O}(a^2) = Z_{O_{\hat{\Gamma}}} \langle \alpha | O_{\hat{\Gamma}} | \beta \rangle^{\text{OS}} + \mathcal{O}(a^2). \quad (2.3)$$

RCs are named, as anticipated, after the twisted basis, in which the Wilson term has its standard form. The operator renormalization pattern in the physical and twisted bases is given in table 1 for both OS and tm formulations at maximal twist. The primed operators refer to the twisted basis while the unprimed ones to the physical basis. Notice that the condition of maximal twist ensures that cut-off effects are of order $\mathcal{O}(a^2)$ [14].

The main point of eq. (2.3) is that the tm and OS determinations of the same continuum matrix element are equal up to discretization effects. Our proposal for the determination of Z_P/Z_S and Z_A , described in the following subsections, is based on this observation.

2.1 Calculation of Z_P/Z_S

The method is based on comparing the amplitude $g_\pi = \langle 0|P|\pi \rangle$, computed both in tm and OS regularizations. We start by considering, in the physical basis, the correlation function

$$C_{PP}(t) \equiv - \sum_{\mathbf{x}} \langle \bar{u}\gamma_5 d(x) \bar{d}\gamma_5 u(0) \rangle , \quad (2.4)$$

which at large time separations behaves like

$$C_{PP}(t) \simeq \frac{|g_\pi|^2}{2m_\pi} [\exp(-m_\pi t) + \exp(-m_\pi(T-t))] \quad (2.5)$$

From table 1 we see that in the twisted basis this corresponds to $C_{S'S'}(t)$ (for OS regularization) and $C_{P'P'}(t)$ (for tm regularization):

$$[C_{PP}(t)]^{\text{cont}} = Z_P^2 [C_{P'P'}(t)]^{\text{tm}} + \mathcal{O}(a^2) = Z_S^2 [C_{S'S'}(t)]^{\text{OS}} + \mathcal{O}(a^2) , \quad (2.6)$$

which implies

$$[g_\pi]^{\text{cont}} = Z_P [g_\pi]^{\text{tm}} + \mathcal{O}(a^2) = Z_S [g_\pi]^{\text{OS}} + \mathcal{O}(a^2) . \quad (2.7)$$

The ratio Z_P/Z_S is extracted from the above equation in the chiral limit. As our simulations are performed at finite quark masses, an extrapolation of our Z_P/Z_S -estimators to zero quark mass is necessary.

2.2 Calculation of Z_A

In order to compute Z_A , we consider the calculation of the charged pseudoscalar meson decay constant f_π , in both OS and tm regularizations. We start with the tm regularization. From the axial Ward identity in the physical basis, we have the well-known result for the pseudoscalar meson decay constant [15]:

$$[f_\pi]^{\text{cont}} = [f_\pi]^{\text{tm}} + \mathcal{O}(a^2) = \left[\frac{2\mu_q g_\pi}{m_\pi^2} \right]^{\text{tm}} + \mathcal{O}(a^2) . \quad (2.8)$$

Note that in this case no RC is needed [15]. Thus the pion decay constant can be extracted from g_π , i.e. from the large time asymptotic behaviour of $[C_{P'P'}(t)]^{\text{tm}}$.

In the OS formulation we consider, besides the correlator C_{PP} of eq. (2.4), also C_{AP} , which is defined by

$$C_{AP}(t) \equiv \sum_{\mathbf{x}} \langle \bar{u}\gamma_0\gamma_5 d(x) \bar{d}\gamma_5 u(0) \rangle . \quad (2.9)$$

Its large time asymptotic behaviour is:

$$C_{AP}(t) \simeq \frac{\xi_{AP}}{2m_\pi} [\exp(-m_\pi t) - \exp(-m_\pi(T-t))] , \quad (2.10)$$

with

$$\xi_{AP} = \langle 0|A_0|\pi \rangle \langle \pi|P|0 \rangle = [f_\pi m_\pi i g_\pi]^{\text{OS}} . \quad (2.11)$$

The last equation may be solved for f_π , which is thus obtained from the correlation functions of eqs. (2.5) and (2.10). With the aid of table 1, this solution is expressed in terms of OS-regularized quantities in the twisted basis (note $\xi_{AP} = i\xi_{A'S'}$) as follows:

$$[f_\pi]^{\text{cont}} = Z_A [f_\pi]^{\text{OS}} + \mathcal{O}(a^2) = Z_A \frac{\xi_{A'S'}}{[g_\pi]^{\text{OS}} [m_\pi]^{\text{OS}}} + \mathcal{O}(a^2) . \quad (2.12)$$

Combining eqs. (2.8) and (2.12), we obtain

$$[f_\pi]^{\text{cont}} = [f_\pi]^{\text{tm}} + \mathcal{O}(a^2) = Z_A [f_\pi]^{\text{OS}} + \mathcal{O}(a^2) , \quad (2.13)$$

from which an estimate of Z_A can be extracted. Again, results obtained at finite quark masses need to be extrapolated to the chiral limit. In ref. [16] the discretization effects affecting the quenched pseudoscalar decay constant in the tm and OS regularizations have been studied in detail. The tm and OS results were compatible within one standard deviation for three values of the lattice spacing, indicating the smallness of $\mathcal{O}(a^2)$ cut-off effects.

2.3 Calculation of Z_V

The determination of the renormalization constant Z_V can be done using *solely* tm quarks. It is based on the comparison of the point-like axial current $A_{\mu,ud}$, which renormalizes with Z_V (see table 1), with the exactly conserved one-point split current $[V_\mu]_{ud}^{\text{1PS}} = [A_\mu]_{ud}^{\text{1PS}}$. The four-divergence of the latter is exactly equal to the sum of the valence quark mass values, $(\mu_1 + \mu_2)$, times the corresponding pseudoscalar density. Therefore, we extract Z_V by solving the equation:

$$Z_V \tilde{\partial}_0 [C_{V'P'}(t)]^{\text{tm}} = (\mu_1 + \mu_2) [C_{P'P'}(t)]^{\text{tm}} \quad (2.14)$$

where $\tilde{\partial}_0$ is the symmetric lattice time derivative, and extrapolating results to the chiral limit.

2.4 Results


In table 2 we give various details of our $N_f = 2$, partially quenched simulations. The smallest sea quark mass corresponds to a pion of about 300 MeV for the two larger β values, while for the smallest coupling the lightest pion weights approximately 400 MeV. The highest sea quark mass is around half the strange quark mass. For the inversions in the valence sector we have made use of the stochastic method (one-end trick of ref. [17]) in order to increase the signal to noise ratio. Propagator sources have been placed at randomly located timeslices. This turned out to be an optimal way to further reduce the autocorrelation time. Our correlation functions are computed for all combinations of valence quark masses appearing in table 2. In order to deal with effectively independent measurements of RC-estimators, we have selected gauge field configurations separated by 20 HMC (length 1/2) trajectories. Within each ensemble of such gauge configurations statistical errors are evaluated using a jackknife procedure. For results involving an extrapolation in the sea quark mass and/or combined fits at several lattice couplings, statistical errors have been estimated with a bootstrap procedure for a 1000 bootstrap events.

	β	$L^3 \times T$	$a\mu_{\text{sea}}$	$a\mu_{\text{val}}$	N_{conf}
A_2	3.80	$24^3 \times 48$	0.0080	{0.0080, 0.0110, 0.0165, 0.0200}	400/240
A_3			0.0110	{0.0080, 0.0110, 0.0165, 0.0200}	400/240
A_4			0.0165	{0.0080, 0.0110, 0.0165, 0.0200}	400/240
B_1	3.90	$24^3 \times 48$	0.0040	{0.0040, 0.0064, 0.0085, 0.0100, 0.0150}	240/240
B_2			0.0064	{0.0040, 0.0064, 0.0085, 0.0100, 0.0150}	240/240
B_3			0.0085	{0.0040, 0.0064, 0.0085, 0.0100, 0.0150}	240/240
B_4			0.0100	{0.0040, 0.0064, 0.0085, 0.0100, 0.0150}	240/240
C_1	4.05	$32^3 \times 64$	0.0030	{0.0030, 0.0060, 0.0080, 0.0120}	130/240
C_2			0.0060	{0.0030, 0.0060, 0.0080, 0.0120}	130/160
C_3			0.0080	{0.0030, 0.0060, 0.0080, 0.0120}	130/160


Table 2. Details of the simulations performed for computing RCs. The number N_{conf} of gauge configurations we analysed is given for both the case of the scale independent RCs discussed in this section (first figure) and the RI-MOM analysis of section 3 (second figure).

Typical plots on the data quality of Z_V , Z_A and Z_P/Z_S are shown in figure 1; in this example we show the case $a\mu_{\text{sea}}^{\text{min}} = a\mu_1 = a\mu_2$ for the three values of the gauge coupling.


Various methods have been implemented in taking the chiral limit (in the sea and valence quark sector). A first method consists in calculating the RCs at fixed sea quark mass for three choices of valence quark pairs, namely with $a\mu_1 = a\mu_2$, or for all pairs of $(a\mu_1, a\mu_2)$ or for $(a\mu_1, a\mu_2) \geq a\mu_{\text{sea}}$ and taking the “valence chiral limit” using a linear fit in terms of the sum of the valence quark masses. Subsequently, the RCs were quadratically extrapolated to the sea quark chiral limit. Note that a quadratic dependence on $a\mu_{\text{sea}}$ is expected from the form of the sea quark determinant, assuming that lattice artefacts on the RCs are not sensitive to spontaneous chiral symmetry breaking. However we have verified that a linear fit in μ_{sea} leads to compatible results, albeit with larger final errors. A second method consists in inverting the order of the two chiral limits. A third method is simply the extraction of the RCs from the subset of data satisfying $\mu_{\text{valence}} = \mu_{\text{sea}}$. This allows reaching the chiral limit with a single, linear extrapolation in the quark mass. In most cases the quality of the fits is very good and the final results, obtained from these different extrapolations, are compatible within one standard deviation. We have opted to quote as final results those produced by the first method, for all pairs of $(a\mu_1, a\mu_2)$, followed by a linear fit in terms of $a^2\mu_{\text{sea}}^2$. In figure 2 (left panel) we show the “valence chiral limit” extrapolation of the three scale independent RCs, computed at the lightest sea quark mass for each gauge coupling. In figure 2 (right panel) we present the chiral limit in the sea sector for all the RCs for each gauge coupling.



(a)



(b)



(c)

Figure 1. The plateaux quality for Z_V (a), Z_A (b) and Z_P/Z_S (c) for three gauge couplings and masses $a\mu_{\text{sea}}^{\text{min}} = a\mu_1 = a\mu_2$. From table 2 we have $a\mu_{\text{sea}}^{\text{min}}(\beta = 3.80) = 0.0080$, $a\mu_{\text{sea}}^{\text{min}}(\beta = 3.90) = 0.0040$ and $a\mu_{\text{sea}}^{\text{min}}(\beta = 4.05) = 0.0030$. Note that in figure(b) the data have been shifted for clarity.




Figure 2. Left panel: valence chiral limit for the three scale independent RCs at $\beta = 3.80$, $a\mu_{\text{sea}} = 0.0080$ (a), $\beta = 3.90$, $a\mu_{\text{sea}} = 0.0040$ (c) and $\beta = 4.05$, $a\mu_{\text{sea}} = 0.0030$ (e). Right panel: sea chiral limit for the same constants at $\beta = 3.80$ (b), $\beta = 3.90$ (d) and $\beta = 4.05$ (f). Note that the data for Z_P/Z_S have been shifted for clarity.

Our final results are collected in table 4 (see rows labelled as “Alt. methods”). For comparison with the RI-MOM results and conclusions on the precision of the methods used, see below. Here we point out that the Z_V value at $\beta = 3.90$ compares nicely with the result of ref. [18], produced from 3- and 2-point correlation functions, in the context of a calculation of the pion form factor.

3 Renormalization constants in the RI-MOM scheme

The non-perturbative determination of the RCs in the RI-MOM scheme [1] is based on the numerical evaluation, in momentum space, of correlation functions of the relevant operators between external quark and/or gluon states. In this work we are interested in the case of the bilinear quark operators $O_\Gamma = \bar{u}\Gamma d$, where $\Gamma = S, P, V, A, T$ stands for $I, \gamma_5, \gamma_\mu, \gamma_\mu\gamma_5, \sigma_{\mu\nu}$ respectively. The fields u and d of the operator O_Γ belong to a twisted doublet of quarks, regularized by the tm action of eq. (2.1), with $r_u = -r_d = 1$.

The relevant Green functions are the quark propagator

$$S_q(p) = a^4 \sum_x e^{-ip \cdot x} \langle q(x) \bar{q}(0) \rangle \quad (q = u, d), \quad (3.1)$$

the forward vertex function

$$G_\Gamma(p) = a^8 \sum_{x,y} e^{-ip \cdot (x-y)} \langle u(x) O_\Gamma(0) \bar{d}(y) \rangle, \quad (3.2)$$

and the amputated Green function¹

$$\Lambda_\Gamma(p) = S_u(p)^{-1} G_\Gamma(p) S_d(p)^{-1}. \quad (3.3)$$

The RI-MOM renormalization scheme consists in imposing that the forward amputated Green function, computed in the chiral limit in the Landau gauge and at a given (large Euclidean) scale $p^2 = \mu^2$, is equal to its tree-level value. In practice, the renormalization condition is implemented by requiring in the chiral limit that

$$Z_q^{-1} Z_\Gamma \mathcal{V}_{\tilde{\Gamma}}(p)|_{p^2=\mu^2} \equiv Z_q^{-1} Z_\Gamma \text{Tr}[\Lambda_{\tilde{\Gamma}}(p) P_{\tilde{\Gamma}}]|_{p^2=\mu^2} = 1, \quad \tilde{\Gamma} = e^{-i\gamma_5\pi/4} \Gamma e^{i\gamma_5\pi/4}, \quad (3.4)$$

where $P_{\tilde{\Gamma}}$ is a Dirac projector satisfying $\text{Tr}[\tilde{\Gamma} P_{\tilde{\Gamma}}] = 1$.² The quark field RC Z_q , which

¹The calculation of $G_\Gamma(p)$ in eq. (3.2) involves the propagator $S_d(0, y)$ which, in the tm approach, equals $\gamma_5 S_u(y, 0)^\dagger \gamma_5$. Similarly, in eq. (3.3) the propagator $S_d(p)$ stands for $\sum_y e^{ip \cdot y} S_d(0, y) = \gamma_5 S_u(p)^\dagger \gamma_5$.

²The appearance of $\tilde{\Gamma}$ in eq. (3.4) is a trivial consequence of the fact that RCs, as we said, are named after the (here unphysical) twisted quark basis while the operator $O_\Gamma = \bar{u}\Gamma d$ is expressed in the physical quark basis.

enters eq. (3.4), is obtained by imposing, in the chiral limit, the condition³

$$Z_q^{-1} \Sigma_1(p)|_{p^2=\mu^2} \equiv Z_q^{-1} \frac{-i}{12} \text{Tr} \left[\frac{\not{p} S_q(p)^{-1}}{p^2} \right]_{p^2=\mu^2} = 1. \quad (3.5)$$

In order to minimize discretization effects, we select momenta with components $p_\nu = (2\pi/L_\nu) n_\nu$ in the intervals

$$n_\nu = \begin{cases} ([0, 2], [0, 2], [0, 2], [0, 3]) \\ ([2, 3], [2, 3], [2, 3], [4, 7]) \end{cases}, \quad \text{for } L = 24 \quad (\beta = 3.8 \text{ and } 3.9) \quad (3.6)$$

and

$$n_\nu = \begin{cases} ([0, 2], [0, 2], [0, 2], [0, 3]) \\ ([2, 5], [2, 5], [2, 5], [4, 9]) \end{cases}, \quad \text{for } L = 32 \quad (\beta = 4.05) \quad (3.7)$$

(L_ν is the lattice size in the direction ν). The time component of the four-momentum is shifted by the constant $\Delta p_4 = \pi/L_4$, in order to account for the use of anti-periodic boundary conditions on the quark fields in the time direction. Furthermore, we have only considered for the final RI-MOM analysis the momenta satisfying

$$\Delta_4(p) \equiv \frac{\sum_\rho \tilde{p}_\rho^4}{\left(\sum_\rho \tilde{p}_\rho^2\right)^2} < 0.28, \quad (3.8)$$

where

$$\tilde{p}_\nu \equiv \frac{1}{a} \sin(ap_\nu), \quad (3.9)$$

which helps in minimizing the contribution of Lorentz non-invariant discretization effects (cfr eq. (3.17)).

In order to improve the statistical accuracy of the RCs, we have averaged the results obtained in the chiral limit from the correlation functions of the operators $O_\Gamma = \bar{u}\Gamma d$ and $O'_\Gamma = \bar{d}\Gamma u$. Similarly, in the case of the quark field RC, we have computed Z_q by averaging the results obtained for the up and down quark propagators.

The RCs, calculated in the chiral limit in the way described above, are automatically improved at $\mathcal{O}(a)$. Actually in the maximal twist situation $\mathcal{O}(a)$ improvement holds for all external momenta p and non-zero (valence and sea) quark mass at the level of the basic quantities $\text{Tr}[\Lambda_\Gamma(p) P_\Gamma]$ and $\text{Tr}[(\not{p} S_q(p)^{-1})/p^2]$ entering in eqs. (3.4)–(3.5). A proof of this statement, which follows from an analysis based on the symmetries of the tm action

³Strictly speaking, the renormalization condition of eq. (3.5) defines the so called RI' scheme. In the original RI-MOM scheme the quark field renormalization condition reads

$$Z_q^{-1} \frac{-i}{48} \text{Tr} \left[\gamma_\mu \frac{\partial S_q(p)^{-1}}{\partial p_\mu} \right]_{p^2=\mu^2} = 1.$$

The two schemes differ in the Landau gauge at the N²LO. In this work, when perturbative predictions are used, the difference between the two schemes has been properly taken into account.

at maximal twist and the $O(4)$ symmetry of the underlying continuum theory, is given in the appendix.⁴

Details of the RI-MOM simulation are collected in table 2. The lattice parameters are those used also in the determination of the scale independent constants with the methods discussed in the previous section. However, a different ensemble of independent gauge configurations has been analyzed in each case. Moreover, the inversions in the valence sector for the RI-MOM study have been performed without using the stochastic approach, but with point-like sources randomly located on the lattice for each gauge configuration.

3.1 Analysis of the twisted mass quark propagator

The necessary Green functions for the RI-MOM determination of RCs of bilinear quark operators are the quark propagator $S_q(p)$ and the amputated vertex $\Lambda_\Gamma(p)$, evaluated in momentum space in the Landau gauge. In this section, we first illustrate the results on the tm quark propagator.

At $\mathcal{O}(a)$, the spin-flavour structure of the lattice artefacts of the quark propagator differs from that of the standard Wilson case, due to the twisted Wilson term in the action. With tm fermions, the explicit breaking of parity at finite lattice spacing allows for the presence of a parity violating contribution in the quark propagator. By neglecting $O(4)$ symmetry violating effects, which only appear at $\mathcal{O}(a^2)$, the inverse quark propagator can be expressed in terms of three scalar form factors, as follows:

$$S_q(p)^{-1} = i \not{p} \Sigma_1(p^2) + \Sigma_2(p^2) - i \gamma_5 \Sigma_3(p^2). \quad (3.10)$$

At large p^2 , Σ_1 and Σ_2 are related to the quark field RC Z_q (see eq. (3.5)), and to the renormalized quark mass $\hat{\mu}_q$ respectively [20]. The parity violating term proportional to Σ_3 represents an $\mathcal{O}(a)$ discretization effect, induced by the twisted Wilson term in the action.⁵ At maximal twist, the form factors $\Sigma_{1,2,3}$ are given at tree-level by

$$\Sigma_1(p^2) = 1, \quad \Sigma_2(p^2) = \mu_q, \quad \Sigma_3(p^2) = \frac{a r_q}{2} p^2. \quad (3.11)$$

In figure 3, the non-perturbatively determined form factors $\Sigma_{1,2,3}$ are plotted against $a^2 \tilde{p}^2$, for $\beta = 3.90$ and $a\mu_{\text{sea}} = 0.0040$. The following observations are in order:

- The form factor Σ_1 exhibits a nice plateau at large p^2 , with a tiny slope which is partly due to a NLO renormalization scale dependence (the quark field anomalous dimension vanishes at LO in the Landau gauge) and partly to $\mathcal{O}(a^2)$ discretization effects. The quark field RC Z_q is obtained from the Σ_1 results at large p^2 , extrapolated to the chiral limit (see eq. (3.5)).

⁴As discussed in the appendix, with the standard Wilson or Clover actions, the RCs determined with the RI-MOM method are also $\mathcal{O}(a)$ -improved. The RC-estimators are however improved only at large momenta, where spontaneous chiral symmetry breaking effects can be neglected [19]. This is sufficient, as the RI-MOM scheme is defined precisely in the high-momentum region. We remark, however, that the $\mathcal{O}(a)$ -improvement of the RI-MOM RCs, with the standard Wilson or Clover actions, does not imply the automatic $\mathcal{O}(a)$ -improvement of the operators as well. The latter can only be achieved, in the Clover case, by implementing a mixing with higher-dimension operators with properly tuned coefficients.

⁵In the standard Wilson case, the analogous $\mathcal{O}(a)$ artifact contributes to the form factor Σ_2 .




Figure 3. The form factors Σ_1 (top), Σ_2 (center) and Σ_3 (bottom), computed at $\beta = 3.90$ with $a\mu_{\text{sea}} = 0.0040$, as a function of $a^2\tilde{p}^2$.

- The form factor Σ_2 also exhibits a plateau at large p^2 , which is expected to be proportional, up to the quark field RC, to the renormalized valence quark mass $\hat{\mu}_q$ in the RI-MOM scheme. Indeed, at variance with the cases of Σ_1 and Σ_3 , a clear separation among the results obtained for Σ_2 at different bare valence quark masses is visible in the plot.
- As expected, the form factor Σ_3 , which represents a pure $\mathcal{O}(a)$ discretization effect, has opposite sign for the up and down quark propagators ($r_u = -r_d = 1$). Its behaviour is quite close to the tree level estimate of eq. (3.11). We find $\Sigma_3 \simeq \pm c_q a p^2 / 2$, with $c_q \simeq 0.8 \div 0.9$.

3.2 Renormalization constants

The RCs in the RI-MOM scheme are determined by closely following the procedure summarised in section 3 above and described in detail in ref. [19]. This procedure involves two main steps: the chiral limit extrapolation of the RCs, at fixed coupling and renormalization scale, and the study of the renormalization scale dependence.

3.2.1 Chiral extrapolations

RI-MOM is a mass independent renormalization scheme. Since in practice the RCs are obtained at non vanishing values of the quark masses, an extrapolation of the results to the chiral limit, both in the valence and the sea quark masses, must be performed.

The validity of the RI-MOM approach relies on the fact that, at large p^2 , Green functions are expected to depend smoothly on the quark masses, since their non-perturbative contributions vanish asymptotically in that limit. However, specific care must be taken in the study of the pseudoscalar Green function \mathcal{V}_P , since the leading $1/p^2$ -suppressed contribution is divergent in the chiral limit, due to the coupling with the Goldstone boson [1, 21, 22]. Moreover, for tm fermions, the explicit breaking of parity at finite lattice spacing induces a coupling of the Goldstone boson also with the scalar operator. Though the latter is suppressed at $\mathcal{O}(a^2)$, its effect may be not negligible at the couplings considered in the present simulations.

In order to subtract the pseudoscalar mass dependence of the amputated vertex functions \mathcal{V}_P and \mathcal{V}_S , at each given p^2 we fit their values at different quark masses to the Ansatz

$$\mathcal{V}_{P(S)}(p^2, \mu_1, \mu_2) = A_{P(S)}(p^2) + B_{P(S)}(p^2) M_{\text{PS}}^2(\mu_1, \mu_2) + \frac{C_{P(S)}(p^2)}{M_{\text{PS}}^2(\mu_1, \mu_2)}, \quad (3.12)$$

where $M_{\text{PS}}(\mu_1, \mu_2)$ is the mass of the pseudoscalar meson composed by valence quarks of mass μ_1 and μ_2 (with $r_1 = -r_2$).⁶ The first term in the fit, i.e. the function $A_{P(S)}(p^2)$, represents the Green function $\mathcal{V}_{P(S)}$ in the (valence) chiral limit, from which the RC

⁶In our preliminary analysis presented in [2], the fitting function used to subtract the Goldstone pole contribution was expressed in terms of the sum of quark masses $\mu_1 + \mu_2$ rather than the pseudoscalar meson mass squared $M_{\text{PS}}(\mu_1, \mu_2)^2$, as in eq. (3.12). Since the latter is simply proportional to $\mu_1 + \mu_2$ at maximal twist, the two fits lead to completely consistent results.

$Z_{P(S)}$ is eventually extracted. The last term in eq. (3.12) accounts for the presence of the Goldstone pole. Using the results of the fit, we can define the subtracted correlators

$$\mathcal{V}_{P(S)}^{\text{sub}}(p^2, \mu_1, \mu_2) = \mathcal{V}_{P(S)}(p^2, \mu_1, \mu_2) - \frac{C_{P(S)}(p^2)}{M_{\text{PS}}^2(\mu_1, \mu_2)}. \quad (3.13)$$

The effect of the subtractions is illustrated in figure 4. We find that, while in the case of \mathcal{V}_P the contribution of the Goldstone pole is clearly visible, the subtraction is practically irrelevant in the case of \mathcal{V}_S , indicating a strong $\mathcal{O}(a^2)$ suppression of the parity violating coupling. We have also verified that, the alternative procedure [23] for the Goldstone pole subtraction, based on the definition

$$\mathcal{V}_{P(S)}^{\text{sub}}(p^2, \mu_1, \mu_2, \mu_3, \mu_4) = \frac{M_{\text{PS}}^2(\mu_1, \mu_2) \mathcal{V}_{P(S)}(p^2, \mu_1, \mu_2) - M_{\text{PS}}^2(\mu_3, \mu_4) \mathcal{V}_{P(S)}(p^2, \mu_3, \mu_4)}{M_{\text{PS}}^2(\mu_1, \mu_2) - M_{\text{PS}}^2(\mu_3, \mu_4)}. \quad (3.14)$$

leads to consistent results in the chiral limit, within our statistical errors.

The dependence of Z_V , Z_A and Z_T , which are not affected by the Goldstone pole contamination, on the valence quark masses, is shown in figure 5, for all three couplings. For illustration purposes, a specific value of the sea quark mass has been chosen in each case. We see that the valence quark mass dependence of the RCs is rather weak and well consistent with a linear behaviour.

In the tm formulation of lattice QCD with $N_f = 2$ flavours of degenerate sea quarks, the fermionic determinant is a function of the sea quark mass squared. Therefore, we expect that the RCs, which by definition are insensitive to the effects of spontaneous chiral symmetry breaking, will also exhibit the same dependence. This is shown in figure 6. We find that all RCs depend very weakly on the sea quark mass squared. We thus obtain our mass independent RCs by performing the sea quark mass extrapolation to the chiral limit linearly in $a^2 \mu_{\text{sea}}^2$. We also checked that chiral extrapolations based on a first or second order polynomial fit in $a \mu_{\text{sea}}$ lead within errors to practically equivalent results.

3.2.2 Renormalization scale dependence and subtraction of the $\mathcal{O}(g^2 a^2)$ effects

Once the RCs have been extrapolated to the chiral limit, we investigate their dependence on the renormalization scale by evolving, at fixed coupling, the RCs to a reference scale $\mu_0 = 1/a$. This is done using

$$Z_{\Gamma}(a, \mu_0) = C_{\Gamma}(\mu_0, \mu) Z_{\Gamma}(a, \mu), \quad (3.15)$$

where the evolution function C_{Γ} is expressed in terms of the beta function $\beta(\alpha)$ and of the anomalous dimension of the relevant operator $\gamma_{\Gamma}(\alpha)$ by

$$C_{\Gamma}(\mu_0, \mu) = \exp \left[\int_{\alpha(\mu_0)}^{\alpha(\mu)} \frac{\gamma_{\Gamma}(\alpha)}{\beta(\alpha)} d\alpha \right]. \quad (3.16)$$

This function is known, in the RI-MOM scheme, at the N²LO for Z_T [24] and at the N³LO for Z_S and Z_P [25]. Since Z_V , Z_A and the ratio Z_P/Z_S are scale independent, they have vanishing anomalous dimensions; thus for these quantities we have $C_{\Gamma}(\mu_0, \mu) = 1$.




Figure 4. The amputated Green functions $\mathcal{V}_P(p^2, \mu_1, \mu_2)$ and $\mathcal{V}_S(p^2, \mu_1, \mu_2)$, evaluated at a scale $a^2 \hat{p}^2 \simeq 1$, as a function of the pseudoscalar meson mass squared. The subtracted Green functions are defined in eq. (3.13). Dashed and solid lines illustrate the results of the fit to eq. (3.12) (with $C = 0$ for $\mathcal{V}_{P(S)}^{sub}$).




Figure 5. Dependence of Z_V , Z_A and Z_T on the valence quark mass, at $a^2\bar{p}^2 \simeq 1$. The solid lines are linear extrapolations to the (valence) chiral limit.




Figure 6. Dependence of the RCs at $a^2\tilde{p}^2 \simeq 1$ on the sea quark mass after valence chiral extrapolation. The solid lines are linear extrapolations in $(a\mu_{\text{sea}})^2$ to the chiral limit.

Γ	$b_\Gamma^{(1)}$	$b_\Gamma^{(2)}$	$c_\Gamma^{(1)}$	$c_\Gamma^{(2)}$	$c_\Gamma^{(3)}$
V	-0.48369852(8)	0	1.5240798(1)	-1/3	-125/288
A	3.57961385(3)	0	0.6999177(1)	-1/3	-125/288
S	0.58345905(5)	-3	2.3547298(2)	-1/4	1/2
P	8.7100837(1)	-3	0.70640549(6)	-1/4	1/2
T	0.51501972(6)	1	0.9724758(2)	-13/36	-161/216

Table 3. Values of the coefficients $b_\Gamma^{(i)}$ and $c_\Gamma^{(i)}$ entering the 1-loop expression (3.17) of the amputated projected Green functions $\mathcal{V}_\Gamma(p)$. Results are presented for the case of the tree-level Symanzik improved gluon action in the Landau gauge and the tm fermionic action at maximal twist.

In the non-perturbative calculation, the RCs evolved to the reference scale $\mu_0 = 1/a$ maintain a dependence on the renormalization scale p^2 at which they have been initially computed. We will keep track of this dependence, which (at large enough p^2) is mostly due to discretization effects, by denoting these RCs as $Z_\Gamma(1/a; a^2 p^2)$, where the first variable indicates the renormalization scale, $\mu_0 = 1/a$, and the second the dependence on the initial momentum.

In figure 7 we show the results for all $Z_\Gamma(1/a; a^2 p^2)$ at $\beta = 3.9$ as a function of $a^2 \tilde{p}^2$ (empty symbols). The residual $a^2 p^2$ -dependence which is observed in these results in the large momentum region is practically linear, suggesting that leading discretization effects are $\mathcal{O}(a^2 p^2)$. As illustrated in the plots, this dependence is particularly pronounced in the case of the pseudoscalar RC Z_P .

In order to reduce the size of discretization errors, we analytically subtract from the quark propagator and the amputated vertex functions the $\mathcal{O}(g^2 a^2)$ contributions, recently computed in lattice perturbation theory [8]. Up to 1-loop and up to $\mathcal{O}(a^2)$, the amputated projected Green functions $\mathcal{V}_\Gamma(p)$, defined in eq. (3.4), have the simple and general expression

$$\mathcal{V}_\Gamma(p)^{\text{pert.}} = 1 + \frac{g^2}{12\pi^2} \left\{ b_\Gamma^{(1)} + b_\Gamma^{(2)} \ln(a^2 p^2) + a^2 \left[p^2 (c_\Gamma^{(1)} + c_\Gamma^{(2)} \ln(a^2 p^2)) + c_\Gamma^{(3)} \frac{\sum_\rho p_\rho^4}{p^2} \right] \right\} + \mathcal{O}(a^4 g^2, g^4). \quad (3.17)$$

The values of the coefficients $b_\Gamma^{(i)}$ and $c_\Gamma^{(i)}$ are collected in table 3 for the lattice action used in the present study, namely the tree-level Symanzik improved gluon action in the Landau gauge and the tm fermionic action at maximal twist.⁷

A result similar to eq. (3.17) also holds for the form factor $\Sigma_1(p)$ of the inverse quark propagator, from which the quark field RC Z_q is evaluated:

$$\Sigma_1(p)^{\text{pert.}} = 1 + \frac{g^2}{12\pi^2} \left\{ b_q^{(1)} + b_q^{(2)} \ln(a^2 p^2) + a^2 \left[p^2 (c_q^{(1)} + c_q^{(2)} \ln(a^2 p^2)) + c_q^{(3)} \frac{\sum_\rho p_\rho^4}{p^2} \right] \right\} + \mathcal{O}(a^4 g^2, g^4). \quad (3.18)$$

⁷The same results are also valid for the standard Wilson fermionic action, except for the exchange of the values of the vector (V) and axial (A) coefficients.

Our definition of this form factor on the lattice, which is equivalent to eq. (3.5) up to terms of $\mathcal{O}(a^2)$, is

$$\Sigma_1(p) \equiv \frac{-i}{12N(p)} \text{Tr} \left[\sum'_\rho (\gamma_\rho S(p)^{-1}) / \tilde{p}_\rho \right], \quad (3.19)$$

where the sum \sum'_ρ only runs over the Lorentz indices for which p_ρ is different from zero and $N(p) = \sum'_\rho 1$. With this definition, the coefficients of eq. (3.18) take the values

$$\begin{aligned} b_q^{(1)} &= -13.02327272(7); & b_q^{(2)} &= 0; & c_q^{(3)} &= \frac{7}{240}; \\ c_q^{(1)} &= 1.14716212(5) + \frac{2.07733285(2)}{N(p)}; & c_q^{(2)} &= -\frac{73}{360} - \frac{157/180}{N(p)}. \end{aligned} \quad (3.20)$$

Using eqs. (3.17) and (3.18), we then define the *corrected* amputated Green functions and the *corrected* form factor Σ_1 as

$$\begin{aligned} \mathcal{V}_\Gamma(p)^{\text{corr.}} &= \mathcal{V}_\Gamma(p) - \frac{g^2}{12\pi^2} a^2 \left[\tilde{p}^2 (c_\Gamma^{(1)} + c_\Gamma^{(2)} \ln(a^2 \tilde{p}^2)) + c_\Gamma^{(3)} \frac{\sum_\rho \tilde{p}_\rho^4}{\tilde{p}^2} \right], \\ \Sigma_1(p)^{\text{corr.}} &= \Sigma_1(p) - \frac{g^2}{12\pi^2} a^2 \left[\tilde{p}^2 (c_q^{(1)} + c_q^{(2)} \ln(a^2 \tilde{p}^2)) + c_q^{(3)} \frac{\sum_\rho \tilde{p}_\rho^4}{\tilde{p}^2} \right], \end{aligned} \quad (3.21)$$

which are free of $\mathcal{O}(g^2 a^2)$ effects. Note that the $\mathcal{O}(a^2)$ terms depend not only on the magnitude, $\sum_\rho p_\rho^2$, but also on the direction of the momentum, p_ρ , as manifested by the presence of $\sum_\rho p_\rho^4$. As a consequence, different renormalization prescriptions, involving different directions of the renormalization scale p_ρ , treat lattice artifacts differently. In the numerical evaluation of the perturbative correction in eq. (3.21), we replaced g^2 with a simple boosted coupling [26], defined as $\tilde{g}^2 = g_0^2 / \langle P \rangle$, where the average plaquette $\langle P \rangle$ is computed non-perturbatively.

The effect of the subtraction (3.21) is also illustrated in figure 7, which shows that the $a^2 p^2$ -dependence of the RCs is significantly reduced by the perturbative correction. By fitting the RCs as

$$Z_\Gamma(1/a; a^2 p^2) = Z_\Gamma(1/a) + \lambda_\Gamma a^2 \tilde{p}^2 \quad (3.22)$$

in the large momentum region $a^2 \tilde{p}^2 \gtrsim 1$ (with λ_Γ just constrained to depend smoothly on g^2 — see eq. (3.24)), we find that the slopes λ_Γ are reduced, after the perturbative subtraction, at the level of 10^{-2} or smaller for Z_V , Z_A , Z_T and Z_q . For Z_S we find that the slope, which is also about 10^{-2} before implementing the correction, increases slightly after the subtraction. For Z_P , on the other hand, the slope is rather large before the subtraction; i.e. $\lambda_P \simeq 5 \cdot 10^{-2}$. As shown in figure 7, the effect of the subtraction is beneficial also in this case, but inadequate for correcting the bulk of the observed $a^2 p^2$ -dependence. This is not completely unexpected: when discretization effects are large, the subtraction of only the leading $\mathcal{O}(g^2 a^2)$ terms may be not sufficient to reduce them to a negligible level. Similar results, with approximately equal values of the slopes λ_Γ , are obtained at all three values of the lattice spacing.

A useful way to investigate the size of lattice artifacts, which are left after the perturbative subtraction, consists in comparing the renormalization scale dependence of




Figure 7. The RCs $Z_\Gamma(1/a; a^2 p^2)$ at $\beta = 3.9$, evaluated at the reference scale $\mu_0 = 1/a$, plotted against the initial renormalization scale $a^2 p^2$. Filled squares (empty circles) are results obtained with (without) the subtraction of the $\mathcal{O}(g^2 a^2)$ discretization effects computed in perturbation theory, see eq. (3.21). The solid lines are linear fits to the data.

the RCs as observed at different lattice spacings. Specifically, at each value of the lattice spacing we fix two common values of the renormalization scale, μ and μ' , and compute the step scaling functions

$$\Sigma_\Gamma(\mu, \mu'; a) = \frac{Z_\Gamma(a, \mu)}{Z_\Gamma(a, \mu')}. \quad (3.23)$$

If not for discretization effects, the step scaling functions should be independent of the




Figure 8. Values of the step scaling functions $\Sigma_\Gamma(\mu, \mu'; a)$ evaluated for $(\mu^2, \mu'^2) = (8.5, 6.0) \text{ GeV}^2$ (red circles) and $(\mu^2, \mu'^2) = (8.5, 7.25) \text{ GeV}^2$ (blue squares) at the three values of the lattice coupling. The solid lines and the empty markers show the results of a linear extrapolation to the continuum limit. For comparison, the corresponding perturbative estimates of the evolution functions $C_\Gamma(\mu, \mu')$ are also shown in the plots (black circles and squares).

cut-off and equal to the evolution functions $C_\Gamma(\mu, \mu')$ of eq. (3.15).

In figure 8 we show the results obtained for the step scaling functions of the quark bilinear operators and of the quark field RCs, as a function of $(a/r_0)^2$, for two common pairs renormalization scales, $(\mu^2, \mu'^2) = (8.5, 6.0) \text{ GeV}^2$ and $(\mu^2, \mu'^2) = (8.5, 7.25) \text{ GeV}^2$, which both lie in the interval of momenta accessible at all values of the lattice spacing. We see from these plots that the dependence of the step scaling functions on the lattice cut-off is tiny in most of the cases, being clearly visible only in the case of Σ_P . Moreover, for the scale independent RCs Z_V and Z_A , a linear extrapolation of the results to the continuum limit leads to values which are perfectly consistent with unity, within the statistical errors. For the other RCs, the continuum limit of the step scaling functions leads to non-perturbative results which are in good agreement with the perturbative determinations of the evolution functions $C_\Gamma(\mu, \mu')$, computed at the N²LO for Z_T and at the N³LO for Z_S , Z_P and Z_q . These results provide evidence that higher order perturbative contributions to the anomalous dimensions of the various operator are not relevant for describing the renormalization scale dependence of the RCs in the region of momenta explored in the present calculation.

In order to account for the residual discretization effects in the calculation of the RCs, we follow two different approaches:

- *Extrapolation method (M1):* after subtraction of the $\mathcal{O}(g^2 a^2)$ terms according to eq. (3.21), we extrapolate the RCs linearly to $a^2 p^2 \rightarrow 0$. Specifically, we fit $Z_\Gamma(1/a; a^2 p^2)$ with eq. (3.22) in the large momentum region, $1.0 \lesssim a^2 \tilde{p}^2 \lesssim 2.2$. The slopes λ_Γ exhibit only a very mild dependence on the coupling. We parameterize this dependence by performing a simultaneous extrapolation at the 3 values of the

coupling and writing the slopes as

$$\lambda_{\Gamma}(g^2) = \lambda_{\Gamma}(g_0^2) + \lambda'_{\Gamma}(g_0^2) \cdot (g^2 - g_0^2), \quad (3.24)$$

where g_0 is the coupling corresponding to the intermediate value $\beta = 3.9$. The intercept of the extrapolation as determined from the fit, and indicated with $Z_{\Gamma}(1/a)$ in eq. (3.22), provides our final estimate of the RC with the extrapolation method. The fit is illustrated, for all RCs, in figure 9.

- *p^2 -window method (M2)*: in this approach we do not attempt any additional subtraction of discretization effects, besides the perturbative one. The final estimates of RCs are obtained by taking directly the results of $Z_{\Gamma}(1/a; a^2 p^2)$ at a fixed value of p^2 (in physical units). In practice, this is done by fitting the RCs to a constant in the small momentum interval $\tilde{p}^2 \in [8.0, 9.5]$ GeV². The idea behind this approach is that, once RCs are combined with bare quantities, so as to construct the physical observables of interest, the residual $\mathcal{O}(a^2)$ effects, which are present in both RCs and bare quantities, will be extrapolated away in the continuum limit.

The two methods are compared below, in a specific example. This consists in the scaling of the pseudoscalar meson mass, composed by two degenerate quarks of fixed (renormalized) mass. As expected, the two methods give different results at fixed lattice cut-off, but the difference disappears in the continuum limit.

3.3 Analysis of systematic errors

Before presenting our final results for the RCs, we list and discuss in some detail the main systematic uncertainties.

- *Finite size effects*: the RCs, determined in the RI-MOM scheme at large momenta, are short distance quantities and, as such, should not be affected by significant finite size effects. In order to verify this expectation, an additional RI-MOM calculation at $\beta = 3.90$ has been performed, on a $32^3 \times 64$ lattice. A comparison of these results with those obtained on the smaller $24^3 \times 48$ lattice is illustrated in figure 10. We find that, for the momentum range in which results on both volumes are available, the differences are smaller than statistical errors.
- *Subtraction of the Goldstone pole and chiral extrapolations*: the chiral extrapolation of the RCs is rather delicate for Z_P (and to a lesser extent for Z_S), due to the presence of the Goldstone pole which has to be subtracted. The uncertainty introduced by the subtraction of the Goldstone pole contribution to Z_S and Z_P can be estimated by comparing the results obtained following two different subtraction approaches: the fit of eq. (3.12) or, alternatively, the procedure based on eq. (3.14). We find that differences are always negligible for Z_S , whereas for Z_P they are at the level of our statistical errors.

As illustrated in figures 5 and 6, the chiral extrapolations in the valence quark mass for the other RCs and in the sea quark mass for all RCs are quite smooth.




Figure 9. The RCs $Z_{\Gamma}(1/a; a^2 p^2)$, evolved to the reference scale $\mu_0 = 1/a$, as a function of the initial renormalization scale $a^2 p^2$. The solid lines are linear fits to the data.

They are also well consistent with the expected leading linear and purely quadratic dependence on the valence and sea quark masses respectively. Therefore, we estimate that the uncertainties associated with these chiral extrapolations are negligible with respect to the statistical errors.




Figure 10. Comparison of the RCs obtained, at $\beta = 3.90$, on $24^3 \times 48$ and $32^3 \times 64$ lattices.

- *Discretization effects:* the analysis of these effects has been presented in section 3.2.2. The leading $\mathcal{O}(g^2 a^2)$ discretization errors have been accounted for, using the analytical results of ref. [8]. With the method denoted as M1 in section 3.2.2, after the perturbative subtraction, residual $\mathcal{O}(a^2 p^2)$ effects are extrapolated away, with the Ansatz (3.22). These contributions are not subtracted, instead, in the




Figure 11. Scaling of the pseudoscalar meson mass squared, computed at a fixed value of the quark mass. The M1 and M2 determinations for $Z_\mu = Z_P^{-1}$ lead to compatible results in the continuum limit.

approach denoted as M2. Discretization effects affecting both RCs and the bare matrix elements will be corrected for, by extrapolating the physical (renormalized) quantity to the continuum limit.

As an example of the results obtained by implementing the two approaches, we show in figure 11 the continuum extrapolation of the pseudoscalar mass squared, computed at a fixed value of the renormalized quark mass μ_R . The renormalization of the latter has been achieved using the estimates M1 and M2 for Z_P given in tables 4 and 5.⁸ Although the two Z_P determinations differ by visible discretization effects at finite lattice spacing, particularly at the two coarsest lattice, the continuum limit results of the pseudoscalar mass squared are consistent within the two determinations. This is evidence that the discretization errors as well as other possible systematic effects in the analysis of the RI-MOM RCs are well under control, at least within our current statistical errors (say at the level of one standard deviation and possibly even less).

3.4 Final results and comparison with perturbation theory

The final results for the bilinear quark operators and the quark field RCs, obtained with the RI-MOM method, are collected in tables 4 and 5 (labelled as RI-MOM (M1) and RI-MOM (M2)). These results are compared with those obtained in section 2 for the scale independent Z_V , Z_A and Z_P/Z_S (Alt. methods). In addition, we give in tables 4 and 5 the predictions of one loop boosted perturbation theory (BPT) [7, 8], obtained with two definitions of the boosted coupling: the first is $\tilde{g}^2 = g_0^2/\langle P \rangle$, also used in the previous section, and the second is based on the one-loop tadpole-improved expansion of $\log\langle P \rangle$ [26].

From tables 4 and 5, we see that the largest deviations between the central values of the RI-MOM (M1) determinations and the alternative determinations of section 2 are at the

⁸Note that in tm lattice QCD we have $Z_\mu = Z_P^{-1}$.

β	Method	Z_V	Z_A	Z_P/Z_S
3.80	RI-MOM (M1)	0.604(07)	0.746(11)	0.580(17)
	RI-MOM (M2)	0.623(05)	0.727(07)	0.653(07)
	Alt. methods	0.5816(02)	0.747(22)	0.498(22)
	BPT	0.61-0.70	0.71-0.77	0.72-0.80
3.90	RI-MOM (M1)	0.624(04)	0.746(06)	0.626(13)
	RI-MOM (M2)	0.634(03)	0.730(03)	0.669(08)
	Alt. methods	0.6103(03)	0.743(18)	0.579(19)
	BPT	0.63-0.71	0.72-0.78	0.74-0.82
4.05	RI-MOM (M1)	0.659(04)	0.772(06)	0.686(12)
	RI-MOM (M2)	0.662(03)	0.758(04)	0.710(07)
	Alt. methods	0.6451(03)	0.746(18)	0.661(21)
	BPT	0.65-0.73	0.74-0.80	0.76-0.83

Table 4. Values of Z_V , Z_A and Z_P/Z_S , obtained with the RI-MOM methods M1 and M2 (see text for details), as well as with the methods described in section 2 (labelled as “Alt. method”). The predictions of one loop boosted perturbation theory (BPT) are also shown for comparison.

level of 3-4% or smaller for Z_A and Z_V . These differences are not always accounted for by the errors quoted for each result. When statistically significant, these differences provide an estimate of the systematic uncertainties affecting the calculations. As discussed in the previous section, we expect these uncertainties to be dominated by $O(a^2)$ discretization effects. Nevertheless, we do not include these differences in the final errors. The reason is that such estimates of discretization effects, though meaningful at finite lattice spacing, are of no practical significance when the continuum limit of a physical quantity is eventually computed.

The comparison of the Z_P/Z_S results, obtained with the RI-MOM and the alternative method, deserves a further comment. The difference between the two results is about 15% for the coarsest lattice and 4% for the finest one. The rather big difference noticed in the former case may be attributed not only to discretization effects, but also to an imprecise tuning of the twist angle to maximal twist at this coupling. This is expected to influence substantially the OS pseudoscalar density.

As shown in tables 4 and 5, the non-perturbative results also lie in the range of one-loop BPT estimates, with the exception of Z_S and, more notably, of Z_P . We emphasise that the accuracy of Z_P is crucial in the determination of quark masses, since the bare quark mass, computed with maximally twisted fermions, renormalizes with of Z_P^{-1} ; see for example ref. [27].

β	Method	$Z_S^{\text{RI}}(1/a)$	$Z_P^{\text{RI}}(1/a)$	$Z_T^{\text{RI}}(1/a)$	$Z_q^{\text{RI}}(1/a)$
3.80	RI-MOM (M1)	0.603(15)	0.338(10)	0.743(09)	0.755(07)
	RI-MOM (M2)	0.668(06)	0.437(04)	0.719(06)	0.755(05)
	BPT	0.68-0.75	0.49-0.60	0.68-0.75	0.69-0.76
3.90	RI-MOM (M1)	0.615(09)	0.377(06)	0.743(05)	0.758(04)
	RI-MOM (M2)	0.669(05)	0.447(05)	0.724(03)	0.755(03)
	BPT	0.70-0.76	0.52-0.62	0.70-0.76	0.71-0.77
4.05	RI-MOM (M1)	0.645(06)	0.440(06)	0.763(06)	0.782(05)
	RI-MOM (M2)	0.678(04)	0.480(04)	0.752(04)	0.778(03)
	BPT	0.72-0.78	0.55-0.65	0.72-0.78	0.73-0.79

Table 5. Values of Z_S, Z_P and Z_T , obtained with the RI-MOM methods M1 and M2 (see text for details). The predictions of one loop boosted perturbation theory (BPT) are also shown for comparison.

β	Z_V	Z_A	$Z_S^{\overline{\text{MS}}}(2 \text{ GeV})$	$Z_P^{\overline{\text{MS}}}(2 \text{ GeV})$	$Z_T^{\overline{\text{MS}}}(2 \text{ GeV})$	$Z_q^{\overline{\text{MS}}}(2 \text{ GeV})$
3.80	0.604(07)	0.746(11)	0.734(18)	0.411(12)	0.733(09)	0.745(07)
3.90	0.624(04)	0.746(06)	0.713(10)	0.437(07)	0.743(05)	0.751(04)
4.05	0.659(03)	0.772(06)	0.699(06)	0.477(06)	0.777(06)	0.780(05)

Table 6. Final results for the RCs of bilinear quark operators and the quark field, obtained with the RI-MOM method. The scheme dependent RCs are given in the $\overline{\text{MS}}$ scheme at scale $\mu=2 \text{ GeV}$.

In most phenomenological applications, the renormalization scheme of choice is $\overline{\text{MS}}$ and the preferred reference scale is 2 GeV . For this reason, we provide in table 6 the values of the scheme dependent RCs Z_S, Z_P, Z_T and Z_q in the $\overline{\text{MS}}$ scheme at the scale of 2 GeV . The conversion from the RI-MOM scheme at $\mu = 1/a$ to $\overline{\text{MS}}$ at $\mu = 2 \text{ GeV}$ has been performed using renormalization group-improved perturbation theory at the N^2LO for Z_T and at the N^3LO for Z_S, Z_P and Z_q . For completeness, we also report in the table the results for the scale independent Z_V and Z_A . Note, however, that the Ward identity determination of Z_V , given in tables 4 and 5, has a much better statistical accuracy.

It should be remarked that in table 6 we only quote the results obtained from RI-MOM renormalization conditions with the method M1, since they are in general affected by smaller discretization effects and should hence be preferably used in computations at fixed gauge coupling without continuum extrapolation.⁹ The case of figure 11 is an exception

⁹Anyway, the values of the RCs in the $\overline{\text{MS}}$ scheme at the scale of 2 GeV obtained with the method M2

in this sense, because there an accidental cancellation (enhancement) of the cutoff effects coming from the bare pseudoscalar meson mass M_{ps} and from Z_P computed with the method M2 (M1) leads to a similar scaling behaviour of the two sets of data corresponding to the M2 and M1 determinations of Z_P . Moreover, with respect to the method M1, the method M2 tends to give results that are affected by smaller statistical uncertainties, since the latter procedure does not involve the extrapolation to $a^2p^2 = 0$ of eq. (3.22).

4 Conclusions

In this paper we have presented a non-perturbative determination of the RCs of bilinear quark operators for the tree-level Symanzik improved gauge action and the $N_f = 2$ twisted mass fermion action at maximal twist, which guarantees automatic $\mathcal{O}(a)$ -improvement. The values of these constants do not depend on whether tm, OS or Wilson fermions are used. The RCs of the five bilinear quark operators and the quark field have been determined with the RI-MOM method. We have also obtained an independent estimate of Z_V , based on a Ward identity, and have introduced a new method for Z_A and Z_S/Z_P , based on a combined use of standard tm and OS fermions. The main results for the RCs are collected in tables 4, 5 and 6, where they are also compared with the predictions of one-loop perturbation theory. We find that the differences between perturbative and non-perturbative determinations are large in the cases of Z_S and Z_P . This result is of particular relevance to the lattice determinations of quark masses and the chiral condensate.

A $\mathcal{O}(a)$ -improvement of RI-MOM renormalization constants

In this appendix we prove that:

1. at maximal twist the RI-MOM “form factors” from which RCs are extracted [1] are automatically $\mathcal{O}(a)$ improved at all p^2 values;
2. at generic values of the twist angle, form factor improvement is guaranteed only at “large” p^2 values; i.e. where spontaneous chiral breaking effects can be neglected.

In both cases the resulting renormalization constants will be improved. In the first case this is obvious because they are extracted from ratios of automatically improved form factors. In the second case the desired conclusion follows from the observation that RCs are short-distance quantities (insensitive to infrared effects), evaluated from large p^2 correlators. In the ensuing sections we give a detailed proof of these two statements.

A.1 Improvement at maximal twist

In the RI-MOM scheme the RC Z_Γ^{fh} of the bilinear operator

$$O_\Gamma^{\text{fh}} = \bar{q}^f \Gamma q^h, \tag{A.1}$$

can easily be obtained for each $a = a(\beta)$ from their M1-method counterparts and the RI-MOM results in tables 4 and 5 with the help of the relation $Z_\Gamma^{\text{MS}}(2 \text{ GeV})|_{\text{M2}}/Z_\Gamma^{\text{RI}}(1/a)|_{\text{M2}} = Z_\Gamma^{\text{MS}}(2 \text{ GeV})|_{\text{M1}}/Z_\Gamma^{\text{RI}}(1/a)|_{\text{M1}}$.

is defined by the condition

$$\lim_{\mu_{\text{val}}^{f,h}, \mu_{\text{sea}} \rightarrow 0} Z_q^{-1} Z_{\Gamma}^{\text{fh}} \text{Tr}[\Lambda_{\Gamma}^{\text{fh}}(p) P_{\Gamma}] \Big|_{p^2=\mu^2} = 1, \quad (\text{A.2})$$

where $f, h = 1, 2$ are flavour indices, and P_{Γ} is the projector onto the Dirac structure of the bilinear. The other quantities entering eq. (A.2) are defined by

$$\lim_{\mu_{\text{val}}^f, \mu_{\text{sea}} \rightarrow 0} Z_q \frac{-i}{12} \frac{\text{Tr}[\not{p} S_q^f(p)^{-1}]}{p^2} \Big|_{p^2=\mu^2} = 1 \quad (\text{A.3})$$

with

$$\Lambda_{\Gamma}^{\text{fh}}(p) = S_q^f(p)^{-1} G_{\Gamma}^{\text{fh}}(p) S_q^h(p)^{-1}, \quad (\text{A.4})$$

$$S_q^f(p) = a^4 \sum_x e^{-ip \cdot x} \langle q^f(x) \bar{q}^f(0) \rangle, \quad (\text{A.5})$$

$$G_{\Gamma}^{\text{fh}}(p) = a^8 \sum_{x,y} e^{-ip \cdot (x-y)} \langle q^f(x) O_{\Gamma}^{\text{fh}}(0) \bar{q}^h(y) \rangle. \quad (\text{A.6})$$

Two useful observations are: (i) the “form factors” $\mathcal{V}_q(p) = \text{Tr}[\not{p} S^f(p)^{-1}]$ and $\mathcal{V}_{\Gamma}^{\text{fh}}(p) = \text{Tr}[\Lambda_{\Gamma}^{\text{fh}}(p) P_{\Gamma}]$ are H(4) invariant by construction. In the continuum limit they approach quantities which are invariant under both O(4) and parity. (ii) Z_q carries no flavour index f since, owing to its definition (A.3), it depends quadratically on r_q .

The proof that our estimators of the RI-MOM RCs are O(a)-improved at all quark masses and external momenta in maximally twisted QCD follows closely the argument of ref. [14] (subsequently refined in ref. [28, 29]) for automatic improvement of correlation functions. It is based on the exact invariance of the maximally twisted QCD action under the transformation $\mathcal{P} \times \mathcal{D}_d \times (\mu_q \rightarrow -\mu_q)$, where μ_q is the twisted quark mass. For completeness we recall the definition of the discrete transformations \mathcal{P} (continuum parity) and \mathcal{D}_d in the physical quark basis

$$\mathcal{P} : \begin{cases} U_0(x) \rightarrow U_0(x_{\mathcal{P}}), & U_k(x) \rightarrow U_k^{\dagger}(x_{\mathcal{P}} - a\hat{k}), & k = 1, 2, 3 \\ q^f(x) \rightarrow \gamma_0 q^f(x_{\mathcal{P}}) \\ \bar{q}^f(x) \rightarrow \bar{q}^f(x_{\mathcal{P}}) \gamma_0 \end{cases} \quad (\text{A.7})$$

$$\mathcal{D}_d : \begin{cases} U_{\mu}(x) \rightarrow U_{\mu}^{\dagger}(-x - a\hat{\mu}) \\ q^f(x) \rightarrow e^{3i\pi/2} q^f(-x) \\ \bar{q}^f(x) \rightarrow e^{3i\pi/2} \bar{q}^f(-x) \end{cases} \quad (\text{A.8})$$

where $x_{\mathcal{P}} = (-\mathbf{x}, x_0)$ and $\hat{\mu}$ is the unit vector in the μ -direction. Actually the proof is also valid in the more general “mixed action” setting, where Osterwalder-Seiler fermions are used as valence quarks. This is because the corresponding action is similarly invariant under $\mathcal{P} \times \mathcal{D}_d \times (M_q \rightarrow -M_q)$, where M_q denotes the set of all quark (sea and valence) twisted mass parameters.

A.1.1 The proof

The proof can be given in three steps:

- Due to $H(4)$ symmetry, the lattice form factors $\mathcal{V}_q(p)$ and $\mathcal{V}_\Gamma^{\text{fh}}(p)$ have no terms of the form $p_0^k + p_1^k + p_2^k + p_3^k$ and $(p_0 p_1 p_2 p_3)^k$, with odd k . Thus they are invariant under parity transformation; i.e. they satisfy the relations

$$\mathcal{V}_q(p) = \mathcal{V}_q(p_{\mathcal{P}}), \quad \mathcal{V}_\Gamma^{\text{fh}}(p) = \mathcal{V}_\Gamma^{\text{fh}}(p_{\mathcal{P}}), \quad p_{\mathcal{P}} = (-\mathbf{p}, p_0). \quad (\text{A.9})$$

- As has been proved in [28, 29], the invariance of maximally twisted QCD under $\mathcal{P} \times \mathcal{D}_d \times (\mu_q \rightarrow -\mu_q)$ implies that $O(a^j)$ cut-off artefacts of lattice correlators arise, in their Symanzik expansion, from the insertion of Symanzik operators of parity $(-1)^j$.
- Subsequently, $O(a^j)$ terms with odd j must then be absent in the Symanzik expansion of parity-even form factors, like $\mathcal{V}_q(p)$ and $\mathcal{V}_\Gamma^{\text{fh}}(p)$.

The Symanzik expansion of the form factors above is straightforwardly obtained from that of the correlators in terms of which they are defined.

A.2 Improvement at a generic value of the twist angle

Since RCs are UV quantities, they should be independent of the QCD infrared structure and properties; in particular they should be unaffected by spontaneous chiral symmetry breaking. Thus, if we compute RCs at zero quark mass and at high momentum scale p^2 such that chiral breaking effects are negligible, we can fully exploit the \mathcal{R}_5 -symmetry of the continuum theory. This will be useful because in turn the lattice action is invariant under $\mathcal{R}_5 \times \mathcal{D}_d$, where

$$\mathcal{R}_5 = \prod_f \mathcal{R}_5^f, \quad \mathcal{R}_5^f : \begin{cases} q^f \rightarrow \gamma_5 q^f \\ \bar{q}^f \rightarrow -\bar{q}^f \gamma_5 \end{cases} \quad (\text{A.10})$$

is a transformation belonging to the $SU(N_f)_L \times SU(N_f)_R$ chiral group in the unitary case (or the graded $SU(N_f^{\text{sea}} + N_f^{\text{val}}|N_f^{\text{val}})_L \times SU(N_f^{\text{sea}} + N_f^{\text{val}}|N_f^{\text{val}})_R$ chiral group in the mixed action case). In eq. (A.10) the product is extended over all flavour species f .

A.2.1 The proof

The proof of the $O(a)$ improvement of RCs in lattice QCD at generic values of the twist angle, e.g. of the bare twist angle $\omega_0 = \arctan(\mu_q/(m_0 - M_{\text{cr}}))$, runs on lines similar to those of the previous section, with \mathcal{R}_5 replacing \mathcal{P} . One can, in fact, argue as follows.

- RCs constants can be extracted from the large- p^2 behaviour of the $\mathcal{V}_q(p)$ and $\mathcal{V}_\Gamma^{\text{fh}}(p)$ form factors in the chiral limit. In this regime chiral breaking effects can be ignored and $\mathcal{V}_q(p)$ and $\mathcal{V}_\Gamma^{\text{fh}}(p)$ are invariant under \mathcal{R}_5 .
- The invariance of the theory under $\mathcal{R}_5 \times \mathcal{D}_d$ implies that the \mathcal{R}_5 -parity of an operator equals the parity of its naive dimension. Thus, $O(a^j)$ lattice artefacts of correlators or form factors arise in the Symanzik expansion from the insertion of operators of \mathcal{R}_5 -parity equal to $(-1)^j$.

- Therefore $O(a^j)$ terms with odd j cannot be present in the Symanzik expansion of \mathcal{R}_5 -even form factors, like $\mathcal{V}_q(p)$ and $\mathcal{V}_F^{\text{th}}(p)$. In fact, the continuum target theory is invariant under the chiral group (including \mathcal{R}_5), while $O(a^j)$ terms would be odd under \mathcal{R}_5 .

A.3 A final comment

The $\mathcal{O}(a)$ improvement of RI-MOM RCs at any value (including zero) of the twist angle ω_0 hardly comes as a surprise because the RC-estimators computed at generic ω_0 become ω_0 -independent in the chiral limit, provided they are analytic in the complex mass parameter $(m_0 - M_{\text{cr}}) + i\mu_q$, which is certainly the case at sufficiently large p^2 . As a consequence, a unique set of RI-MOM RCs, all free from $\mathcal{O}(a^j)$ (j odd) artifacts, exist for lattice QCD with Wilson quarks and a given pure gauge lattice action. These RCs can in principle be extracted from simulations at arbitrary (and not necessarily all equal) values of the twist angle. Computing the RI-MOM RCs at maximal twist ($|\omega_0| = \pi/2$), as we do in the present study, is just a simple and technically convenient choice. In our case this choice was a quite natural one, as we could then evaluate the necessary correlators on the $N_f = 2$ gauge ensembles which were produced for studying large volume physics.

Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] G. Martinelli, C. Pittori, C.T. Sachrajda, M. Testa and A. Vladikas, *A general method for nonperturbative renormalization of lattice operators*, *Nucl. Phys. B* **445** (1995) 81 [[hep-lat/9411010](#)] [[SPIRES](#)].
- [2] P. Dimopoulos et al., *Renormalisation of quark bilinears with $N_f = 2$ Wilson fermions and tree-level improved gauge action*, *PoS(LATTICE 2007)241* [[arXiv:0710.0975](#)] [[SPIRES](#)].
- [3] EUROPEAN TWISTED MASS collaboration, P. Boucaud et al., *Dynamical twisted mass fermions with light quarks*, *Phys. Lett. B* **650** (2007) 304 [[hep-lat/0701012](#)] [[SPIRES](#)].
- [4] EUROPEAN TWISTED MASS collaboration, P. Boucaud et al., *Dynamical twisted mass fermions with light quarks: simulation and analysis details*, *Comput. Phys. Commun.* **179** (2008) 695 [[arXiv:0803.0224](#)] [[SPIRES](#)].
- [5] EUROPEAN TWISTED MASS collaboration, C. Urbach, *Lattice QCD with two light Wilson quarks and maximally twisted mass*, *PoS(LATTICE 2007)022* [[arXiv:0710.1517](#)] [[SPIRES](#)].
- [6] EUROPEAN TWISTED MASS collaboration, R. Baron et al., *Light meson physics from maximally twisted mass lattice QCD*, [arXiv:0911.5061](#) [[SPIRES](#)].
- [7] S. Aoki, K.-I. Nagai, Y. Taniguchi and A. Ukawa, *Perturbative renormalization factors of bilinear quark operators for improved gluon and quark actions in lattice QCD*, *Phys. Rev. D* **58** (1998) 074505 [[hep-lat/9802034](#)] [[SPIRES](#)].
- [8] M. Constantinou, V. Lubicz, H. Panagopoulos and F. Stylianou, *$O(a^2)$ corrections to the one-loop propagator and bilinears of clover fermions with Symanzik improved gluons*, *JHEP* **10** (2009) 064 [[arXiv:0907.0381](#)] [[SPIRES](#)].

- [9] M. Gockeler et al., *Perturbative and nonperturbative renormalization in lattice QCD*, [arXiv:1003.5756](#) [SPIRES].
- [10] XLF collaboration, K. Jansen, M. Papinutto, A. Shindler, C. Urbach and I. Wetzorke, *Quenched scaling of Wilson twisted mass fermions*, *JHEP* **09** (2005) 071 [[hep-lat/0507010](#)] [SPIRES].
- [11] D. Becirevic et al., *Exploring twisted mass lattice QCD with the clover term*, *Phys. Rev.* **D 74** (2006) 034501 [[hep-lat/0605006](#)] [SPIRES].
- [12] K. Osterwalder and E. Seiler, *Gauge field theories on the lattice*, *Annals Phys.* **110** (1978) 440 [SPIRES].
- [13] R. Frezzotti and G.C. Rossi, *Chirally improving Wilson fermions. II: four-quark operators*, *JHEP* **10** (2004) 070 [[hep-lat/0407002](#)] [SPIRES].
- [14] R. Frezzotti and G.C. Rossi, *Chirally improving Wilson fermions. I: $O(a)$ improvement*, *JHEP* **08** (2004) 007 [[hep-lat/0306014](#)] [SPIRES].
- [15] ALPHA collaboration, R. Frezzotti, P.A. Grassi, S. Sint and P. Weisz, *Lattice QCD with a chirally twisted mass term*, *JHEP* **08** (2001) 058 [[hep-lat/0101001](#)] [SPIRES].
- [16] P. Dimopoulos, H. Simma and A. Vladikas, *Quenched B_K -parameter from Osterwalder-Seiler tmQCD quarks and mass-splitting discretization effects*, *JHEP* **07** (2009) 007 [[arXiv:0902.1074](#)] [SPIRES].
- [17] UKQCD collaboration, M. Foster and C. Michael, *Quark mass dependence of hadron masses from lattice QCD*, *Phys. Rev.* **D 59** (1999) 074503 [[hep-lat/9810021](#)] [SPIRES].
- [18] R. Frezzotti, V. Lubicz and S. Simula, *Electromagnetic form factor of the pion from twisted-mass lattice QCD at $N_f = 2$* , *Phys. Rev.* **D 79** (2009) 074506 [[arXiv:0812.4042](#)] [SPIRES].
- [19] D. Becirevic, V. Gimenez, V. Lubicz, G. Martinelli, M. Papinutto and J. Reyes, *Renormalization constants of quark operators for the non-perturbatively improved Wilson action*, *JHEP* **08** (2004) 022 [[hep-lat/0401033](#)] [SPIRES].
- [20] D. Becirevic, V. Gimenez, V. Lubicz and G. Martinelli, *Light quark masses from lattice quark propagators at large momenta*, *Phys. Rev.* **D 61** (2000) 114507 [[hep-lat/9909082](#)] [SPIRES].
- [21] J.-R. Cudell, A. Le Yaouanc and C. Pittori, *Pseudoscalar vertex, Goldstone boson and quark masses on the lattice*, *Phys. Lett.* **B 454** (1999) 105 [[hep-lat/9810058](#)] [SPIRES].
- [22] J.R. Cudell, A. Le Yaouanc and C. Pittori, *Large pion pole in $Z_S^{\text{MOM}}/Z_P^{\text{MOM}}$ from Wilson action data*, *Phys. Lett.* **B 516** (2001) 92 [[hep-lat/0101009](#)] [SPIRES].
- [23] L. Giusti and A. Vladikas, *RI/MOM renormalization window and Goldstone pole contamination*, *Phys. Lett.* **B 488** (2000) 303 [[hep-lat/0005026](#)] [SPIRES].
- [24] J.A. Gracey, *Three loop anomalous dimension of non-singlet quark currents in the RI' scheme*, *Nucl. Phys.* **B 662** (2003) 247 [[hep-ph/0304113](#)] [SPIRES].
- [25] K.G. Chetyrkin and A. Retey, *Renormalization and running of quark mass and field in the regularization invariant and \overline{MS} schemes at three and four loops*, *Nucl. Phys.* **B 583** (2000) 3 [[hep-ph/9910332](#)] [SPIRES].
- [26] G.P. Lepage and P.B. Mackenzie, *On the viability of lattice perturbation theory*, *Phys. Rev.* **D 48** (1993) 2250 [[hep-lat/9209022](#)] [SPIRES].

- [27] EUROPEAN TWISTED MASS collaboration, B. Blossier et al., *Light quark masses and pseudoscalar decay constants from $N_f = 2$ lattice QCD with twisted mass fermions*, *JHEP* **04** (2008) 020 [[arXiv:0709.4574](#)] [[SPIRES](#)].
- [28] R. Frezzotti, G. Martinelli, M. Papinutto and G.C. Rossi, *Reducing cutoff effects in maximally twisted lattice QCD close to the chiral limit*, *JHEP* **04** (2006) 038 [[hep-lat/0503034](#)] [[SPIRES](#)].
- [29] G. Rossi, R. Frezzotti, G. Martinelli and M. Papinutto, *Cutoff effects in maximally twisted LQCD*, *PoS(LAT2005)285* [[hep-lat/0509168](#)] [[SPIRES](#)].
- [30] R. Frezzotti, G. Martinelli, M. Papinutto and G.C. Rossi, *Reducing cutoff effects in maximally twisted lattice QCD close to the chiral limit*, *JHEP* **04** (2006) 038 [[hep-lat/0503034](#)] [[SPIRES](#)].
- [31] G. Rossi, R. Frezzotti, G. Martinelli and M. Papinutto, *Cutoff effects in maximally twisted LQCD*, *PoS(LAT2005)285* [[hep-lat/0509168](#)] [[SPIRES](#)].