

RESEARCH

Open Access



Pronuclear formation by ICSI using chemically activated ovine oocytes and zona pellucida bound sperm

J. E. Hernández-Pichardo^{1,6}, Y. Ducolomb², S. Romo³, M. E. Kjelland⁴, R. Fierro², F. Casillas⁵ and M. Betancourt^{2*}

Abstract

Background: In order to improve ICSI, appropriate sperm selection and oocyte activation is necessary. The objective of the present study was to determine the efficiency of fertilization using ICSI with chemically activated ovine oocytes and sperm selected by swim up (SU) or swim up + zona pellucida (SU + ZP) binding.

Results: Experiment 1, 4–20 replicates with total 821 *in vitro* matured oocytes were chemically activated with ethanol, calcium ionophore or ionomycin, to determine oocyte activation (presence of one PN). Treatments showed similar results (54, 47, 42 %, respectively) but statistically different ($P < 0.05$) than mechanical activated oocytes in sham, ICSI and sham injection (13, 25, 32 %, respectively) (10–17 replicates; $n = 429$).

Experiment 2: Twelve ejaculates and 28 straws of semen were used (11–19 replicates). Sperm were selected by SU in BSA-TCM 199-H medium. A total of 2,294 fresh sperm and 2,760 from frozen-thawed semen were analyzed after SU or SU + ZP binding. Fresh sperm selected by SU showed acrosome reaction (AR) of 59 %, the sperm selected by SU + ZP binding increased AR to 91 %. In comparison, the AR of frozen-thawed sperm using SU or SU + ZP binding was 77 and 86 %, respectively ($P < 0.05$).

Experiment 3: fertilization in 200 mechanical activated oocytes (17 replicates) was 4 %, but fertilization increased in ethanol activated oocytes after ICSI (12–28 %) (5–6 replicates). When fresh sperm only selected by SU were injected to 123 oocytes, a fertilization rate (28 %) was achieved; in sperm selected by SU + ZP was 25 % (73 oocytes). In comparison, in frozen-thawed sperm selected by SU, fertilization was 13 % (70 oocytes), whereas sperm from SU + ZP binding displayed 12 % (51 oocytes) ($P > 0.05$).

Conclusions: Chemical activation induces higher ovine oocyte activation than mechanical activation. Ethanol slightly displays higher oocyte activation than calcium ionophore and ionomycin. Sperm selection with SU + ZP increased AR/A and AR/D rates in comparison with SU in fresh and frozen-thawed sperm. According to this, in terms of fertilization rates, chemical activation after ICSI increased oocyte PN formation compared to mechanical activation. Also, fresh sperm treated with SU and SU + ZP were significantly different than frozen-thawed sperm, but between sperm treatments no significant differences were obtained.

Keywords: Ethanol, ICSI, Oocyte activation, Ovine, Pronucleus, Zona pellucida

* Correspondence: bet@xanum.uam.mx

²División de Ciencias Biológicas y de la Salud, Departamento de Ciencias de la Salud, Universidad Autónoma Metropolitana-Iztapalapa, San Rafael Atlixco 186, CP 09340 Ciudad de México, Mexico

Full list of author information is available at the end of the article



© The Author(s). 2016 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Background

In recent decades, assisted reproductive techniques (ARTs) in mammals, have been developed for its use in super-ovulation, embryo transfer (ET), *in vitro* fertilization (IVF), intracytoplasmic sperm injection (ICSI) and cloning by somatic cell nuclear transfer. Despite the great advances in the protocols of these ARTs, their efficiency still remains low [1].

Among ARTs, ICSI is the current procedure for avoiding polyspermy and increasing the number of produced zygotes, as well as allowing for the study of several aspects of fertilization such as gamete interaction, sperm decondensation, oocyte activation [2] and gene transfer [3, 4]. In human, this technique has been developed with great success [5]; however, its application in livestock is still not well established, mainly due to the different reproductive physiology among species of economically important domestic animals such as cattle, pigs, horses and sheep. In general, embryo development (ED) using the ICSI procedure is lower than that of IVF [2, 6].

Although ovine oocyte fertilization by means of the injection of immobilized sperm has been done before, and normal lambs have been born [7, 8], ICSI success is markedly restricted due to sperm head decondensation failure and low male pronuclei (PN) formation [9]. In turn, several studies have tried to solve this issue in sheep using sperm pretreatments before performing ICSI. In order to induce sperm head decondensation and male PN formation, sperm have been pre-treated with lysophosphatidylcholine (LPC) [10], triton X-100 [11], dithiothreitol (DTT), sodium dodecyl sulphate (SDS) or by freezing and thawing without cryoprotectants [9]. These treatments are directed at removing the sperm membranes and dissolving the nuclear proteins allowing the head decondensation.

In human, physiological strategies are being explored to investigate the selection of sperm that are more capable of achieving successful ICSI, compared to the subjective way that the embryologists currently use [12], increasing fertilization. A strategy for improving ICSI is using sperm attached to the zona pellucida (ZP) [13, 14]. Studies [12, 14, 15] have concluded that the ZP has the ability to select functionally normal sperm and even those with a superior quality. Further, the attachment to the ZP during fertilization is necessary for inducing the acrosome reaction (AR), which is an important factor for ICSI success.

Other important factor is the oocyte activation which is the resumption of meiosis II during fertilization increasing calcium ion levels in the cytoplasm. Without this step, the inseminated oocyte prevents the decondensation of the sperm head. Notably, artificial stimuli can mimic the action of sperm during fertilization. Chemical activation of oocytes has been necessary with ICSI

because mechanical activation alone, by means of sperm injection, is sometimes not sufficient for inducing oocyte activation [16, 17]. To improve results, appropriate sperm treatment and oocyte activation is necessary to achieve fertilization [17].

Oocyte activation has not been shown to be crucial in several species such as mice, hamsters and rabbits; however, it is essential in pigs, cattle and sheep [18]. The chemical activation of ovine oocytes after ICSI has been performed using: ionomycin (ION) [17, 19, 20], ION+ 6 dimethylaminopurine (6-DMAP) [4, 9, 17, 21, 22], ethanol (ETL) [4, 21], DTT [9] and calcium ionophore (CAI) [16]. However, even with such chemical treatments, the percent of sheep blastocysts produced after ICSI and cultured *in vitro* was around 20 %. Moreover most investigations have evaluated ED but not fertilization, determined by PN formation, to assure that the zygotes are diploid. This condition is essential for the normal development of the embryos [23]. When oocytes are artificially activated, a high probability that the resultant blastocysts will be parthenogenic exists [17, 24, 25]. Therefore, the objective of the present study was to determine the efficiency of fertilization using ICSI with chemically activated ovine oocytes with sperm selected by swim-up (SU) or SU plus binding to ZP (SU + ZP).

Methods

The ovaries were collected from a slaughterhouse, "El Rojo". The aforementioned facility has the animal health federal law authorization under the number 150810056630. Except for the maturation and development media, which were prepared in a commercial laboratory (In Vitro S.A., México), all chemicals were purchased from Sigma Chemical Company (ST Louis, MO USA). All incubation conditions were performed at 38.5 °C in an atmosphere with 5 % CO₂, 95 % air, and humidity at saturation (NUAIRE, USA). *In vitro* maturation (IVM) and ICSI procedures were performed under mineral oil (Fisher Scientific, USA). All media were stored at 4 °C for not more than three wk and supplemented 24 to 48 h before use.

Experimental design

The experimental design consisted of three separate experiments. Experiment 1 was conducted to identify the percent activation of oocytes matured *in vitro* only exposed to chemical activation with ETL ($n = 393$), CAI ($n = 350$), ION ($n = 78$) or mechanical activation through ICSI ($n = 200$), sham injection (false injection) ($n = 78$) or sham (only manipulation of the gametes without injection) ($n = 151$), and evaluated by the PN formation.

Experiment 2 was conducted to determine the viability and sperm acrosome status after selection by SU or SU + ZP of fresh ($n = 1,194$, SU; $n = 1,100$ SU + ZP) and frozen-thawed sperm ($n = 718$, SU; $n = 922$ SU + ZP).

Experiment 3 was conducted to determine fertilization evaluated by the formation of two PN and without the presence of a sperm head when performing ICSI with fresh or frozen-thawed sperm selected by SU or SU + ZP. After ICSI, oocyte chemical activation with 7 % ETL for 5 min ($n = 123$ SU; $n = 73$ SU + ZP and $n = 70$ SU; $n = 51$ SU + ZP respectively) and mechanical activation ($n = 200$) were performed.

Oocyte collection and maturation

Ovaries were collected from adult ewes and transported, less than three h, to the laboratory in 0.157 mol/L NaCl solution at 37 °C with antibiotics (10,000 UI/mL ampicillin, 10,000 µg/mL streptomycin and 25 µg/mL, amphotericin (In Vitro S.A., México). Ovaries were rinsed three times in the fresh solution described above. Follicular fluid was aspirated from 2 to 6 mm follicles punctured with an 18-gauge needle fixed to a 10 mL disposable syringe containing 1 mL of modified Tyrode's medium supplemented with 10 mmol/L sodium lactate, 10 mmol/L HEPES and 0.1 % polyvinilic alcohol (PVA) (TL-HEPES-PVA) with a pH of 7.3–7.4, supplemented with 200 UI/mL heparine. Follicular fluid was pooled in a 50 mL conical tube (PISA, México). Cumulus cell-oocyte complexes (COCs) with compact cumulus mass and oocytes with uniform cytoplasm were selected and washed three times in 500 µL drops of maturation medium TCM-199 with Earle's salts, 26.2 mmol/L sodium bicarbonate and L glutamine (In Vitro S.A., México), supplemented with 0.1 % PVA, 3.05 mmol/L D-glucose, 0.91 mmol/L sodium pyruvate, 0.57 mmol/L cysteine and 10 ng/mL epidermal growth factor [26].

For oocyte maturation, sterile four well plates were used (Thermo-Scientific Nunc, Rochester NY). First, 500 µL of maturation medium with 0.5 µg/mL LH, 0.5 µg/mL FSH and 10 % (v/v) fetal calf serum (FBS) (In Vitro S.A., México) was added to each well. Next, 20–30 COCs were placed in the well with the maturation medium and incubated for 24 h [19]. After IVM, COCs were placed in 500 µL of TCM 199-HEPES (TCM 199-H) with 300 IU of hyaluronidase for three min, then washed three times in TCM 199-H [13]. Maturation was determined by identification of the first polar body (PB) using an inverted microscope (Nikon Eclipse TE200, Japan).

Oocyte activation

For experiment 1, mature denuded oocytes (only with a PB) were distributed randomly among the following groups to perform activation; mechanical activation: (1) Sham: oocytes manipulated as in ICSI without injection, (2) ICSI, (3) Sham injection: oocyte pierced without sperm insertion, and for chemical activation without sperm injection: (4) 7 % ETL for 5 min, (5) 50 µmol/L CAI for 10 min, and (6) 5 µmol/L ION for 5 min. After

activation, oocytes were washed three times in modified Tris-buffered Medium (mTBM) consisting of 13.1 mmol/L NaCl, 3 mmol/L KCl, 7.5 mmol/L $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 20 mmol/L Tris, 11 mmol/L glucose, 5 mmol/L sodium pyruvate, supplemented with 0.4 % bovine serum albumin (BSA) fraction V, and 2.5 mmol/L caffeine benzoate and placed in 200 mL of medium and incubated during 18 h.

Assessment of pronucleus formation

After incubation, oocytes were stained with 10 µg/mL bisbenzimidazole (Hoechst 33258) in cold ETL for 15 min, protected from the light. Further, oocytes were placed on a slide in a microdrop of PBS-glycerol (1:9) [9], and examined under an epifluorescence microscope at 400× (Eclipse E600, Nikon, Japan).

Non-activated oocytes were determined by the identification of the metaphase II, whereas activated oocytes presented a PN.

Sperm preparation

Semen was collected from a creole ram of proven fertility, located in the animal facilities at the Universidad Autónoma Metropolitana-Xochimilco, as approved under the regulations of the Committee for Care and Use of Animals. One hundred µL of fresh or frozen semen were added to 2 mL of TCM 199-H supplemented with 3 mg/mL BSA (BSA-TCM 199-H), and incubated for 1 h for SU to help capacitation. Afterwards, 500 µL of medium were gently removed from the top of the suspension, then 3 mL of BSA-TCM 199-H were added and centrifuged at 200 ×g for 10 min [9, 27]. The pellet was suspended to give a final concentration of 1×10^6 sperm/mL [28].

Exposure of sperm to mature oocytes ZP

In each assay, four oocytes in metaphase II were placed in 400 µL drops of BSA-TCM-H and 1 % gentamycin, followed by the addition of 100 µL of the sperm suspension and incubated for 1 h. The oocytes were transferred to a 300 µL drop of TCM-H medium and washed by repeated aspirations with a fine glass pipette to eliminate the non-bound ZP sperm. Two oocytes with sperm tightly bound to the ZP were placed in 20 µL drops of the same medium and covered with mineral oil. The sperm bound to the ZP were removed by repeated aspiration of the oocytes through a fine glass pipette, for ICSI [13, 29], while the other two oocytes were placed in 100 µL drops to obtain the sperm for viability and AR analysis.

Viability and acrosomal status of sperm after SU and ZP binding

Fresh and frozen-thawed sperm were selected by SU or SU + ZP binding. To determine their viability and acrosome

status, propidium iodide (PI) and fluorescein isothiocyanate with peanut agglutinin (FITC-PNA) double staining was developed [30, 31]. From each sperm suspension sample, 100 μ L were obtained. Next, 5 μ L of FITC-PNA and 5 μ L of PI were added, followed by incubation for 5 min. Ten μ L of the sperm suspension were mixed with 10 μ L 1.6 % paraformaldehyde in a slide and analyzed under a fluorescence microscope (Eclipse E600, Nikon, Japan) at 400 \times . The functional status of the sperm was classified according to the following staining patterns: live sperm with intact acrosome: cells with FITC-PNA stain at the acrosome, without PI stain at the post-acrosomal region. Live sperm with AR: cells without FITC-PNA acrosome stain and without PI nuclear stain. Dead sperm with intact acrosome: cells with FITC-PNA stain in the acrosome region, and with PI nuclear stain. Dead sperm with AR: without FITC-PNA stain in the acrosome region, and with PI nuclear stain.

Intracytoplasmic sperm injection

For ICSI, a 50 μ L drop of TCM 199-H, supplemented with 2 % FBS and 1 % gentamycin containing five *in vitro* matured denuded oocytes, and one 10 μ L drop of sperm suspension with 10 % polivinilpirrolidone (PVP) (1:1 dilution) were added to a 6 cm diameter Petri dish [8, 16]. Microinjection was performed as previously reported by Shirazi et al. [9], using an inverted microscope at 200 \times (Nikon Eclipse TE200, Japan). The microinjection pipette was immersed in the sperm suspension-PVP and one sperm was immobilized by scoring the midpiece and aspirating the tail-first, and then injected into the ooplasm through the ZP. The first PB was either in the 6 or 12 o'clock position, and the injection pipette was in the 3 o'clock position. During the injection, cytoplasm was aspirated to ensure that the oolema was punctured. After ICSI, oocytes were washed three times in BSA-TCM 199-H. Mechanical and chemical activated oocytes were incubated in 100 μ L of mTBM for 18 h. A total of 517 oocytes were injected (5–17 replicates).

Oocyte chemical activation after ICSI

For experiment 3, 10–20 ICSI inseminated oocytes were activated in 100 μ L drops of 7 % ETL (Baker Absolute

alcohol 9000-02) in TCM 199-H at room temperature for 5 min [32]. After activation, oocytes were washed and incubated in mTBM during 18 h.

Nuclear staining

Eighteen h after ICSI, oocytes were stained with 10 μ g/mL Hoechst 33258 as mentioned, to determine the following stages: 1) non-activated: oocytes in metaphase II and without a decondensed sperm head; 2) activated oocytes: formation of the female PN and without a decondensed sperm head; and 3) fertilized oocytes: oocytes with two PN and without the presence of the sperm head [9, 19, 33].

Statistical analysis

With regard to the oocyte activation, 4–20 replicates were performed. For sperm viability and acrosome status, 11–19 replicates were performed for fresh and frozen-thawed semen. To assess fertilization 5–6 replicates were performed for each group. ANOVA and Bonferroni multiple post hoc test were used for comparing the results of the following variables: oocyte activation, viability and acrosome status of fresh sperm compared to frozen-thawed sperm, and fertilization with mechanical and chemical activation, but also between sperm treatments [34], considering significant differences between the groups when $P < 0.05$.

Results

Oocyte activation

The mechanical activation of oocytes with sham injection was higher than sham ($P < 0.05$). However, no differences were observed between sham injection and ICSI. Chemical activation with ETL, CAI and ION was higher than the mechanical activation ($P < 0.05$). When compared to chemical activation with ETL, the three chemical activation treatments had a similar proportion of PN with a slightly increase in the ETL group, shown in Table 1. Based on these results, oocytes subjected to ICSI were hereafter treated only with 7 % ETL for 5 min.

Table 1 Activation procedures in sheep oocytes matured *in vitro*

	Mechanical activation			Chemical activation		
	Sham	ICSI	Sham Injection	ETL	CAI	ION
N	13	17	10	20	19	4
n	151	200	78	393	350	78
Activated oocytes,%	2,0	4,9	25	21,1	16,4	3,3
	13 \pm 1.6 ^a	25 \pm 0.5 ^{ab}	32 \pm 3.4 ^b	54 \pm 1.3 ^c	47 \pm 1.3 ^c	42 \pm 5.0 ^c

Sham: oocytes manipulated as in ICSI but no injection; Sham injection: oocyte pierced but no sperm insertion; ETL: 7 % Ethanol 5 min; CAI: 50 μ mol/L Calcium lonophore 10 min; ION: 5 μ mol/L lonomicine 5 min. Mechanically activated oocytes: Oocytes in metaphase II. Activated oocytes: Presence of one pronucleus

N number of replicates; n number of oocytes examined

Percentage data are presented as mean \pm SD

^{a,b,c}Values with different letters indicate statistically significant differences ($P < 0.05$)

Viability and acrosome reaction

Independently of whether the sperm were alive or dead, the proportion with AR was greater in those selected with SU + ZP than SU only (control) ($P < 0.05$), either in fresh or frozen-thawed semen. These results indicate that if ICSI is performed using sperm selection through ZP binding, the probability of injecting sperm with AR is very high, independently of the viability and fresh or frozen-thawed condition, shown in Table 2.

ICSI

The oocytes treated with ETL had an increased fertilization rate with ICSI (presence of two PN without the sperm head) compared to mechanical activation. Fertilization was increased when fresh semen was used compared to frozen-thawed semen, independent of the sperm selection procedure ($P < 0.05$). The comparison of the four groups of oocytes exposed to chemical activation with ETL were significant different than mechanical activation ($P < 0.05$), shown in Table 3. Also, a significant difference in fertilization was observed between the fresh and frozen-thawed sperm but between selection (SU and SU + ZP) no significant differences were obtained (Table 3). The percentage of activated oocytes was greater than percentage of fertilized oocytes, indicating that the proportion of sperm decondensation is still lower when the oocytes are activated.

Discussion

The chemical activation of the sheep oocytes during ICSI is necessary due to the low percentages of sperm decondensation and PN formation. Oocyte activation implies an increase of intracellular free calcium, causing the degradation of the maturation promoting factor (MPF), which is essential for the resumption of the second meiotic division and the formation of both PN [25, 35].

ETL is among the most utilized chemical activators through the formation of inositol 1,4,5-triphosphate and its union to the membrane receptors in the endoplasmic reticulum releasing of intracellular calcium [25, 36, 37].

Other chemical activators include CAI and ION, which induce the liberation of intracellular calcium stored in the endoplasmic reticulum, evident by the activation of several proteolytic pathways calcium dependent that drive the destruction of cyclin B, reducing the activity of MPF [25, 36, 38].

In regard to experiment 1 the present study demonstrated that chemical activation induced a higher female PN formation than that produced by mechanical activation. The results obtained with ETL oocyte activation were similar to those reported by Shirazi et al. (50 %) [36] who studied the effect of sheep oocyte maturation on parthenogenetic development. In other mammalian domestic species, oocyte activation with ETL generally leads to a lower incidence of pronuclear formation. Despite this, Wang et al. [39] activated cow oocytes and reported 42 % activation. Additionally, Yi and Park [40] observed an increase in the activation of pig oocytes with ETL exposure (34 %).

Also, in the present study, the use of CAI resulted in an oocyte activation of 47 %. However, little is known about sheep CAI oocyte activation. Nevertheless, studies in cattle reported 22 % [39]. Oocyte activation using ION resulted in 42 % PN formation, which is similar to that reported by Shirazi et al. [36], who evaluated activation through ED. These results support the use of these chemical activators with sheep oocytes, which are essential in ICSI.

In the present study, mechanical activation by ICSI displayed 25 % PN formation, similar to that reported by Gómez et al. [8, 33] with 31 % (determined through ED) and 38 % (resumption of second meiotic division), while Shirazi et al. [17] observed a PN formation of 5 % only. The effect of the sham injection resulted in 32 % of the oocytes undergoing PN formation. The results of this study are superior to those reported by Gómez et al. [10], with 22 %.

The sham group showed that 13 % of the oocytes were activated, which is similar to those by Catt and Rhodes [41]. This could be due to the use of hyaluronidase

Table 2 Viability and acrosome status of sperm selected by swim-up or swim-up and oocyte ZP binding

		Fresh Semen, %					Frozen-thawed semen, %			
		AR/A	AR/D	NAR/A	NAR/D		AR/A	AR/D	NAR/A	NAR/D
SU	N = 12					N = 14				
	n = 1,194	52,8 44 ± 1.8 ^a	17,9 15 ± 0.8 ^a	34,0 28 ± 8.3 ^a	14,7 12 ± 0.6 ^a	n = 1410	71,8 51 ± 0.9 ^{a*}	37,3 26 ± 0.9 ^{a*}	17,5 12 ± 0.3 ^{a*}	14,4 10 ± 0.5 ^a
SU + ZP	N = 11					N = 19				
	n = 1,100	64,8 59 ± 0.9 ^b	34,7 32 ± 1.1 ^b	40 4 ± 0.3 ^b	65 6 ± 0.2 ^b	n = 1350	92,2 68 ± 1.8 ^{b*}	23,7 18 ± 0.9 ^{b*}	14,3 11 ± 0.5 ^{a*}	48 4 ± 0.2 ^{b*}

SU swim-up, ZP Zona pellucida, AR/A acrosome reaction/alive, AR/D acrosome reaction/dead, NAR/A no acrosome reaction/alive, NAR/D no acrosome reaction/dead, N number of replicates, n number of sperm examined

Percentage data are presented as mean ± SD

^{a,b}Values in the same column with different letters are significantly different ($P < 0.05$)

*Within rows, among the variables corresponding to fresh and frozen-thawed semen counterparts, significant difference was observed ($P < 0.05$)

Table 3 Effect of oocyte activation, semen type and sperm selection technique for ICSI on fertilization percentage

Semen type	Frozen-thawed semen	Fresh		Frozen-thawed	
	Mechanical activation	Activated with 7 % Ethanol			
Sperm Treatment	"Swim-up"	"Swim-up"	"Swim-up" + Zona pellucida	"Swim-up"	"Swim-up" + Zona pellucida
N	17	5	6	5	5
n	200	123	73	70	51
Activated oocytes, %	4,9 25 ± 2.7	7,5 61 ± 1.3*	3,6 49 ± 1.6*	4,0 57 ± 3.2*	3,1 61 ± 6.3*
Fertilized oocytes, %	8 4 ± 0.2	3,5 28 ± 1.7*	18 25 ± 1.7*	9 13 ± 1.2*▲	6 12 ± 0.9▲

Activated oocytes: Formation of female pronucleus and no decondensed sperm head; Fertilized oocytes: Oocytes with two pronuclei without sperm head

N number of replicates, n Number of oocytes examined

Percentage data are presented as mean ± SD

*Indicates significant differences with respect to mechanical activated oocytes ($P < 0.05$)

▲Indicates significant differences with respect to SU fresh ($P < 0.05$)

during the removal of cumulus cells, since this enzyme is capable of producing oocyte activation in mice, cattle and human itself [32]. With regard to the activation using sham injection or ICSI, our results indicate that the injection by itself induces sheep oocyte activation and not necessarily the presence of a sperm.

Several factors influence oocyte sensitivity to artificial activation and this can explain the differences between the results obtained in the present study and those reported previously. Factors influencing oocyte activation include: differences among species [37], *in vivo* or *in vitro* maturation, presence or absence of cumulus cells [32, 36] and type of stimuli [32].

For improving ICSI results, new methods of sperm selection are being developed in human, as injecting sperm previously bound to the ZP of oocytes. To our knowledge, this is the first study that evaluates the acrosome status and viability of sperm from fresh and frozen-thawed ovine sperm after being in contact with the *in vitro* matured oocyte ZP of the same species. It is important to consider that the use of ZP does not just select sperm, but also induces AR, which has been reported to be essential for decondensation of sperm chromatin in ooplasm [9]. In experiment 2 the percentage of sperm with AR after the ZP binding was 91 and 86 % for fresh and frozen-thawed sperm, respectively, which are similar to those reported with fresh goats semen, [42] pigs [43] and human [44]. Notably, a significant difference was observed between the SU + ZP and SU selection procedures. Among the factors that affect the trigger of the AR through ZP contact, is the state of oocyte maturation; immature porcine oocytes in contact with fresh semen for 10 min had an AR of 34 %, while mature oocytes 42 % underwent AR, similar tendency was observed when frozen-thawed sperm was used with immature (37 %) and mature oocytes (45 %) [45].

In experiment 3, a significant increase in the percentage of activated and fertilized oocytes was observed when stimulated with ETL after ICSI, similar results were obtained

by Shirazi et al. [17], besides, they reported that mechanical stimulation (ICSI or sham injection) is not sufficient to activate ovine oocytes for PN formation, indicating that chemical activation is essential for ICSI in sheep.

Oocyte treatment with ETL after ICSI increased the oocyte activation percentage from 49 to 61 %, and 12 to 28 % for fertilization, compared to mechanical activated (25 and 4 %, respectively). The fertilization rate with fresh sperm was only 50 % with respect to the proportion of the activated oocytes; while that in frozen-thawed sperm was not higher than 25 %. If the evaluation of these oocytes was through the ED, more than 50 % could be parthenotes. Similar results have been reported in cattle oocytes treated with ETL after ICSI with frozen-thawed sperm heads, with 52 % of activation, but only 27 % of fertilization, being 56 % parthenote embryos [46]. In equine oocytes with the same treatment, 56 % activation, 16 % fertilization, and 40 % parthenotes have been obtained; indicating that even when oocytes are activated, the sperm decondensation may not occur, but artificial activation can induce parthenogenetic development in oocytes with ICSI [47].

In the present study, an increase in fertilization was not observed using sperm previously bound to oocyte ZP and with an AR higher than 85 %. This finding is in agreement with that reported by Gómez et al. [10], in which an increase in fertilization was not observed (22 %) after ICSI with AR sperm (86 %) compared to that obtained with untreated sperm (33 %). These results indicate that there is not a correlation between the percentage of sperm with AR and the number of oocytes fertilized, suggesting that it is not necessary to induce AR before ICSI in sheep to obtain fertilization. In human, Black et al. [13] did not observe a significant difference in fertilization between ICSI performed with sperm bound to the ZP and conventional ICSI; however, higher quality embryos and a greater number of pregnancies were produced.

A significant difference in the fertilization rate was observed between oocytes fertilized with frozen-thawed

and fresh sperm. This is consistent with Gómez et al. [10], reporting higher fertilization rates after ICSI with fresh sperm than frozen-thawed sperm. Also Gómez et al. [10] results could be due to acrosome-membrane damage which may occur during the freeze-thaw process; thereby predisposing sperm to further membrane changes or some other deleterious effect on sperm DNA. In addition, in human sperm, it has been reported that the freeze-thaw process produces an oxidative stress due to the generation of oxygen free radicals and an alteration of the sperm antioxidant system [48]. The aforementioned oxidative stress can cause the peroxidation of membrane lipids, alterations of proteins and DNA fragmentation [49] which decreases sperm fertilization capacity.

Conclusions

Chemical activation induces higher ovine oocyte activation than mechanical activation. ETL slightly displays higher oocyte activation than CAI and ION. Sperm selection with SU + ZP increased AR/A and AR/D rates in comparison with SU in fresh and frozen-thawed sperm. According to this, in terms of fertilization rates, chemical activation after ICSI increased oocyte PN formation compared to mechanical activation. Fresh sperm selected by SU and SU + ZP were significantly different than frozen-thawed sperm, but between both selection procedures no differences were obtained. Therefore, this study demonstrates that chemical activation is required to activate oocytes after injection and acrosome reaction is not essential for fertilization. In turn, fresh sperm was more efficient in terms of fertilization capacity when performing ICSI.

Acknowledgments

The authors thank the slaughterhouse "El Rojo", Estado de México for the donation of ovaries. This publication represents partial fulfillment of the requirements for the degree of Doctor in Biología Experimental for José Ernesto Hernández-Pichardo.

Funding

This study was not financially supported.

Availability of data and materials

The authors declare that they do not apply new software and databases.

Authors' contributions

Direction of the study: MB, YD, RF and SR. Established the experimental design: JEHP and MB. Managed the animals and performed the experiments: JEHP. Analyzed the results and wrote the manuscript: JEHP, MB, MK, YD, SR and FC. All authors verify reading and approving the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Author details

¹División de Ciencias Biológicas y de la Salud, Departamento de Producción Agrícola y Animal, Universidad Autónoma Metropolitana-Xochimilco, Ciudad de México, Mexico. ²División de Ciencias Biológicas y de la Salud, Departamento de Ciencias de la Salud, Universidad Autónoma

Metropolitana-Iztapalapa, San Rafael Atlixco 186, CP 09340 Ciudad de México, Mexico. ³Departamento de Ciencias Pecuarías, Facultad de Estudios Superiores Cuautitlán, UNAM, Ciudad de México, Estado de México, Mexico. ⁴Conservation, Genetics & Biotech, LLC, Valley City, ND, USA. ⁵División de Ciencias Biológicas y de la Salud, Departamento de Biología de la Reproducción, Universidad Autónoma Metropolitana-Iztapalapa, Ciudad de México, Mexico. ⁶Doctorado en Biología Experimental, Universidad Autónoma Metropolitana-Iztapalapa, Ciudad de México, Mexico.

Received: 12 May 2016 Accepted: 14 October 2016

Published online: 08 November 2016

References

1. Franco MM, Prickett AR, Oakey RJ. Minireview. The role of CCCTC-Binding Factor (CTCF) in genomic imprinting, development, and reproduction. *Biol Reprod*. 2014;91:1–9.
2. García-Roselló E, García-Mengual E, Coy P, Alfonso J, Silvestre MA. Intracytoplasmic sperm injection in livestock species: an update. *Reprod Domest Anim*. 2009;44:143–51.
3. Bevacqua RJ, Pereyra-Bonnet F, Fernandez-Martin R, Salamone DF. High rates of bovine blastocyst development after ICSI-mediated gene transfer assisted by chemical activation. *Theriogenology*. 2010;74:922–31.
4. Pereyra-Bonnet F, Gibbons A, Cueto M, Sipowicz P, Fernández-Martín R, Salamone D. Efficiency of sperm-mediated gene transfer in the ovine by laparoscopic insemination, *in vitro* fertilization and ICSI. *J Reprod Dev*. 2011;57:188–96.
5. Gerris JMR. Single embryo transfer and IVF/ICSI outcome: a balanced appraisal. *Hum Reprod Update*. 2005;11:105–21.
6. Casillas F, Ducolomb Y, Lemus AE, Cuello C, Betancourt M. Porcine embryo production following *in vitro* fertilization and intracytoplasmic sperm injection from vitrified immature oocytes matured with a granulosa cell co-culture system. *Cryobiology*. 2015;71:299–305.
7. Catt JW. Intracytoplasmic sperm injection (ICSI) and related technology. *Anim Reprod Sci*. 1996;42:239–50.
8. Gómez MC, Catt JW, Evans G, Maxwell WMC. Cleavage, development and competence of sheep embryos fertilized by intracytoplasmic sperm injection and *in vitro* fertilization. *Theriogenology*. 1998;49:1143–54.
9. Shirazi A, Derakhshan-Horeh M, Pilvarian AA, Ahmadi E, Nazari H, Heidari B. Effect of pre-treatment of ovine sperm on male pronuclear formation and subsequent embryo development following intracytoplasmic sperm injection. *Reprod Domest Anim*. 2011;46:87–94.
10. Gómez MC, Catt JW, Evans G, Maxwell WMC. Effect of culture, incubation and acrosome reaction of fresh and frozen-thawed ram spermatozoa for *in vitro* fertilization and intracytoplasmic sperm injection. *Reprod Fertil Dev*. 1997;9:665–73.
11. Pereyra-Bonnet F, Gibbons A, Cueto M, Naim P, Fernández-Martín F, Salamone D. Transgénesis mediada por espermatozoides en ovejas prepúberes. VII Simp Int Reprod Anim. IRAC. 2007;213.
12. De-Yi L. Could using the zona pellucida bound sperm for intracytoplasmic sperm injection (ICSI) enhance the outcome of ICSI? *Asian J Androl*. 2011;13:197–8.
13. Black M, de Liu Y, Bourne H, Baker HW. Comparison of outcomes of conventional intracytoplasmic sperm injection and intracytoplasmic sperm injection using sperm bound to the zona pellucida of immature oocytes. *Fertil Steril*. 2010;93:672–4.
14. Henkel R. Sperm preparation: state-of-the art-physiological aspect and application of advanced sperm preparation methods. *Asian J Androl*. 2012;14:260–9.
15. Parmegiani L, Cognigni GE, Filicori M. New Advances in Intracytoplasmic Sperm Injection (ICSI). *Advances in Embryo Transfer*. In: Bin Wu, editor. InTech. [On line]. 2015. <http://cdn.intechopen.com/pdfs-wm/32057.pdf>.
16. Gómez MC, Catt JW, Evans G, Maxwell WMC. Sheep oocyte activation after intracytoplasmic sperm injection (ICSI). *Reprod Fertil Dev*. 1998;10:197–205.
17. Shirazi A, Ostad-Hosseini S, Ahmadi E, Heidari B, Shams-Esfandabadi N. *In vitro* developmental competence of ICSI-derived activated ovine embryos. *Theriogenology*. 2009;71:342–8.
18. Yong HY, Pyo BS, Hong JY, Kang SK, Lee BC, Lee ES, et al. A modified method for ICSI in the pig: injection of head membrane-damaged sperm using a 3–4 µm diameter injection pipette. *Hum Reprod*. 2003;18:2390–6.
19. Catalá MG, Izquierdo D, Rodríguez-Prado M, Hammami S, Paramio MT. Effect of oocyte quality on blastocyst development after *in vitro* fertilization (IVF)

- and intracytoplasmic sperm injection (ICSI) in a sheep model. *Fert Steril*. 2012;97:1004–8.
20. Hosseini SM, Asgari V, Ostadhosseini S, Hajian M, Piryaei A, Najarasi M, et al. Potential applications of sheep oocytes as affected by vitrification and *in vitro* aging. *Theriogenology*. 2012;77:1741–53.
 21. Loi P, Ledda S, Fulka J, Cappai P, Moor RM. Development of parthenogenetic and cloned ovine embryos: effect of activation protocols. *Biol Reprod*. 1998;58:1177–87.
 22. Pereyra-Bonnet F, Fernández-Martín R, Olivera R, Jarazo J, Vichera G, Gibbons A, et al. A unique method to produce transgenic embryos in ovine, porcine, feline, bovine and equine species. *Reprod Fertil Dev*. 2008;20:741–9.
 23. Nicolí A, Capodanno F, Rondini I, Valli B, Villani MT, Morini D, et al. Pronuclear morphology evaluation *in vitro* fertilization (IVF) intracytoplasmic sperm injection (ICSI) cycles: a retrospective clinical review. *J Ovarian Res*. 2013;6:1–6.
 24. Jiménez-Macedo AR, Anguita B, Izquierdo D, Mogas T, Paramio MT. Embryo development of prepubertal goat oocytes fertilised by intracytoplasmic sperm injection (ICSI) according to oocyte diameter. *Theriogenology*. 2006;66:1065–72.
 25. Kharche SD, Birade HS. Parthenogenesis and activation of mammalian oocytes for *in vitro* embryo production: A review. *Adv Biosci Biotechnol*. 2013;4:170–82.
 26. Fernández RF, Ducolomb Y, Romo S, Casas E, Salazar Z, Betancourt M. Viability, maturation and embryo development *in vitro* of immature porcine and ovine oocytes vitrified in different devices. *Cryobiology*. 2012;64:261–6.
 27. Paramio MT, Izquierdo D. Current status of *in vitro* embryo production in sheep and goats. *Reprod Domest Anim*. 2014;49:37–8.
 28. McPartlin LA, Suarez SS, Czaya CA, Hinrichs K, Bedford-Guaus SJ. Hyperactivation of stallion sperm is required for successful *in vitro* fertilization of equine oocytes. *Biol Reprod*. 2009;81:199–206.
 29. Liu F, Qiu Y, Zou Y, Deng ZH, Yang H, Liu YD. Use of zona pellucida-bound sperm for intracytoplasmic sperm injection produces higher embryo quality and implantation than conventional intracytoplasmic sperm injection. *Fertil Steril*. 2011;95:815–8.
 30. Ortgies F, Klewitz J, Görgens A, Martinsson G, Sieme H. Effect of procaine, pentoxifylline and trolox on capacitation and hyperactivation of stallion spermatozoa. *Andrologia*. 2012;44:130–8.
 31. Varesi S, Vernocchi V, Faustini M, Luvoni GC. Morphological and acrosomal changes of canine spermatozoa during epididymal transit. *Acta Vet Scand*. 2013;55:17.
 32. Vallejo J, Gómez-Piquer V, Tarín JJ. Inducción de la partenogénesis en ovocitos de mamíferos. *Rev Iberoam Fert*. 2003;20:177–87.
 33. Gómez MC, Catt SL, Gillan L, Catt JW, Evans G, Maxwell WMC. Time course of pronuclear formation and fertilization after insemination *in vitro* and intracytoplasmic sperm injection of *in vitro* matured sheep oocytes. *Zygote*. 1998;6:261–70.
 34. Wayne WD, Chad LC. *Biostatistics. A foundation for Analysis in the Health Sciences*. 10th ed. USA: Wiley; 2013.
 35. Tasripoo K, Srisakwattana K, Nualchuen W, Sophon S. Effects of various activators on bovine embryonic development following intracytoplasmic sperm injection. *Iran J Appl Anim Sci*. 2012;2:167–73.
 36. Shirazi A, Bahiraei A, Ahmadi E, Nazari H, Heidari B, Borjian S. The Effect of the duration of *in vitro* maturation (IVM) on parthenogenetic development of ovine oocytes. *Avicenna J Med Biotechnol*. 2009;1:181–91.
 37. Barbosa FC, Gasparini DL, Rigatto ML, Pria BID, De Lima NJF, Myakawa TP, et al. Artificial activation of bovine and equine oocytes with cycloheximide, roscovitine, strontium, or 6-dimethylaminopurine in low or high calcium concentrations. *Zygote*. 2013;22:387–94.
 38. Hosseini SM, Hajian M, Moulavi F, Shahverdi AH, Nasr-Esfahani MH. Optimized combined electrical–chemical parthenogenetic activation for *in vitro* matured bovine oocytes. *Anim Reprod Sci*. 2008;108:122–33.
 39. Wang ZG, Wang W, Yu SD, Xu ZX. Effects of different activation protocols on preimplantation development, apoptosis and ploidy of bovine parthenogenetic embryos. *Anim Reprod Sci*. 2008;105:292–301.
 40. Yi YJ, Park CS. Parthenogenetic development of porcine oocytes treated by ethanol, cycloheximide, cytochalasin B and 6-dimethylaminopurine. *Anim Reprod Sci*. 2005;86:297–304.
 41. Catt JW, Rhodes SL. Comparative intracytoplasmic sperm injection (ICSI) in human and domestic species. *Reprod Fertil Dev*. 1995;7:161–7.
 42. Somanath PR, Suraj K, Gandhi KK. Caprine sperm acrosome reaction: promotion by progesterone and homologous zona pellucida. *Small Rumin Res*. 2000;37:279–86.
 43. Fazeli AR, Holt C, Steenweg W, Bevers MM, Holt WW, Colenbrander B. Development of a sperm hemizona binding assay for boar semen. *Theriogenology*. 1995;44:17–27.
 44. Franken DR, Bastiaan HS, Oehninger SC. Physiological induction of the acrosome reaction in human sperm: Validation of a microassay using minimal volumes of solubilized, homologous zona pellucida. *J Assist Reprod Genet*. 2000;17:156–61.
 45. Rath D, Töpfer-Petersen E, Michelmann H-W, Schwartz P, Ebeling S. Zona pellucida characteristics and sperm-binding patterns of *in vivo* and *in vitro* produced porcine oocytes inseminated with differently prepared spermatozoa. *Theriogenology*. 2005;63:352–62.
 46. Li X, Hamano KI, Qian XQ, Funauchi K, Furudate M, Minato Y. Oocyte activation and parthenogenetic development of bovine oocytes following intracytoplasmic sperm injection. *Zygote*. 1999;7:233–7.
 47. Li X, Morris LHA, Allen WR. Effects of different activation treatments on fertilization of horse oocytes by intracytoplasmic sperm injection. *J Reprod Fertil*. 2000;119:253–60.
 48. Gadea J, Molla M, Selles E, Marco MA, Garcia-Vazquez FA, Gardon JC. Reduced glutathione content in human sperm is decreased after cryopreservation: Effect of the addition of reduced glutathione to the freezing and thawing extenders. *Cryobiology*. 2011;62:40–6.
 49. Thomson L, Fleming S, Aitken R, De Lullis G, Zieschang JA, Clark A. Cryopreservation-induced human sperm DNA damage is predominantly mediated by oxidative stress rather than apoptosis. *Hum Reprod*. 2009;24:2061–70.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

