

Hindawi Publishing Corporation  
Journal of Inequalities and Applications  
Volume 2008, Article ID 397340, 15 pages  
doi:10.1155/2008/397340

## Research Article

# The Reverse Hölder Inequality for the Solution to $p$ -Harmonic Type System

Zhenhua Cao,<sup>1</sup> Gejun Bao,<sup>1</sup> Ronglu Li,<sup>1</sup> and Haijing Zhu<sup>2</sup>

<sup>1</sup> Department of Mathematics, Harbin Institute of Technology, Harbin 150001, China

<sup>2</sup> College of Mathematics and Physics, Shan Dong Institute of Light Industry, Jinan 250353, China

Correspondence should be addressed to Gejun Bao, baogi@hit.edu.cn

Received 6 July 2008; Revised 9 September 2008; Accepted 5 November 2008

Recommended by Shusen Ding

Some inequalities to  $A$ -harmonic equation  $A(x, du) = d^*v$  have been proved. The  $A$ -harmonic equation is a particular form of  $p$ -harmonic type system  $A(x, a + du) = b + d^*v$  only when  $a = 0$  and  $b = 0$ . In this paper, we will prove the Poincaré inequality and the reverse Hölder inequality for the solution to the  $p$ -harmonic type system.

Copyright © 2008 Zhenhua Cao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Recently, amount of work about the  $A$ -harmonic equation for the differential forms has been done. In fact, the  $A$ -harmonic equation is an important generalization of the  $p$ -harmonic equation in  $\mathbb{R}^n$ ,  $p > 1$ , and the  $p$ -harmonic equation is a natural extension of the usual Laplace equation (see [1] for the details). The reverse Hölder inequalities have been widely studied and frequently used in analysis and related fields, including partial differential equations and the theory of elasticity (see [2]). In 1999, Nolder gave the reverse Hölder inequality for the solution to the  $A$ -harmonic equation in [3], and different versions of Caccioppoli estimates have been established in [4–6]. In 2004, D’Onofrio and Iwaniec introduced the  $p$ -harmonic type system in [7], which is an important extension of the conjugate  $A$ -harmonic equation. In 2007, Ding proved the following inequality in [8].

**Theorem A.** *Let  $(u, v)$  be a pair of solutions to  $A(x, g + du) = h + d^*v$  in a domain  $\Omega \subset \mathbb{R}^n$ . If  $g \in L^p(B, \Lambda^L)$  and  $h \in L^q(B, \Lambda^L)$ , then  $du \in L^p(B, \Lambda^L)$  if and only if  $d^*v \in L^q(B, \Lambda^L)$ . Moreover, there exist constants  $C_1, C_2$  independent of  $u$  and  $v$ , such that*

$$\begin{aligned} \|d^*v\|_{q,B}^q &\leq C_1 (\|h\|_{q,B}^q + \|g\|_{p,B}^p + \|du\|_{p,B}^p), \\ \|du\|_{p,B}^p &\leq C_2 (\|h\|_{q,B}^q + \|g\|_{p,B}^p + \|d^*v\|_{q,B}^q) \quad \forall B \subset \sigma B \subset \Omega. \end{aligned} \quad (1.1)$$

In this paper, we will prove the Poincaré inequality (see Theorem 2.5) and the reverse Hölder inequality for the solution to the  $p$ -harmonic type system (see Theorem 3.5). Now let us see some notions and definitions about the  $p$ -harmonic type system.

Let  $e_1, e_2, \dots, e_n$  denote the standard orthogonal basis of  $\mathbb{R}^n$ . For  $l = 0, 1, \dots, n$ , we denote by  $\Lambda^l = \Lambda^l(\mathbb{R}^n)$  the linear space of all  $l$ -vectors, spanned by the exterior product  $e_I = e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_l}$  corresponding to all ordered  $l$ -tuples  $I = (i_1, i_2, \dots, i_l)$ ,  $1 \leq i_1 < i_2 < \dots < i_l \leq n$ . The Grassmann algebra  $\Lambda = \oplus \Lambda^l$  is a graded algebra with respect to the exterior products. For  $\alpha = \sum \alpha_I e_I \in \Lambda$  and  $\beta = \sum \beta_I e_I \in \Lambda$ , then its inner product is obtained by

$$\langle \alpha, \beta \rangle = \sum \alpha_I \beta_I, \quad (1.2)$$

with the summation over all  $I = (i_1, i_2, \dots, i_l)$  and all integers  $l = 0, 1, \dots, n$ . The Hodge star operator  $*$ :  $\Lambda \rightarrow \Lambda$  is defined by the rule

$$\begin{aligned} *1 &= e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_n}, \\ \alpha \wedge * \beta &= \beta \wedge * \alpha = \langle \alpha, \beta \rangle (*1) \quad \forall \alpha, \beta \in \Lambda. \end{aligned} \quad (1.3)$$

Hence, the norm of  $\alpha \in \Lambda$  can be given by

$$|\alpha|^2 = \langle \alpha, \alpha \rangle = *(\alpha \wedge * \alpha) \in \Lambda_0 = \mathbb{R}. \quad (1.4)$$

Throughout this paper,  $\Omega \subset \mathbb{R}^n$  is an open subset, for any constant  $\sigma > 1$ ,  $Q$  denotes a cube such that  $Q \subset \sigma Q \subset \Omega$ , where  $\sigma Q$  denotes the cube whose center is as same as  $Q$  and  $\text{diam}(\sigma Q) = \sigma \text{diam} Q$ . We say  $\alpha = \sum \alpha_I e_I \in \Lambda$  is a differential  $l$ -form on  $\Omega$ , if every coefficient  $\alpha_I$  of  $\alpha$  is Schwartz distribution on  $\Omega$ . The space spanned by differential  $l$ -form on  $\Omega$  is denoted by  $D'(\Omega, \Lambda^l)$ . We write  $L^p(\Omega, \Lambda^l)$  for the  $l$ -form  $\alpha = \sum \alpha_I dx_I$  on  $\Omega$  with  $\alpha_I \in L^p(\Omega)$  for all ordered  $l$ -tuple  $I$ . Thus  $L^p(\Omega, \Lambda^l)$  is a Banach space with the norm

$$\|\alpha\|_{p,\Omega} = \left( \int_{\Omega} |\alpha|^p dx \right)^{1/p} = \left( \int_{\Omega} \left( \sum_I |\alpha_I|^2 \right)^{p/2} dx \right)^{1/p}. \quad (1.5)$$

Similarly  $W^{k,p}(\Omega, \Lambda^l)$  denotes those  $l$ -forms on  $\Omega$  with all coefficients in  $W^{k,p}(\Omega)$ . We denote the exterior derivative by

$$d : D'(\Omega, \Lambda^l) \longrightarrow D'(\Omega, \Lambda^{l+1}), \quad \text{for } l = 0, 1, 2, \dots, n, \quad (1.6)$$

and its formal adjoint operator (the Hodge codifferential operator)

$$d^* : D'(\Omega, \Lambda^l) \longrightarrow D'(\Omega, \Lambda^{l-1}). \quad (1.7)$$

The operators  $d$  and  $d^*$  are given by the formulas

$$d\alpha = \sum_I d\alpha_I \wedge dx_I, \quad d^* = (-1)^{nl+1} *d*. \quad (1.8)$$

The following two definitions appear in [7].

*Definition 1.1.* The Hodge system holds:

$$A(x, a + du) = b + d^*v, \quad (1.9)$$

where  $a \in L^p(\Omega, \Lambda^l)$  and  $b \in L^q(\Omega, \Lambda^l)$ , is a  $p$ -harmonic type system if  $A$  is a mapping from  $\Omega \times \Lambda^l$  to  $\Lambda^l$  satisfying

- (1)  $x \rightarrow A(x, \xi)$  is measurable in  $x \in \Omega$  for every  $\xi \in \Lambda^l$ ;
- (2)  $\xi \rightarrow A(x, \xi)$  is continuous in  $\xi \in \Lambda^l$  for almost every  $x \in \Omega$ ;
- (3)  $A(x, t\xi) = t^{p-1}A(x, \xi)$  for every  $t \geq 0$ ;
- (4)  $K \langle A(x, \xi) - A(x, \zeta), \xi - \zeta \rangle \geq |\xi - \zeta|^2(|\xi| + |\zeta|)^{p-2}$ ;
- (5)  $|A(x, \xi) - A(x, \zeta)| \leq K|\xi - \zeta|(|\xi| + |\zeta|)^{p-2}$

for almost every  $x \in \Omega$  and all  $\xi, \zeta \in \Lambda^l$ , where  $K \geq 1$  is a constant. It should be noted that  $A(x, *) : \Omega \times \Lambda^l \rightarrow \Lambda^l$  is invertible and its inverse denoted by  $A^{-1}$  satisfies similar conditions as  $A$  but with Hölder conjugate exponent  $q$  in place of  $p$ .

*Definition 1.2.* If (1.9) is a  $p$ -harmonic type system, then we say the equation

$$d^*A(x, a + du) = d^*b \quad (1.10)$$

is a  $p$ -harmonic type equation.

The following definition appears in [9].

*Definition 1.3.* A differential form  $u$  is a weak solution for (1.10) in  $\Omega$  if  $u$  satisfies

$$\int_{\Omega} \langle A(x, a + du), d\varphi \rangle + \langle d^*b, \varphi \rangle \equiv 0 \quad (1.11)$$

for every  $\varphi \in W^{k,p}(\Omega, \Lambda^{l-1})$  with compact support.

We can find that if we let  $a = 0$  and  $b = 0$ , then the  $p$ -harmonic type system

$$A(x, a + du) = b + d^*v \quad (1.12)$$

becomes

$$A(x, du) = d^*v. \quad (1.13)$$

It is the conjugate  $A$ -harmonic equation, where the mapping  $A : \Omega \times \Lambda^l \rightarrow \Lambda^l$  satisfies the following conditions:

$$|A(x, \xi)| \leq a|\xi|^{p-1}, \quad \langle A(x, \xi), \xi \rangle \geq |\xi|^p. \quad (1.14)$$

If we let  $A(x, \xi) = |\xi|^{p-2}\xi$ , then the conjugate  $A$ -harmonic equation becomes the form

$$|du|^{p-2}du = d^*v. \quad (1.15)$$

It is the conjugate  $p$ -harmonic equation.

So we can see that the conjugate  $p$ -harmonic equation and the conjugate  $A$ -harmonic equation are the specific  $p$ -harmonic type system.

*Remark 1.4.* It should be noted that the mapping  $A(x, *)$  in  $p$ -harmonic system  $A(x, a + du) = b + d^*v$ , is invertible. If we denote its inverse by  $A^{-1}(x, *)$ , then the mapping  $A^{-1}(x, *) : \Lambda^l \rightarrow \Lambda^l$  satisfies similar conditions as  $A$  but with Hölder conjugate exponent  $q$  in place of  $p$ .

## 2. The Poincaré inequality

In this section, we will introduce the Poincaré inequality for the differential forms.

Now first let us see a lemma, which can be found in [9, Section 4] for the details.

**Lemma 2.1.** *Let  $\mathbb{D}$  be a bounded, convex domain in  $\mathbb{R}^n$ . To each  $\mathbf{y} \in \mathbb{D}$  there corresponds a linear operator  $K_{\mathbf{y}} : C^\infty(\mathbb{D}, \Lambda^l) \rightarrow C^\infty(\mathbb{D}, \Lambda^{l-1})$  defined by*

$$(K_{\mathbf{y}}\omega)(x; \xi_1, \dots, \xi_{l-1}) = \int_0^1 t^{l-1} \omega(tx + \mathbf{y} - t\mathbf{y}; x - \mathbf{y}, \xi_1, \dots, \xi_{l-1}) dt, \quad (2.1)$$

and the decomposition

$$\omega = d(K_{\mathbf{y}}\omega) + K_{\mathbf{y}}(d\omega) \quad (2.2)$$

holds at any point  $\mathbf{y} \in \mathbb{D}$ .

We construct a homotopy operator  $T : C^\infty(\mathbb{D}, \Lambda^l) \rightarrow C^\infty(\mathbb{D}, \Lambda^{l-1})$  by averaging  $K_{\mathbf{y}}$  over all points  $\mathbf{y} \in \mathbb{D}$ :

$$T\omega = \int_{\mathbb{D}} \varphi(\mathbf{y}) K_{\mathbf{y}}\omega d\mathbf{y}, \quad (2.3)$$

where  $\varphi$  form  $C^\infty(\mathbb{D})$  is normalized so that  $\int \varphi(\mathbf{y}) d\mathbf{y} = 1$ . It is obvious that  $\omega = d(K_{\mathbf{y}}\omega) + K_{\mathbf{y}}(d\omega)$  remains valid for the operator  $T$ :

$$\omega = d(T\omega) + T(d\omega). \quad (2.4)$$

We define the  $l$ -forms  $\omega_{\mathbb{D}} \in D'(\mathbb{D}, \Lambda^l)$  by  $\omega_{\mathbb{D}} = |\mathbb{D}|^{-1} \int_{\mathbb{D}} \omega(\mathbf{y}) d\mathbf{y}$  for  $l = 0$  and  $\omega_{\mathbb{D}} = d(T\omega)$  for  $l = 1, 2, \dots, n$ , and all  $\omega \in W^{1,p}(D, \Lambda^l)$ ,  $1 < p < \infty$ .

The following definition can be found in [9, page 34].

**Definition 2.2.** For  $\omega \in D'(\mathbb{D}, \Lambda^l)$ , the vector valued differential form

$$\nabla\omega = \left( \frac{\partial\omega}{\partial x_1}, \dots, \frac{\partial\omega}{\partial x_n} \right) \quad (2.5)$$

consists of differential forms  $\partial\omega/\partial x_i \in D'(\mathbb{D}, \Lambda^l)$ , where the partial differentiation is applied to coefficients of  $\omega$ .

The proof of [9, Proposition 4.1] implies the following inequality.

**Lemma 2.3.** *For any  $\omega \in L^p(\mathbb{D}, \Lambda^l)$ , it holds that*

$$\|\nabla T\omega\|_{p, \mathbb{D}} \leq C(n, p)\|\omega\|_{p, \mathbb{D}} \quad (2.6)$$

for any ball or cube  $\mathbb{D} \in \mathbb{R}^n$ .

The following Poincaré inequality can be found in [2].

**Lemma 2.4.** *If  $u \in W_0^{1,p}(\Omega)$ , then there is a constant  $C = C(n, p) > 0$  such that*

$$\left(\frac{1}{|B|} \int_B |u|^{p\chi} dx\right)^{1/p\chi} \leq Cr \left(\frac{1}{|B|} \int_B |\nabla u|^p dx\right)^{1/p}, \quad (2.7)$$

whenever  $B = B(x_0, r)$  is a ball in  $\mathbb{R}^n$ , where  $n \geq 2$  and  $\chi = 2$  for  $p \geq n$ ,  $\chi = np/(n-p)$  for  $p < n$ .

**Theorem 2.5.** *Let  $u \in D'(\mathbb{D}, \Lambda^l)$ , and  $du \in L^p(\mathbb{D}, \Lambda^{l+1})$ . Then,  $u - u_{\mathbb{D}}$  is in  $L^{p\chi}(\mathbb{D}, \Lambda^l)$  and*

$$\left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |u - u_{\mathbb{D}}|^{p\chi} dx\right)^{1/p\chi} \leq C(n, p, l) \text{diam}(\mathbb{D}) \left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |du|^p dx\right)^{1/p} \quad (2.8)$$

for any ball or cube  $\mathbb{D} \in \mathbb{R}^n$ , where  $\chi = 2$  for  $p \geq n$  and  $\chi = np/(n-p)$  for  $1 < p < n$ .

*Proof.* We know  $T(du) = u - u_{\mathbb{D}}$ . Now we suppose  $u - u_{\mathbb{D}} = T(du) = \sum_I u_I dx_I$ , where  $I = (i_1, \dots, i_{l+1})$  take over all  $l+1$ -tuples. So we have

$$\nabla T(du) = \left(\frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n}\right) = \left(\sum_I \frac{\partial u_I}{\partial x_1} dx_I, \dots, \sum_I \frac{\partial u_I}{\partial x_n} dx_I\right). \quad (2.9)$$

So we have

$$\begin{aligned} \left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |u - u_{\mathbb{D}}|^{p\chi} dx\right)^{1/p\chi} &= \left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} \left|\sum_I u_I dx_I\right|^{p\chi} dx\right)^{1/p\chi} \\ &= \left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} \left(\sum_I |u_I|^2\right)^{p\chi/2} dx\right)^{1/p\chi}. \end{aligned} \quad (2.10)$$

By the inequality

$$\left(\sum_{i=1}^n (a_i)^2\right)^{1/2} \leq \sum_{i=1}^n a_i \leq n^{1/2} \left(\sum_{i=1}^n (a_i)^2\right)^{1/2} \quad (2.11)$$

for any  $a_i \geq 0$ , and the Minkowski inequality, we have

$$\left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} \left(\sum_I |u_I|^2\right)^{pX/2} dx\right)^{1/pX} \leq \sum_I \left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |u_I|^{pX} dx\right)^{1/pX}. \quad (2.12)$$

According to the Poincaré inequality, we have

$$\sum_I \left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |u_I|^{pX} dx\right)^{1/pX} \leq C_1(n, p) \text{diam}(\mathbb{D}) \sum_I \left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |\nabla u_I|^p dx\right)^{1/p}. \quad (2.13)$$

Combining (2.10), (2.12), and (2.13), we can obtain

$$\left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |u - u_{\mathbb{D}}|^{pX} dx\right)^{1/pX} \leq C_1(n, p) \text{diam}(\mathbb{D}) \sum_I \left(\frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |\nabla u_I|^p dx\right)^{1/p}. \quad (2.14)$$

By (2.9) we have

$$\begin{aligned} \|\nabla T du\|_{p, \mathbb{D}} &= \left\| \left( \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n} \right) \right\|_{p, \mathbb{D}} \\ &= \left( \int_{\mathbb{D}} \left| \left( \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n} \right) \right|^p dx \right)^{1/p} \\ &= \left( \int_{\mathbb{D}} \left( \sum_{i=1}^n \left| \frac{\partial u}{\partial x_i} \right|^2 \right)^{p/2} dx \right)^{1/p} \\ &= \left( \int_{\mathbb{D}} \left( \sum_{i=1}^n \sum_I \left| \frac{\partial u_I}{\partial x_i} \right|^2 \right)^{p/2} dx \right)^{1/p} \\ &= \left( \int_{\mathbb{D}} \left( \sum_I \sum_{i=1}^n \left| \frac{\partial u_I}{\partial x_i} \right|^2 \right)^{p/2} dx \right)^{1/p}. \end{aligned} \quad (2.15)$$

Combining (2.11) and (2.15), then we have

$$\begin{aligned}
\|\nabla T du\|_{p,\mathbb{D}} &= \left( \int_{\mathbb{D}} \left( \sum_I \sum_{i=1}^n \left| \frac{\partial u_I}{\partial x_i} \right|^2 \right)^{p/2} dx \right)^{1/p} \\
&\geq (C_n^{(l+1)})^{-1/2} \left( \int_{\mathbb{D}} \left( \sum_I \left( \sum_{i=1}^n \left| \frac{\partial u_I}{\partial x_i} \right|^2 \right)^{1/2} \right)^p dx \right)^{1/p} \\
&\geq (C_n^{(l+1)})^{-1/2} \left( \int_{\mathbb{D}} \sum_I \left( \sum_{i=1}^n \left| \frac{\partial u_I}{\partial x_i} \right|^2 \right)^{p/2} dx \right)^{1/p} \\
&\geq (C_n^{(l+1)})^{-1/2} (C_n^{(l+1)})^{-(p-1)/p} \sum_I \left( \int_{\mathbb{D}} \left( \sum_{i=1}^n \left| \frac{\partial u_I}{\partial x_i} \right|^2 \right)^{p/2} dx \right)^{1/p} \\
&\geq (C_2(n,p,l))^{-1} \sum_I \left( \int_{\mathbb{D}} |\nabla u_I|^p dx \right)^{1/p},
\end{aligned} \tag{2.16}$$

where  $C_2(n,p,l) = (C_n^{(l+1)})^{1/2+(p-1)/p}$ . Now combining (2.14), (2.16), and (2.6), we can get

$$\begin{aligned}
\left( \frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |u - u_{\mathbb{D}}|^p dx \right)^{1/p} &\leq C_1(n,p) \text{diam}(\mathbb{D}) \sum_I \left( \frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |\nabla u_I|^p dx \right)^{1/p} \\
&\leq C_1(n,p,l) C_2(n,p,l) \left( \frac{1}{|\mathbb{D}|} \right)^{1/p} \|\nabla T du\|_{p,\mathbb{D}} \\
&\leq C_3(n,p,l) \text{diam}(\mathbb{D}) \left( \frac{1}{|\mathbb{D}|} \int_{\mathbb{D}} |du|^p dx \right)^{1/p}.
\end{aligned} \tag{2.17}$$

□

### 3. The reverse Hölder inequality

In this section, we will prove the reverse Hölder inequality for the solution of the  $p$ -harmonic type system. Before we prove the reverse Hölder inequality, let us first see some lemmas.

**Lemma 3.1.** *If  $f, g \geq 0$  and for any nonnegative  $\eta \in C_0^\infty(\Omega)$ , it holds*

$$\int_{\Omega} \eta f dx \leq \int_{\Omega} g dx, \tag{3.1}$$

then for any  $h \geq 0$ :

$$\int_{\Omega} \eta f h dx \leq \int_{\Omega} g h dx. \tag{3.2}$$

*Proof.* Let  $\mu$  be a measure in  $X$ ,  $f$  be a nonnegative  $\mu$ -measurable function in a measure space  $X$ , using the standard representation theorem, we have

$$\int_X f^q d\mu = q \int_0^\infty t^{q-1} \mu(x : f(x) > t) dt \quad (3.3)$$

for any  $0 < t < q$ . Now, we let  $\mu(E) = \int_E \eta f dx$  and  $\nu(E) = \int_E g dx$  then, we can obtain

$$\int_\Omega \eta f h dx = \int_0^\infty \int_{h>t} \eta f dx dt \leq \int_0^\infty \int_{h>t} g dx dt = \int_\Omega g h dx. \quad (3.4)$$

So Lemma 3.1 is proved.  $\square$

**Lemma 3.2.** *If  $(u, v)$  is a pair of solution to the  $p$ -harmonic type system (1.9), then it holds*

$$\int_\Omega |\eta da|^p dx \leq C \int_\Omega |(a + du)d\eta|^p dx \quad (3.5)$$

for any nonnegative  $\eta \in C_0^\infty(\Omega)$  and where  $C = (C_n^{l+1})^p$ .

*Proof.* Since  $(u, v)$  is a pair of solutions to  $A(x, a + du) = b + d^*v$ , it is also the solution to  $A^{-1}(x, b + d^*v) = a + du$ , where  $A^{-1}(x, *)$  is the inverse  $A(x, *)$ . Now, we suppose that  $da = \sum_I \omega_I dx_I$  and let  $\varphi_1 = -\sum_I \eta \text{sign}(\omega_I) dx_I$ . By using  $\varphi = \varphi_1$  and  $d\varphi_1 = \sum_I \text{sign}(\omega_I) d\eta \wedge dx_I$  in (1.11), we can obtain

$$\int_\Omega \langle A^{-1}(x, b + d^*v), d\varphi_1 \rangle + \langle da, \varphi_1 \rangle dx \equiv 0. \quad (3.6)$$

That is,

$$\int_\Omega \left\langle da, \sum_I \eta \text{sign}(\omega_I) dx_I \right\rangle dx = \int_\Omega \left\langle A^{-1}(x, b + d^*v), -\sum_I \text{sign}(\omega_I) d\eta \wedge dx_I \right\rangle dx. \quad (3.7)$$

In other words,

$$\int_\Omega \sum_I \eta |\omega_I| dx = \int_\Omega \left\langle A^{-1}(x, b + d^*v), -\sum_I \text{sign}(\omega_I) d\eta \wedge dx_I \right\rangle dx. \quad (3.8)$$

By the elementary inequality

$$\left( \sum_{i=1}^n a_i^2 \right)^{1/2} \leq \sum_{i=1}^n |a_i|, \quad (3.9)$$



we have

$$\begin{aligned} \int_{\Omega} \eta |da| dx &= \int_{\Omega} \eta \left( \sum_I \omega_I^2 \right)^{1/2} dx \leq \int_{\Omega} \sum_I \eta |\omega_I| dx \\ &= \int_{\Omega} \left\langle A^{-1}(x, b + d^*v), -\sum_I \text{sign}(\omega_I) d\eta \wedge dx_I \right\rangle dx. \end{aligned} \quad (3.10)$$

Using the inequality

$$|\langle a, b \rangle| \leq |a| |b|, \quad (3.11)$$

(3.10) becomes

$$\begin{aligned} \int_{\Omega} \eta |da| dx &\leq \int_{\Omega} |A^{-1}(x, b + d^*v)| \left| \sum_I \text{sign}(\omega_I) d\eta \wedge dx_I \right| \\ &\leq \int_{\Omega} |A^{-1}(x, b + d^*v)| \sum_I |\text{sign}(\omega_I)| |d\eta| dx \\ &= C_n^{l+1} \int_{\Omega} |A^{-1}(x, b + d^*v)| |d\eta| dx \\ &= C_n^{l+1} \int_{\Omega} |a + du| |d\eta| dx, \end{aligned} \quad (3.12)$$

where  $I$  takes over all  $(l+1)$ -tuples for  $d\eta \in \Lambda^{l+1}$ , thus it has  $C_n^{l+1}$  numbers at most. Now we let  $f = |da|$  and  $g = C_n^{l+1} |a + du| |d\eta|$ . In the subset  $\{x : f\eta = g\}$ , we have

$$\int_{\{x: f\eta=g\}} |d\eta| dx \leq \int_{\{x: f\eta=g\}} |(a + du)d\eta|^p dx. \quad (3.13)$$

In the subset  $\{x : f\eta \neq g\}$ , let  $h = (|f\eta|^p - |g|^p) / (f\eta - g)$ , then we easily obtain  $h > 0$ . So by Lemma 3.1, we have

$$\int_{\{x: f\eta \neq g\}} h f \eta dx \leq \int_{\{x: f\eta \neq g\}} h g dx. \quad (3.14)$$

That is to say

$$\int_{\{x: f\eta \neq g\}} h (f\eta - g) dx \leq 0, \quad (3.15)$$

that is,

$$\int_{\{x:f\eta \neq g\}} |f\eta|^p dx \leq \int_{f\eta \neq g} |g|^p dx. \quad (3.16)$$

Combining (3.13) and (3.16), we have

$$\int_{\Omega} |f\eta|^p dx \leq \int_{\Omega} |g|^p dx, \quad (3.17)$$

that is,

$$\int_{\Omega} |\eta da|^p dx \leq \int_{\Omega} |C_n^{l+1}(a + du)d\eta|^p dx. \quad (3.18)$$

So Lemma 3.2 is proved.

The following lemma appears in [2]. □

**Lemma 3.3.** *Suppose that  $0 < q < p < s \leq \infty$ ,  $\xi \in \mathbb{R}$ , and that  $B = B(x_0, r)$  is a ball. If a nonnegative function  $v \in L^p(B, d\mu)$  satisfies*

$$\left( \frac{1}{\mu(\lambda B')} \int_{\lambda B'} v^s d\mu \right)^{1/s} \leq C(1 - \lambda)^{\xi} \left( \frac{1}{\mu(B')} \int_{B'} v^p d\mu \right)^{1/p} \quad (3.19)$$

for each ball  $B' = B(x_0, r')$  with  $r' \leq r$  and for all  $0 < \lambda < 1$ , then

$$\left( \frac{1}{\mu(\lambda B)} \int_{\lambda B} v^s d\mu \right)^{1/s} \leq C(1 - \lambda)^{\xi/\theta} \left( \frac{1}{\mu(B)} \int_B v^q d\mu \right)^{1/q} \quad \forall 0 < \lambda < 1. \quad (3.20)$$

Here  $C > 0$  is a constant depending on  $p, q, s$  and  $\theta \in (0, 1)$  is such that  $1/p = \theta/q + (1 - \theta)/s$ .

The following lemma appears in [10].

**Lemma 3.4.** *Let  $(u, v)$  be a pair of solutions of the  $p$ -harmonic type system on domain  $\Omega$ , then we have a constant  $C$  only depending on  $K, n, p$ , and  $l$ , such that*

$$\|\eta du\|_{p,\Omega} \leq C(\|(u - c)d\eta\|_{p,\Omega} + \|\eta a\|_{p,\Omega}), \quad (3.21)$$

where  $c$  is any closed form (i.e.,  $dc = 0$ ) and for any  $\eta \in C_0^\infty(\Omega)$ . Also we have a constant  $C'$  only depending on  $K$ ,  $n$ ,  $q$ , such that

$$\|\eta d^*v\|_{q,\Omega} \leq C'(\|(v - c')d\eta\|_{q,\Omega} + \|\eta b\|_{q,\Omega}), \quad (3.22)$$

where  $c'$  is any coclosed form (i.e.,  $d^*c' = 0$ ) and  $q$  is the conjugate exponent of  $p$ .

**Theorem 3.5.** *If  $(u, v)$  is a pair of solutions to the  $p$ -harmonic type system, then there exists a constant  $C > 0$  dependent on  $K$ ,  $p$ ,  $n$ , and  $l$ , such that*

$$\begin{aligned} \left(\frac{1}{|Q|} \int_Q (|u - u_Q| + \|a\|_{\infty,Q})^s dx\right)^{1/s} &\leq C(1 - \sigma^{-1})^{-t\chi/p(\chi-1)} (\text{diam } Q + 1)^{\chi/(\chi-1)} \\ &\times \left(\frac{1}{|\sigma Q|} \int_{\sigma Q} (|u - u_{\sigma Q}| + \|a\|_{\infty,\sigma Q})^t dx\right)^{1/t} \end{aligned} \quad (3.23)$$

for any  $0 < s$ ,  $t < \infty$ ,  $\sigma > 1$  and all cubes with  $Q \subset \sigma Q \subset \Omega$ , where  $\chi > 1$  is the Poincaré constant.

*Proof.* Suppose that the center of  $Q$  is  $x_0$  and  $\text{diam } Q = r$ ,  $0 < \lambda = \sigma^{-1} < 1$ . Let

$$r_m = \lambda + (1 - \lambda)2^{-m}, \quad m = 0, 1, 2, \dots \quad (3.24)$$

Then  $r_m$  is decreasing and  $\lambda < r_m < 1$ . So we have  $u_Q|_{r_m Q} = u_{r_m Q}$ , for any  $m \in 0, 1, 2, \dots$ . Let  $\eta_m \in C_0^\infty(r_m Q)$  be a nonnegative function such that  $\eta_m = 1$  in  $r_{m+1}Q$ ,  $0 \leq \eta_m \leq 1$  in  $r_m Q - r_{m+1}Q$ .  $|d\eta_m| \leq (1 - \lambda)^{-1}2^m r^{-1}$ . Given any  $t \geq 0$  and let  $\omega_m = (|u - u_Q| + \|a\|_{\infty,Q})^{1+t/p} \eta_m$ , then we have

$$du_m = \left(1 + \frac{t}{p}\right) (|u - u_Q| + \|a\|_{\infty,Q})^{t/p} \eta_m d|u - u_Q| + (|u - u_Q| + \|a\|_{\infty,Q})^{1+t/p} d\eta_m. \quad (3.25)$$

By the Minkowski inequality, we can obtain

$$\begin{aligned} \left(\int_{r_m Q} |du_m|^p dx\right)^{1/p} &\leq \left(\int_{r_m Q} (|u - u_Q| + \|a\|_{\infty,Q})^{p+t} |d\eta_m|^p dx\right)^{1/p} \\ &+ \frac{(p+t)}{p} \left(\int_{r_m Q} |d|u - u_Q||^p (|u - u_Q| + \|a\|_{\infty,Q})^t |\eta_m|^p dx\right)^{1/p}. \end{aligned} \quad (3.26)$$

We assume that  $u - u_Q = \sum_I a_I dx_I$ , then we have  $|u - u_Q| = (\sum_I a_I^2)^{1/2}$ . If  $u - u_Q$  is zero, then we have  $|d|u - u_Q|| = 0 = |\nabla T(du)|$ . If  $u - u_Q$  is not equal zero, and the proof of (2.15) implies

that  $|\nabla T du| = (\sum_I \sum_{i=1}^n |\partial a_I / \partial x_i|^2)^{1/2}$

$$\begin{aligned}
 |d|u - u_Q|| &= |\nabla|u - u_Q|| = \left| \left( \frac{\partial|u - u_Q|}{\partial x_1}, \dots, \frac{\partial|u - u_Q|}{\partial x_n} \right) \right| \\
 &= \left( \sum_{i=1}^n \left| \frac{\partial|u - u_Q|}{\partial x_i} \right|^2 \right)^{1/2} = \left( \sum_{i=1}^n \left| \frac{\partial|u - u_Q|}{\partial x_i} \right|^2 \right)^{1/2} \\
 &= \left( \sum_{i=1}^n \left| \frac{\partial(\sum_I a_I^2)^{1/2}}{\partial x_i} \right|^2 \right)^{1/2} = \left( \sum_{i=1}^n \frac{1}{\sum_I a_I^2} \left| \sum_I a_I \frac{\partial a_I}{\partial x_i} \right|^2 \right)^{1/2} \quad (3.27) \\
 &\leq \left( \sum_{i=1}^n \frac{1}{\sum_I a_I^2} \sum_I a_I^2 \sum_I \left( \frac{\partial a_I}{\partial x_i} \right)^2 \right)^{1/2} = \left( \sum_{i=1}^n \sum_I \left( \frac{\partial a_I}{\partial x_i} \right)^2 \right)^{1/2} \\
 &= \left( \sum_{i=1}^n \sum_I \left| \frac{\partial a_I}{\partial x_i} \right|^2 \right)^{1/2} = |\nabla T(du)| = |\nabla(u - u_Q)|.
 \end{aligned}$$

So we have

$$|d|u - u_Q|| \leq |\nabla T(du)|. \quad (3.28)$$

For any  $\eta \in C_0^\infty(\Omega)$ , according to (2.6), we have

$$\|\eta \nabla T dw\|_{p, \mathbb{D}} \leq C(n, p) \max_{x \in \mathbb{D}}(\eta) \|dw\|_{p, \mathbb{D}}. \quad (3.29)$$

By the similar method as Lemma 3.1, we can prove the following inequality:

$$\begin{aligned}
 &\left( \int_{r_m Q} |d|u - u_Q||^p (|u - u_Q| + \|a\|_{\infty, Q})^t |\eta_m|^p dx \right)^{1/p} \\
 &\leq \left( \int_{r_m Q} |\eta_m|^p |\nabla T(du)|^p (|u - u_Q| + \|a\|_{\infty, Q})^t dx \right)^{1/p} \quad (3.30) \\
 &\leq C(n, p) \max_{x \in \mathbb{D}}(\eta_m^p) \left( \int_{r_m Q} |\eta_m|^p |du|^p (|u - u_Q| + \|a\|_{\infty, Q})^t dx \right)^{1/p}
 \end{aligned}$$

for any  $\eta \in C_0^\infty(\Omega)$ . By Lemma 3.1 and (3.21), we can obtain

$$\begin{aligned}
& \left( \int_{r_m Q} |\eta_m|^p |du|^p (|u - u_Q| + \|a\|_{\infty, Q})^t dx \right)^{1/p} \\
& \leq 2C \left( \int_{r_m Q} |\eta_m|^p |a|^p (|u - u_Q| + \|a\|_{\infty, Q})^t dx \right)^{1/p} \\
& \quad + 2C \left( \int_{r_m Q} |d\eta_m|^p (|u - u_Q| + \|a\|_{\infty, Q})^{p+t} dx \right)^{1/p} \\
& \leq 2C \left( \int_{r_m Q} |\eta_m|^p \|a\|_{\infty, Q}^p (|u - u_Q| + \|a\|_{\infty, Q})^t dx \right)^{1/p} \\
& \quad + 2C \left( \int_{r_m Q} |d\eta_m|^p (|u - u_Q| + \|a\|_{\infty, Q})^{p+t} dx \right)^{1/p} \\
& \leq 2C \left( \int_{r_m Q} |\eta_m|^p (|u - u_Q| + \|a\|_{\infty, Q})^{p+t} dx \right)^{1/p} \\
& \quad + 2C \left( \int_{r_m Q} |d\eta_m|^p (|u - u_Q| + \|a\|_{\infty, Q})^{p+t} dx \right)^{1/p}.
\end{aligned} \tag{3.31}$$

Combining (3.26), (3.30), and (3.31), by the values of  $\eta_m$ , we have

$$\left( \int_{r_m Q} |du_m|^p dx \right)^{1/p} \leq C_1(p+t)(1 + (1-\lambda)^{-1}2^m r^{-1}) \left( \int_{r_m Q} (|u - u_Q| + \|a\|_{\infty, Q})^{p+t} dx \right)^{1/p}. \tag{3.32}$$

For  $\eta_m = 1$  in  $r_{m+1}Q$  and  $0 \leq \eta_m \leq 1$  in  $r_m Q - r_{m+1}Q$ , and as we have  $|r_m|/r_{m+1} = |\lambda + (1-\lambda)2^{-m}|/(\lambda + (1-\lambda)2^{-m-1}) \leq 2$ , so we have  $|r_m Q|/|r_{m+1}Q| \leq 2^n$ . By the Poincaré inequality, we know

$$\begin{aligned}
& \left( \frac{1}{|r_{m+1}Q|} \int_{r_{m+1}Q} (|u - u_Q| + \|a\|_{\infty, Q})^{X(p+t)} dx \right)^{1/pX} \\
& \leq \frac{1}{|r_{m+1}Q|} \int_{r_m Q} (\eta_m^{pX} |u - u_Q| + \|a\|_{\infty, Q})^{X(p+t)} dx)^{1/pX} \\
& \leq \left( \frac{1}{|r_{m+1}Q|} \int_{r_m Q} |u_m|^{pX} dx \right)^{1/pX} \\
& \leq 2^n \left( \frac{1}{|r_m Q|} \int_{r_m Q} |u_m|^{pX} dx \right)^{1/pX} \\
& \leq C_2 r_m r \left( \frac{1}{|r_m Q|} \int_{r_m Q} |du_m|^p dx \right)^{1/p} \\
& \leq C_3 r_m r (p+t)(1 + (1-\lambda)^{-1}2^m r^{-1}) \left( \int_{r_m Q} (|u - u_Q| + \|a\|_{\infty, Q})^{p+t} dx \right)^{1/p} \\
& \leq C_3(p+t)(1-\lambda)^{-1}2^m(1+r) \left( \int_{r_m Q} (|u - u_Q| + \|a\|_{\infty, Q})^{p+t} dx \right)^{1/p}.
\end{aligned} \tag{3.33}$$

Now we set  $\kappa = p + t$ , then by computation, we obtain

$$\begin{aligned} \left( \frac{1}{|r_{m+1}Q|} \int_{r_{m+1}Q} (|u - u_Q| + \|a\|_{\infty, Q})^{\kappa \chi} dx \right)^{1/\kappa \chi} &\leq (C_3)^{p/\kappa} \kappa^{p/\kappa} (1 - \lambda)^{-p/\kappa} 2^{pm/\kappa} (r + 1)^{p/\kappa} \\ &\times \left( \frac{1}{|r_m Q|} \int_{r_m Q} (|u - u_Q| + \|a\|_{\infty, Q})^{\kappa} dx \right)^{1/\kappa}. \end{aligned} \quad (3.34)$$

Since this inequality holds for all  $\kappa > p$ , it can be applied with  $\kappa = \kappa_m = p\chi^m$ . And we can easily prove  $((1/|Q|) \int_Q |f|^p dx)^{1/p}$  is increasing with  $p$  and its limit is  $\text{ess sup}_Q |f|$ . So by iterating we arrive at the desired inequality for  $q = p$ :

$$\begin{aligned} &\text{ess sup}_{\lambda Q} (|u - u_Q| + \|a\|_{\infty, Q}) \\ &\leq \lim_{m \rightarrow \infty} \left( \frac{1}{|r_m Q|} \int_{r_m Q} (|u - u_Q| + \|a\|_{\infty, Q})^{\kappa_m \chi} dx \right)^{1/\kappa_m \chi} \\ &\leq C_4 ((1 - \lambda)^{-1} (r + 1))^{\sum_{i=0}^{\infty} \chi^{-m}} \prod_{m=0}^{\infty} 2^{m\chi^{-m}} \prod_{m=0}^{\infty} (p\chi^m)^{\chi^{-m}} \\ &\quad \times \left( \frac{1}{|Q|} \int_Q (|u - u_Q| + \|a\|_{\infty, Q})^p dx \right)^{1/p} \\ &\leq C_5 (1 - \lambda)^{-\chi/(\chi-1)} (r + 1)^{\chi/(\chi-1)} \left( \frac{1}{|Q|} \int_Q (|u - u_Q| + \|a\|_{\infty, Q})^p dx \right)^{1/p}. \end{aligned} \quad (3.35)$$

We can observe that the constants  $C_5$  and  $\chi$  are independent of  $x_0$  and  $r$  in (3.35), thus (3.35) holds not only in the cube  $Q = Q(x_0, r)$  but also in each ball inside  $Q$ . By Lemma (3.5) we can obtain

$$\begin{aligned} \left( \frac{1}{|\lambda Q|} \int_{\lambda Q} (|u - u_Q| + \|a\|_{\infty, Q})^s dx \right)^{1/s} &\leq C_5 (1 - \lambda)^{-\theta \chi/(\chi-1)} (r + 1)^{\chi/(\chi-1)} \\ &\quad \times \left( \frac{1}{|Q|} \int_Q (|u - u_Q| + \|a\|_{\infty, Q})^t dx \right)^{1/t} \end{aligned} \quad (3.36)$$

for any  $0 < t < p < s \leq \infty$ , where  $\theta = t(s - p)/p(s - t)$ . So we have  $\theta \leq t/p$  for any  $0 < t < p < s \leq \infty$ . Since  $((1/|Q|) \int_Q |f|^p dx)^{1/p}$  is increasing with  $p$ ,

$$\begin{aligned} \left( \frac{1}{|\lambda Q|} \int_{\lambda Q} (|u - u_Q| + \|a\|_{\infty, Q})^s dx \right)^{1/s} &\leq C_5 (1 - \lambda)^{-t\chi/p(\chi-1)} (r + 1)^{\chi/(\chi-1)} \\ &\quad \times \left( \frac{1}{|Q|} \int_Q (|u - u_Q| + \|a\|_{\infty, Q})^t dx \right)^{1/t} \end{aligned} \quad (3.37)$$

for any  $0 < s < \infty$  and  $1 < p < t < \infty$ . Combining (3.36) and (3.37), we have

$$\begin{aligned} \left( \frac{1}{|\Omega|} \int_{\Omega} (|u - u_{\Omega}| + \|a\|_{\infty, \Omega})^s dx \right)^{1/s} &\leq C_6 (1 - \lambda)^{-tX/p(X-1)} (r + 1)^{X/(X-1)} \\ &\times \left( \frac{1}{|\sigma Q|} \int_{\sigma Q} (|u - u_{\sigma Q}| + \|a\|_{\infty, \sigma Q})^t dx \right)^{1/t} \end{aligned} \quad (3.38)$$

for any  $0 < s, t < \infty$  and  $\sigma > 1$  such that  $\sigma Q \subset \Omega$ . Theorem 3.5 is proved.  $\square$

### Acknowledgment

This work is supported by the NSF of China (Grants nos. 10671046 and 10771044).

### References

- [1] S. Ding, "Some examples of conjugate  $p$ -harmonic differential forms," *Journal of Mathematical Analysis and Applications*, vol. 227, no. 1, pp. 251–270, 1998.
- [2] J. Heinonen, T. Kilpeläinen, and O. Martio, *Nonlinear Potential Theory of Degenerate Elliptic Equations*, Oxford Mathematical Monographs, Oxford University Press, New York, NY, USA, 1993.
- [3] C. A. Nolder, "Hardy-Littlewood theorems for  $A$ -harmonic tensors," *Illinois Journal of Mathematics*, vol. 43, no. 4, pp. 613–632, 1999.
- [4] G. Bao, " $A_r(\lambda)$ -weighted integral inequalities for  $A$ -harmonic tensors," *Journal of Mathematical Analysis and Applications*, vol. 247, no. 2, pp. 466–477, 2000.
- [5] S. Ding, "Weighted Caccioppoli-type estimates and weak reverse Hölder inequalities for  $A$ -harmonic tensors," *Proceedings of the American Mathematical Society*, vol. 127, no. 9, pp. 2657–2664, 1999.
- [6] X. Yuming, "Weighted integral inequalities for solutions of the  $A$ -harmonic equation," *Journal of Mathematical Analysis and Applications*, vol. 279, no. 1, pp. 350–363, 2003.
- [7] L. D’Onofrio and T. Iwaniec, "The  $p$ -harmonic transform beyond its natural domain of definition," *Indiana University Mathematics Journal*, vol. 53, no. 3, pp. 683–718, 2004.
- [8] S. Ding, "Local and global norm comparison theorems for solutions to the nonhomogeneous  $A$ -harmonic equation," *Journal of Mathematical Analysis and Applications*, vol. 335, no. 2, pp. 1274–1293, 2007.
- [9] T. Iwaniec and A. Lutoborski, "Integral estimates for null Lagrangians," *Archive for Rational Mechanics and Analysis*, vol. 125, no. 1, pp. 25–79, 1993.
- [10] Z. Cao, G. Bao, and R. Li, "The Caccioppoli estimate for the solution to the  $p$ -harmonic type system," to appear in the Proceedings of the 6th International Conference on Differential Equations and Dynamical Systems.