

RESEARCH

Open Access

Sum of squared logarithms - an inequality relating positive definite matrices and their matrix logarithm

Mircea Birsan^{1,2}, Patrizio Neff^{1*} and Johannes Lankeit¹

*Correspondence:

patrizio.neff@uni-due.de
¹Lehrstuhl für Nichtlineare Analysis und Modellierung, Fakultät für Mathematik, Universität Duisburg-Essen, Essen, Germany
Full list of author information is available at the end of the article**Abstract**Let $y_1, y_2, y_3, a_1, a_2, a_3 \in (0, \infty)$ be such that $y_1 y_2 y_3 = a_1 a_2 a_3$ and

$$y_1 + y_2 + y_3 \geq a_1 + a_2 + a_3, \quad y_1 y_2 + y_2 y_3 + y_1 y_3 \geq a_1 a_2 + a_2 a_3 + a_1 a_3.$$

Then

$$(\log y_1)^2 + (\log y_2)^2 + (\log y_3)^2 \geq (\log a_1)^2 + (\log a_2)^2 + (\log a_3)^2.$$

This can also be stated in terms of real positive definite 3×3 -matrices P_1, P_2 : If their determinants are equal, $\det P_1 = \det P_2$, then

$$\operatorname{tr} P_1 \geq \operatorname{tr} P_2 \quad \text{and} \quad \operatorname{tr} \operatorname{Cof} P_1 \geq \operatorname{tr} \operatorname{Cof} P_2 \quad \implies \quad \| \log P_1 \|_F^2 \geq \| \log P_2 \|_F^2,$$

where \log is the principal matrix logarithm and $\|P\|_F^2 = \sum_{i,j=1}^3 P_{ij}^2$ denotes the Frobenius matrix norm. Applications in matrix analysis and nonlinear elasticity are indicated.**MSC:** 26D05; 26D07**Keywords:** matrix logarithm; elementary symmetric polynomials; inequality; characteristic polynomial; positive definite matrices; means

1 Introduction

Convexity is a powerful source for obtaining new inequalities; see, e.g., [1, 2]. In applications coming from nonlinear elasticity, we are faced, however, with variants of the squared logarithm function; see the last section. The function $(\log(x))^2$ is neither convex nor concave. Nevertheless, the sum of squared logarithms inequality holds. We will proceed as follows: In the first section, we will give several equivalent formulations of the inequality, for example, in terms of the coefficients of the characteristic polynomial (Theorem 1), in terms of elementary symmetric polynomials (Theorem 3), in terms of means (Theorem 5) or in terms of the Frobenius matrix norm (Theorem 7). A proof of the inequality will be given in Section 2, and some counterexamples for slightly changed variants of the inequality are discussed in Section 3. In the last section, an application of the sum of squared loga-

rhythms inequality in matrix analysis and in the mathematical theory of nonlinear elasticity is indicated.

2 Formulations of the problem

All theorems in this section are equivalent.

Theorem 1 For $n = 2$ or $n = 3$ let $P_1, P_2 \in \mathbb{R}^{n \times n}$ be positive definite real matrices. Let the coefficients of the characteristic polynomials of P_1 and P_2 satisfy

$$\operatorname{tr} P_1 \geq \operatorname{tr} P_2 \quad \text{and} \quad \operatorname{tr} \operatorname{Cof} P_1 \geq \operatorname{tr} \operatorname{Cof} P_2 \quad \text{and} \quad \det P_1 = \det P_2.$$

Then

$$\|\log P_1\|_F^2 \geq \|\log P_2\|_F^2.$$

For $n = 3$, we will now give equivalent formulations of this statement. The case $n = 2$ can be treated analogously. For its proof, see Remark 15. By orthogonal diagonalization of P_1 and P_2 , the inequalities can be rewritten in terms of the eigenvalues y_1, y_2, y_3 and a_1, a_2, a_3 , respectively.

Theorem 2 Let the real numbers $a_1, a_2, a_3 > 0$ and $y_1, y_2, y_3 > 0$ be such that

$$\begin{aligned} y_1 + y_2 + y_3 &\geq a_1 + a_2 + a_3, \\ y_1 y_2 + y_2 y_3 + y_1 y_3 &\geq a_1 a_2 + a_2 a_3 + a_1 a_3, \\ y_1 y_2 y_3 &= a_1 a_2 a_3. \end{aligned} \tag{1}$$

Then

$$(\log y_1)^2 + (\log y_2)^2 + (\log y_3)^2 \geq (\log a_1)^2 + (\log a_2)^2 + (\log a_3)^2. \tag{2}$$

The elementary symmetric polynomials, see, e.g., [3, p.178]

$$\begin{aligned} e_0(y_1, y_2, y_3) &= 1, \\ e_1(y_1, y_2, y_3) &= y_1 + y_2 + y_3, \\ e_2(y_1, y_2, y_3) &= y_1 y_2 + y_1 y_3 + y_2 y_3, \\ e_3(y_1, y_2, y_3) &= y_1 y_2 y_3 \end{aligned}$$

are known to have the Schur-concavity property (i.e., $-e_k$ is Schur-convex) [1, 4]; see (16). It is possible to express the problem in terms of these elementary symmetric polynomials as follows.

Theorem 3 Let $a_1, a_2, a_3 > 0$ and $y_1, y_2, y_3 > 0$ satisfy

$$\begin{aligned} e_1(y_1, y_2, y_3) &\geq e_1(a_1, a_2, a_3), & e_2(y_1, y_2, y_3) &\geq e_2(a_1, a_2, a_3), \\ e_3(y_1, y_2, y_3) &= e_3(a_1, a_2, a_3). \end{aligned}$$

Then

$$e_1((\log y_1)^2, (\log y_2)^2, (\log y_3)^2) \geq e_1((\log a_1)^2, (\log a_2)^2, (\log a_3)^2).$$

Because $y_1 y_2 y_3 = a_1 a_2 a_3 > 0$, we have

$$y_1 y_2 + y_2 y_3 + y_1 y_3 \geq a_1 a_2 + a_2 a_3 + a_1 a_3 \Leftrightarrow \frac{1}{y_1} + \frac{1}{y_2} + \frac{1}{y_3} \geq \frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3}.$$

Thus, we obtain the following theorem.

Theorem 4 *Let the real numbers $a_1, a_2, a_3 > 0$ and $y_1, y_2, y_3 > 0$ be such that*

$$\begin{aligned} y_1 + y_2 + y_3 &\geq a_1 + a_2 + a_3, \\ \frac{1}{y_1} + \frac{1}{y_2} + \frac{1}{y_3} &\geq \frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3}, \\ y_1 y_2 y_3 &= a_1 a_2 a_3. \end{aligned} \tag{3}$$

Then

$$(\log y_1)^2 + (\log y_2)^2 + (\log y_3)^2 \geq (\log a_1)^2 + (\log a_2)^2 + (\log a_3)^2. \tag{4}$$

The conditions (3) are also simple expressions in terms of arithmetic, harmonic and geometric and quadratic mean

$$\begin{aligned} A(y_1, y_2, y_3) &= \frac{y_1 + y_2 + y_3}{3}, & H(y_1, y_2, y_3) &= \frac{3}{\frac{1}{y_1} + \frac{1}{y_2} + \frac{1}{y_3}}, \\ G(y_1, y_2, y_3) &= \sqrt[3]{y_1 y_2 y_3}, & Q(y_1, y_2, y_3) &= \sqrt{\frac{1}{3}(y_1^2 + y_2^2 + y_3^2)}. \end{aligned}$$

Theorem 5 *Let $a_1, a_2, a_3 > 0$ and $y_1, y_2, y_3 > 0$. Then $A(y_1, y_2, y_3) \geq A(a_1, a_2, a_3)$, $H(a_1, a_2, a_3) \geq H(y_1, y_2, y_3)$ ('reverse!') and $G(y_1, y_2, y_3) = G(a_1, a_2, a_3)$ imply*

$$Q(\log y_1, \log y_2, \log y_3) \geq Q(\log a_1, \log a_2, \log a_3).$$

We denote by

$$a_i =: d_i^2, \quad y_i =: x_i^2$$

and arrive at

Theorem 6 *Let the real numbers d_i and x_i be such that $d_1, d_2, d_3 > 0$, $x_1, x_2, x_3 > 0$ and*

$$\begin{aligned} x_1^2 + x_2^2 + x_3^2 &\geq d_1^2 + d_2^2 + d_3^2, \\ x_1^2 x_2^2 + x_2^2 x_3^2 + x_1^2 x_3^2 &\geq d_1^2 d_2^2 + d_2^2 d_3^2 + d_1^2 d_3^2, \\ x_1 x_2 x_3 &= d_1 d_2 d_3. \end{aligned} \tag{5}$$

Then

$$(\log x_1)^2 + (\log x_2)^2 + (\log x_3)^2 \geq (\log d_1)^2 + (\log d_2)^2 + (\log d_3)^2. \tag{6}$$

If we again view x_i and d_i as eigenvalues of positive definite matrices, an equivalent formulation of the problem can be given in terms of their Frobenius matrix norms:

Theorem 7 For $n \in \{2, 3\}$, let $P_1, P_2 \in \mathbb{R}^{n \times n}$ be positive definite real matrices. Let

$$\|P_1\|_F^2 \geq \|P_2\|_F^2 \quad \text{and} \quad \|P_1^{-1}\|_F^2 \geq \|P_2^{-1}\|_F^2 \quad \text{and} \quad \det P_1 = \det P_2.$$

Then

$$\|\log P_1\|_F^2 \geq \|\log P_2\|_F^2.$$

Let us reconsider the formulation from Theorem 5. If we denote

$$c_i := \log a_i, \quad z_i := \log y_i,$$

from $H(a_1, a_2, a_3) \geq H(y_1, y_2, y_3)$, we obtain

$$e^{-z_1} + e^{-z_2} + e^{-z_3} \geq e^{-c_1} + e^{-c_2} + e^{-c_3}.$$

Theorem 8 Let the real numbers c_1, c_2, c_3 and z_1, z_2, z_3 be such that

$$\begin{aligned} e^{z_1} + e^{z_2} + e^{z_3} &\geq e^{c_1} + e^{c_2} + e^{c_3}, \\ e^{-z_1} + e^{-z_2} + e^{-z_3} &\geq e^{-c_1} + e^{-c_2} + e^{-c_3}, \\ z_1 + z_2 + z_3 &= c_1 + c_2 + c_3. \end{aligned} \tag{7}$$

Then

$$z_1^2 + z_2^2 + z_3^2 \geq c_1^2 + c_2^2 + c_3^2. \tag{8}$$

In order to prove Theorem 8, one can assume without loss of generality that

$$z_1 + z_2 + z_3 = c_1 + c_2 + c_3 = 0. \tag{9}$$

Thus, we have the equivalent formulation

Theorem 9 Let the real numbers $\bar{c}_1, \bar{c}_2, \bar{c}_3$ and $\bar{z}_1, \bar{z}_2, \bar{z}_3$ be such that

$$\begin{aligned} e^{\bar{z}_1} + e^{\bar{z}_2} + e^{\bar{z}_3} &\geq e^{\bar{c}_1} + e^{\bar{c}_2} + e^{\bar{c}_3}, \\ e^{-\bar{z}_1} + e^{-\bar{z}_2} + e^{-\bar{z}_3} &\geq e^{-\bar{c}_1} + e^{-\bar{c}_2} + e^{-\bar{c}_3}, \\ \bar{z}_1 + \bar{z}_2 + \bar{z}_3 &= \bar{c}_1 + \bar{c}_2 + \bar{c}_3 = 0. \end{aligned} \tag{10}$$

Then

$$\bar{z}_1^2 + \bar{z}_2^2 + \bar{z}_3^2 \geq \bar{c}_1^2 + \bar{c}_2^2 + \bar{c}_3^2. \tag{11}$$

Let us prove that Theorem 8 can be reformulated as Theorem 9. Indeed, let us assume that Theorem 9 is valid and show that the statement of Theorem 8 also holds true. We denote by s the sum $s = z_1 + z_2 + z_3 = c_1 + c_2 + c_3$ and we designate

$$\bar{z}_i = z_i - \frac{s}{3}, \quad \bar{c}_i = c_i - \frac{s}{3} \quad (i = 1, 2, 3).$$

Then the real numbers \bar{z}_i and \bar{c}_i satisfy the hypotheses of Theorem 9 and we obtain $\bar{z}_1^2 + \bar{z}_2^2 + \bar{z}_3^2 \geq \bar{c}_1^2 + \bar{c}_2^2 + \bar{c}_3^2$. This inequality is equivalent to

$$\sum_{i=1}^3 \left(z_i - \frac{s}{3} \right)^2 \geq \sum_{i=1}^3 \left(c_i - \frac{s}{3} \right)^2,$$

which, by virtue of the condition $(7)_3$, reduces to

$$z_1^2 + z_2^2 + z_3^2 \geq c_1^2 + c_2^2 + c_3^2.$$

Thus, Theorem 8 is also valid.

By virtue of the logical equivalence

$$(A \wedge B \Rightarrow C) \Leftrightarrow (\neg C \Rightarrow \neg A \vee \neg B)$$

for any statements A, B, C , we can formulate the inequality (11) (*i.e.*, Theorem 9) in the following equivalent manner.

Theorem 10 *Let the real numbers c_1, c_2, c_3 and z_1, z_2, z_3 be such that*

$$z_1 + z_2 + z_3 = c_1 + c_2 + c_3 = 0 \quad \text{and} \quad z_1^2 + z_2^2 + z_3^2 < c_1^2 + c_2^2 + c_3^2. \tag{12}$$

Then one of the following inequalities holds:

$$\begin{aligned} e^{z_1} + e^{z_2} + e^{z_3} &< e^{c_1} + e^{c_2} + e^{c_3} \quad \text{or} \\ e^{-z_1} + e^{-z_2} + e^{-z_3} &< e^{-c_1} + e^{-c_2} + e^{-c_3}. \end{aligned} \tag{13}$$

We use the statement of Theorem 10 for the proof.

Before continuing, let us show that our new inequality is not a consequence of majorization and Karamata's inequality [5]. Consider $z = (z_1, \dots, z_n) \in \mathbb{R}_+^n$ and $c = (c_1, \dots, c_n) \in \mathbb{R}_+^n$ arranged already in decreasing order $z_1 \geq z_2 \geq \dots \geq z_n$ and $c_1 \geq c_2 \geq \dots \geq c_n$. If

$$\sum_{i=1}^k z_i \geq \sum_{i=1}^k c_i \quad (1 \leq k \leq n-1), \quad \sum_{i=1}^n z_i = \sum_{i=1}^n c_i, \tag{14}$$

we say that z majorizes c , denoted by $z \succ c$. The following result is well known [6, p.89], [4, 5]. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is convex, then

$$z \succ c \Rightarrow \sum_{i=1}^n f(z_i) \geq \sum_{i=1}^n f(c_i). \tag{15}$$

A function $g : \mathbb{R}^n \mapsto \mathbb{R}$ which satisfies

$$z \succ c \Rightarrow g(z_1, \dots, z_n) \geq g(c_1, \dots, c_n) \tag{16}$$

is called Schur-convex. In Theorem 8, the convex function to be considered would be $f(t) = t^2$. Do conditions (7) (upon rearrangement of $z, c \in \mathbb{R}_+^3$ if necessary) yield already majorization $z \succ c$? This is not the case, as we explain now. Let the real numbers $z_1 \geq z_2 \geq z_3$ and $c_1 \geq c_2 \geq c_3$ be such that

$$\begin{aligned} e^{z_1} + e^{z_2} + e^{z_3} &\geq e^{c_1} + e^{c_2} + e^{c_3}, \\ e^{-z_1} + e^{-z_2} + e^{-z_3} &\geq e^{-c_1} + e^{-c_2} + e^{-c_3}, \\ z_1 + z_2 + z_3 &= c_1 + c_2 + c_3. \end{aligned} \tag{17}$$

These conditions do not imply the majorization $z \succ c$,

$$z_1 \geq c_1, \quad z_1 + z_2 \geq c_1 + c_2, \quad z_1 + z_2 + z_3 = c_1 + c_2 + c_3. \tag{18}$$

Therefore, our inequality (i.e., $z_1^2 + z_2^2 + z_3^2 \geq c_1^2 + c_2^2 + c_3^2$) does not follow from majorization in disguise.

Indeed, let

$$z_1 = \frac{1}{2} + \frac{0.95}{2\sqrt{3}}, \quad z_2 = \frac{1}{2} + \frac{0.85}{2\sqrt{3}}, \quad z_3 = -1 - \frac{0.9}{\sqrt{3}}$$

and

$$c_1 = \frac{1}{2} + \frac{1}{2\sqrt{3}}, \quad c_2 = -\frac{1}{2} + \frac{1}{2\sqrt{3}}, \quad c_3 = -\frac{1}{\sqrt{3}}.$$

Then we have $z_1 > z_2 > z_3$ and $c_1 > c_2 > c_3$, together with

$$\begin{aligned} e^{z_1} + e^{z_2} + e^{z_3} &= 4.49497\dots > 3.57137\dots = e^{c_1} + e^{c_2} + e^{c_3}, \\ e^{-z_1} + e^{-z_2} + e^{-z_3} &= 5.50607\dots > 3.47107\dots = e^{-c_1} + e^{-c_2} + e^{-c_3}, \\ z_1 + z_2 + z_3 &= c_1 + c_2 + c_3 = 0, \end{aligned}$$

but the majorization inequalities (18) are not satisfied, since $z_1 < c_1$.

3 Proof of the inequality

Of course, we may assume without loss of generality that $c_1 \geq c_2 \geq c_3$ and $z_1 \geq z_2 \geq z_3$ (and the same for a_i, d_i, x_i, y_i).

The proof begins with the crucial lemma.

Lemma 11 *Let the real numbers $a \geq b \geq c$ and $x \geq y \geq z$ be such that*

$$a + b + c = x + y + z = 0, \quad a^2 + b^2 + c^2 = x^2 + y^2 + z^2. \tag{19}$$

Then the inequality

$$e^a + e^b + e^c \leq e^x + e^y + e^z \tag{20}$$

is satisfied if and only if the relation

$$a \leq x \tag{21}$$

holds, or equivalently, if and only if

$$c \leq z \tag{22}$$

holds.

Proof Let us denote by $r := \sqrt{\frac{2}{3}(a^2 + b^2 + c^2)} > 0$. Then, from (19), it follows

$$\begin{aligned} b + c = -a, \quad b^2 + c^2 &= \frac{3}{2}r^2 - a^2, \\ y + z = -x, \quad y^2 + z^2 &= \frac{3}{2}r^2 - x^2, \end{aligned}$$

and we find

$$\begin{aligned} b &= \frac{1}{2}(-a + \sqrt{3(r^2 - a^2)}), & c &= \frac{1}{2}(-a - \sqrt{3(r^2 - a^2)}), \\ y &= \frac{1}{2}(-x + \sqrt{3(r^2 - x^2)}), & z &= \frac{1}{2}(-x - \sqrt{3(r^2 - x^2)}). \end{aligned} \tag{23}$$

In view of (19) and $a \geq b \geq c$, $x \geq y \geq z$, one can show that

$$a, x \in \left[\frac{r}{2}, r \right], \quad b, y \in \left[-\frac{r}{2}, \frac{r}{2} \right], \quad c, z \in \left[-r, -\frac{r}{2} \right]. \tag{24}$$

Indeed, let us verify the relations (24). We have

$$\begin{aligned} \frac{r}{2} \leq a \leq r &\Leftrightarrow \frac{1}{6}(a^2 + b^2 + c^2) \leq a^2 \leq \frac{2}{3}(a^2 + b^2 + c^2) \\ &\Leftrightarrow b^2 + c^2 \leq 5a^2 \quad \text{and} \quad a^2 \leq 2(b^2 + c^2) \\ &\Leftrightarrow b^2 + (a + b)^2 \leq 5a^2 \quad \text{and} \quad (b + c)^2 \leq 2(b^2 + c^2) \\ &\Leftrightarrow 4a^2 - 2ab - 2b^2 \geq 0 \quad \text{and} \quad b^2 + c^2 \geq 2bc \\ &\Leftrightarrow 2(a - b)(2a + b) \geq 0 \quad \text{and} \quad (b - c)^2 \geq 0, \end{aligned}$$

which hold true since $a \geq b$ and $2a + b \geq a + b + c = 0$. Similarly, we have

$$\begin{aligned} -\frac{r}{2} \leq b \leq \frac{r}{2} &\Leftrightarrow b^2 \leq \frac{r^2}{4} \Leftrightarrow 4b^2 \leq \frac{2}{3}(a^2 + b^2 + c^2) \Leftrightarrow 5b^2 \leq a^2 + c^2 \\ &\Leftrightarrow 5b^2 \leq a^2 + (a+b)^2 \Leftrightarrow 2a^2 + 2ab - 4b^2 \geq 0 \\ &\Leftrightarrow 2(a-b)(a+2b) \geq 0, \end{aligned}$$

which holds true since $a \geq b$ and $a + 2b \geq a + b + c = 0$. Also, we have

$$\begin{aligned} -r \leq c \leq -\frac{r}{2} &\Leftrightarrow r^2 \geq c^2 \geq \frac{r^2}{4} \Leftrightarrow \frac{2}{3}(a^2 + b^2 + c^2) \geq c^2 \geq \frac{1}{6}(a^2 + b^2 + c^2) \\ &\Leftrightarrow 2(a^2 + b^2) \geq c^2 \quad \text{and} \quad 5c^2 \geq a^2 + b^2 \\ &\Leftrightarrow 2(a^2 + b^2) \geq (a+b)^2 \quad \text{and} \quad 5(a+b)^2 \geq a^2 + b^2 \\ &\Leftrightarrow (a-b)^2 \geq 0 \quad \text{and} \quad 4a^2 + 10ab + 4b^2 \geq 0 \\ &\Leftrightarrow (a-b)^2 \geq 0 \quad \text{and} \quad 2(a+2b)(2a+b) \geq 0, \end{aligned}$$

which hold true since $a + 2b \geq a + b + c = 0$ and $2a + b \geq a + b + c = 0$. One can show in the same way that $x \in [\frac{r}{2}, r]$, $y \in [-\frac{r}{2}, \frac{r}{2}]$, $z \in [-r, -\frac{r}{2}]$, so that (24) has been verified.

We prove now that the inequality (21) holds if and only if (22) holds. Indeed, using (23)_{2,4} and (24) we get

$$\begin{aligned} c \leq z &\Leftrightarrow -a - \sqrt{3(r^2 - a^2)} \leq -x - \sqrt{3(r^2 - x^2)} \\ &\Leftrightarrow \frac{a}{r} + \sqrt{3\left(1 - \left(\frac{a}{r}\right)^2\right)} \geq \frac{x}{r} + \sqrt{3\left(1 - \left(\frac{x}{r}\right)^2\right)} \Leftrightarrow a \leq x, \end{aligned}$$

since the function $t \mapsto t + \sqrt{3(1 - t^2)}$ is decreasing for $t \in [\frac{1}{2}, 1]$.

Let us prove next that the inequalities (20) and (21) are equivalent. To accomplish this, we introduce the function $f : [\frac{r}{2}, r] \rightarrow \mathbb{R}$ by

$$f(x) = e^x + e^{(-x + \sqrt{3(r^2 - x^2)})/2} + e^{(-x - \sqrt{3(r^2 - x^2)})/2}. \tag{25}$$

Taking into account (23) and (24)₁, the inequality (20) can be written equivalently as

$$f(a) \leq f(x), \tag{26}$$

which is equivalent to

$$a \leq x,$$

since the function f defined by (25) is monotone increasing on $[\frac{r}{2}, r]$, as we show next. To this aim, we denote by

$$\cos \varphi := \frac{x}{r} \in \left[\frac{1}{2}, 1\right], \quad \text{i.e. } \varphi := \arccos\left(\frac{x}{r}\right) \in \left[0, \frac{\pi}{3}\right].$$

Then the function (25) can be written as

$$f(x) = h(r, \varphi), \quad \text{where } h : (0, \infty) \times \left[0, \frac{\pi}{3}\right] \rightarrow \mathbb{R}, \tag{27}$$

$$h(r, \varphi) = e^{r \cos \varphi} + e^{r \cos(\varphi+2\pi/3)} + e^{r \cos(\varphi-2\pi/3)}.$$

We have to show that $h(r, \varphi)$ is decreasing with respect to $\varphi \in [0, \frac{\pi}{3}]$. We compute the first derivative

$$\frac{\partial}{\partial \varphi} h(r, \varphi) = -r \left[e^{r \cos \varphi} \sin \varphi + e^{r \cos(\varphi+2\pi/3)} \sin\left(\varphi + \frac{2\pi}{3}\right) + e^{r \cos(\varphi-2\pi/3)} \sin\left(\varphi - \frac{2\pi}{3}\right) \right]. \tag{28}$$

The function (28) has the same sign as the function

$$F(r, \varphi) := \frac{1}{r} e^{-r \cos \varphi} \frac{\partial}{\partial \varphi} h(r, \varphi), \tag{29}$$

i.e., the function $F : (0, \infty) \times [0, \frac{\pi}{3}] \rightarrow \mathbb{R}$ given by

$$F(r, \varphi) = -\sin \varphi - e^{-r\sqrt{3} \sin(\varphi+\pi/3)} \sin\left(\varphi + \frac{2\pi}{3}\right) - e^{r\sqrt{3} \sin(\varphi-\pi/3)} \sin\left(\varphi - \frac{2\pi}{3}\right). \tag{30}$$

In order to show that $F(r, \varphi) \leq 0$ for all $(r, \varphi) \in (0, \infty) \times [0, \frac{\pi}{3}]$, we remark that $\lim_{r \searrow 0} F(r, \varphi) = 0$ for fixed $\varphi \in [0, \frac{\pi}{3}]$ and we compute

$$\begin{aligned} \frac{\partial}{\partial r} F(r, \varphi) &= \sqrt{3} \left[e^{-r\sqrt{3} \sin(\varphi+\pi/3)} \sin\left(\varphi + \frac{\pi}{3}\right) \sin\left(\varphi + \frac{2\pi}{3}\right) \right. \\ &\quad \left. - e^{r\sqrt{3} \sin(\varphi-\pi/3)} \sin\left(\varphi - \frac{\pi}{3}\right) \sin\left(\varphi - \frac{2\pi}{3}\right) \right] \\ &= \sqrt{3} \left[e^{-r\sqrt{3} \sin(\varphi+\pi/3)} \frac{1}{2} \left(-\cos(2\varphi + \pi) + \cos \frac{\pi}{3} \right) \right. \\ &\quad \left. - e^{r\sqrt{3} \sin(\varphi-\pi/3)} \frac{1}{2} \left(-\cos(2\varphi - \pi) + \cos \frac{-\pi}{3} \right) \right] \\ &= \frac{\sqrt{3}}{2} \left(\cos 2\varphi + \frac{1}{2} \right) \left[e^{-r\sqrt{3} \sin(\varphi+\pi/3)} - e^{r\sqrt{3} \sin(\varphi-\pi/3)} \right] \leq 0, \end{aligned}$$

since $\varphi \in [0, \frac{\pi}{3}]$ implies $\cos 2\varphi \geq -\frac{1}{2}$ and $-\sin(\varphi + \frac{\pi}{3}) \leq \sin(\varphi - \frac{\pi}{3})$.

Consequently, the function $F(r, \varphi)$ is decreasing with respect to r and for any $(r, \varphi) \in (0, \infty) \times [0, \frac{\pi}{3}]$ we have that

$$F(r, \varphi) \leq \lim_{r \searrow 0} F(r, \varphi) = 0. \tag{31}$$

From (29) and (31), it follows that $h(r, \varphi)$ is decreasing with respect to $\varphi \in [0, \frac{\pi}{3}]$. This means that $f(x)$ is increasing as a function of $x \in [\frac{r}{2}, r]$, i.e., the relation (26) is indeed equivalent to $a \leq x$ and the proof is complete. \square

Consequence 12 Let the real numbers $a \geq b \geq c$ and $x \geq y \geq z$ be such that

$$a + b + c = x + y + z = 0, \quad a^2 + b^2 + c^2 = x^2 + y^2 + z^2.$$

Then one of the following inequalities holds:

$$e^a + e^b + e^c \leq e^x + e^y + e^z, \tag{32}$$

or

$$e^{-a} + e^{-b} + e^{-c} \leq e^{-x} + e^{-y} + e^{-z}. \tag{33}$$

The inequalities (32) and (33) are satisfied simultaneously if and only if $a = x$, $b = y$ and $c = z$.

Proof According to Lemma 11, the inequality (32) is equivalent to

$$a \leq x, \tag{34}$$

while the inequality (33) is equivalent to

$$-a \leq -x. \tag{35}$$

Since one of the relations (34) and (35) must hold, we have proved that one of the inequalities (32) and (33) is satisfied. They are simultaneously satisfied if and only if both (34) and (35) hold true, i.e., $a = x$ (and consequently $b = y$, $c = z$). \square

Consequence 13 Let the real numbers $a \geq b \geq c$ and $x \geq y \geq z$ be such that

$$a + b + c = x + y + z = 0, \quad a^2 + b^2 + c^2 = x^2 + y^2 + z^2$$

$$\text{and } e^a + e^b + e^c = e^x + e^y + e^z.$$

Then we have $a = x$, $b = y$ and $c = z$.

Proof Since by hypothesis $e^a + e^b + e^c = e^x + e^y + e^z$ holds, we can apply Lemma 11 to deduce $a \leq x$ and $c \leq z$.

On the other hand, by virtue of the inverse inequality $e^x + e^y + e^z \leq e^a + e^b + e^c$ and Lemma 11, we obtain $x \leq a$ and $z \leq c$. In conclusion, we get $a = x$, $c = z$ and $b = y$. \square

Proof of Theorem 10 In order to prove (13), we define the real numbers

$$t_i = kz_i \quad (i = 1, 2, 3), \quad \text{where } k = \sqrt{\frac{c_1^2 + c_2^2 + c_3^2}{z_1^2 + z_2^2 + z_3^2}} > 1. \tag{36}$$

Then we have

$$t_1 + t_2 + t_3 = c_1 + c_2 + c_3 = 0 \quad \text{and} \quad t_1^2 + t_2^2 + t_3^2 = c_1^2 + c_2^2 + c_3^2. \tag{37}$$

If we apply the Consequence 12 for the numbers $c_1 \geq c_2 \geq c_3$ and $t_1 \geq t_2 \geq t_3$, then we obtain that

$$\begin{aligned} e^{t_1} + e^{t_2} + e^{t_3} &\leq e^{c_1} + e^{c_2} + e^{c_3} \quad \text{or} \\ e^{-t_1} + e^{-t_2} + e^{-t_3} &\leq e^{-c_1} + e^{-c_2} + e^{-c_3}. \end{aligned} \tag{38}$$

In what follows, let us show that

$$e^{z_1} + e^{z_2} + e^{z_3} < e^{t_1} + e^{t_2} + e^{t_3}. \tag{39}$$

Using the notations $\rho := \sqrt{\frac{2}{3}(z_1^2 + z_2^2 + z_3^2)}$ and

$$\cos \zeta := \frac{z_1}{\rho} \in \left[\frac{1}{2}, 1 \right], \quad \text{i.e., } \zeta := \arccos\left(\frac{z_1}{\rho}\right) \in \left[0, \frac{\pi}{3} \right],$$

we have $k\rho := \sqrt{\frac{2}{3}(t_1^2 + t_2^2 + t_3^2)}$ and $\cos \zeta = \frac{t_1}{k\rho}$. With the help of the function h defined in (27), we can write the inequality (39) in the form

$$\begin{aligned} e^{\rho \cos \zeta} + e^{\rho \cos(\zeta + 2\pi/3)} + e^{\rho \cos(\zeta - 2\pi/3)} &< e^{k\rho \cos \zeta} + e^{k\rho \cos(\zeta + 2\pi/3)} + e^{k\rho \cos(\zeta - 2\pi/3)}, \quad \text{or} \\ h(\rho, \zeta) &< h(k\rho, \zeta), \quad \forall (\rho, \zeta) \in (0, \infty) \times \left[0, \frac{\pi}{3} \right], \quad k > 1. \end{aligned} \tag{40}$$

The relation (40) asserts that the function h defined in (27) is increasing with respect to the first variable $r \in (0, \infty)$. To show this, we compute the derivative

$$\frac{\partial}{\partial r} h(r, \varphi) = e^{r \cos \varphi} \cos \varphi + e^{r \cos(\varphi + 2\pi/3)} \cos\left(\varphi + \frac{2\pi}{3}\right) + e^{r \cos(\varphi - 2\pi/3)} \cos\left(\varphi - \frac{2\pi}{3}\right). \tag{41}$$

By virtue of the Chebyshev's sum inequality, we deduce from (41) that

$$\frac{\partial}{\partial r} h(r, \varphi) > 0. \tag{42}$$

Indeed, the Chebyshev's sum inequality [6, 2.17] asserts that: if $a_1 \geq a_2 \geq \dots \geq a_n$ and $b_1 \geq b_2 \geq \dots \geq b_n$ then

$$n \sum_{k=1}^n a_k b_k \geq \left(\sum_{k=1}^n a_k \right) \left(\sum_{k=1}^n b_k \right).$$

In our case, we derive the following result: for any real numbers x, y, z such that $x + y + z = 0$, the inequality

$$xe^x + ye^y + ze^z \geq \frac{1}{3}(x + y + z)(e^x + e^y + e^z) = 0, \tag{43}$$

holds true, with equality if and only if $x = y = z = 0$.

Applying the result (43) to the function (41), we deduce the relation (42). This means that $h(r, \varphi)$ is an increasing function of r , i.e. the inequality (40) holds, and hence, we have proved (39).

One can show analogously that the inequality

$$e^{-z_1} + e^{-z_2} + e^{-z_3} < e^{-t_1} + e^{-t_2} + e^{-t_3} \tag{44}$$

is also valid. From (38), (39) and (44), it follows that the assertion (13) holds true. Thus, the proof of Theorem 10 is complete. \square

Since the statements of the Theorems 8 and 10 are equivalent, we have proved also the inequality (8).

Remark 14 The inequality (8) becomes an equality if and only if $z_i = c_i$, $i = 1, 2, 3$.

Proof Indeed, assume that $z_1^2 + z_2^2 + z_3^2 = c_1^2 + c_2^2 + c_3^2$. Then we can apply the Consequence 12 and we deduce that

$$e^{z_1} + e^{z_2} + e^{z_3} \leq e^{c_1} + e^{c_2} + e^{c_3} \quad \text{or} \quad e^{-z_1} + e^{-z_2} + e^{-z_3} \leq e^{-c_1} + e^{-c_2} + e^{-c_3}. \tag{45}$$

Taking into account (7)_{1,2} in conjunction with (45), we find

$$e^{z_1} + e^{z_2} + e^{z_3} = e^{c_1} + e^{c_2} + e^{c_3} \quad \text{or} \quad e^{-z_1} + e^{-z_2} + e^{-z_3} = e^{-c_1} + e^{-c_2} + e^{-c_3}. \tag{46}$$

By virtue of (46), we can apply the Consequence 13 to derive $z_1 = c_1$, and consequently $z_2 = c_2$, $z_3 = c_3$. \square

Let us prove the following version of the inequality (6) for two pairs of numbers d_1, d_2 and x_1, x_2 :

Remark 15 If the real numbers $d_1 \geq d_2 > 0$ and $x_1 \geq x_2 > 0$ are such that

$$x_1^2 + x_2^2 \geq d_1^2 + d_2^2 \quad \text{and} \quad x_1 x_2 = d_1 d_2 = 1, \tag{47}$$

then the inequality

$$(\log x_1)^2 + (\log x_2)^2 \geq (\log d_1)^2 + (\log d_2)^2 \tag{48}$$

holds true. Note that the additional condition

$$\frac{1}{x_1^2} + \frac{1}{x_2^2} \geq \frac{1}{d_1^2} + \frac{1}{d_2^2}$$

is automatically fulfilled.

Proof Since $x_1 x_2 = d_1 d_2 = 1$ and $d_1 \geq d_2 > 0$, $x_1 \geq x_2 > 0$, we have $x_1 \geq 1$, $d_1 \geq 1$ and

$$\log x_1 = -\log x_2 \geq 0, \quad \log d_1 = -\log d_2 \geq 0,$$

so that the inequality (48) is equivalent to $\log x_1 \geq \log d_1$, *i.e.*, we have to show that $x_1 \geq d_1$.

Indeed, if we insert $x_2 = \frac{1}{x_1}$ and $d_2 = \frac{1}{d_1}$ into the inequality (47)₁ then we find

$$x_1^2 + \frac{1}{x_1^2} \geq d_1^2 + \frac{1}{d_1^2},$$

which means that $x_1 \geq d_1$ since the function $t \mapsto t^2 + \frac{1}{t^2}$ is increasing for $t \in [1, \infty)$. This completes the proof. \square

Alternative proof of Remark 15 Let $x_3 = d_3 = 1$. Then (47) implies $x_1^2 + x_2^2 + x_3^2 \geq d_1^2 + d_2^2 + d_3^2$ and $x_1x_2x_3 = d_1d_2d_3 = 1$ as well as

$$x_1^2x_2^2 + x_2^2x_3^2 + x_1^2x_3^2 = 1 + x_2^2 + x_1^2 \geq 1 + d_2^2 + d_1^2 = d_1^2d_2^2 + d_2^2d_3^2 + d_1^2d_3^2, \tag{49}$$

because $x_1^2x_2^2 = 1 = d_1^2d_2^2$, and Theorem 6 provides the assertion. \square

4 Some counterexamples for weakened assumptions

Example 16 Unlike in the 2D case in Remark 15, for two triples of numbers the second condition (18)₂ of Theorem 2, namely $y_1y_2 + y_2y_3 + y_1y_3 \geq a_1a_2 + a_2a_3 + a_1a_3$, cannot be removed. Let

$$y_1 = e^6, \quad y_2 = 1, \quad y_3 = e^{-6}, \quad a_1 = e^4, \quad a_2 = e^4, \quad a_3 = e^{-8}.$$

Then $y_1y_2y_3 = a_1a_2a_3 = 1$ and

$$y_1 + y_2 + y_3 > e^6 \geq e^2e^4 > 3e^4 > a_1 + a_2 + a_3,$$

but

$$\begin{aligned} (\log y_1)^2 + (\log y_2)^2 + (\log y_3)^2 &= 36 + 0 + 36 \\ &< 16 + 16 + 64 = (\log a_1)^2 + (\log a_2)^2 + (\log a_3)^2. \end{aligned}$$

Example 17 The condition $y_1y_2y_3 = a_1a_2a_3$ cannot be weakened to $y_1y_2y_3 \geq a_1a_2a_3$. Indeed, let $y_2 = y_3 = a_1 = a_2 = 1, y_1 = e, a_3 = e^{-2}$. Then

$$\begin{aligned} y_1 + y_2 + y_3 &= e + 1 + 1 \geq 1 + 1 + e^{-2} = a_1 + a_2 + a_3, \\ y_1y_2 + y_1y_3 + y_2y_3 &= e + e + 1 \geq 1 + e^{-2} + e^{-2} = a_1a_2 + a_1a_3 + a_2a_3, \\ y_1y_2y_3 &= e \geq e^{-2} = a_1a_2a_3. \end{aligned}$$

But nevertheless

$$(\log y_1)^2 + (\log y_2)^2 + (\log y_3)^2 = 1 + 0 + 0 < 0 + 0 + 4 = (\log a_1)^2 + (\log a_2)^2 + (\log a_3)^2.$$

A counterexample for the two variable case can be constructed analogously.

Example 18 Even with an analogous condition, the inequality (4) does not hold for $n = 4$ numbers (without further assumptions). Indeed, let

$$y_1 = e, \quad y_2 = y_3 = e^7, \quad y_4 = e^{-15}, \quad a_1 = a_2 = e^6, \quad a_3 = e^7, \quad a_4 = e^{-19}.$$

Then $y_1 y_2 y_3 y_4 = a_1 a_2 a_3 a_4 = 1$. Also,

$$y_1 + y_2 + y_3 + y_4 = e + e^7 + e^7 + e^{-15} > 0 + e^7 + 2e^6 + e^{-19} = a_1 + a_2 + a_3 + a_4.$$

Furthermore,

$$y_1 y_2 + y_1 y_3 + y_1 y_4 + y_2 y_3 + y_2 y_4 + y_3 y_4 = e^8 + e^8 + e^{-14} + e^{14} + e^{-8} + e^{-8}$$

and

$$a_1 a_2 + a_1 a_3 + a_1 a_4 + a_2 a_3 + a_2 a_4 + a_3 a_4 = e^{12} + e^{13} + e^{-13} + e^{13} + e^{-13} + e^{-12}.$$

Since $e^2 > 2e + 1$, we have $e^{14} > e^{13} + e^{13} + e^{12}$ and, therefore,

$$y_1 y_2 + y_1 y_3 + y_1 y_4 + y_2 y_3 + y_2 y_4 + y_3 y_4 \geq a_1 a_2 + a_1 a_3 + a_1 a_4 + a_2 a_3 + a_2 a_4 + a_3 a_4.$$

Nevertheless, for the sum of squared logarithms, the ‘reverse’ inequality

$$\begin{aligned} (\log y_1)^2 + (\log y_2)^2 + (\log y_3)^2 + (\log y_4)^2 &= 1 + 49 + 49 + 225 = 324 \\ &< 482 = 36 + 36 + 49 + 361 \\ &= (\log a_1)^2 + (\log a_2)^2 + (\log a_3)^2 + (\log a_4)^2 \end{aligned}$$

holds true.

Example 19 The inequality (4) does not remain true either, if the function $\log(y)$ is replaced by its linearization $(y - 1)$. Indeed, let $y_1 = 9, y_2 = 5, y_3 = \frac{1}{45}, a_1 = 10, a_2 = 1, a_3 = \frac{1}{10}$. Then

$$y_1 + y_2 + y_3 > 14 > 11.1 = a_1 + a_2 + a_3$$

and

$$y_1 y_2 + y_1 y_3 + y_2 y_3 > 45 \geq 11.1 = a_1 a_2 + a_1 a_3 + a_2 a_3.$$

But

$$\begin{aligned} (y_1 - 1)^2 + (y_2 - 1)^2 + (y_3 - 1)^2 &= 64 + 16 + \left(\frac{44}{45}\right)^2 < 81 < 9^2 + 0 + \left(\frac{9}{10}\right)^2 \\ &= (a_1 - 1)^2 + (a_2 - 1)^2 + (a_3 - 1)^2. \end{aligned}$$

5 Conjecture for arbitrary n

The structure of the inequality in dimensions $n = 2$ and $n = 3$ and extensive numerical sampling strongly suggest that the inequality holds for all $n \in \mathbb{N}$ if the n corresponding conditions are satisfied. More precisely, in terms of the elementary symmetric polynomials, we expect the following:

Conjecture 20 *Let $n \in \mathbb{N}$ and $y_i, a_i > 0$ for $i = 1, \dots, n$. If for all $i = 1, \dots, n - 1$ we have*

$$e_i(y_1, \dots, y_n) \geq e_i(a_1, \dots, a_n) \quad \text{and} \quad e_n(y_1, \dots, y_n) = e_n(a_1, \dots, a_n),$$

then

$$\sum_{i=1}^n (\log y_i)^2 \geq \sum_{i=1}^n (\log a_i)^2.$$

6 Applications

The investigation in this paper has been motivated by some recent applications. The new sum of squared logarithms inequality is one of the fundamental tools in deducing a novel optimality result in matrix analysis and the conditions in the form (3) had been deduced in the course of that work. Optimality in the matrix problem suggested the sum of squared logarithms inequality. Indeed, based on the present result in [7], it has been shown that for all invertible $Z \in \mathbb{C}^{3 \times 3}$ and for any definition of the matrix logarithm as possibly multivalued solution $X \in \mathbb{C}^{3 \times 3}$ of $\exp X = Z$ it holds

$$\begin{aligned} \min_{Q^*Q=I} \|\log Q^*Z\|_F^2 &= \|\log U_p^*Z\|_F^2 = \|\log H\|_F^2, \\ \min_{Q^*Q=I} \|\text{sym} \log Q^*Z\|_F^2 &= \|\text{sym} \log U_p^*Z\|_F^2 = \|\log H\|_F^2, \end{aligned} \tag{50}$$

where $\text{sym} X = \frac{1}{2}(X + X^*)$ is the Hermitian part of $X \in \mathbb{C}^{3 \times 3}$ and U_p is the unitary factor in the polar decomposition of Z into unitary and Hermitian positive definite matrix H

$$Z = U_p H. \tag{51}$$

This result (50) generalizes the fact that for any complex logarithm and for all $z \in \mathbb{C} \setminus \{0\}$

$$\min_{\vartheta \in (-\pi, \pi]} |\log_{\mathbb{C}} [e^{-i\vartheta} z]|^2 = |\log_{\mathbb{R}} |z||^2, \quad \min_{\vartheta \in (-\pi, \pi]} |\Re \log_{\mathbb{C}} [e^{-i\vartheta} z]|^2 = |\log_{\mathbb{R}} |z||^2. \tag{52}$$

The optimality result (50) can now also be viewed as another characterization of the unitary factor in the polar decomposition. In addition, in a forthcoming contribution [8], we use (50) to calculate the geodesic distance of the isochoric part of the deformation gradient $\frac{F}{\det F^{\frac{1}{3}}} \in \text{SL}(3, \mathbb{R})$ to $\text{SO}(3, \mathbb{R})$ in the canonical left-invariant Riemannian metric on $\text{SL}(3, \mathbb{R})$, to the effect that

$$\text{dist}_{\text{geod}}^2 \left(\frac{F}{\det F^{\frac{1}{3}}}, \text{SO}(3, \mathbb{R}) \right) = \|\text{dev}_3 \log \sqrt{F^T F}\|_F^2, \tag{53}$$

where $\text{dev}_3 X = X - \frac{1}{3}(\text{tr } X)I$ is the orthogonal projection of $X \in \mathbb{R}^{3 \times 3}$ to trace free matrices. Thereby, we provide a rigorous geometric justification for the preferred use of the Hencky-strain measure $\|\log \sqrt{F^T F}\|_F^2$ in nonlinear elasticity and plasticity theory [9].

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed fully to all parts of the manuscript. Notably all ideas have emerged by continuous discussions among them.

Author details

¹Lehrstuhl für Nichtlineare Analysis und Modellierung, Fakultät für Mathematik, Universität Duisburg-Essen, Essen, Germany. ²Department of Mathematics, University 'A.I. Cuza' of Iași, Iași, Romania.

Acknowledgements

The first author (MB) was supported by the German state grant: 'Programm des Bundes und der Länder für bessere Studienbedingungen und mehr Qualität in der Lehre'.

Received: 21 January 2013 Accepted: 28 March 2013 Published: 12 April 2013

References

1. Guan, K: Schur-convexity of the complete elementary symmetric functions. *J. Inequal. Appl.* **2006**, Article ID 67624 (2006). doi:10.1155/JIA/2006/67624
2. Roventa, I: A note on Schur-concave functions. *J. Inequal. Appl.* **159**, 1-9 (2012)
3. Steele, JM: *The Cauchy-Schwarz Master Class: an Introduction to the Art of Mathematical Inequalities*. Cambridge University Press, Cambridge (2004)
4. Khan, AR, Latif, N, Pečarić, J: Exponential convexity for majorization. *J. Inequal. Appl.* **105**, 1-13 (2012)
5. Karamata, J: Sur une inégalité relative aux fonctions convexes. *Publ. Math. Univ. Belgrad* **1**, 145-148 (1932)
6. Hardy, GH, Littlewood, JE, Pólya, G: *Inequalities*. The University Press, Cambridge (1934)
7. Neff, P, Nagatsukasa, Y, Fischle, A: The unitary polar factor $Q = U_p$ minimizes $\|\text{Log}(Q^* Z)\|^2$ and $\|\text{sym, Log}(Q^* Z)\|^2$ in the spectral norm in any dimension and the Frobenius matrix norm in three dimensions (2013, submitted)
8. Neff, P, Eidel, B, Osterbrink, F, Martin, R: The isotropic Hencky strain energy $\|\log U\|^2$ measures the geodesic distance of the deformation gradient $F \in \text{GL}^+(n)$ to $\text{SO}(n)$ in the unique left-invariant Riemannian metric on $\text{GL}^+(n)$ which is also right $O(n)$ -invariant (2013, in preparation)
9. Hencky, H: Über die Form des Elastizitätsgesetzes bei ideal elastischen Stoffen. *Z. Techn. Physik* **9**, 215-220 (1928)

doi:10.1186/1029-242X-2013-168

Cite this article as: Biršan et al.: Sum of squared logarithms - an inequality relating positive definite matrices and their matrix logarithm. *Journal of Inequalities and Applications* 2013 **2013**:168.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com