

RESEARCH

Open Access

The relationship between sperm viability and DNA fragmentation rates

Mary K Samplaski¹, Apostolos Dimitromanolakis², Kirk C Lo^{1,3}, Ethan D Grober¹, Brendan Mullen², Alaina Garbens¹ and Keith A Jarvi^{1,3,4*}

Abstract

Background: In humans, sperm DNA fragmentation rates have been correlated with sperm viability rates. Reduced sperm viability is associated with high sperm DNA fragmentation, while conversely high sperm viability is associated with low rates of sperm DNA fragmentation. Both elevated DNA fragmentation rates and poor viability are correlated with impaired male fertility, with a DNA fragmentation rate of > 30% indicating subfertility. We postulated that in some men, the sperm viability assay could predict the sperm DNA fragmentation rates. This in turn could reduce the need for sperm DNA fragmentation assay testing, simplifying the infertility investigation and saving money for infertile couples.

Methods: All men having semen analyses with both viability and DNA fragmentation testing were identified via a prospectively collected database. Viability was measured by eosin-nigrosin assay. DNA fragmentation was measured using the sperm chromosome structure assay. The relationship between DNA fragmentation and viability was assessed using Pearson's correlation coefficient.

Results: From 2008-2013, 3049 semen analyses had both viability and DNA fragmentation testing. A strong inverse relationship was seen between sperm viability and DNA fragmentation rates, with $r = -0.83$. If viability was $\leq 50\%$ ($n = 301$) then DNA fragmentation was $\geq 30\%$ for 95% of the samples. If viability was $\geq 75\%$ ($n = 1736$), then the DNA fragmentation was $\leq 30\%$ for 95% of the patients. Sperm viability correlates strongly with DNA fragmentation rates.

Conclusions: In men with high levels of sperm viability $\geq 75\%$, or low levels of sperm viability $\leq 30\%$, DFI testing may be not be routinely necessary. Given that DNA fragmentation testing is substantially more expensive than vitality testing, this may represent a valuable cost-saving measure for couples undergoing a fertility evaluation.

Keywords: DNA fragmentation, Viability, Prediction, Semen

Background

Both elevated DNA fragmentation and poor viability are known to be associated with male factor infertility. These two conditions are linked, as DNA fragmentation is one of the final steps before spermatozoa death [1] and conversely, DNA breaks are one of the primary instigators of sperm apoptosis [2-4]. The literature has demonstrated a clear link between DNA fragmentation and sperm viability [5], and in groups of men with high levels of DNA fragmentation, high levels of necrospemia are also seen [5].

The "Georgetown Male Factor Infertility Study" was the first study to establish DNA fragmentation fertility data in humans [6]. Data from this study were used to establish the thresholds DNA fragmentation and fertility data. For male fertility, the sperm chromatin structure assay was used to define > 30% DNA fragmentation as "significant lack of", 15-30% DNA fragmentation as 'reasonable' and < 15% DNA fragmentation for 'high' fertility status [6]. Subsequent studies have also demonstrated that a sperm DNA fragmentation rate of > 30% is correlated with impaired fertility outcomes. A meta-analysis of four studies, with a total of 1962 men evaluating the relationship between sperm DNA fragmentation and spontaneous and intrauterine insemination pregnancies found that men

* Correspondence: kjarvi@mtsinai.on.ca

¹Division of Urology, Department of Surgery, Mount Sinai Hospital, University of Toronto, Toronto, ON, Canada

³Institute of Medical Sciences, University of Toronto, Toronto, ON, Canada

Full list of author information is available at the end of the article

with a DNA fragmentation < 30% were more likely to achieve a pregnancy or live birth ($p = 0.0001$) [6-8]. The second study looked at the relationship between DNA fragmentation and in vitro fertilization pregnancies, and found that couples were ~2x more likely to become pregnant if the sperm DNA fragmentation rate was < 30% [7,9]. These studies and others, support the notion that semen samples with $\geq 30\%$ sperm DNA fragmentation by sperm chromatin structure assay have reduced fertility [10].

Necrostermia is defined as the percentage of dead sperm in a semen analysis sample > 42%, and is usually measured by assessing the membrane integrity of the cells, as determined by a dye exclusion assay [11,12]. In spite of its relatively straightforward definition, the causes and impact of necrostermia on male fertility are still relatively poorly understood. Necrostermia is commonly seen in men with spinal cord injuries [12], infections [13], chronic medical conditions [5], and after exposure to toxic substances [14]. While intuitively it makes sense that high levels of necrostermia are associated with poorer reproductive outcomes, there is a paucity of literature to support this.

One of the primary disadvantages of DNA fragmentation testing is its cost. Compared with other male fertility testing, DNA fragmentation testing is costly, as vitality testing costs approximately \$2 and DNA fragmentation testing costs approximately \$250, based on internal and internet quotes [15]. These costs generally are transferred to the patient and can add further financial burden to a couple looking to conceive.

We sought to determine the relationship between sperm viability and DNA fragmentation, with the hypothesis that for some patients, sperm DNA fragmentation rates could be predicted from the sperm viability rates. This may eliminate the need for sperm DNA fragmentation testing in a subset of men, representing a cost-saving measure for some couples undergoing a fertility evaluation.

Methods

All men presenting for a fertility evaluation from 2008-2013, and having semen analyses with both viability and DNA fragmentation testing were identified. This data was reviewed in a retrospective manner. The collection of data and the analysis of the data in this database were approved by the Research Ethics Board of the Mount Sinai Hospital with reference number 05-0161-E (collection of data) and 07-0032-E (analysis of data) respectively. The date of the approval was October 18, 2005 and October 30, 2007. All participants have signed the IRB approved informed consent form.

Semen samples were collected at Mount Sinai Hospital, at least 48 hours, but not more than 7 days, after the time of last ejaculation. Semen samples were collected between

2 and 5 days from the last ejaculation. Semen samples were assessed for volume and then analyzed for sperm count, sperm concentration and motility following the 2001 WHO criteria. These criteria are based on microscopic high-power evaluation of 200 sperm for intactness of membranes of acrosome, head, neck, midpiece and tail.

Viability was assessed within 30 minutes of ejaculation. It was measured by eosin-nigrosin assay, by dissolving 1 g of eosin with 1 g of fresh sperm and then 3 g of nigrosin [16]. The percentage of viable sperm (sperm head unstained indicating living sperm) and non-viable sperm (sperm head stained indicating dead spermatozoa) was assessed by counting a minimum of 100 spermatozoa. Replicate counts of 100 sperm on each of two slides were performed. These were then repeated if >5% difference was found.

DNA fragmentation testing was performed on a frozen prepared semen sample using the sperm chromosome structure assay, as previously described [17]. Samples were treated for 30 seconds with 400 μL of a solution of 0.1% Triton X-100, 0.15 M NaCl, and 0.08 N HCl (pH 1.2). After 30 seconds, 1.2 mL of staining buffer (6 $\mu\text{g}/\text{mL}$ acridine orange, 37 mM citric acid, 126 mM Na_2HPO_4 , 1 mM disodium EDTA, and 0.15 M NaCl, pH 6.0) was admixed to the test tube, and the sample was analyzed by flow cytometry. After excitation by a 488-nm wavelength light source, acridine orange bound to double-stranded DNA fluoresced green (515 to 530 nm) and acridine orange bound to single-stranded DNA fluoresced red (630 nm or more). Three minutes after acridine orange staining, samples were analyzed in a flow cytometry activated cell sorter (Caliburflow cytometer, Becton Dickinson, San Jose, California). A minimum of 5000 cells were analyzed by a flow cytometry activated cell sorter scan interfaced with a data handler. The proportion of cells exhibiting an abnormal emission of red fluorescence, reflecting the percentage of sperm with denatured DNA, was recorded.

The relationship between DNA fragmentation and viability was assessed using Pearson's product moment correlation coefficient. P-values and confidence intervals were obtained by standard methods, assuming normality of the

Table 1 Semen analysis DNA fragmentation rates when grouped by viability

Sperm viability	DNA fragmentation rate (mean +/- standard deviation)
0-20%	70.07 +/- 34.39%
21-40%	58.03 +/- 28.28%
41-60%	38.19 +/- 17.95%
61-80%	20.40 +/- 10.16%
81-100%	12.40 +/- 5.61%

Higher sperm DNA fragmentation rates were associated with lower sperm viability rates.




Figure 1 Sperm DNA fragmentation versus necrospemia. A strong inverse relationship ($p < 0.001$) was seen between viability and DNA fragmentation. If viability was $\geq 75\%$, then the DNA fragmentation was $< 30\%$ for 95% of the patients. If viability was $\leq 50\%$ then DNA fragmentation was $\geq 30\%$ for 95% of the samples.

data. Statistical analysis was performed in R Foundation for Statistical Computing, version 2.15.2.

Results

A total of 2695 men underwent semen analysis testing with both DNA fragmentation and viability assays. Some men had multiple semen analyses with DNA fragmentation and viability testing. 2438 men had 1 test, 191 men had 2 tests, 37 men had 3 tests, 15 men had 4 tests, 3 men had 5 tests, 6 men had 6 tests, and 1 man had 7 tests. A total of 3049 semen samples with both DNA fragmentation and viability testing were performed.

Of the 3049 semen samples analyzed, 47 (1.5%) had sperm viability of 0-20%, 113 (3.7%) had viability of 21-40%, 450 (14.8%) had viability of 41-60%, 1920 (63%) had viability of 61-80%, and 519 (17%) had viability of 81-100%.

Higher sperm DNA fragmentation rates were associated with lower sperm viability rates (see Table 1), and samples with the lowest sperm viability 0-20% had the highest sperm DNA fragmentation rates ($70.07 \pm 34.39\%$), and samples with the highest sperm viability rates, 81-100%, were associated with the lowest sperm DNA fragmentation rates ($12.40 \pm 5.61\%$).

433 (16.1%) samples had a sperm viability of $\leq 58\%$, the lower limit of normal according to the 2010 WHO laboratory manual guidelines [18]. For these samples, the mean DNA fragmentation was $48.65 \pm 25.59\%$. For the 2262 samples with viability $> 58\%$, the mean DNA fragmentation was $19.57 \pm 10.58\%$ ($p < 0.001$).

A strong inverse relationship ($p < 0.001$) was seen between sperm viability and DNA fragmentation, with Pearson’s product-moment correlation coefficient $r = -0.83$ (Figure 1), $t = -80.69$, $df = 3047$, $p < 0.001$ and a 95% confidence interval of -0.836 to -0.814 . If viability was very high

($\geq 80\%$, $n = 1104$) then DNA fragmentation was consistently $< 30\%$ (100% sensitivity to predict DNA fragmentation $< 30\%$). If viability was $\geq 75\%$ ($n = 1736$), then the DNA fragmentation was $< 30\%$ for 95% of the patients (Table 2). For samples with very low viability (viability $\leq 35\%$, $n = 91$) then DFI was always $\geq 30\%$. If viability was $\leq 50\%$ then DNA fragmentation was $\geq 30\%$ for 95% of the samples ($n = 310$). The results of the semen analysis testing with respect to vitality and DFI are seen in Tables 3 and 4.

Discussion

Both elevated sperm DNA fragmentation and poor sperm viability are linked to male infertility. While it is unclear in some cases which process comes first, the two are linked and the literature has demonstrated a clear link between DNA fragmentation and sperm viability [5]. Both apoptosis and necrosis result in DNA fragmentation, either by an active mechanism through apoptotic endonuclease activation or passively as in necrosis [19,20]. DNA fragmentation may also be seen in mature, viable sperm, although the mechanisms that trigger this degradation have not been fully elucidated [21,19]. After ejaculation, the incidence of sperm DNA

Table 2 Number of men having viability testing with DNA fragmentation \geq or $< 30\%$ (95% confidence interval)

Viability	DNA fragmentation rate (95% confidence interval)	Number of men (%)
$\geq 75\%$	$< 30\%$	1736 (56.9%)
74-49%		1004 (32.9%)
$\leq 50\%$	$\geq 30\%$	310 (10.2%)

If sperm viability was $\geq 75\%$ ($n = 1736$), then the DNA fragmentation was $< 30\%$ for 95% of the patients. If sperm viability was $\leq 50\%$ then DNA fragmentation was $\geq 30\%$ for 95% of the samples ($n = 310$).

Table 3 Semen analysis parameters as grouped by viability

Viability	Ejaculate volume	Concentration	Motility	Normal morphology	Total motile sperm count
≥ 75%	2.82 +/- 1.49 mL	64.23 +/- 56.05	31.89 +/- 12.62%	20.93 +/- 11.34%	58.15 +/- 61.38
74-49%	3.00 +/- 1.72 mL	54.55 +/- 58.81	22.88 +/- 11.45%	16.27 +/- 10.89%	37.44 +/- 51.82
≤ 50%	2.55 +/- 1.79 mL	38.74 +/- 44.57	8.23 +/- 7.12%	12.04 +/- 9.34%	10.31 +/- 17.88

fragmentation increases, both with duration since ejaculation [19,22] and with temperatures of 37°C or greater [23,24], but there is great inter-individual variability in these increases [19].

While reported correlations between DNA fragmentation and sperm concentration in subfertile men have varied [25-27], the literature has demonstrated a clear link between sperm DNA fragmentation and sperm viability [5]. For groups of men with high levels of sperm DNA fragmentation, high levels of necrospermia are also seen [5]. In addition, increases in DNA fragmentation rates following incubation of human spermatozoa have been shown to correlate strongly with sperm viability loss [28].

Elevated sperm DNA fragmentation rates have been positively correlated with impaired fertility, including longer times to natural conception [8], impaired embryo cleavage [29], impaired implantation rates [30], higher miscarriage rates [29], and increased risk of pregnancy loss after both in vitro fertilization and intracytoplasmic sperm injection [31]. Using the sperm chromosome structure assay, couples with a sperm DNA fragmentation of <40% have been shown to have an odds ratio of 10× greater probability of pregnancy via natural intercourse than those with DNA fragmentation >40% [8]. Likewise, intrauterine insemination patients have been shown to be 8.7× more likely to have a live birth if the DNA fragmentation is ≤27% [7]. Finally, a DNA fragmentation rate of ≥30% has been associated with increased spontaneous abortion rates [32]. In addition to these individual studies, two large meta-analyses using the sperm chromatin structure assay have demonstrated the clear relationship between DNA fragmentation and pregnancy. The first of these was a meta-analysis of four studies, with a total of 1962 men, found that men with a DNA fragmentation <30% were more likely to achieve a pregnancy or live birth either spontaneously or via intrauterine insemination (p = 0.0001) [6-8]. The second found that couples were ~2× more likely to become

pregnant via in vitro fertilization if their DNA fragmentation was <30% [7,9]. Because of studies like this, sperm DNA fragmentation testing has been used increasingly as an adjunct to the standard sperm parameters [33-35].

Interestingly, while a DNA fragmentation cutoff of 30% is commonly used, there are limited studies looking at DNA fragmentation rates >30%. We identified a single study which compared reproductive outcomes in men with sperm having DNA fragmentation rates of ≤15% compared with >50% [33]. They found that using sperm selected by movement and morphology characteristic for intracytoplasmic sperm injection, couples with high sperm DNA fragmentation rates had similar fertilization and clinical pregnancy rates compared with sperm with low DNA fragmentation rates [33]. There are otherwise no studies looking at sperm DNA fragmentation rates >30%, but the literature clearly demonstrates that a DNA fragmentation of >30% negatively correlates with male reproductive outcomes.

Necrospermia is defined as a high percentage of dead sperm, as determined in our study by dye exclusion [11,12]. Living sperm have an intact cytoplasmic membrane, which is the basis for viability assays such as or dye exclusion testing, which tests sperms' ability to resist the absorption of certain dyes, including eosin, nigrosin, or trypan blue [16]. According to the 2010 WHO laboratory manual for the examination and processing of human semen, the lower reference limit for viability (membrane-intact spermatozoa) is 58% (5th centile, 95% CI 55-63) [18].

The incidence of necrospermia in the fertile and infertile populations is poorly defined. A 2003 study of 4108 infertile men identified a prevalence of 0.7% [36], and a 2004 study estimated the prevalence of epididymal necrospermia at 0.5% in healthy men undergoing infertility work-up [37]. Both of these studies used a viability of 40% dead sperm as their cutoff definition of necrospermia, whereas we used the WHO 2010 guidelines which state that 42% dead sperm is the cutoff for

Table 4 Semen analysis parameters as grouped by DNA fragmentation

DNA fragmentation	Ejaculate volume	Concentration	Motility	Normal morphology	Total motile sperm count
< 30%	2.84 +/- 1.53 mL	62.06 +/- 56.23	30.36 +/- 12.74%	20.06 +/- 11.31%	56.67 +/- 10.96
≥ 30%	2.93 +/- 1.84 mL	49.38 +/- 57.05	16.5 +/- 11.3%	14.2 +/- 10.71%	26.21 +/- 43.44

necropermia. Using a cutoff of 42%, we found an incidence of necropermia of 16.1% in our population of infertile males, relatively high compared with the prior reported rates. For these samples, the mean DNA fragmentation was $48.65 \pm 25.59\%$. For the 2262 samples with viability $> 58\%$, the mean DNA fragmentation was $19.57 \pm 10.58\%$ ($p < 0.001$). Thus our data clearly support the notion that men with WHO defined necropermia have higher rates of DNA fragmentation. However, in spite of its relatively straightforward definition and diagnosis, the impact and etiology of necropermia in men with infertility is still relatively poorly understood.

This is the first study to use vitality as a predictor for the level of sperm DNA fragmentation. Given our findings, in men with sperm vitality of $\geq 75\%$, routine DNA fragmentation testing is unlikely to provide any additional information, as in these men sperm DNA fragmentation is very likely to be low. Likewise, for men with sperm vitality of $\leq 50\%$ routine DNA fragmentation testing may not be required as $> 95\%$ will have sperm DNA integrity of $> 30\%$. Based on these estimates, only 32.9% of men in our series undergoing both viability and DNA fragmentation testing would have gained additional information from DNA fragmentation testing (Table 2). In the majority of men, viability testing may predict sperm DNA fragmentation rates, allowing the couple to avoid the sperm DNA fragmentation assay and may represent a valuable cost-saving measure. While a formal cost-analysis was not performed, this may represent a valuable cost-savings measure for couples undergoing fertility evaluation, as viability testing costs approximately \$2 and DNA fragmentation testing costs approximately \$250, based on internal and internet quotes [15].

Conclusions

Sperm viability correlates strongly with DNA fragmentation rates and is predictive of sperm DNA fragmentation rates. In men with high levels of sperm viability $\geq 75\%$, or low levels of sperm viability $\leq 30\%$, DFI testing may not provide additional information. Given that DNA fragmentation testing is approximately 100× more expensive than viability testing (\$250 versus \$2), this may represent a valuable cost-saving measure for couples undergoing a fertility evaluation.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MS wrote the manuscript with AG and was responsible for primary data collection and analysis. AD performed the data and statistical analysis. KL and EG contributed patients from their practices as well as contributing constructively to the message of the manuscript. AG was involved in writing the manuscript as well as data collection. BM contributed patients, as well as conceived the original idea for the manuscript. KJ contributed patients from his practice, was involved in manuscript design and revisions and data analysis. All authors were involved in manuscript revisions and have approved the final manuscript.

Author details

¹Division of Urology, Department of Surgery, Mount Sinai Hospital, University of Toronto, Toronto, ON, Canada. ²Department of Pathology, Mount Sinai Hospital, University of Toronto, Toronto ON, Canada. ³Institute of Medical Sciences, University of Toronto, Toronto, ON, Canada. ⁴Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, Toronto, ON, Canada.

Received: 29 December 2014 Accepted: 27 April 2015

Published online: 14 May 2015

References

- Irvine DS, Macleod IC, Templeton AA, Masterton A, Taylor A. A prospective clinical study of the relationship between the computer-assisted assessment of human semen quality and the achievement of pregnancy in vivo. *Hum Reprod.* 1994;9(12):2324–34.
- Shen HM, Dai J, Chia SE, Lim A, Ong CN. Detection of apoptotic alterations in sperm in subfertile patients and their correlations with sperm quality. *Hum Reprod.* 2002;17(5):1266–73.
- Barroso G, Morshedi M, Oehninger S. Analysis of DNA fragmentation, plasma membrane translocation of phosphatidylserine and oxidative stress in human spermatozoa. *Hum Reprod.* 2000;15(6):1338–44.
- Gandini L, Lombardo F, Paoli D, Caponecchia L, Familiari G, Verlengia C, et al. Study of apoptotic DNA fragmentation in human spermatozoa. *Hum Reprod.* 2000;15(4):830–9.
- Brahem S, Jellad S, Iballa S, Saad A, Mehdi M. DNA fragmentation status in patients with necrozoospermia. *Syst Biol Reprod Med.* 2012;58(6):319–23. doi:10.3109/19396368.2012.710869.
- Evenson DP, Jost LK, Marshall D, Zinaman MJ, Clegg E, Purvis K, et al. Utility of the sperm chromatin structure assay as a diagnostic and prognostic tool in the human fertility clinic. *Hum Reprod.* 1999;14(4):1039–49.
- Bungum M, Humaidan P, Spano M, Jepsen K, Bungum L, Giwercman A. The predictive value of sperm chromatin structure assay (SCSA) parameters for the outcome of intrauterine insemination, IVF and ICSI. *Hum Reprod.* 2004;19(6):1401–8. doi:10.1093/humrep/deh280.
- Spano M, Bonde JP, Hjollund HI, Kolstad HA, Cordelli E, Leter G. Sperm chromatin damage impairs human fertility. The Danish First Pregnancy Planner Study Team. *Fertil Steril.* 2000;73(1):43–50.
- Henkel R, Hajimohammad M, Stalf T, Hoogendijk C, Mehnert C, Menkveld R, et al. Influence of deoxyribonucleic acid damage on fertilization and pregnancy. *Fertil Steril.* 2004;81(4):965–72. doi:10.1016/j.fertnstert.2003.09.044.
- Evenson DP, Wixon R. Clinical aspects of sperm DNA fragmentation detection and male infertility. *Theriogenology.* 2006;65(5):979–91. doi:10.1016/j.theriogenology.2005.09.011.
- Wilton LJ, Temple-Smith PD, Baker HW, de Kretser DM. Human male infertility caused by degeneration and death of sperm in the epididymis. *Fertil Steril.* 1988;49(6):1052–8.
- Mallidis C, Lim TC, Hill ST, Skinner DJ, Brown DJ, Johnston WL, et al. Necropermia and chronic spinal cord injury. *Fertil Steril.* 2000;74(2):221–7.
- Nduwayo L, Barthelemy C, Lansac J, Tharanne MJ, Leconte P. Management of necropermia. *Contracept Fertil Sex.* 1995;23(11):682–5.
- Lerda D, Rizzi R. Study of reproductive function in persons occupationally exposed to 2,4-dichlorophenoxyacetic acid (2,4-D). *Mutat Res.* 1991;262(1):47–50.
- Health OCFWs. Sperm Chromatin Structure Assay. <http://www.ohsu.edu/xd/health/services/women/services/fertility/fertility-services/lab-services/sperm-chromatin-structure.cfm>. Accessed April 5, 2014.
- Moskovtsev SI, Librach CL. Methods of sperm vitality assessment. *Methods Mol Biol.* 2013;927:13–9. doi:10.1007/978-1-62703-038-0_2.
- Zini A, Kamal K, Phang D, Willis J, Jarvi K. Biologic variability of sperm DNA denaturation in infertile men. *Urology.* 2001;58(2):258–61.
- WHO laboratory manual for the examination and processing of human semen. 2010. <http://www.who.int/reproductivehealth/publications/infertility/9789241547789/en/>. Accessed March 30, 2014 2014.
- Gosalvez J, Cortes-Gutierrez EI, Nunez R, Fernandez JL, Caballero P, Lopez-Fernandez C, et al. A dynamic assessment of sperm DNA fragmentation versus sperm viability in proven fertile human donors. *Fertil Steril.* 2009;92(6):1915–9. doi:10.1016/j.fertnstert.2008.08.136.
- Arends MJ, Morris RG, Wyllie AH. Apoptosis. The role of the endonuclease. *Am J Pathol.* 1990;136(3):593–608.
- Jaattela M. Programmed cell death: many ways for cells to die decently. *Ann Med.* 2002;34(6):480–8.

22. Gosalvez J, Cortes-Gutierrez E, Lopez-Fernandez C, Fernandez JL, Caballero P, Nunez R. Sperm deoxyribonucleic acid fragmentation dynamics in fertile donors. *Fertil Steril*. 2009;92(1):170–3. doi:10.1016/j.fertnstert.2008.05.068.
23. Cortes-Gutierrez EI, Crespo F, Gosalvez A, Davila-Rodriguez MI, Lopez-Fernandez C, Gosalvez J. DNA fragmentation in frozen sperm of *Equus asinus*: Zamorano-Leones, a breed at risk of extinction. *Theriogenology*. 2008;69(8):1022–32. doi:10.1016/j.theriogenology.2008.02.002.
24. Lopez-Fernandez C, Crespo F, Arroyo F, Fernandez JL, Arana P, Johnston SD, et al. Dynamics of sperm DNA fragmentation in domestic animals II. The stallion. *Theriogenol*. 2007;68(9):1240–50. doi:10.1016/j.theriogenology.2007.08.029.
25. Irvine DS, Twigg JP, Gordon EL, Fulton N, Milne PA, Aitken RJ. DNA integrity in human spermatozoa: relationships with semen quality. *J Androl*. 2000;21(1):33–44.
26. Sun JG, Jurisicova A, Casper RF. Detection of deoxyribonucleic acid fragmentation in human sperm: correlation with fertilization in vitro. *Biol Reprod*. 1997;56(3):602–7.
27. Lopes S, Sun JG, Jurisicova A, Meriano J, Casper RF. Sperm deoxyribonucleic acid fragmentation is increased in poor-quality semen samples and correlates with failed fertilization in intracytoplasmic sperm injection. *Fertil Steril*. 1998;69(3):528–32.
28. Muratori M, Maggi M, Spinelli S, Filimberti E, Forti G, Baldi E. Spontaneous DNA fragmentation in swim-up selected human spermatozoa during long term incubation. *J Androl*. 2003;24(2):253–62.
29. Morris ID, Ilott S, Dixon L, Brison DR. The spectrum of DNA damage in human sperm assessed by single cell gel electrophoresis (Comet assay) and its relationship to fertilization and embryo development. *Hum Reprod*. 2002;17(4):990–8.
30. Brahem S, Mehdi M, Elghezal H, Saad A. Semen processing by density gradient centrifugation is useful in selecting sperm with higher double-strand DNA integrity. *Andrologia*. 2011;43(3):196–202. doi:10.1111/j.1439-0272.2010.01050.x.
31. Zini A, Sigman M. Are tests of sperm DNA damage clinically useful? Pros and cons. *J Androl*. 2009;30(3):219–29.
32. Virro MR, Larson-Cook KL, Evenson DP. Sperm chromatin structure assay (SCSA) parameters are related to fertilization, blastocyst development, and ongoing pregnancy in in vitro fertilization and intracytoplasmic sperm injection cycles. *Fertil Steril*. 2004;81(5):1289–95. doi:10.1016/j.fertnstert.2003.09.063.
33. Dar S, Grover SA, Moskovtsev SI, Swanson S, Baratz A, Librach CL. In vitro fertilization-intracytoplasmic sperm injection outcome in patients with a markedly high DNA fragmentation index (>50%). *Fertil Steril*. 2013;100(1):75–80. doi:10.1016/j.fertnstert.2013.03.011.
34. Cohen-Bacrie P, Belloc S, Menezo YJ, Clement P, Hamidi J, Benkhalifa M. Correlation between DNA damage and sperm parameters: a prospective study of 1,633 patients. *Fertil Steril*. 2009;91(5):1801–5. doi:10.1016/j.fertnstert.2008.01.086.
35. Zini A, Boman JM, Belzile E, Ciampi A. Sperm DNA damage is associated with an increased risk of pregnancy loss after IVF and ICSI: systematic review and meta-analysis. *Hum Reprod*. 2008;23(12):2663–8.
36. Fang S, Baker HW. Male infertility and adult polycystic kidney disease are associated with necrospemia. *Fertil Steril*. 2003;79(3):643–4.
37. Correa-Perez JR, Fernandez-Pelegrina R, Aslanis P, Zavos PM. Clinical management of men producing ejaculates characterized by high levels of dead sperm and altered seminal plasma factors consistent with epididymal necrospemia. *Fertil Steril*. 2004;81(4):1148–50. doi:10.1016/j.fertnstert.2003.09.047.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

