

## RESEARCH ARTICLE

## Open Access

# Food intake during the previous 24 h as a percentage of usual intake: a marker of hypoxia in infants with bronchiolitis: an observational, prospective, multicenter study

François Corrard<sup>1\*</sup>, France de La Rocque<sup>1</sup>, Elvira Martin<sup>1</sup>, Claudie Wollner<sup>1^</sup>, Annie Elbez<sup>1</sup>, Marc Koskas<sup>1,2</sup>, Alain Wollner<sup>1</sup>, Michel Boucherat<sup>1</sup> and Robert Cohen<sup>1,2,3</sup>

## Abstract

**Background:** Hypoxia associated with bronchiolitis is not always easy to assess on clinical grounds alone. The aim of this study was to determine the value of food intake during the previous 24 hours (bottle and spoon feeding), as a percentage of usual intake (24h FI), as a marker of hypoxia, and to compare its diagnostic value with that of usual clinical signs.

**Methods:** In this observational, prospective, multicenter study, 18 community pediatricians, enrolled 171 infants, aged from 0 to 6 months, with bronchiolitis (rhinorrhea + dyspnea + cough + expiratory sounds). Infants with risk factors (history of prematurity, chronic heart or lung disorders), breast-fed infants, and infants having previously been treated for bronchial disorders were excluded.

The 24h FI, subcostal, intercostal, supracostal retractions, nasal flaring, respiratory rate, pauses, cyanosis, rectal temperature and respiratory syncytial virus test results were noted. The highest stable value of transcutaneous oxygen saturation (SpO<sub>2</sub>) was recorded. Hypoxia was noted if SpO<sub>2</sub> was below 95% and verified.

**Results:** 24h FI  $\geq$  50% was associated with a 96% likelihood of SpO<sub>2</sub>  $\geq$  95% [95% CI, 91–99]. In univariate analysis, 24h FI < 50% had the highest odds ratio (13.8) for SpO<sub>2</sub> < 95%, compared to other 24h FI values and other clinical signs, as well as providing one of the best compromises between specificity (90%) and sensitivity (60%) for identifying infants with hypoxia. In multivariate analysis with adjustment for age, SpO<sub>2</sub> < 95% was related to the presence of intercostal retractions (OR = 9.1 [95% CI, 2.4–33.8%]) and 24h FI < 50% (OR = 10.9 [95% CI, 3.0–39.1%]). Hospitalization (17 infants) was strongly related to younger age, 24h FI and intercostal retractions.

**Conclusion:** In practice, the measure of 24 h FI may be useful in identifying hypoxia and deserves further study.

**Keywords:** Bronchiolitis, Hypoxia, Feeding, Infant, Out-patient, Intercostal retraction, Subcostal retraction, Supracostal retractions, Respiratory syncytial virus

\* Correspondence: [francois.corrard@free.fr](mailto:francois.corrard@free.fr)

<sup>^</sup>Deceased

<sup>1</sup>ACTIV (Association Clinique et Thérapeutique Infantile du Val de marne), 27 rue d'Inkermann, 94100, Saint Maur des fossés, France

Full list of author information is available at the end of the article

## Background

Bronchiolitis is the most common viral infection of the lower respiratory tract in infants under 1 year of age [1]. It affects more than 400 000 infants every year in France [2] and is responsible for tens of thousands of hospital admissions. Respiratory syncytial virus (RSV) is usually responsible for the first episode, but other viruses may also be involved. Bronchiolitis is usually mild, resolving spontaneously within a few days, but some infants, especially the very young, develop severe forms with hypoxia and may need hospitalization, primarily for supplemental oxygen administration. Treatment is solely symptomatic, and recent studies have shown that physiotherapy is not beneficial [3,4]. Diagnosis of this infection is often easy, but its severity (hypoxia due to broncho-bronchiolar obstruction) may be difficult to assess. Suckling is a muscular effort that must be coordinated with breathing, and this is more difficult when breathing is restricted by airway obstruction, therefore, reduced feeding might represent a marker of severity.

The amount of food ingested in the past 24 hours is relatively easy to measure accurately, whether taken exclusively from a (graduated) bottle or partly by spoon.

Hypoxia can be diagnosed by non-invasive transcutaneous measurement of oxygen saturation (SpO<sub>2</sub>). Although many inexpensive devices are available, few are suitable for young infants, who require specific, often costly probes that can be difficult to disinfect. In addition, isolated SpO<sub>2</sub> values may not be predictive of subsequent outcome.

The aims of this study were to determine whether food intake in the previous 24 hours, expressed as a percentage of usual intake (24h FI), might serve as an initial screening tool for hypoxia in infants with bronchiolitis, and to analyze the relation between both 24h FI and standard clinical signs and the decision to hospitalize the infant. This tool might be useful for telephone triage or in settings where the history can be used to guide further evaluation.

## Patients and methods

This observational, prospective, multicenter study included infants aged from 0–6 months diagnosed with bronchiolitis (rhinorrhea + cough + dyspnea + expiratory breath sounds) during the three winter months (November to January) when the infection is most frequent in Europe. Infants were excluded if they had risk factors (history of prematurity, chronic heart or lung disease), were breast-fed (even partially), or had previously received treatment for a bronchial disorder (bronchodilators, corticosteroids, physiotherapy). Infants cared for outside the home by child minders could be enrolled, provided the parents knew the precise amount of food ingested in the previous 24 hours. Food intake was calculated as the sum of bottle and spoon feeding. The infants were recruited by 18 community pediatricians in the Paris region.

## Data collection

The pediatrician noted the infant's age, the duration (days) of wheezing, any fever or loose stools in the previous 24 hours, the usual type of feeding, the number of meals taken with a bottle or spoon, and the amount of formula-milk usually drunk, per meal, then calculated the total volume of milk taken over the past 24 hours. If the baby was partly spoon-fed, the pediatrician noted any change in the amount ingested during the previous 24 hours compared to normal.

Quiet, non crying infants were examined for retractions (subcostal, intercostal, suprasternal, and nasal flaring), the respiratory rate (recorded for one minute; rapid if >50/min), cyanosis, and respiratory pauses (during the examination or reported by the parents). SpO<sub>2</sub> was measured only after collecting this information, in order not to interfere with the clinical examination. Rectal temperature was routinely measured.

The pediatricians used a Hellcor pulse oximeter to measure SpO<sub>2</sub>. The prolonged measurements took into account the pulsatile nature of arterial flow (displayed on the device), as well as the highest SpO<sub>2</sub> value recorded (sources of error tend to give lower results) and its stability. Hypoxia was defined as an SpO<sub>2</sub> below 95%. All results below < 95% were verified by a second measurement, and only the highest value was recorded.

RSV was sought routinely (immuno-chromatographic screening test VRSTOP+<sup>®</sup> (ALL.DIAG SA, Strasbourg, France) with nasal swabs).

The doctor recorded the decision regarding immediate hospitalization. The hospital reports were recovered later to determine whether the child had received specific hospital care (oxygen, infusion, or gastric gavage).

## Statistical analysis

The sensitivity, specificity, positive predictive values (PPV), negative predictive values (NPV), positive likelihood ratios (LR<sup>+</sup>) and negative likelihood ratios (LR<sup>-</sup>) were calculated with their 95% confidence intervals, taking an SpO<sub>2</sub> of < 95% as reference, and ROC curves were computed. Means were compared between groups by using a t-test with an unequal variance option if necessary, and percentages were compared by using the chi<sup>2</sup> test or Fisher's exact test, as appropriate. Significance was assumed at  $p < 0.05$ . Univariate and multivariate analyses (logistic regression) were used to identify factors associated with SpO<sub>2</sub> < 95% after adjustment for age (< or ≥ 2 months), and odds ratios (OR) were calculated with their 95%CI. Stata SE 9.1 statistical software was used.

## Ethics

A poster placed in the waiting room invited parents to participate in the study and informed them that they were free to refuse their participation. They were informed that the study was anonymous and would have no influence

on the care received by their child. Furthermore, before enrolment, the investigators informed the parents about the purpose of the study and requested their oral consent. During the period in which this observational study was performed, French legislation did not require ethical approval or other authorizations for research not involving unusual or additional procedures relative to usual practice.

## Results

During three winter periods, from late 2006 to early 2009, the participating pediatricians recruited 171 infants with a mean age of  $1.6 \pm 3.7$  months (median 4 months). The infants were seen between the first and sixth day after the onset of chest sounds (average 2 days,  $\pm 1.5$  days), as reported by the parents.

During the previous 24 hours, 22% of the infants had been febrile ( $\geq 38^\circ\text{C}$ ) (21% were febrile at the time of the examination), and 14% had had three or more loose stools.

Feeding consisted exclusively of formula-milk in 87% of infants, while the remaining 13% of infants, all aged more than 4 months, were partly spoon-fed. Food intake in the previous 24 hours (24h FI), as a percentage of the usual amount, was less than 50% in 14% of cases,  $\geq 50\%$  to  $< 70\%$  in 26% of cases, and  $\geq 70\%$  in 60% of cases.

Nasal flaring and suprasternal, intercostal or subcostal retractions were present in respectively 2%, 15%, 25% and 30% of infants.

The respiratory frequency was  $\geq 50/\text{min}$  in 43% of infants and  $\geq 60/\text{min}$  in 23% of infants.

Cyanosis was noted in 5 infants, one of whom also had breathing pauses.

Mean oxygen saturation was  $98\% \pm 3\%$  (80% to 100%; median 98%). Fifteen infants (9%) had  $\text{SpO}_2 < 95\%$  and 2 infants had  $\text{SpO}_2 < 90\%$ .

RSV was tested for in 104 infants, of whom 42% were positive.

Seventeen of the 171 infants were hospitalized immediately after the examination. Ten of the hospitalized infants received specific hospital care, consisting of oxygen in 9 cases, gastric gavage in 6 cases and infusion in 5 cases.

### Relation between $\text{SpO}_2$ and 24h FI

The closer food intake in the previous 24 hours was to the usual amount, the closer  $\text{SpO}_2$  was to normal: higher 24h FI values were associated with higher  $\text{SpO}_2$  (Figure 1). The average  $\text{SpO}_2$  in infants whose food intake fell by more than 50% was 95.5% [95% CI, 93.6-97.4%], compared to 98.1% [95% CI, 97.7-98.3%] in the infants whose food intake remained higher than 50% of normal ( $p < 0.001$ ). In the latter group,  $\text{SpO}_2$  was normal ( $\geq 95\%$ ) in 96% of cases [95% CI, 91-99%].

When food intake during the previous 24 hours was 50% or more of the usual amount, the likelihood that

saturation would be normal ( $\text{SpO}_2 \geq 95\%$ ) was 96% [95% CI, 91%-99%].

### Relation between $\text{SpO}_2$ , clinical status and food intake

The positive predictive values (PPV) of 24h FI for  $\text{SpO}_2 < 95\%$  were highest with cutoffs of  $< 40\%$  and  $< 50\%$  (Table 1). However, 24h FI  $< 50\%$  had the highest specificity (90%) and moderate sensitivity (60%) (Figure 2), and also the highest odds ratio of all 24h FI cutoffs in univariate analysis (Table 1).

In multivariate analysis, the only variables significantly associated with  $\text{SpO}_2 < 95\%$  after adjustment for age were intercostal retraction (OR = 9.1 [95% CI 2.4 to 33.8]) and 24h FI  $< 50\%$  (OR = 10.9 [95% CI 3.0 to 39.1]).

### 24h FI and hospitalization


The 17 hospitalized infants had significantly lower  $\text{SpO}_2$  values ( $94\% \pm 5\%$  versus  $98\% \pm 2\%$ ,  $p < 0.005$ ) and significantly lower 24h FI values ( $56\% \pm 24\%$  versus  $76 \pm 19\%$ ,  $p < 0.0003$ ) than the non-hospitalized infants.

Nine of the 17 hospitalized infants (Table 2) had both 24h FI  $< 50\%$  and  $\text{SpO}_2 < 95\%$ , while 24h FI was  $\geq 50\%$  in 90% of the non-hospitalized infants ( $p < 0.001$ ). Six hospitalized infants had 24h FI  $> 50\%$  and no obvious signs of respiratory distress, but all were very young ( $< 2$  months). Four of these six infants did not require specific hospital care. The fifth child had a 24h FI value very close to the cutoff (52%), with rapid breathing, intercostal retractions and an  $\text{SpO}_2$  of 92%. The sixth child had a 24h FI of 67% and an  $\text{SpO}_2$  of 92%, but also had high fever ( $39.6^\circ\text{C}$ ) during the previous 24 hours and rapid breathing (55 breaths/min), probably due to fever. This infant did not require specific hospital care.

In multivariate analysis, the decision to hospitalize was very strongly associated with  $\text{SpO}_2 < 95\%$  ( $p < 0.0001$ ) and age  $< 2$  months ( $p = 0.001$ ), but not with the other parameters. However, when saturation was excluded, the decision to hospitalize was strongly associated with age  $< 2$  months (OR = 14.7 [95% CI 3.1 to 69.8]), 24h FI  $< 50\%$  (OR = 10.6 [95% CI 3.0 to 37.3]) and intercostal retractions (OR = 3.4 [95% CI 1.0 to 11.4]). Thus, if  $\text{SpO}_2$  measurement had not been available, the decision to hospitalize could have been based on age  $< 2$  months, 24h FI  $< 50\%$ , and intercostal retractions.

## Discussion

This study suggests that recent changes in food intake by infants with bronchiolitis closely reflect their oxygenation status. The 24h FI cutoff most strongly associated with hypoxia (odds ratio = 13.8 [4.3 to 44.1]) was 50% of the usual amount, and this cutoff also had high negative predictive value. In the study population, there were 14 infants who took less than 50% of their usual food intake, and only one of them was not hypoxic. This 50% cutoff




also represented the best compromise (moderate sensitivity, high specificity) for identifying infants with SpO<sub>2</sub> < 95%. In addition, multivariate analysis showed that this parameter remained correlated with SpO<sub>2</sub> < 95%, even after adjustment for age and intercostal retractions, while the other clinical and biological parameters showed no significant correlation with SpO<sub>2</sub>. This cutoff (50%) would

have two advantages. First, if food intake during the past 24 hours remains more than half the usual amount, the likelihood of normal oxygenation status is 96% (or at least 91%, the lower limit of the confidence interval), indicating that bronchiolar obstruction is well tolerated and that pulse oximetry is probably unnecessary in the absence of associated gravity signs. In contrast, food intake below

**Table 1 Sensitivity, specificity, PPV, NPV, LR<sup>+</sup>, LR<sup>-</sup> and odds ratios of clinical signs of hypoxia (SpO<sub>2</sub> < 95%) in infants with bronchiolitis**

	Sensitivity % [95% IC]	Specificity % [95% IC]	PPV % [95% IC]	NPV % [95% IC]	LR <sup>+</sup> n [95% IC]	LR <sup>-</sup> n [95% IC]	Odds ratio n [95% IC]
Suprasternal retraction	50 [21–79]	87 [81–92]	24 [9–45]	96 [91–98]	4 [2–8]	0,6 [0,3 - 1]	6,9 [2,0 - 23,7]
Intercostal retraction	73 [45–92]	80 [73–86]	26 [14–42]	97 [92–99]	3,6 [2–6]	0,3 [0,1 - 78]	10,8 [3,2 - 36,3]
Subcostal retraction	47 [21–73]	72 [64–79]	14 [6–27]	93 [87–97]	1,7 [0,9 - 3]	0,7 [0,5 - 1,2]	2,3 [0,8 - 6,6]
Polypnea (≥ 50/min)	87 [60–98]	62 [53–70]	19 [10–30]	98 [93–100]	2,3 [1,7 - 3]	0,2 [0,06 - 0,8]	10,5 [2,3 - 48,2]
24h FI < 70%	87 [60–98]	65 [56–72]	19 [11–31]	98 [93–100]	2,4 [1,8 - 3,2]	0,2 [0,06 - 0,8]	11,8 [2,6 - 54,2]
24h FI < 60%	67 [38–88]	78 [70–84]	23 [11 - 38]	96 [91–99]	3 [1,9 - 4,7]	0,4 [0,2 - 0,8]	6,9 [2,2 - 21,7]
<b>24h FI &lt; 50%</b>	<b>60</b> [32–84]	<b>90</b> [84–94]	<b>38</b> [19–59]	<b>96</b> [91–99]	<b>6,1</b> [3–11]	<b>0,4</b> [0,2 - 0,8]	<b>13,8</b> [4,3 - 44,1]
24h FI < 40%	33 [12–62]	96 [92–99]	46 [17–77]	94 [89–97]	8,5 [2,9 - 24]	0,7 [0,5 - 0,99]	12,2 [3,2 - 47,2]

24h FI food intake in previous 24h as a % of usual intake; PPV positive predictive value ; NPV negative predictive value ; LR<sup>+</sup> positive likelihood ratio ; LR<sup>-</sup> negative likelihood ratio; [95% CI] 95% confidence interval.



50% of normal calls for pulse oximetry and medical attention.

This 50% cutoff has a certain tolerance: if 24h FI is 40%, then the negative predictive value only falls from 96% to 94% and the odds ratio from 13.8 to 12.2. The clinical implications therefore remain valid.

24h FI might be used as a screen to further testing. This might be useful for telephone triage or other settings where the history can be used to guide further evaluation, as monitoring a bronchiolitic child in the home.

**Table 2 Factors associated with hospitalization for bronchiolitis**

	Hospitalized (n = 17)		Non hospitalized (n = 154)		p
	n	%	n	%	
Age < 2 months	13	76%	7	5%	< 0.001
24h FI < 50%	9	53%	15/150 (*)	10%	< 0.001
Intercostal retraction	9	53%	33/151 (*)	22%	< 0.005
1 or more of these 3 factors	14	82%	49/147 (*)	33%	< 0.001
SpO <sub>2</sub> < 95%	11	65%	4	3%	< 0.001

\* missing data.

Food intake has already been included in severity scores and guidelines, but only as “reduced or normal” [2,5-7], or “good, reduced or poor” [8-10]. The November 2006 Scottish guideline [11] mentioned a < 50% reduction in fluid intake in the previous 24 hours as a gravity sign, but this was based on expert opinion and not on published research.

Physiological studies of milk intake by infants with bronchiolitis [12] have shown that the number of sucking and swallowing actions is the same as in healthy infants, but that milk flow during these two movements is reduced. In addition, swallowing movements are followed less by expiration and more by inspiration, and periods of apnea are twice as frequent after swallowing. Suckling is a muscular effort that must be coordinated with breathing, and this is more difficult when breathing is restricted by airway obstruction, when the child is tired, and when the muscles lack oxygen [13].

Parents are generally aware of the amount of food their infant usually consumes, and can accurately recall food intake in the previous 24 hours, based on the amount prepared and the amount left, especially when their child is sick. Indeed, parents prepare the same

quantity of milk (powder scoops and bottle graduations) each day for a given meal. The amount left in the bottle by healthy infants does not vary much, but increases during bronchiolitis. Thus, in practice, 24-hour percentage food intake is easy to calculate.

In theory, SpO<sub>2</sub> would be an accurate marker of hypoxia, which is directly related through the hemoglobin dissociation curve. Hypoxia is the main complication of bronchiolitis. However, SpO<sub>2</sub> is tricky to measure, requiring a considerable period of calm, and a correctly recorded arterial pulse. Correct sensor positioning may take several attempts. In addition, pediatric sensors are fragile, difficult to disinfect correctly, and expensive for single use. In the range of values generally measured in this setting (85-100%), a small change in SpO<sub>2</sub> is associated with a large change in PaO<sub>2</sub>. The result is also observer-dependent and may vary over a 30-minute interval [14]. This is why we verified all values below < 95% by a second measurement, and only used the highest value.

The 95% SpO<sub>2</sub> cutoff is commonly used to define hypoxia, and oxygen therapy is recommended if SpO<sub>2</sub> is < 90% for a prolonged period in an infant not at risk (term birth, no heart or lung disease and age more than 3 months) [5].

Among the three types of retraction (suprasternal, intercostal and subcostal), intercostal retractions were most closely associated with SpO<sub>2</sub> < 95%. Wang [13] reported that the appreciation of retractions is observer-dependent, even when the observer is a healthcare professional, and that the appreciation may vary between two separate observations made 10 to 30 minutes apart. The degree of retractions is difficult to quantify, especially in borderline situations. In addition, they are difficult for parents to assess without specific training. Retractions are included in most scores, but these are not always easy to use in daily practice. Measurement of 24 hours percentage food intake seems to be far simpler for parents.

Hospitalization was associated with hypoxia and with age < 2 months (French guidelines recommend hospitalization of all bronchiolitic infants less than 6 weeks old). Hypoxia was associated with 24h FI < 50% and intercostal retractions. In the physician's office, if SpO<sub>2</sub> cannot be measured, these two signs, especially 24h FI < 50%, in addition to age, can help to decide whether hospitalization is necessary.

This study may have certain limitations. First, the same pediatrician assessed all the data (24h FI, clinical signs of respiratory impairment, oxymetry) and took the decision regarding hospitalization. We tried to limit the subjective element by establishing the order of data collection and applying precautions during oxymetry.

Second, these results are applicable to patients presenting to community practices. Further studies of sicker infants (emergency departments, hospital inpatients) would be necessary to determine whether 24h FI remains valid in this

population. It should be noted that most studies of severe adverse outcomes in this setting have involved inpatients, which limits their relevance to other situations [5]. Finally, we excluded breast-feeding infants, as the amount of milk ingested could not easily be estimated, notably from the suckling time [15].

## Conclusion

We studied the semiological value of a clinical sign for assessing the severity of bronchiolitis in previously healthy, mainly bottle-fed, full-term infants presenting to a pediatrician's office.

Food intake during the previous 24 hours, as a percentage of usual intake, was predictive of oxygenation status (hypoxia or normoxia), and is easy for parents to memorize and measure. Apart from in very young infants (less than 6 weeks old), this parameter could serve as an initial screening tool and to monitor bronchiolitic child in the home.

If an infant has ingested at least half the usual amount of food during the previous 24 hours, hypoxia is unlikely; in contrast, if food intake falls below half the usual amount, medical attention is required.

French version of the manuscript : "see Additional file 1".

## Additional file

**Additional file 1: Bronchiolite et prise alimentaire des dernières 24 heures: un outil de dépistage de l'hypoxie (The French version of the manuscript).**

## Abbreviations

24h FI: Food intake during the previous 24 hours as a percentage of usual intake; ACTIV: Association Clinique et Thérapeutique Infantile du Val de marne, a research institute on pediatric community acquired infections; CI: Confidence interval; LR<sup>+</sup>: Positive likelihood ratio; LR<sup>-</sup>: Negative likelihood ratio; NPV: Negative Predictive Value; OR: Odds ratio; PPV: Positive predictive value; RSV: Respiratory Syncytial Virus; SpO<sub>2</sub>: Saturation of hemoglobin with oxygen as measured by pulse oxymetry.

## Competing interests

All the authors declare that no financial support for the submitted work was received from anyone other than their employer; that they have no financial relationships with commercial entities that might have an interest in the submitted work in the previous 3 years; that they have no spouses, partners, or children with financial relationships that may be relevant to the submitted work; and that they have no non-financial interests that may be relevant to the submitted work.

This work was submitted by ACTIV. The funders of ACTIV had no role in the study design, data collection or analysis, the decision to publish, or the preparation of the manuscript.

## Authors' contributions

Designed the study: FC FLR EM CW AE MK AW MB RC. MB designed the data base. Enrolled patients: FC CW AE MK AW RC. Analysed the data: FC FLR EM MK RC. Contributed to the writing of the paper: FC EM MK RC. Agree with the manuscript's results and conclusions: FC FLR EM AE MK AW MB RC. All authors read and approved the final manuscript.

## Authors' information

François Corrard is a French pediatric and President of a research institute on pediatric community acquired infections (ACTIV). His main research interests

are epidemiologic studies and clinical trials in community acquired infections, including pneumococcal diseases, the rhinopharyngeal flora, and vaccines. He has published more than 10 papers in English.

#### Acknowledgments

This work is dedicated to the memory of Claudie Wollner. We thank the other investigating pediatricians (Christian Copin, Patrice Deberdt, Solange Duriez, Geneviève Granat, Eric Sandrin, Anne-Sylvestre Michot, Mohammed Benani, Allegra Brami, Jean Pierre Henon, Sydney Sebban, Marie Héléne A'kou and Eric Osika), Philippe d'Athis for his statistical expertise and Nicole Beydon for proof-reading the manuscript. None of the physicians listed received any financial compensation for their contributions. We thank all the patients and their parents who participated in this study. We also thank Corinne Levy, (ACTIV), Manuela Pereira (ACTIV), Sadia Tortorelli (ACTIV) and Stéphane Béchet (ACTIV) for their technical assistance.

#### Author details

<sup>1</sup>ACTIV (Association Clinique et Thérapeutique Infantile du Val de marne), 27 rue d'Inkermann, 94100, Saint Maur des fossés, France. <sup>2</sup>Physiology Lung Function Department Armand-Trousseau Hospital, Paris, France. <sup>3</sup>Department of Microbiology, CHI Créteil, 40 avenue de Verdun, Créteil, France.

Received: 25 March 2012 Accepted: 8 January 2013

Published: 11 January 2013

#### References

1. Carraro S, Zanconato S, Baraldi E: **Bronchiolitis: from empiricism to scientific evidence.** *Minerva Pediatr* 2009, **61**(2):217–225.
2. Stagnara J, Balagny E, Cossalter B, Dommerges JP, Dournel C, Drahi E, Gauchez H, Guillot F, Javault D, Lagardere B, et al: **Management of bronchiolitis in the infant. Recommendations. Long text.** *Arch Pediatr* 2001, **8**(1):115–235.
3. Gajdos V, Katsahian S, Beydon N, Abadie V, de Pontual L, Larrar S, Epaud R, Chevallier B, Bailleux S, Mollet-Boudjemline A, et al: **Effectiveness of chest physiotherapy in infants hospitalized with acute bronchiolitis: a multicenter, randomized, controlled trial.** *PLoS Med* 2010, **7**(9): e1000345.
4. Roquéi Figuls M, Giné-Garriga M, Granados Rugeles C, Perrotta C: **Chest physiotherapy for acute bronchiolitis in paediatric patients between 0 and 24 months old.** *Cochrane Database Syst Rev* 2012, **2**:CD004873. doi:10.1002/14651858.CD004873.pub4.
5. American Academy of Pediatrics, Subcommittee on diagnosis and management of bronchiolitis, Clinical practice guideline: **Diagnosis and management of bronchiolitis.** *Pediatrics* 2006, **118**(4):1774–1793.
6. Perlstein PH, Kotagal UR, Bolling C, Steele R, Schoettker PJ, Atherton HD, Farrell MK: **Evaluation of an evidence-based guideline for bronchiolitis.** *Pediatrics* 1999, **104**(6):1334–1341.
7. Shaw KN, Bell LM, Sherman NH: **Outpatient assessment of infants with bronchiolitis.** *Am J Dis Child* 1991, **145**(2):151–155.
8. Barben J, Hammer J: **Treatment of acute bronchiolitis in newborns.** *Pediatrics* 2003, **14**:22–23. http://wwwswiss-paediatricsorg/paediatrica/vol14/n6/bronchiolite-frhtml.
9. Dawson K, Kennedy D, Asher I, Cooper D, Cooper P, Francis P, Henry R, Le Souef P, Martin J, Masters B, et al: **The management of acute bronchiolitis. Thoracic Society of Australia and New Zealand.** *J Paediatr Child Health* 1993, **29**(5):335–337.
10. Simoes EA: **Respiratory syncytial virus infection.** *Lancet* 1999, **354**(9181):847–852.
11. Baumer JH: **SIGN guideline on bronchiolitis in infants.** *Arch Dis Child Educ Pract Ed* 2007, **92**(5):ep149-151.
12. Pinnington LL, Smith CM, Ellis RE, Morton RE: **Feeding efficiency and respiratory integration in infants with acute viral bronchiolitis.** *J Pediatr* 2000, **137**(4):523–526.
13. Verges S, Bachasson D, Wuyam B: **Effect of acute hypoxia on respiratory muscle fatigue in healthy humans.** *Respir Res* 2010, **11**:109.

14. Wang EE, Law BJ, Stephens D, Langley JM, MacDonald NE, Robinson JL, Dobson S, McDonald J, Boucher FD, de Carvalho V, et al: **Study of interobserver reliability in clinical assessment of RSV lower respiratory illness: a Pediatric Investigators Collaborative Network for Infections in Canada (PICNIC) study.** *Pediatr Pulmonol* 1996, **22**(1):23–27.
15. Riordan J, Gill-Hopple K, Angeron J: **Indicators of effective breastfeeding and estimates of breast milk intake.** *J Hum Lact* 2005, **21**(4):406–412.

doi:10.1186/1471-2431-13-6

**Cite this article as:** Corrard et al.: Food intake during the previous 24 h as a percentage of usual intake: a marker of hypoxia in infants with bronchiolitis: an observational, prospective, multicenter study. *BMC Pediatrics* 2013 **13**:6.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

