

POSTER PRESENTATION

Open Access

Prevalence of HIV, HBV and chlamydia infections in Cameroonian University context: case of the University of Dschang, in the Western region

Martin Sanou Sobze*, Jean Hubert Donfack, James-Francis Onohiol, Joseph Fokam, Ghislaine Bruna Djeunang Dongho, Patrick Stephane Nkamedjie Pete, Vittorio Colizzi, Gianluca Russo

From International Symposium HIV and Emerging Infectious Diseases 2014
Marseille, France. 21-23 May 2013

Introduction

In sub-Saharan Africa HIV infection remains largely epidemic, whereas HBV infection is highly endemic (>8%). In Cameroon, the prevalence of HIV is 4.3%. Concerning HBV and chlamydia infections, their prevalence are both $\geq 10\%$. Teenagers and young adults, including university students, are the population groups mostly affected by STDs. Epidemiological data on these infections, among university students could be helpful to implement specific prevention strategies.

Methods

A descriptive study was performed in May 2013 among 624 students from the University of Dschang, Cameroon. Participants were screened for HIV, HBV and Chlamydia infections. Data was collected by a standard questionnaire and analyzed by Epi Info.

Results

The average age of participants was 23.3 years ($\sigma = 3.2$) with a female predominance (58.7%). The observed prevalence of HIV, HBV and Chlamydia infection was 1.1% (7/624), 2.8% (5/176) and 2.0% (2/100) respectively. 83.2% of participants were sexually active. Concerning sexual risky behaviors, participants reported having multi partners (14.8%), using condom regularly (36.4%), occasionally (58.6%) or never (5.0%). 100%, 62.6% and 52.2% reported to be aware on HIV, HBV and Chlamydia infections respectively. In addition, only 5.5% and 21.3% of the participants were aware of their HBV and Chlamydia status respectively, versus 64.4% for HIV. The excessive cost of

HBV and Chlamydia tests has been identified as the major barrier to testing (87.6%).

Conclusion

Among college Cameroonian students the prevalence of HIV, HBV and Chlamydia infections seems to be relatively low if compared to general population. However, having multiple sexual partners in addition to non-systematic use of condoms during sexual intercourse represents risky behaviors among students. Awareness campaigns and screening facilitation on HBV and chlamydia infections among students need to be strengthened.

Published: 23 May 2014

doi:10.1186/1471-2334-14-S2-P3

Cite this article as: Sobze et al.: Prevalence of HIV, HBV and chlamydia infections in Cameroonian University context: case of the University of Dschang, in the Western region. *BMC Infectious Diseases* 2014 14(Suppl 2):P3.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Department of Biomedical Sciences, Faculty of Sciences, University of Dschang, Dschang, Cameroon



© 2014 Sobze et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.