



Genetic evolution of uveal melanoma guides the development of an inflammatory microenvironment

Gülçin Gezgin¹ · Mehmet Dogrusöz¹ · T. Huibertus van Essen¹ · Wilhelmina G. M. Kroes² · Gregorius P. M. Luyten¹ · Pieter A. van der Velden¹ · Vonn Walter³ · Robert M. Verdijk⁴ · Thorbald van Hall⁵ · Sjoerd H. van der Burg⁵ · Martine J. Jager¹

Received: 5 September 2016 / Accepted: 20 March 2017
© The Author(s) 2017. This article is an open access publication

Abstract Uveal melanoma (UM) is characterized by a number of genetic aberrations that follow a certain chronology and are tightly linked to tumor recurrence and survival. Loss of chromosome 3, bi-allelic loss of BAP1 expression, and gain in chromosome 8q have been associated with metastasis formation and death, while loss of chromosome 3 has been associated with the influx of macrophages and T cells. We used a set of genetically-classified UM to study immune infiltration in the context of their genetic evolution. We show in two independent cohorts that lack of BAP1 expression is associated with an increased density of CD3⁺ T cells and CD8⁺ T cells. The presence of extra copies of chromosome 8q in disomy 3 tumors with a normal BAP1 expression is associated with an increased influx of macrophages (but not T cells). Therefore, we propose that the genetic evolution of UM is associated with changes in the inflammatory phenotype. Early changes resulting in gain of chromosome 8q may activate macrophage infiltration, while sequential loss of BAP1 expression seems to drive T cell infiltration in UM.

Keywords BAP1 · Infiltration · Macrophages · T cells · Lymphocytes · Chromosome

Abbreviations

<i>BAP1</i>	BRCA1-associated protein 1
<i>CCL2</i>	Chemokine (C-C motif) ligand 2 (Monocyte chemoattractant protein-1)
<i>CCL3</i>	Chemokine (C-C motif) ligand 3 (Macrophage inflammatory protein 1 α)
<i>CCL5</i>	Chemokine (C-C motif) ligand 5 (RANTES)
<i>CXCL10</i>	Chemokine (C-X-C motif) ligand 10 (Interferon gamma-induced protein 10)
<i>CXCL12</i>	Chemokine (C-X-C motif) ligand 12 (Stromal cell-derived factor 1)
<i>dPCR</i>	Digital polymerase chain reaction
<i>HLA</i>	Human leukocyte antigen
<i>IF</i>	Immunofluorescence
<i>IHC</i>	Immunohistochemistry
<i>NF-κB</i>	Nuclear factor κ B
<i>SNP</i>	Single-nucleotide polymorphism
<i>TCGA</i>	The Cancer Genome Atlas
<i>UCH</i>	Ubiquitin-carboxy-terminal hydrolase
<i>UCHL1</i>	Ubiquitin carboxyl-terminal hydrolase L1
<i>UCHL2</i>	Ubiquitin carboxyl-terminal hydrolase L2
<i>UM</i>	Uveal melanoma
<i>VEGFA</i>	Vascular Endothelial Growth Factor A

✉ Martine J. Jager
M.J.Jager@lumc.nl

¹ Department of Ophthalmology, Leiden University Medical Center, PO Box 9600, 2300 RC Leiden, The Netherlands

² Department of Clinical Genetics, Leiden University Medical Center, Leiden, The Netherlands

³ Department of Biochemistry, Penn State Milton S. Hershey Medical Center, Hershey, PA, USA

⁴ Department of Pathology, Section Ophthalmic Pathology, Erasmus MC University Medical Center, Rotterdam, The Netherlands

⁵ Department of Clinical Oncology, Leiden University Medical Center, Leiden, The Netherlands

Introduction

Uveal melanoma (UM) is the most common primary intraocular malignancy in Caucasian adults and may lead to metastatic disease in up to 50% of patients [1, 2]. Current treatments are hardly ever effective against metastases [3], and hence, most research efforts are focused on the

development of targeted therapies or immunotherapeutic approaches, such as treatments with immune checkpoint inhibitors, vaccination, or adoptive T cell therapy [4–8].

Some UM express increased levels of human leukocyte antigen (HLA) class I and are infiltrated by macrophages and lymphocytes, and this is known as an inflammatory phenotype [9–12]. The presence of this inflammatory phenotype has been correlated with a specific genetic aberration, which is the loss of one copy of chromosome 3 (monosomy 3) [13]. Other chromosomal abnormalities frequently occur in chromosomes 1, 6, and 8 [14–17]. Following an initiating mutation in either GNAQ or GNA11, gain of 8q is thought to be one of the earliest genetic aberrations, followed by loss of one chromosome 3 [18, 19]. Gain of 8q and monosomy 3 and are both associated with the development of UM metastases and a poor prognosis [16, 20]. Similarly, gene expression analysis has been used to divide UM into two major classes, 1 and 2, which are good predictors of prognosis [21, 22]. Moreover, we recently showed that, in our hands, class II tumors can be subdivided into IIa and IIb: while class IIa tumors are composed of highly homogeneous tumor cells and class IIb tumors contain a larger percentage of non-tumor cells which are likely to be immune cells. Interestingly, class IIa and IIb tumors differed in their numbers of chromosome 8q copies [19].

Chromosome 3 contains the gene for *BRCA1-associated protein 1* (BAP1). In UM, inactivating hemizygous mutations in this gene have been found [23, 24], which are associated with loss of BAP1 protein expression and a high metastatic risk [23–26]. However, monosomy 3 and loss of BAP1 may occur independently, as tumors with a normal chromosome 3 status with lack of BAP1 expression have been identified, as well as tumors with monosomy 3 that still express BAP1. Such atypical tumors show high-risk clinico-pathological features and convey an increased metastatic risk [25]. BAP1 is a member of the ubiquitin-carboxy-terminal hydrolase (UCH) family [27], and is also known as ubiquitin carboxyl-terminal hydrolase L2 (UHL2). Another member of the UCH family, ubiquitin carboxyl-terminal hydrolase L1 (UHL1), is associated with suppressed production of pro-inflammatory chemokines and cytokines in keratinocytes [28]. We, therefore, hypothesized that loss of BAP1 expression might be related to macrophage and/or T cell infiltration.

To assess whether genetic alterations affect immune cell infiltration in UM, we studied the presence and type of tumor-infiltrating immune cells in UM subtypes consisting of typical cases of UM (e.g. disomy 3 tumors with BAP1 expression and monosomy 3 with loss of BAP1 expression) and atypical cases of UM (e.g. disomy 3 tumors with loss of BAP1 expression and monosomy 3 tumors with expression of BAP1). In addition, we studied disomy 3/BAP1-positive cases with and without extra copies of chromosome 8q.

Our data show that loss of BAP1 protein expression is predominantly related to T cell infiltration in UM, while early gain of chromosome 8q is associated with macrophage infiltration.

Materials and methods

Patient selection

Tumor tissue specimens were obtained from 84 UM patients. Patients underwent primary enucleation for UM between 1999 and 2008 at the Leiden University Medical Center (Leiden, The Netherlands). Part of the tumor was snap frozen using 2-methyl butane, and DNA and RNA were isolated. The remaining tumor was fixed in 4% neutral-buffered formalin for 48 h and subsequently embedded in paraffin. This study was approved by the Medical Ethics Committee of the Leiden University Medical Center. Tumor material was handled according to the Dutch National Ethical Guidelines ('Code for Proper Secondary Use of Human Tissue'), and the tenets of the Declaration of Helsinki (World Medical Association of Declaration 2013; ethical principles for medical research involving human subjects). In addition, 80 patients from The Cancer Genome Atlas (TCGA) Project on UM were included in this study as an independent validation cohort.

Immunohistochemistry and fluorescent immunostaining of the Leiden Cohort

4- μ m serial sections from paraffin-embedded tissue were cut and used for immunostaining. Immunohistochemistry (IHC) of BAP1 was performed on 74 tumors, as described previously [24]. Tumors were scored by two independent investigators as BAP1-positive or -negative based on nuclear staining. Immunofluorescence (IF) staining for T cells and macrophages was performed on 43 tumors as described previously [29, 30]. T cell types were detected by primary antibodies: anti-CD3 (ab828, rabbit polyclonal; Abcam, Cambridge, MA, United States of America) and anti-CD8 (4B11, mouse monoclonal IgG2b; Novocastra, Valkenswaard, The Netherlands). To visualize the T cells, the following secondary antibodies were used: goat-anti-rabbit IgG Alexa 546 and goat-anti-mouse IgG2b Alexa 647 (Molecular Probes, Invitrogen, Breda, The Netherlands). Counts of intratumoral CD3⁺ and CD8⁺ T cells were represented as the number of cells per square millimeter. For IF staining of CD68⁺ macrophages, we used the primary mouse anti-human macrophage CD68 antibody (clone 514H12; ab49777; Abcam, Cambridge, United Kingdom), and as secondary antibody AlexaFluor IgG2a

(488) goat-anti-mouse. The amount of CD68⁺ expression was determined in pixels per square millimeter.

DNA and gene expression analysis

DNA and gene expression analysis were performed on 54 tumor specimens from Leiden, in which the BAP1 status was known. The QIAmp DNA Mini kit was used to isolate DNA for single-nucleotide polymorphism (SNP) analysis (Qiagen, Venlo, The Netherlands). SNP analysis was then performed with the Affymetrix 250K_NSIP microarray and Affymetrix Cytoscan HD chip (Affymetrix, Santa Clara, California, United States of America) to detect aberrations of chromosome 3 as described previously [20]. Information on chromosome 8q was obtained by digital polymerase chain reaction (dPCR) [20]. A threshold of >2.1 was defined as having extra copies of chromosome 8q. The RNeasy mini kit was used to isolate RNA for gene expression analyses (Qiagen, Venlo, The Netherlands). Gene expression levels of *CD3* and *CD8* (T cells), *CD68* (macrophages), and pro-inflammatory cytokines, specifically *macrophage inflammatory protein 1 α* (*CCL3*), *vascular endothelial growth factor A* (*VEGFA*), *stromal cell-derived factor 1* (*CXCL12*), *CCL7*, *CSF-1*, *monocyte chemoattractant protein-1* (*CCL2*), *RANTES* (*CCL5*), *interferon gamma-induced protein 10* (*CXCL10*), *CCR7*, and *CXCR4* were obtained using the Illumina HT-12 v4 chip (Illumina, San Diego, California, United States of America). The pro-inflammatory cytokines were selected based on our previous papers [31–33]. We could validate the probes for *CD3*, *CD8* and *CD68* in 24 tumors in which gene expression levels had been determined with an Illumina HT12 v4 array and in which the number of infiltrating cells was analyzed by IF. In addition, data on RNA sequencing and Affymetrix SNP 6.0 array from 80 samples of UM were obtained from the TCGA Research Network: <http://cancergenome.nih.gov/>. Copy numbers for 8q were determined by Affymetrix SNP 6.0 array and analyzed with the GISTIC 2.0 algorithm [34, 35]. Copy numbers >2 were categorized as extra copies of chromosome 8q. *BAP1*, *CD68*, *CD3*, and *CD8* expression were obtained by RNA sequencing and quantified as $\log_2(\text{RSEM}+1)$. *BAP1* expression was dichotomized into *BAP1*-positive and *BAP1*-negative expression at the median.

Statistical analysis

Analyses were performed using SPSS version 20.0.0 (IBM SPSS Statistics, IBM Corporation, Armonk, New York, United States of America) and graphs were made using GraphPad Prism version 5.0 for Windows (GraphPad Software, La Jolla, California, United States of America, <http://www.graphpad.com>). The Mann–Whitney *U* test was

applied for continuous parameters. Correlation analyses were performed with Spearman's rho correlation test. A *P* value <0.05 was considered statistically significant.

Results

Loss of BAP1-protein expression is associated with an inflammatory phenotype in UM

As expression of UCHL1, a member of the UCH family, has been associated with suppression of pro-inflammatory cytokines and chemokines in keratinocytes [28], we determined whether loss of expression of another UCH-member, BAP1 might be correlated to the development of an inflammatory phenotype in UM. We quantified the number of tumor-infiltrating macrophages and T cells by immunofluorescence staining of tissue sections in a cohort of 20 BAP1-immunopositive tumors and 23 BAP1-immunonegative tumors from the Leiden cohort, and found that BAP1-negative tumors contained significantly higher numbers of CD3⁺ T cells (*P*=0.002), CD8⁺ T cells (*P*=0.003), and CD68⁺ macrophages (*P*<0.001) (Fig. 1a–b).

To corroborate our findings, we analyzed gene expression array data from 54 cases of UM in which BAP1 expression as well as chromosome 3 and 8q status had been determined. We first correlated potentially-useful probes by comparing IF staining data for CD68⁺ macrophages, CD3⁺ and CD8⁺ T cells with their corresponding CD68, CD3, and CD8 probes used for RNA gene expression analysis in a group of 24 patients for whom both tests were available. The probes with the highest correlation were used for further analyses (Table 1), since we assumed that these probes represent the corresponding protein expression most accurately. We subsequently compared the presence of specific probes with BAP1 expression: a higher expression of *CD3* (*P*=0.016), *CD8* (*P*=0.015), and *CD68* (*P*=0.002) was observed in tumors that did not express BAP1 (*n*=24) compared to tumors that did express BAP1 (*n*=30; Fig. 1c), corroborating our results obtained with IF staining (Fig. 1a, b). These results indicate that loss of BAP1 expression in UM is associated with a higher T cell and macrophage infiltration.

Since most monosomy 3 tumors lack BAP1 expression, it is difficult to determine whether the effects observed are due to loss of BAP1 expression or due to loss of one chromosome 3. We therefore, focused on atypical cases, and observed that within the disomy 3 tumors, loss of BAP1 protein expression was associated with increased numbers of CD3⁺ T cells (*P*=0.036), CD8⁺ T cells (*P*=0.018) and CD68⁺ macrophages (*P*=0.018). Within the monosomy 3 tumors, we noticed that the expression of BAP1 protein was associated with significantly lower




Fig. 1 Presence of T cells (**a**), and macrophages (**b**) as determined by IF staining or by gene expression (**c**) was compared between BAP1-positive and BAP1-negative tumors from the Leiden cohort

Table 1 Correlation of the values of different probes obtained with an Illumina gene expression array with immunohistochemical data in the Leiden cohort

T cell type	Probe name	Probe number	R	P value
CD68 (n=24)	CD68 probe 1	ILMN_1714861	0.490	0.015
	CD68 probe 2	ILMN_2359907	0.372	0.073
	CD68 probe 3	ILMN_2267914	0.252	0.235
CD3 (n=24)	CD3D probe 1	ILMN_2261416	0.760	<0.001
	CD3D probe 2	ILMN_2325837	0.748	<0.001
	CD3E	ILMN_1739794	0.523	0.009
	CD3G	ILMN_1717197	0.622	0.001
CD8 (n=24)	CD8A probe 1	ILMN_1768482	0.685	<0.001
	CD8A probe 2	ILMN_1760374	0.500	0.013
	CD8A probe 3	ILMN_2353732	0.744	<0.001
	CD8B probe 1	ILMN_1748601	0.126	0.558
	CD8B probe 2	ILMN_2354191	0.388	0.061

The highest correlated probes (in bold) were used for further analysis
P values were obtained by the Spearman's Rho correlation test
n number of patients, *R* correlation coefficient

numbers of CD3⁺ T cells ($P=0.034$) and CD8⁺ T cells ($P=0.034$), but not of CD68⁺ macrophages ($P=0.11$; Fig. 2a-b). This shows that in disomy 3 as well as monosomy 3 tumors, the loss of BAP1 protein is associated with more infiltrating T cells.

In addition, we scrutinized an independent cohort that was available from the TCGA for a potential association between the BAP1-status and infiltration with T cells and macrophages. *BAP1*-negative tumors showed a higher expression of *CD3* ($P<0.001$) and *CD8* ($P<0.001$), but not for *CD68* ($P=0.149$) than *BAP1*-positive tumors (Fig. 3), which confirms our finding that loss of BAP1 is associated with a higher density of T cells in UM.

Immune cell infiltration of tumors is governed by the local production of cytokines and chemokines. We analyzed whether the expression of several cytokines and chemokines involved in T cell and macrophage chemotaxis differed between 24 BAP1-positive and 30 BAP1-negative tumors from the Leiden cohort [31, 32]. Gene expression of *CCL5* and *CXCL10*, important chemokines in T cell chemotaxis, was higher in the BAP1-negative tumors ($P<0.001$; $P=0.005$ and $P=0.065$; Table 2a).

Altogether, these results show that the production of immune cell-attracting chemokines and infiltration of different types of immune cells in UM are associated with the loss of BAP1 expression, independent of chromosome 3 loss.




Fig. 2 Comparison of T cells (a) and macrophages (b) as determined by IF staining between BAP1-positive and BAP1-negative tumors from the Leiden cohort that were either disomic or monosomic for chromosome 3 according to the Mann–Whitney *U* test

Influence of chromosome 8q gain on the inflammatory microenvironment

Gain in the copy number of chromosome 8q is considered an early event in UM development, occurring before loss of chromosome 3 [18, 19]. Seventeen cases of UM from Leiden were available in which the tumor showed gain in chromosome 8q but did not show loss of chromosome 3 or lack of BAP1 protein expression. These cases allowed us to study the influence of gain of chromosome 8q on

immune cell infiltration. Tumors that carried additional copies of chromosome 8q had an increased expression of *CD68* ($P=0.006$), but not of *CD3* and *CD8* (Fig. 4). Again, using the TCGA data as an independent cohort, we analyzed whether there was an association between gain of chromosome 8q and the presence of macrophages and T cells. *BAP1*-positive tumors with extra copies of 8q had an increased expression of *CD68* ($P<0.001$) (Fig. 3) but not of *CD3* or *CD8* compared to *BAP1*-positive tumors with a normal chromosome 8q. Since these tumors had no loss of chromosome 3 or aberrant expression of BAP1, these data confirm our finding in the Leiden cohort that the early gain in chromosome 8q is responsible for macrophage infiltration but not for T cell infiltration.

In addition, we determined whether, in the disomy3/BAP1-positive tumors from the Leiden cohort, extra copies of chromosome 8q affect the expression of pro-inflammatory cytokines. The myeloid-cell attracting chemokines *CCL3* and *CCL2* showed a higher expression in disomy3/BAP1-positive tumors ($P=0.002$ and $P=0.059$, respectively) that carried extra copies of chromosome 8q than did disomy3/BAP1-positive tumors with two copies of chromosome 8q (Table 2b). This was not the case for the expression of typical T cell attracting chemokines, such as *CCL5*, *CXCL12*, and *CXCL10*.


Discussion

We show that the presence of extra copies of chromosome 8q in UM is associated with macrophage infiltration, while loss of BAP1 protein expression, with or without loss of chromosome 3, is associated with T cell infiltration in UM.

The chromosomal evolution of aggressive UM is thought to start with a mutation in *GNAQ/GNA11* [36, 37], followed by gain of chromosome 8q that precedes a potential loss of one copy of chromosome 3 and/or mutation in the *BAP1* gene [20, 38]. We show that in UM with disomy 3 and expression of BAP1, the presence of additional copies of chromosome 8q is highly associated with the increased expression of macrophage-attracting chemokines and a stronger macrophage infiltration. In this subgroup, no effect was found with respect to the production of chemokines associated with T cell infiltration. This phenomenon could not be assessed in monosomy 3 tumors as more than 90% of monosomy 3 UM carry extra copies of 8q. Thus, a gain in copy number of chromosome 8q is associated with an increase in macrophage infiltration.

One might expect that this influx is initiated by activation of the *c-Myc* gene, a proto-oncogene located on chromosome 8q24, which is upregulated in many types of cancer and has been studied in UM [18, 39, 40]. It has previously been suggested that *c-Myc* may be involved in the

Fig. 3 Comparison of T cells and macrophages as determined by gene expression between tumors from the TCGA cohort with a normal and abnormal chromosome 8q status in cases that had low (*BAP1*-negative) or high (*BAP1*-positive) *BAP1* expression as determined by RNA sequencing



activation of inflammatory mediators in the tumor microenvironment [41]. However, we previously observed the opposite, i.e., an association between a high *c-Myc* expression and a low inflammatory phenotype, making it unlikely that *c-Myc* is the relevant factor [42].

Monosomy 3 and loss of *BAP1*-protein expression are strongly correlated in UM, but we observed several atypical cases which allowed us to separately assess the contribution of chromosome 3 and *BAP1*-protein expression on the magnitude and type of immune cell infiltration and to pinpoint that it was the loss of *BAP1* which was associated with the higher expression of T cell-attracting chemokines and a stronger T cell infiltration in UM. The previous studies have reported that tumor suppressor proteins can be involved in the processes and pathways of tumor-promoting inflammation by interacting with transcription factors such as nuclear factor- κ B (NF- κ B) [43]. NF- κ B regulates genes which are involved in inflammation and immune responses. A close family member of *BAP1* is *UCHL1*. Similar to *BAP1*, *UCHL1* functions as a tumor suppressor protein [44] and was recently shown to suppress the NF- κ B pathway, thereby negatively affecting the production of type 1 interferon and pro-inflammatory cytokines and chemokines, including *CCL5* [28]. We, therefore,

hypothesize that *BAP1* may have a similar function as *UCHL1* and that loss of *BAP1* alleviates the suppression pathways leading to activation of NF- κ B, resulting in the production of cytokines and chemokines that attract tumor-specific T cells into UM.

Obvious correlations between genetic changes and the development of an immune infiltrate are not easy to find. Loss of function of several tumor suppressor genes (*p53*, *PTEN*) due to genetic aberrations is known to be associated with inflammation [45]. Interestingly, loss of *BAP1* in an unusual cutaneous tumor, the atypical Spitz nevus, was associated with a higher presence of T cells [46].

When looking at UM, one of the chemokines that was higher in *BAP1*-negative than *BAP1*-positive tumors was *CCL5*; another chemokine that was almost significantly higher in *BAP1*-negative tumors was *CXCL10*. *CCL5* and *CXCL10* play a role in the recruitment of T cells [47]. An influence of *BAP1* on NF- κ B and the additional release of pro-inflammatory multifunctional chemokines might explain why both macrophages and T cells are found in *BAP1*-negative UM, but this requires further research.

Another chemokine involved in T cell recruitment is *CXCL12*, which is the ligand for chemokine receptor *CXCR4*. Previously, it has been described that *CXCR4* is

Table 2 Expression of pro-inflammatory chemokines and receptors in BAP1-positive and BAP1-negative UM (a), and in disomy 3/BAP1-positive tumors, with and without 8q gain (b) in the Leiden cohort

(a)	BAP1-positive (n = 24)	BAP1-negative (n = 30)	P value ⁺
CCL3 (MIP-1 α)	6.7 (6.5–8.5)	6.7 (6.3–8.0)	0.321
VEGFA probe 1	6.8 (6.5–7.4)	6.8 (6.5–7.3)	0.651
VEFGA probe 2	6.4 (6.3–6.8)	6.4 (6.2–6.8)	0.403
CXCL12 (SDF-1) probe 1	7.1 (6.6–8.6)	7.3 (6.5–9.4)	0.689
CXCL12 (SDF-1) probe 2	6.6 (6.3–7.1)	6.6 (6.3–7.6)	0.175
CXCL12 (SDF-1) probe 3	6.5 (6.2–7.1)	6.5 (6.0–7.3)	0.848
CCL7	6.2 (6.0–6.5)	6.3 (5.9–6.7)	0.088
CSF-1	6.4 (6.3–6.7)	6.5 (6.3–6.7)	0.130
CCL2 (MCP-1)	7.2 (6.4–9.5)	7.1 (6.4–9.1)	0.614
CCL5 (RANTES) probe 1	6.9 (6.4–10.7)	7.5 (6.5–12.1)	<0.001
CCL5 (RANTES) probe 2	7.7 (6.6–12.4)	8.8 (7.0–14.4)	0.005
CXCL10 (IP-10)	6.8 (6.4–10.5)	7.4 (6.4–10.4)	0.065
CCR7	6.4 (6.1–7.0)	6.3 (6.0–7.2)	0.903
CXCR4 probe 1	6.5 (6.2–7.2)	6.7 (6.2–8.3)	0.038
CXCR4 probe 2	6.5 (6.1–6.9)	6.4 (6.2–6.8)	0.010
CXCR4 probe 3	6.6 (6.2–7.5)	6.8 (6.2–8.2)	0.216
(b)	D3/BAP1+/n8q (n = 9)	D3/BAP1+/8q gain (n = 8)	P value*
CCL3 (MIP-1 α)	6.6 (6.5–6.8)	6.8 (6.7–8.1)	0.002
VEGFA probe 1	6.7 (6.5–6.9)	6.9 (6.6–7.4)	0.059
VEFGA probe 2	6.4 (6.3–6.6)	6.5 (6.3–6.8)	0.236
CXCL12 (SDF-1) probe 1	7.1 (6.7–7.9)	7.1 (6.6–8.6)	0.743
CXCL12 (SDF-1) probe 2	6.5 (6.3–6.8)	6.5 (6.4–6.8)	0.743
CXCL12 (SDF-1) probe 3	6.5 (6.3–6.7)	6.5 (6.2–7.0)	0.888
CCL7	6.1 (6.0–6.3)	6.3 (6.1–6.4)	0.277
CSF-1	6.4 (6.3–6.6)	6.4 (6.3–6.6)	0.606
CCL2 (MCP-1)	6.8 (6.4–7.7)	7.3 (6.8–9.1)	0.059
CCL5 (RANTES) probe 1	6.8 (6.4–7.0)	6.9 (6.5–7.5)	0.236
CCL5 (RANTES) probe 2	7.4 (6.6–8.2)	7.9 (7.0–8.9)	0.277
CXCL10 (IP-10)	6.7 (6.4–7.4)	6.8 (6.5–8.4)	0.321
CCR7	6.3 (6.1–6.6)	6.4 (6.2–6.7)	0.606
CXCR4 probe 1	6.4 (6.2–6.7)	6.5 (6.3–6.9)	0.114
CXCR4 probe 2	6.5 (6.1–6.6)	6.5 (6.2–6.9)	0.370
CXCR4 probe 3	6.5 (6.2–6.9)	6.6 (6.3–7.3)	0.321

The boldface indicates a significant difference

D3 disomy 3, BAP1+ positive BAP1 protein expression, n8q normal chromosome 8q status, 8q gain chromosome 8q gain, n number of patients; median (range)

⁺P value comparison of chemokine expression in BAP1-positive and BAP1-negative tumors

^{*}P value comparison of chemokine expression in tumors with D3/BAP1+/n8q and tumors with D3/BAP1+/8q gain

involved in the migration of UM cells to the liver [48, 49]. In contrast, another group reported that the expression of CXCR4 was, indeed, correlated with lymphocyte infiltration, but had no prognostic relevance in UM patients [33]. We observed discrepant results for CXCR4 in our Leiden cohort, with one probe showing a higher expression in BAP1-negative tumors, and another one a lower expression (Table 2a).

Previously, it had been shown that tumor-intrinsic active β -catenin signaling restrains tumor-infiltration by T cells, resulting in the escape of tumors from immune surveillance [50]. The β -catenin protein is encoded by the gene *CTNNB1*, which, like *BAP1*, is located on chromosome 3p21. Hence, loss of chromosome 3 may also reduce β -catenin expression. In contrast to *BAP1* [24, 25], however, there is no evidence that the other allele of *CTNNB1* is frequently mutated in UM and thus that the loss of




Fig. 4 Comparison of T cells and macrophages as determined by gene expression between tumors from the Leiden cohort with a normal and abnormal chromosome 8q status in cases that were disomic for chromosome 3 and BAP1-positive as determined by IHC staining

β -catenin signalling is underlying T cell infiltration in UM. In our cohort, we found no correlation between *CTNBL1* expression and the amount of CD8⁺ T cells. Furthermore, the number of β -catenin-positively staining tumor cells in UM is around 10% [51], making its expression an unlikely explanation for the absence of T cell infiltration in most UM.

Our current findings show that alterations in copy numbers or mutations in certain genes can drive a specific type of immune response. As the most common treatment of UM is irradiation and not enucleation, we wondered whether local treatments might affect immune infiltration in UM. No tumors in either the Leiden cohort or the TCGA cohort had received prior irradiation. A previous study from our group showed that more T cells were present in secondarily enucleated eyes after prior irradiation compared to primarily enucleated eyes [52]. As irradiation influenced our chromosome testing, we could not always analyze the chromosome status in previously irradiated tumors [53]. We do not yet know how the type of inflammation or irradiation influences the patient's response to immunotherapy, which at this moment has not been very successful in UM.

In conclusion, we provide evidence that the magnitude and type of immune cell infiltration observed in the subgroup of inflamed UM co-evolve with the sequential genetic changes occurring in UM. The initial infiltration

by macrophages is related to a gain in the copy number of chromosome 8q, while additional T cell infiltration is correlated to a loss of functional BAP1-protein expression.

Acknowledgements This project was supported by Algemene Nederlandse Vereniging Ter Voorkoming van Blindheid (ANVVB), Stichting Blinden Penning, Landelijke Stichting voor Blinden en Slechtienden (LSBS), Novartis Foundation, Stichting Nederlands Oogheelkundig Onderzoek (SNOO), Rotterdamse Stichting Blindenbelangen, and the European Union Horizon 2020 program UM Cure (Project number: 667787).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Kujala E, Makitie T, Kivela T (2003) Very long-term prognosis of patients with malignant uveal melanoma. *Invest Ophthalmol Vis Sci* 44:4651–4659
- Kaliki S, Shields CL (2016) Uveal melanoma: relatively rare but deadly cancer. *Eye (Lond)*. doi:10.1038/eye.2016.275
- Augsburger JJ, Correa ZM, Shaikh AH (2009) Effectiveness of treatments for metastatic uveal melanoma. *Am J Ophthalmol* 148:119–127. doi:10.1016/j.ajo.2009.01.023
- Hodi FS, O'Day SJ, McDermott DF et al (2010) Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 363:711–723. doi:10.1056/NEJMoa1003466
- Larkin J, Chiarion-Sileni V, Gonzalez R et al (2015) Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. *N Engl J Med* 373:23–34. doi:10.1056/NEJMoa1504030
- Weber JS, D'Angelo SP, Minor D et al (2015) Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* 16:375–384. doi:10.1016/S1470-2045(15)70076-8
- Verdegaal EM (2016) Adoptive cell therapy: a highly successful individualized therapy for melanoma with great potential for other malignancies. *Curr Opin Immunol* 39:90–95. doi:10.1016/j.coi.2016.01.004
- Bol KF, van den Bosch T, Schreiber G et al (2016) Adjuvant Dendritic Cell Vaccination in High-Risk Uveal Melanoma. *Ophthalmology* 123:2265–2267. doi:10.1016/j.ophtha.2016.06.027
- de la Cruz PO Jr, Specht CS, McLean IW (1990) Lymphocytic infiltration in uveal malignant melanoma. *Cancer* 65:112–115
- Whelchel JC, Farah SE, McLean IW, Burnier MN (1993) Immunohistochemistry of infiltrating lymphocytes in uveal malignant melanoma. *Invest Ophthalmol Vis Sci* 34:2603–2606
- de Waard-Siebinga I, Hilders CG, Hansen BE, van Delft JL, Jager MJ (1996) HLA expression and tumor-infiltrating immune cells in uveal melanoma. *Graefes Arch Clin Exp Ophthalmol* 234:34–42

12. Ericsson C, Seregard S, Bartolazzi A, Levitskaya E, Ferrone S, Kiessling R, Larsson O (2001) Association of HLA class I and class II antigen expression and mortality in uveal melanoma. *Invest Ophthalmol Vis Sci* 42:2153–2156
13. Maat W, Ly LV, Jordanova ES, de Wolff-Rouendaal D, Schalijs-Delfos NE, Jager MJ (2008) Monosomy of chromosome 3 and an inflammatory phenotype occur together in uveal melanoma. *Invest Ophthalmol Vis Sci* 49:505–510. doi:10.1167/iops.07-0786
14. Prescher G, Bornfeld N, Hirche H, Horsthemke B, Jockel KH, Becher R (1996) Prognostic implications of monosomy 3 in uveal melanoma. *The Lancet* 347:1222–1225
15. Coupland SE, Lake SL, Zeschnigk M, Damato BE (2013) Molecular pathology of uveal melanoma. *Eye (Lond)* 27:230–242. doi:10.1038/eye.2012.255
16. Damato B, Duke C, Coupland SE, Hiscott P, Smith PA, Campbell I, Douglas A, Howard P (2007) Cytogenetics of uveal melanoma: a 7-year clinical experience. *Ophthalmology* 114:1925–1931. doi:10.1016/j.ophtha.2007.06.012
17. Cassoux N, Rodrigues MJ, Plancher C et al (2014) Genome-wide profiling is a clinically relevant and affordable prognostic test in posterior uveal melanoma. *Br J Ophthalmol* 98:769–774. doi:10.1136/bjophthalmol-2013-303867
18. Singh AD, Tubbs R, Biscotti C, Schoenfeld L, Trizzoi P (2009) Chromosomal 3 and 8 status within hepatic metastasis of uveal melanoma. *Arch Pathol Lab Med* 133:1223–1227. doi:10.1043/1543-2165-133.8.1223
19. de Lange MJ, van Pelt SI, Versluis M et al (2015) Heterogeneity revealed by integrated genomic analysis uncovers a molecular switch in malignant uveal melanoma. *Oncotarget* 6:37824–37835. doi:10.18632/oncotarget.5637
20. Versluis M, de Lange MJ, van Pelt SI, Ruivenkamp CA, Kroes WG, Cao J, Jager MJ, Luyten GP, van der Velden PA (2015) Digital PCR validates 8q dosage as prognostic tool in uveal melanoma. *PLoS One* 10:e0116371. doi:10.1371/journal.pone.0116371
21. Tschentscher F, Husing J, Holter T et al (2003) Tumor classification based on gene expression profiling shows that uveal melanomas with and without monosomy 3 represent two distinct entities. *Cancer Res* 63:2578–2584
22. Onken MD, Worley LA, Ehlers JP, Harbour JW (2004) Gene expression profiling in uveal melanoma reveals two molecular classes and predicts metastatic death. *Cancer Res* 64:7205–7209. doi:10.1158/0008-5472.can-04-1750
23. Harbour JW, Onken MD, Roberson ED et al (2010) Frequent mutation of BAP1 in metastasizing uveal melanomas. *Science* 330:1410–1413. doi:10.1126/science.1194472
24. Koopmans AE, Verdijk RM, Brouwer RW et al (2014) Clinical significance of immunohistochemistry for detection of BAP1 mutations in uveal melanoma. *Mod Pathol* 27:1321–1330. doi:10.1038/modpathol.2014.43
25. Kalirai H, Dodson A, Faqir S, Damato BE, Coupland SE (2014) Lack of BAP1 protein expression in uveal melanoma is associated with increased metastatic risk and has utility in routine prognostic testing. *Br J Cancer* 111:1373–1380. doi:10.1038/bjc.2014.417
26. van Essen TH, van Pelt SI, Versluis M et al (2014) Prognostic parameters in uveal melanoma and their association with BAP1 expression. *Br J Ophthalmol* 98:1738–1743. doi:10.1136/bjophthalmol-2014-305047
27. Fang Y, Fu D, Shen XZ (2010) The potential role of ubiquitin c-terminal hydrolases in oncogenesis. *Biochim Biophys Acta* 1806:1–6. doi:10.1016/j.bbcan.2010.03.001
28. Karim R, Tummers B, Meyers C et al (2013) Human papillomavirus (HPV) upregulates the cellular deubiquitinase UCHL1 to suppress the keratinocyte's innate immune response. *PLoS Pathog* 9:e1003384. doi:10.1371/journal.ppat.1003384
29. Bronkhorst IH, Ly LV, Jordanova ES, Vrolijk J, Versluis M, Luyten GP, Jager MJ (2011) Detection of M2-macrophages in uveal melanoma and relation with survival. *Invest Ophthalmol Vis Sci* 52:643–650. doi:10.1167/iops.10-5979
30. Bronkhorst IH, Vu TH, Jordanova ES, Luyten GP, Burg SH, Jager MJ (2012) Different subsets of tumor-infiltrating lymphocytes correlate with macrophage influx and monosomy 3 in uveal melanoma. *Invest Ophthalmol Vis Sci* 53:5370–5378. doi:10.1167/iops.11-9280
31. Ly LV, Bronkhorst IH, van Beelen E, Vrolijk J, Taylor AW, Versluis M, Luyten GP, Jager MJ (2010) Inflammatory cytokines in eyes with uveal melanoma and relation with macrophage infiltration. *Invest Ophthalmol Vis Sci* 51:5445–5451. doi:10.1167/iops.10-5526
32. Bronkhorst IH, Jehs TM, Dijkgraaf EM, Luyten GP, van der Velden PA, van der Burg SH, Jager MJ (2014) Effect of hypoxic stress on migration and characteristics of monocytes in uveal melanoma. *JAMA Ophthalmol*. 132: 614–621. doi:10.1001/jamaophthalmol.2014.43
33. van den Bosch T, Koopmans AE, Vaarwater J, van den Berg M, de Klein A, Verdijk RM (2013) Chemokine receptor CCR7 expression predicts poor outcome in uveal melanoma and relates to liver metastasis whereas expression of CXCR4 is not of clinical relevance. *Invest Ophthalmol Vis Sci* 54:7354–7361. doi:10.1167/iops.13-12407
34. McCarroll SA, Kuruvilla FG, Korn JM, Cawley S, Nemesh J, Wysoker A, Shapero MH, de Bakker PIW, Maller JB, Kirby A, Elliott AL, Parkin M, Hubbell E, Webster T, Mei R, Veitch J, Collins PJ, Handsaker R, Lincoln S, Nizzari M, Blume J, Jones KW, Rava R, Daly MJ, Gabriel SB, Altshuler D (2008) Integrated detection and population-genetic analysis of SNPs and copy number variation. *Nat Genet* 40(10):1166–1174
35. Mermel CH, Schumacher SE, Hill B, Meyerson ML, Beroukhi R, Getz G (2011) GISTIC2.0 facilitates sensitive and confident localization of the targets of focal somatic copy-number alteration in human cancers. *Genome Biol* 12(4):R41
36. Van Raamsdonk CD, Griewank KG, Crosby MB et al (2010) Mutations in GNA11 in uveal melanoma. *N Engl J Med* 363:2191–2199. doi:10.1056/NEJMoa1000584
37. Onken MD, Worley LA, Long MD, Duan S, Council ML, Bowcock AM, Harbour JW (2008) Oncogenic mutations in GNAQ occur early in uveal melanoma. *Invest Ophthalmol Vis Sci* 49:5230–5234. doi:10.1167/iops.08-2145
38. Singh N, Singh AD, Hide W (2015) Inferring an Evolutionary Tree of Uveal Melanoma From Genomic Copy Number Aberrations. *Invest Ophthalmol Vis Sci* 56:6801–6809. doi:10.1167/iops.15-16822
39. Royds JA, Sharrard RM, Parsons MA, Lawry J, Rees R, Cottam D, Wagner B, Rennie IG (1992) C-myc oncogene expression in ocular melanomas. *Graefes Arch Clin Exp Ophthalmol* 230:366–371
40. Parrella P, Caballero OL, Sidransky D, Merbs SL (2001) Detection of c-myc amplification in uveal melanoma by fluorescent in situ hybridization. *Invest Ophthalmol Vis Sci* 42:1679–1684
41. Borrello MG, Degl'Innocenti D, Pierotti MA (2008) Inflammation and cancer: the oncogene-driven connection. *Cancer Lett* 267:262–270. doi:10.1016/j.canlet.2008.03.060
42. De Waard-Siebinga I, Blom DJ, Griffioen M, Schrier PI, Hoogendoorn E, Beverstock G, Danen EH, Jager MJ (1995) Establishment and characterization of an uveal-melanoma cell line. *Int J Cancer* 62:155–161
43. Mantovani A, Allavena P, Sica A, Balkwill F (2008) Cancer-related inflammation. *Nature* 454:436–444. doi:10.1038/nature07205

44. Li L, Tao Q, Jin H et al (2010) The tumor suppressor UCHL1 forms a complex with p53/MDM2/ARF to promote p53 signaling and is frequently silenced in nasopharyngeal carcinoma. *Clin Cancer Res* 16:2949–2958. doi:[10.1158/1078-0432.ccr-09-3178](https://doi.org/10.1158/1078-0432.ccr-09-3178)
45. Yang L, Karin M (2014) Roles of tumor suppressors in regulating tumor-associated inflammation. *Cell Death Differ* 21:1677–1686. doi:[10.1038/cdd.2014.131](https://doi.org/10.1038/cdd.2014.131)
46. Wiesner T, Murali R, Fried I, Cerroni L, Busam K, Kutzner H, Bastian BC (2012) A distinct subset of atypical Spitz tumors is characterized by BRAF mutation and loss of BAP1 expression. *Am J Surg Pathol* 36:818–830. doi:[10.1097/PAS.0b013e3182498be5](https://doi.org/10.1097/PAS.0b013e3182498be5)
47. Liu J, Li F, Ping Y et al (2015) Local production of the chemokines CCL5 and CXCL10 attracts CD8 + T lymphocytes into esophageal squamous cell carcinoma. *Oncotarget* 6:24978–24989. doi:[10.18632/oncotarget.4617](https://doi.org/10.18632/oncotarget.4617)
48. Li H, Alizadeh H, Niederkorn JY (2008) Differential expression of chemokine receptors on uveal melanoma cells and their metastases. *Invest Ophthalmol Vis Sci* 49:636–643. doi:[10.1167/iovs.07-1035](https://doi.org/10.1167/iovs.07-1035)
49. Li H, Yang W, Chen PW, Alizadeh H, Niederkorn JY (2009) Inhibition of chemokine receptor expression on uveal melanomas by CXCR4 siRNA and its effect on uveal melanoma liver metastases. *Invest Ophthalmol Vis Sci* 50:5522–5528. doi:[10.1167/iovs.09-3804](https://doi.org/10.1167/iovs.09-3804)
50. Spranger S, Bao R, Gajewski TF (2015) Melanoma-intrinsic beta-catenin signalling prevents anti-tumour immunity. *Nature* 523:231–235. doi:[10.1038/nature14404](https://doi.org/10.1038/nature14404)
51. Zuidervaart W, Pavey S, van Nieuwpoort FA, Packer L, Out C, Maat W, Jager MJ, Gruis NA, Hayward NK (2007) Expression of Wnt5a and its downstream effector beta-catenin in uveal melanoma. *Melanoma Res* 17:380–386. doi:[10.1097/CMR.0b013e3282f1d302](https://doi.org/10.1097/CMR.0b013e3282f1d302)
52. Vu TH, Bronkhorst IH, Versluis M, Marinkovic M, van Duinen SG, Vrolijk J, Luyten GP, Jager MJ (2013) Analysis of inflammatory cells in uveal melanoma after prior irradiation. *Invest Ophthalmol Vis Sci* 54:360–369. doi:[10.1167/iovs.12-9424](https://doi.org/10.1167/iovs.12-9424)
53. Dogrusoz M, Kroes WG, van Duinen SG, Creutzberg CL, Versluis M, Bleeker JC, Marinkovic M, Luyten GP, Jager MJ (2015) Radiation treatment affects chromosome testing in uveal melanoma. *Invest Ophthalmol Vis Sci* 56:5956–5964. doi:[10.1167/iovs.15-17092](https://doi.org/10.1167/iovs.15-17092)