

POSTER PRESENTATION

Open Access

Chronic cobalt exposure modifies nociception and orofacial inflammatory pain

T Alexa^{1*}, A Dondas¹, A Luca¹, VA Grigoras², OC Mungiu³, CR Bohotin³*From* The European Headache and Migraine Trust International Congress London, UK. 20-23 September 2012

Introduction

Cobalt chloride (CoCl₂) is a widely used hypoxia-mimetic; preconditioning with cobalt has been reported to exert protection against ischemic injury as well as decreasing pro-inflammatory cytokine levels (Shukla et al. 2011). Exposure to moderate hypoxia may lead to neurological phenomena such as paraesthesia, numbness or pain, possibly as a consequence of the modifications that appear in small DRG neurones. (Gruss et al. 2006).

Purpose

to investigate the effects of chronic (CoCl₂) administration on nociception and orofacial formalin-induced pain in mice.

Methods

Sixteen Swiss male mice were divided into 2 groups: control (n=8, i.p. saline) and chronic (CoCl₂) (n=8, 12.5 mg/kg, i.p. 21 days). Tail flick (TF), hot plate (HP), mechanical (electronic von Frey aesthesiometer) and thermal (Hargreaves method) withdrawal thresholds were measured before cobalt/saline administration and thereafter every two days. 24 hours after the last dose, each group was evaluated for the nociceptive tests (TF&HP) and heat/mechanical hyperalgesia. Results were compared with the baseline using paired Student's t test. After the nociceptive tests, the mice were injected with 20 μ l of 5% formalin into the upper right lip in order to assess the orofacial pain. The results were compared with saline group using unpaired Student's t test.

Results

Chronic (CoCl₂) administration, significantly increased nociception in TF (p<0.001) and HP (p<0.001) tests and

decreased the mechanical withdrawal threshold (p<0.02). On orofacial formalin pain, (CoCl₂) tends to exert a pronociceptive effect on the first phase (p=0.09) and produces a significant antinociceptive effect (p=0.04) on the inflammatory phase.

Conclusion

Taken together, our results demonstrate that chronic cobalt exposure increases acute nociception (TF, HP and first phase of the orofacial formalin pain) but has an antinociceptive effect on the inflammatory pain induced by orofacial formalin injection.

Acknowledgements

research supported by UEFISCSU Romania, project PN-II-ID-PCE-2011-3-0875.

Author details

¹Centre for the Study and Therapy of Pain, University of Medicine and Pharmacy "Grigore T Popa" Iasi, Romania. ²Department of Neurology, Rehabilitation Hospital, University of Medicine and Pharmacy "Grigore T Popa" Iasi, Romania. ³Centre for the Study and Therapy of Pain, Department of Oncology, University of Medicine and Pharmacy "Grigore T Popa" Iasi, Romania.

Published: 21 February 2013

References

1. Gruss M, Ettore G, Stehr AJ, Henrich M, Hempelmann G, Scholz A: Moderate hypoxia influences excitability and blocks dendrotoxin sensitive K⁺ currents in rat primary sensory neurones. *Mol Pain* 2006, **2**:12.
2. Shukla D, Saxena S, Purushothaman J, Shrivastava K, Singh M, Shukla S, Malhotra VK, Mustoori S, Bansal A: Hypoxic preconditioning with cobalt ameliorates hypobaric hypoxia induced pulmonary.

doi:10.1186/1129-2377-14-S1-P87

Cite this article as: Alexa et al: Chronic cobalt exposure modifies nociception and orofacial inflammatory pain. *The Journal of Headache and Pain* 2013 **14**(Suppl 1):P87.

¹Centre for the Study and Therapy of Pain, University of Medicine and Pharmacy "Grigore T Popa" Iasi, Romania
Full list of author information is available at the end of the article