



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: February 26, 2011

ACCEPTED: April 26, 2011

PUBLISHED: May 11, 2011

Stable causality of Black Saturns

Sebastian J. Szybka

*Astronomical Observatory, Jagellonian University,
Orla 171, 30-244 Kraków, Poland*

E-mail: sebastian.szybka@uj.edu.pl

ABSTRACT: We prove that the Black Saturns are stably causal on the closure of the domain of outer communications.

KEYWORDS: Black Holes, Differential and Algebraic Geometry

ARXIV EPRINT: [1102.3942](https://arxiv.org/abs/1102.3942)

JHEP05(2011)052

Contents

1	Introduction	1
2	Stable causality	2

1 Introduction

The Black Saturn solution of Elvang and Figueras is a five dimensional black hole with disjoint event horizons with topology $\mathbb{R} \times S^3$ and $\mathbb{R} \times S^1 \times S^2$ [1, 2]. It describes a spherical black hole¹ surrounded by a black ring.

In this article, we address the question of causality violations in the Black Saturn solution. The absence of closed causal curves is one of the desired properties of a solution to the Einstein equations. Such property should be stable against small perturbations. Therefore, the closed causal curves are not allowed in any Lorentzian metric that is sufficiently near the original one. This leads to the notion of *stable causality* as introduced by Hawking [3]. Technically, a spacetime $(M, g_{\mu\nu})$ is *stably causal* if and only if there exists a differentiable function f on M such that $\nabla^\mu f$ is a past directed timelike vector field [3, 4]. The non-unique function f may be interpreted as a cosmic time that increases along every causal curve.

The natural guess for the Black Saturn is that the generalized Weyl t coordinate is a cosmic time in the domain of outer communications (d.o.c.). Under this hypothesis, the problem of *stable causality* of the d.o.c. was reduced in [1] to the study of the $g_{\psi\psi}$ component of the metric. Namely, stable causality will result from the following inequality in the generalized Weyl coordinates

$$g(\nabla t, \nabla t) = g^{tt} = -\frac{g_{\psi\psi}}{G_y} < 0, \tag{1.1}$$

where G_y is a non-negative function which is zero only on the axis of the rotation of the Killing field ∂_ψ . In other words, it is sufficient to show that $g_{\psi\psi}$ vanishes as fast as G_y on the axis of the rotation of ∂_ψ and that $g_{\psi\psi} > 0$ in the remaining part of the d.o.c.

Numerical evidence for positivity of $g_{\psi\psi}$ *in the plane of the ring* (as defined in [2, section 3.8]) was already signalled in the original paper of Elvang and Figueras [2]. This numerical evidence was extended in [1] to the part of spacetime covered by the generalized Weyl coordinates away from points where ∂_ψ vanishes. Before our work, all analytical proofs of *stable causality* were restricted to the situation when the Komar angular momentum of the spherical component of the horizon is equal to zero: under this restriction, the positivity of $g_{\psi\psi}$ in the plane of the ring was established in [2], and *stable causality* of the d.o.c. was shown in [1].

¹The adjective “spherical” refers to the topology of the horizon.

In this article, we prove that the d.o.c. of the general Black Saturn solution is *stably causal*. In addition, we show that the event horizons may be included to the domain of *stable causality*, hence the Black Saturns are *stably causal* on the closure of the d.o.c.²

The calculations presented here³ involve manipulations of huge algebraic expressions and were done with MATHEMATICA. We use the same notation and definitions of auxiliary functions as in [1].

2 Stable causality

In the generalized Weyl coordinates $(t, \psi, \varphi, \rho, z)$ the d.o.c. corresponds to

$$\{\rho > 0\} \cup \{\rho = 0, z \notin [a_5, a_4] \cup [a_3, a_2]\},$$

where $a_{i=1,\dots,5}$ are parameters. Since $g_{\psi\psi} = g_{\psi\psi}(\rho, z, a_1, \dots, a_5)$, then we would like to prove *stable causality* for a union of the following sets in \mathbb{R}^7

$$\begin{aligned} \tilde{V}_a &= \{a_1 < a_5 < a_4 < a_3 < a_2\}, \\ \tilde{V}_I &= \{\rho > 0\} \cap \tilde{V}_a, \\ \tilde{V}_{II} &= \{\rho = 0, z < a_1\} \cap \tilde{V}_a, \\ \tilde{V}_{III} &= \{\rho = 0, a_1 \leq z < a_5\} \cap \tilde{V}_a, \\ \tilde{V}_{IV} &= \{\rho = 0, a_4 < z < a_3\} \cap \tilde{V}_a, \\ \tilde{V}_\psi &= \{\rho = 0, a_2 < z\} \cap \tilde{V}_a. \end{aligned} \tag{2.1}$$

Hence, we have $\tilde{V}_{d.o.c.} = \tilde{V}_I \cup \tilde{V}_{II} \cup \tilde{V}_{III} \cup \tilde{V}_{IV} \cup \tilde{V}_\psi$. We are interested in non-degenerate solutions, so the parameters are restricted to \tilde{V}_a . The “plane of the ring” corresponds to $\tilde{V}_{II} \cup \tilde{V}_{III} \cup \tilde{V}_{IV}$, while \tilde{V}_ψ is the intersection of the rotation axis of ∂_ψ with the d.o.c. The event horizons of the black ring and the spherical component coincide with $(\{\rho = 0\} \setminus \tilde{V}_{d.o.c.}) \cap \tilde{V}_a$. We would like to show that $g_{\psi\psi} > 0$ on $\tilde{V}_I \cup \tilde{V}_{II} \cup \tilde{V}_{III} \cup \tilde{V}_{IV}$. Moreover, since G_y vanishes as ρ^2 on the axis of ∂_ψ [1], then it is necessary to check that

$$\lim_{\rho \rightarrow 0^+} \frac{g_{\psi\psi}}{\rho^2} > 0$$

on \tilde{V}_ψ .

It turns out to be convenient to view $g_{\psi\psi}$ as a function of $\rho, \mu_1, \dots, \mu_5$, where $\mu_i = \sqrt{\rho^2 + (z - a_i)^2} - (z - a_i)$. In this parametrization, the translational symmetry of z, a_i is explicit. The analogues of the sets⁴ (2.1) are now in \mathbb{R}^6

$$\begin{aligned} V_\mu &= \{\rho \leq \mu_1 \leq \mu_5 \leq \mu_4 \leq \mu_3 \leq \mu_2\}, \\ V_I &= \{\rho > 0\} \cap V_\mu, \end{aligned} \tag{2.2}$$

²In fact, our proof remains valid for a family of the Black Saturn solutions with conical singularities on the axes of the periodic Killing fields $\partial_\psi, \partial_\varphi$.

³The MATHEMATICA code is available at <http://th.if.uj.edu.pl/~szybka/BScasuality>.

⁴These sets are not equivalent to (2.1) because $V_4 \subset V_{IV}, V_2 \subset V_\psi$, where V_4, V_2 correspond to the sets that are defined in the old parametrization as $\{\rho = 0, z = a_4\} \cap \tilde{V}_a, \{\rho = 0, z = a_2\} \cap \tilde{V}_a$, respectively.

$$\begin{aligned}
V_{II} &= \{\rho = 0, \mu_1 > 0\} \cap V_\mu, \\
V_{III} &= \{\rho = 0, \mu_1 = 0, \mu_5 > 0\} \cap V_\mu, \\
V_{IV} &= \{\rho = 0, \mu_4 = 0, \mu_3 > 0\} \cap V_\mu, \\
V_\psi &= \{\rho = 0, \mu_2 = 0\} \cap V_\mu,
\end{aligned}$$

and, of course, $V_{d.o.c.} \subset V_I \cup V_{II} \cup V_{III} \cup V_{IV} \cup V_\psi$. It follows from the definition of μ_i that if $\mu_i = \mu_j$ for $i \neq j$, then $\rho = \mu_i = \mu_j = 0$. This parametrization turns out to be more helpful in completing the proof.

The numerator and the denominator of $g_{\psi\psi}$, when written as polynomials in ρ, μ_i , and some c_1, c_2, q contain tens of thousands monomials [1, 2]. One may check with a direct MATHEMATICA calculation that some non-trivial factors from the numerator and the denominator cancel and the original form of $g_{\psi\psi}$ may be simplified to

$$g_{\psi\psi} = \frac{\mu_4\mu_5 A^2 - \mu_3 B^2}{\mu_1\mu_4 H_x F}, \quad (2.3)$$

where

$$\begin{aligned}
A &= (\mu_2 p_1 (\mu_5 p_2 + c_2 q \mu_1 \mu_3 p_3) + c_1 \mu_3 p_4 (-q \rho^2 p_5 + c_2 \mu_1 \mu_4 p_6)), \\
B &= (\mu_5 p_1 (q \mu_1 p_2 - c_2 \mu_4 \rho^2 p_3) + c_1 \mu_2 \mu_4 p_4 (\mu_1 \mu_5 p_5 + c_2 q \mu_3 p_6)), \\
p_1 &= (\mu_3 - \mu_1)(\mu_1 \mu_4 + \rho^2), \\
p_2 &= (\mu_2 - \mu_4)(\mu_1 \mu_2 + \rho^2)(\mu_2 \mu_3 + \rho^2), \\
p_3 &= (\mu_2 - \mu_1)(\mu_2 \mu_5 + \rho^2), \\
p_4 &= \mu_1(\mu_5 - \mu_1), \\
p_5 &= (\mu_2 - \mu_1)(\mu_2 - \mu_4)(\mu_2 \mu_3 + \rho^2), \\
p_6 &= (\mu_1 \mu_2 + \rho^2)(\mu_2 \mu_5 + \rho^2),
\end{aligned} \quad (2.4)$$

and $p_i \geq 0$. The functions H_x, F are non-negative and they were defined in [2]. It follows from the analysis in [1] that zeros of $H_x F$ exist only for $\rho = 0$ and they cancel with the zeros of the numerator of $g_{\psi\psi}$.

The parameters c_1, c_2, q depend only on a_i and do not depend on ρ, z . However, if one assumes that ρ, μ_i are independent variables, then c_1, c_2, q are finite continuous functions of ρ, μ_i and are given by

$$c_1^2 = \frac{(\mu_3 - \mu_1)(\mu_4 - \mu_1)\mu_5(\mu_1\mu_3 + \rho^2)(\mu_1\mu_4 + \rho^2)}{\mu_1\mu_3\mu_4(\mu_5 - \mu_1)(\mu_1\mu_5 + \rho^2)}, \quad (2.5)$$

$$q = \frac{c_1 c_2 \mu_4 (\mu_2 - \mu_1)(\mu_1 \mu_2 + \rho^2)}{c_1 \mu_1 (\mu_2 - \mu_4)(\mu_2 \mu_4 + \rho^2) + c_2 \mu_2 (\mu_4 - \mu_1)(\mu_1 \mu_4 + \rho^2)}, \quad (2.6)$$

where we imposed (2.5) on c_1 in the formula for q . The formula for c_2 in terms of ρ, μ_i is too long to be usefully cited here. It may be derived from equations (4.2), (5.1) in [1], but it is not necessary for our calculations.

The formulas (2.5), (2.6) are explicitly valid in $V_I \cup V_{II}$. They are also valid in the remaining part of the d.o.c. provided the limit $\rho \rightarrow 0^+$ is carefully taken. If $\mu_i(\rho = 0) = 0$, then taking this limit should be preceded by the substitution $\mu_i \rightarrow \rho^2 \hat{\mu}_i$, where $\hat{\mu}_i > 0$ [1].

The simplification (2.3) is a significant one. Even if one does not substitute formulas for c_1, c_2, q , the original expression for $g_{\psi\psi}$ written as a rational function contained 106995 monomials. This number was reduced to 2344 in the simplified formula.

In order to present the proof in a concise form, we introduce the following operators. Let N , D denote operators acting on a rational functions that return polynomials: a numerator or a denominator, respectively. The result is not unique and the action of N , D is given only up to an overall factor. However, this non-uniqueness is not important for our problem. We also define substitution operators S_q , respectively $S_{c_1^2}$, that return the rational function which is obtained after q , respectively c_1^2 , has been substituted in the original expression using (2.6), respectively (2.5). One should note that S_q and $S_{c_1^2}$ do not commute with N and D in general. The formula for c_2 , in contrast to the formula for c_1 , contains a square root that cannot be eliminated by taking c_2^2 . We prefer to preserve the polynomial form of the evaluated expressions, hence we will substitute only c_1^2 and avoid substituting c_2 .

The calculations described below were done with MATHEMATICA. We present them here in a brief form. It follows from the smoothness of the Black Saturns [1] that the expressions evaluated below are regular.

We start with the analysis of $g_{\psi\psi}$ on V_I ($\rho > 0$).⁵ The denominator of $g_{\psi\psi}$ is given by $\mu_1\mu_4H_xF$ and it is a positive function on V_I . The numerator of $g_{\psi\psi}$ is equal to $\Xi_+\Xi_-$, where

$$\Xi_{\pm} = \sqrt{\mu_4\mu_5}A \pm \sqrt{\mu_3}B. \tag{2.7}$$

If $\Xi_+\Xi_- > 0$, then $g_{\psi\psi} > 0$ on V_I , as desired. Firstly, we check at a random point⁶ $P \in V_I$ that $\Xi_{\pm}|_P > 0$. Since both Ξ_+ and Ξ_- are continuous in ρ , μ_i , then it is sufficient to show that they cannot vanish. The functions Ξ_{\pm} are linear in q , as may be seen from (2.4), (2.7). We substitute q into Ξ_{\pm} and examine the numerators of the resulting expressions. We would like to show that none of them ($NS_q\Xi_{\pm}$) has zeros. By inspection, we find that $NS_q\Xi_{\pm}$ are quadratic in c_2 . Since c_2 is real, a negative discriminant of $NS_q\Xi_{\pm}$ with respect to c_2 would imply that none of the equations $NS_q\Xi_{\pm} = 0$ has a solution. We calculate these discriminants Δ_{\pm} and they turn out to be fourth order in c_1 . Next, we substitute c_1^2 into Δ_{\pm} using (2.5) and taking $c_1^4 = (c_1^2)^2$, $c_1^3 = c_1c_1^2$. With a help of MATHEMATICA we have derived

$$S_{c_1^2}\Delta_{\pm} = w_{\pm} \frac{\mu_1\mu_2^2\mu_4\mu_5^{5/2}(\mu_1 - \mu_2)^2(\mu_3 - \mu_1)(\mu_4 - \mu_1)(\mu_2 - \mu_4)^2(\mu_1\mu_2 + \rho^2)^2(\mu_1\mu_4 + \rho^2)^2}{(\mu_5 - \mu_1)(\mu_1\mu_5 + \rho^2)^2}.$$

The factors multiplying w_{\pm} are strictly positive, and w_{\pm} are complicated polynomials in ρ , $\sqrt{\mu_i}$. These polynomials are linear in c_1 (with non-vanishing coefficients in front of c_1 as it will follow from our further analysis). We check at a random point $P' \in V_I$ that $w_{\pm}|_{P'} < 0$, hence if w_{\pm} have no zeros, then $\Delta_{\pm} < 0$. Let c_1^{\pm} be solutions to the equations $w_{\pm} = 0$. A MATHEMATICA calculation reveals that $c_1^+ = -c_1^-$. We set $U = (c_1^{\pm})^2 - S_{c_1^2}c_1^2$ and calculate

$$U = \frac{\mu_5(\mu_1^2 + \rho^2)^2}{4\mu_1^2\mu_3\mu_4(\mu_1 - \mu_5)^2(\mu_1\mu_5 + \rho^2)^2} \frac{\hat{U}}{\tilde{U}},$$

where \hat{U} , \tilde{U} are complicated polynomials⁷ in ρ , μ_i with signs unknown so far. The coefficient in front of \hat{U}/\tilde{U} is strictly positive. Now, we succeeded in making the signs of \hat{U} , \tilde{U}

⁵ $V_I = \{\rho > 0\} \cap V_{\mu}$ as indicated in (2.2), but for the sake of brevity we will remind only first part of the definitions.

⁶We impose the equation satisfied by c_2 in such checks, but this is actually not necessary.

⁷The polynomial \tilde{U} is a full square.

explicit by writing them in terms of the new positive functions

$$\Delta_{51} = \mu_5 - \mu_1, \quad \Delta_{45} = \mu_4 - \mu_5, \quad \Delta_{34} = \mu_3 - \mu_4, \quad \Delta_{23} = \mu_2 - \mu_3.$$

The coefficients in \hat{U} , \tilde{U} in front of ρ , μ_1 , Δ_{ij} turn out⁸ to be positive and belong to

$$[9, 13705432344] \cap \mathbb{Z}, \quad [1, 137075730] \cap \mathbb{Z},$$

respectively. Since ρ , μ_1 , Δ_{ij} are strictly positive, then it follows that \hat{U} , \tilde{U} are strictly positive and the equation $U = 0$ does not have solutions. Therefore, $c_1^\pm \neq S_{c_1^2} c_1$ and the polynomials w_\pm cannot vanish. This means that the discriminants of $NS_q \Xi_\pm$ in respect to c_2 are negative ($\Delta_\pm < 0$) and there are no real c_2 that would satisfy any of the equations $\Xi_\pm = 0$. Finally, this implies that $g_{\psi\psi} > 0$ on V_I (for $\rho > 0$), as desired. To complete the proof it is sufficient to repeat the analysis above in the remaining part of the d.o.c.

For $\rho = 0$ the denominator of $g_{\psi\psi}$ (given by $\mu_1 \mu_4 H_x F$) is not strictly positive any more. However, it follows from the Black Saturns' smoothness [1] that whenever the denominator of $g_{\psi\psi}$ vanishes the numerator of $g_{\psi\psi}$ (equal to $\Xi_+ \Xi_-$) vanishes as well and the limit $\lim_{\rho \rightarrow 0^+} g_{\psi\psi}$ is finite. Moreover, the continuity of $g_{\psi\psi}$ implies that this limit is non-negative, possibly zero.

On V_{II} ($\rho = 0$, $\mu_1 > 0$) the argument proceeds along the same lines as for V_I . Intermediate expressions have different form, but the reasoning is analogous. We have found

$$U = \frac{\mu_1^2}{4\mu_3\mu_4\mu_5(\mu_1 - \mu_5)^2} \frac{\hat{U}}{\tilde{U}}.$$

The coefficients in the polynomials \hat{U} , \tilde{U} in front of μ_1 , Δ_{ij} range in

$$[9, 7882] \cap \mathbb{Z}, \quad [9, 184] \cap \mathbb{Z},$$

respectively. Hence, $U > 0$ and none of the expressions $NS_q \Xi_\pm$ vanishes, and $g_{\psi\psi} > 0$ on V_{II} .

In order to study $g_{\psi\psi}$ on V_{III} ($\rho = \mu_1 = 0$, $\mu_5 > 0$) we substitute $\mu_1 = \rho^2 \hat{\mu}_1$ into Ξ_\pm (the numerator of $g_{\psi\psi}$ is given by $\Xi_+ \Xi_-$). It turns out that ρ^4 factors out in each term Ξ_\pm . On the other hand, ρ^8 factors in $\mu_1 \mu_4 H_x F$ (the denominator of $g_{\psi\psi}$). We set

$$\check{\Xi}_\pm = \lim_{\rho \rightarrow 0^+} \frac{\Xi_\pm}{\rho^4}$$

and repeat the proof for $\check{\Xi}_\pm$ as in the case $\rho > 0$. We have

$$U = \frac{1}{4\hat{\mu}_1^2 \mu_3 \mu_4 \mu_5 (1 + \hat{\mu}_1 \mu_5)^2} \frac{\hat{U}}{\tilde{U}}.$$

Finally, the coefficients in the polynomials \hat{U} , \tilde{U} in front of the strictly positive functions $\hat{\mu}_1$, Δ_{ij} are again greater than zero and in

$$[9, 8714] \cap \mathbb{Z}, \quad [9, 184] \cap \mathbb{Z},$$

⁸We have $\mu_1 > \rho$, but Δ_{ij} do not have to form monotonically increasing sequence like μ_i .

respectively. Therefore, $g_{\psi\psi} > 0$ on V_{III} , as expected.

We continue our analysis on V_{IV} ($\rho = \mu_4 = 0, \mu_3 > 0$). Here, $\mu_1 = \rho^2 \hat{\mu}_1, \mu_5 = \rho^2 \hat{\mu}_5, \mu_4 = \rho^2 \hat{\mu}_4$. We note that $0 < \hat{\mu}_1 < \hat{\mu}_5 < \hat{\mu}_4$. The calculations are similar to the calculations for V_{III} . A factor ρ^{16} appears both in the numerator and the denominator of $g_{\psi\psi}$, thus we define

$$\check{\Xi}_{\pm} = \lim_{\rho \rightarrow 0^+} \frac{\Xi_{\pm}}{\rho^8},$$

and apply our standard analysis to $\check{\Xi}_{\pm}$. The formula for U is

$$U = \frac{\hat{\mu}_5}{4\hat{\mu}_1^2 \hat{\mu}_4 \mu_2^2 \mu_3^3 (\hat{\mu}_1 - \hat{\mu}_5)^2 (1 + \hat{\mu}_1 \mu_2)^2} \hat{U}.$$

In the final step we introduce $\Delta_{51} = \rho^2 \hat{\Delta}_{51}, \Delta_{45} = \rho^2 \hat{\Delta}_{45}$ and the coefficients in \hat{U} in front of strictly positive functions $\hat{\mu}_1, \hat{\Delta}_{51}, \hat{\Delta}_{45}, \Delta_{34}, \Delta_{23}$ are greater than zero and belong to $[9, 5852] \cap \mathbb{Z}$. Hence, $g_{\psi\psi} > 0$ on V_{IV} , as desired.

In the case of the rotation axis of the periodic Killing field ∂_{ψ} (the set V_{ψ} , where $\rho = \mu_i = 0$), the analysis is slightly more involved. We substitute $\mu_i = \rho^2 \hat{\mu}_i$ into Ξ_{\pm} . Then, we have verified that $\Xi_{\pm} \sim \rho^{15}$ for generic c_1 and q . However, a MATHEMATICA calculation reveals that for our choice of c_1^2 and q (the equations (2.5), (2.6)) the leading terms vanish and we have at least $\Xi_{\pm} \sim \rho^{16}$. Therefore, we drop the leading terms in Ξ_{\pm} and analyse the remaining higher order terms. We denote them with Ξ'_{\pm} . Next, we set

$$\check{\Xi}_{\pm} = \lim_{\rho \rightarrow 0^+} \frac{\Xi'_{\pm}}{\rho^{16}},$$

and apply our standard procedure to $\check{\Xi}_{\pm}$. We have found that

$$S_{c_1^2} \Delta_{\pm} = - \frac{4\hat{\mu}_1 \hat{\mu}_2^2 \hat{\mu}_4 \hat{\mu}_5^3 (\hat{\mu}_1 - \hat{\mu}_2)^2 (\hat{\mu}_3 - \hat{\mu}_1)^3 (\hat{\mu}_4 - \hat{\mu}_1) (\hat{\mu}_2 - \hat{\mu}_4)^2}{\hat{\mu}_5 - \hat{\mu}_1},$$

which is strictly negative because $0 < \hat{\mu}_1 < \hat{\mu}_5 < \hat{\mu}_4 < \hat{\mu}_3 < \hat{\mu}_2$. Thus, we have $\Xi_+ \Xi_- \sim \rho^{32}$. The denominator of $g_{\psi\psi}$ (given by $\mu_1 \mu_4 H_x F$) behaves like ρ^{30} and $g_{\psi\psi}$ vanishes like ρ^2 . This, together with positivity of $g_{\psi\psi}$ for $\rho > 0$ and continuity of $g_{\psi\psi}$, implies that

$$\lim_{\rho \rightarrow 0^+} \frac{g_{\psi\psi}}{\rho^2} > 0$$

on the axis of the rotation of ∂_{ψ} and the inequality (1.1) holds. It completes the proof of *stable causality* of the Black Saturns' d.o.c.

The generalized Weyl coordinates cover also the sets

$$\begin{aligned} V_{br} &= \{\rho = 0, \mu_5 = 0, \mu_4 > 0\} \cap V_{\mu}, \\ V_{sb} &= \{\rho = 0, \mu_3 = 0, \mu_2 > 0\} \cap V_{\mu}, \end{aligned}$$

which have been ignored in our analysis so far. The set $V_{BR} = V_{br} \cup V_4$ corresponds to the event horizon of a black ring and the set $V_{SB} = V_{sb} \cup V_2$ corresponds to the event horizon

of a spherical black hole.⁹ The closure of $V_{d.o.c.}$ is given by $V_{d.o.c.} \cup V_{BR} \cup V_{SB}$. Since $V_4 \subset V_{IV}$, $V_2 \subset V_\psi$, it remains to apply our analysis on V_{br} , V_{sb} .

On V_{br} ($\rho = \mu_5 = 0$, $\mu_4 > 0$) we substitute $\mu_1 = \rho^2 \hat{\mu}_1$, $\mu_5 = \rho^2 \hat{\mu}_5$ into Ξ_\pm . Then, it turns out that ρ^7 factors out of Ξ_\pm and ρ^{14} factors out of the denominator of $g_{\psi\psi}$ ($\mu_1 \mu_4 H_x F$). We define

$$\check{\Xi}_\pm = \lim_{\rho \rightarrow 0^+} \frac{\Xi_\pm}{\rho^7},$$

and calculate the discriminants Δ_\pm of $NS_q \check{\Xi}_\pm$ in respect to c_2 . Next, we substitute c_1^2 into Δ_\pm and obtain

$$S_{c_1^2} \Delta_\pm = -w_\pm \frac{\hat{\mu}_1 \hat{\mu}_5^3 \mu_2^4 \mu_3^4 \mu_4 (1 + \hat{\mu}_1 \mu_2)^2 (1 + \hat{\mu}_1 \mu_4)^2 (\mu_2 - \mu_4)^2}{\hat{\mu}_5 - \hat{\mu}_1},$$

where the coefficient behind w_\pm is strictly greater than zero. This time w_\pm are polynomials in ρ , $\hat{\mu}_1$, $\hat{\mu}_5$, μ_4 , μ_3 , μ_2 . They become explicitly positive if written in terms of $\hat{\mu}_1$, $\hat{\Delta}_{51}$, Δ_{45} , Δ_{34} , Δ_{23} with coefficients in the range $[3, 16] \cap \mathbb{Z}$. Then, $S_{c_1^2} \Delta_\pm < 0$ and $g_{\psi\psi} > 0$ on V_{br} .

The analysis of positivity of $g_{\psi\psi}$ on V_{sb} ($\rho = \mu_3 = 0$, $\mu_2 > 0$) mimics the calculations on V_{br} . There are the following changes. We substitute $\mu_1 = \rho^2 \hat{\mu}_1$, $\mu_5 = \rho^2 \hat{\mu}_5$, $\mu_4 = \rho^2 \hat{\mu}_4$, $\mu_3 = \rho^3 \hat{\mu}_4$ into Ξ_\pm . We set

$$\check{\Xi}_\pm = \lim_{\rho \rightarrow 0^+} \frac{\Xi_\pm}{\rho^{12}},$$

and obtain

$$S_{c_1^2} \Delta_\pm = -\frac{4\hat{\mu}_1 \hat{\mu}_4 \hat{\mu}_5^3 \mu_2^4 (1 + \hat{\mu}_1 \mu_2)^2 (1 + \hat{\mu}_3 \mu_2) (1 + \hat{\mu}_4 \mu_2) (1 + \hat{\mu}_5 \mu_2) (\hat{\mu}_3 - \hat{\mu}_1)^3 (\hat{\mu}_4 - \hat{\mu}_1)}{\hat{\mu}_5 - \hat{\mu}_1},$$

which is strictly negative. Therefore, $g_{\psi\psi} > 0$ on V_{sb} .

In summary, our analysis applied to V_{br} , V_{sb} establishes that $g_{\psi\psi} > 0$ there. The strict positivity of $g_{\psi\psi}$ remains valid on $V_4 \subset V_{IV}$, $V_2 \subset V_\psi$, as we know from the analysis of the sets V_{IV} , V_ψ . Therefore, $g_{\psi\psi} > 0$ holds on both event horizons V_{BR} , V_{SB} . This implies that the inequality (1.1) is satisfied on the event horizons. They may be included to the Black Saturns' domain of *stable causality* and the Black Saturns are *stably causal* on the closure of the d.o.c.

We stress that we have not assigned a particular value to c_2 in our calculations so all results hold also for the Black Saturns with conical singularities on the axes of the rotation of the periodic Killing fields ∂_ψ , ∂_φ .

Acknowledgments

We are grateful to Piotr Chruściel and Michał Eckstein for useful discussions. Our calculations were carried out with MATHEMATICA. This research was supported by the Foundation for Polish Science.

⁹The sets V_4 , V_2 were defined in the footnote on page 3.

Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] P.T. Chruściel, M. Eckstein and S.J. Szybka, *On smoothness of Black Saturns*, *JHEP* **11** (2010) 048 [[arXiv:1007.3668](#)] [[SPIRES](#)].
- [2] H. Elvang and P. Figueras, *Black Saturn*, *JHEP* **05** (2007) 050 [[hep-th/0701035](#)] [[SPIRES](#)].
- [3] S. Hawking, *The existence of cosmic time functions*, *Proc. Roy. Soc. Lond. A* **308** (1969) 433 [[SPIRES](#)].
- [4] S.W. Hawking and G.F.R. Ellis, *The large-scale structure of space-time*, Cambridge University Press, Cambridge U.K. (1973) [[SPIRES](#)].