

RESEARCH ARTICLE

Open Access

Assessment of antiproliferative and antiplasmodial activities of five selected Apocynaceae species

Siu Kuin Wong¹, Yau Yan Lim^{1*}, Noor Rain Abdullah², Fariza Juliana Nordin²**Abstract**

Background: Studies have shown that the barks and roots of some Apocynaceae species have anticancer and antimalarial properties. In this study, leaf extracts of five selected species of Apocynaceae used in traditional medicine (*Alstonia angustiloba*, *Calotropis gigantea*, *Dyera costulata*, *Kopsia fruticosa* and *Vallisneria spiralis*) were assessed for antiproliferative (APF) and antiplasmodial (APM) activities, and analysed for total alkaloid content (TAC), total phenolic content (TPC) and radical-scavenging activity (RSA). As *V. glabra* leaf extracts showed wide spectrum APF and APM activities, they were further screened for saponins, tannins, cardenolides and terpenoids.

Methods: APF and APM activities were assessed using the sulphorhodamine B and lactate dehydrogenase assays, respectively. TAC, TPC and RSA were analysed using Dragendorff precipitation, Folin-Ciocalteu and DPPH assays, respectively. Screening for saponins, tannins, cardenolides and terpenoids were conducted using the frothing, ferric chloride, Kedde and vanillin-H₂SO₄ tests, respectively.

Results: Leaf extracts of *A. angustiloba*, *C. gigantea* and *V. glabra* displayed positive APF activity. Dichloromethane (DCM) extract of *C. gigantea*, and DCM and DCM:MeOH extracts of *V. glabra* showed strong APF activity against all six human cancer cell lines tested. DCM extract of *A. angustiloba* was effective against three cancer cell lines. Against MCF-7 and MDA-MB-231 cell lines, DCM extract of *C. gigantea* was stronger than standard drugs of xanthorrhizol, curcumin and tamoxifen. All five species were effective against K1 strain of *Plasmodium falciparum* and three species (*C. gigantea*, *D. costulata* and *K. fruticosa*) were effective against 3D7 strain. Against K1 strain, all four extracts of *V. glabra* displayed effective APM activity. Extracts of *D. costulata* were effective against 3D7 strain. Selectivity index values of extracts of *A. angustiloba*, *C. gigantea* and *V. glabra* suggested that they are potentially safe for use to treat malaria. Extracts of *K. fruticosa* had the highest TAC while *D. costulata* had the highest TPC and RSA. Phytochemical screening of extracts of *V. glabra* also showed the presence of terpenoids, tannins and saponins.

Conclusions: Leaf extracts of *C. gigantea* and *V. glabra* showed great promise as potential candidates for anticancer drugs as they inhibited the growth of all six cancer cell lines. Against K1 strain of *P. falciparum*, all four extracts of *V. glabra* displayed effective APM activity. The wide spectrum APF and APM activities of *V. glabra* are reported for the first time and this warrants further investigation into its bioactive compounds.

Background

The family Apocynaceae consists of about 250 genera and 2000 species of tropical trees, shrubs and vines [1,2]. A characteristic feature of the family is that almost all species produce milky sap. Leaves are simple, opposite and whorled. Flowers are large and colourful.

In traditional medicine, Apocynaceae species are used to treat gastrointestinal ailments, fever, malaria, pain and diabetes [2]. Of the five species studied, roots and leaves of *Calotropis gigantea* (L.) Aiton are used to treat skin and liver diseases, leprosy, dysentery, worms, ulcers, tumours and earaches [3]. Its latex has been reported to have wound healing properties [4]. *Kopsia fruticosa* (Ker.) A. DC. is used to treat sore and syphilis, and has cholinergic effects [5]. Leaves and barks of *Dyera costulata* Hook have been used for treating fever, inflammation and pain [6]. Stems, leaves and latex of *Alstonia*

* Correspondence: lim.yau.yan@sci.monash.edu.my¹School of Science, Monash University Sunway Campus, Bandar Sunway, 46150 Petaling Jaya, Selangor, Malaysia

Full list of author information is available at the end of the article

angustiloba Miq. have been used for gynaecological problems and skin sores in Indonesia [7]. Leaves are externally applied to treat headache in Malaysia [8]. *Vallisneria spiralis* Kuntze is well known in Thailand because the scent of its flowers is similar to that of pandan leaves and aromatic rice [9]. Its use in traditional medicine has not been reported, and its bioactivity and phytochemistry have yet to be studied. Species of Apocynaceae have also been reported to possess anticancer properties [2,10]. Species having cytotoxic activity include those of *Allamanda* [11], *Alstonia* [12,13], *Cerbera* [14,15], *Nerium* [16,17], *Plumeria* [18] and *Tabernaemontana* [19]. Species of Apocynaceae, notably those of *Alstonia*, are also known to have antimalarial properties [20-22].

Prompted by the anticancer and antimalarial properties of Apocynaceae, leaf extracts of five selected species used in traditional medicine were assessed for antiproliferative (APF) activity against six human cancer cell lines and for antiplasmodial (APM) activity against two strains of *Plasmodium falciparum*. Their extracts were also analysed for total alkaloid content, total phenolic content and radical-scavenging activity. Having wide spectrum APF and APM activities, leaf extracts of *V. glabra* were further screened for saponins, tannins, cardenolides and terpenoids. Information from the screening will serve as a useful guide to further work on isolating compounds with APF and APM activities.

Methods

Plant materials

Species studied were *A. angustiloba*, *C. gigantea*, *D. costulata*, *K. fruticosa* and *V. glabra*. Leaf samples of *A. angustiloba*, *C. gigantea* and *V. glabra* were collected from Puchong (3°2'42"N; 101°37'12"E), Sunway (3°4'30"N; 101°36'8" E) and Kepong (3°12'14"N; 101°37'50"E) in Selangor, Malaysia, respectively. Those of *D. costulata* and *K. fruticosa* were collected from the Forest Research Institute Malaysia (3°14'6"N; 101°37'58"E). Identification of species was based on documented descriptions and illustrations [1,2]. With brief descriptions of their morphology and location of collection, the voucher specimens of these species (WSK01, WSK02, WSK03, WSK04 and WSK05, respectively) were deposited in the herbarium of Monash University Sunway Campus in Malaysia.

Extraction of leaves

For crude extraction, fresh leaves of each species (40 g) were cut into small pieces and freeze-dried overnight. Dried samples were blended and extracted with 250 ml of methanol (MeOH) three times for 1 h each time. Samples were filtered and the solvent was removed using a rotary evaporator (Eyela). The dried crude extracts were stored at -20°C for further analysis. For

sequential extraction, fresh leaves of each species (40 g) were freeze-dried, ground and extracted successively with hexane (HEX), dichloromethane (DCM), DCM: MeOH (1:1) and MeOH (HmbG Chemicals). For each solvent, the suspension of ground leaves in 250-300 ml of solvent was shaken for 1 h on the orbital shaker. After filtering, the samples were extracted two more times for each solvent. Solvents were removed with a rotary evaporator to obtain the dried extracts, which were stored at -20°C for further analysis.

Antiproliferative activity

Antiproliferative (APF) activity of extracts (25 µg/ml) was initially screened for growth inhibitory activity against three human cancer cell lines (MCF-7, MDA-MB-231 and HeLa) using the sulphorhodamine B (SRB) assay [23]. Growth inhibitory activity with less than 50% cell growth was considered positive while that with more than 50% cell growth was considered negative. Extracts with positive growth inhibition were further tested against six human cancer lines (MCF-7, MDA-MB-231, HeLa, HT-29, SKOV-3 and HepG2) using six different extract concentrations. Human cancer cell lines were obtained from the American Type Culture Collection (ATCC, Manassas, VA, USA). The cells were seeded 24 h prior to treatment in 96-well plates at densities of 10,000-20,000 cells/well. Each cell line was designated one plate. Initial cell population of each cell lines prior to addition of extracts was determined by fixing with trichloroacetic acid (TCA) (Sigma). Extracts were dissolved in dimethyl sulphoxide (DMSO) (Sigma) and serially diluted from 8-25 µg/ml. Control cultures were treated with the same volume of DMSO. The concentration of DMSO was kept within 1% to avoid any interference with cell viability. After the addition of extracts, the plates were incubated for 48 h. After incubation, the cells were fixed with 50 µl of cold 50% TCA and incubated for 1 h at 4°C. The plates were then washed with tap water and air dried. Cells were stained with 100 µl of 0.4% SRB solution (Sigma) diluted with 1% acetic acid followed by incubation for 10 min at room temperature. Unbound dye was removed by washing with 1% acetic acid. Bound stain was then solubilised with 200 µl of 10 mM trizma base (Sigma). Absorbance of each well at 505 nm was obtained using a microplate reader. Dose-response curves were constructed to obtain GI₅₀ or concentration of extract that causes growth inhibition (GI) of cells by 50%. GI₅₀ was calculated using the formula $[(T_z - T_i)/(T_c - T_i)] \times 100 = 50$ where T_z is the absorbance of cells treated with extracts or drugs at the end of incubation, T_i is the absorbance of cells prior to treatment with extracts or drugs and T_c is the absorbance of untreated cells at the end of incubation [24]. IC₅₀ or inhibition concentration

at which there is a 50% reduction in cells was obtained using the formula $T_z/T_c \times 100 = 50$. Activity is considered to be effective when GI_{50} value ≤ 20 $\mu\text{g/ml}$ [25].

Antiplasmodial activity

Antiplasmodial (APM) activity of extracts was assessed *in vitro* in human blood using the lactate dehydrogenase assay with slight modifications [26]. Chloroquine-resistant K1 and chloroquine-sensitive 3D7 strains of *Plasmodium falciparum* were tested. Standard drugs of artemisinin (Sigma) and mefloquine (Sigma) were used as positive controls. Extracts dissolved in DMSO (10 mg/ml) were diluted with deionised water to 320 $\mu\text{g/ml}$. The solution was serially diluted two-fold six times to give seven different concentrations. Aliquots of each concentration (10 μl) were transferred into 96-well microtiter plates. Parasitised red blood cell suspensions (1% parasitaemia, 190 μl) were added to each well. Parasitised and non-parasitised red blood cells were used as negative controls. The plates were incubated for 24 h at 37°C in a candle jar and were subsequently chilled at -20°C to lyse the red blood cells. The plates were then allowed to cool to room temperature, and 20 μl of blood suspension was dispensed into a new microtiter plate containing 100 μl MALSTAT™ reagent (Flow Inc.), and 20 μl of nitroblue tetrazolium (Sigma) and phenazine ethosulphate (Sigma) mixture. Absorbance was measured with a plate reader at 630 nm. Percentage inhibition at each concentration was determined and the mean of EC_{50} values of parasite viability was calculated using probit analysis. EC_{50} or effective concentration is the extract concentration that kills 50% of malaria parasites. Activity is effective if EC_{50} value ≤ 10 $\mu\text{g/ml}$ [27]. The selectivity index (SI) for APM activity was calculated based on the ratio of cytotoxicity (IC_{50}) on HepG2 and MCF-7 cells to APM activity (EC_{50}) on chloroquine-resistant K1 strain.

Analysis of TAC, TPC and RSA

Total alkaloid content

Total alkaloid content (TAC) of extracts was determined using the Dragendorff precipitation assay [28]. For each species, extracts (15 mg) were dissolved in 1 ml of distilled water that was acidified to pH 2.0-2.5 with 0.01 M HCl. Analysis was conducted in triplicate. Alkaloids were then precipitated with 0.4 ml of Dragendorff reagent. After washing with 0.5 ml of distilled water to remove traces of the reagent, the precipitate was later treated with 0.4 ml of 1% sodium sulphide, resulting in a brownish-black precipitate. Precipitates formed at each stage were recovered by centrifugation at 14,000 rpm for 1 min. The resulting precipitate was dissolved in 0.2 ml of concentrated nitric acid and diluted to 1 ml with distilled water. Addition of 2.5 ml of 3% thiourea

to 0.5 ml aliquots of this solution resulted in a yellow colored complex. Absorbance was measured at 435 nm and TAC was expressed as boldine equivalent in milligram per gram of extract. The calibration equation for boldine (Sigma) was $y = 1.068x$ ($R^2 = 0.9959$) where y is absorbance and x is mg/ml of boldine. Dragendorff reagent was prepared by dissolving 0.8 g of bismuth nitrate (Sigma) in 40 ml of distilled water and 10 ml of glacial acetic acid. The resulting solution was mixed with 20 ml of 40% potassium iodide.

Total phenolic content

Total phenolic content (TPC) of extracts was determined using the Folin-Ciocalteu (FC) assay [29-31]. Extracts (300 μl in triplicate) were introduced into test tubes followed by 1.5 ml of FC reagent (Fluka) at 10 times dilution and 1.2 ml of sodium carbonate (Fluka) at 7.5% w/v. The tubes were allowed to stand for 30 min in the dark before absorbance was measured at 765 nm. TPC was expressed as gallic acid (GA) equivalent in milligram per gram of extract. The calibration equation for GA (Fluka) was $y = 0.0111x - 0.0148$ ($R^2 = 0.9998$) where y is absorbance and x is mg/ml of GA.

Radical-scavenging activity

Radical-scavenging activity (RSA) of extracts was determined using the 1,1-diphenyl-2-picrylhydrazyl (DPPH) assay [29-31]. Different dilutions of extracts (1 ml in triplicate) were added to 2 ml of DPPH (Sigma). The concentration of DPPH used was 5.9 mg in 100 ml of methanol. Absorbance was measured at 517 nm after 30 min. RSA was calculated as IC_{50} , the concentration of extract to scavenge 50% of the DPPH radical. RSA was then expressed as ascorbic acid equivalent antioxidant capacity (AEAC) using the equation: $AEAC$ (mg ascorbic acid/g) = $IC_{50}(\text{ascorbate})/IC_{50}(\text{sample}) \times 10^5$. IC_{50} of ascorbic acid used for calculation of AEAC was 0.00387 mg/ml.

Phytochemical screening

Qualitative phytochemical screening for saponins, tannins, cardenolides and terpenoids from leaf extracts of *V. glabra* was carried out using standard phytochemical procedures [32,33]. The frothing test was used for saponins, the ferric chloride test for tannins, the Kedde test for cardenolides and the vanillin- H_2SO_4 test for terpenoids.

Results and Discussion

APF activity

Initial screening of leaf extracts of five Apocynaceae species against three human cancer cell lines (MCF-7, MDA-MB-231 and HeLa) showed that DCM, DCM:MeOH and MeOH extracts of *A. angustiloba*, *C. gigantea* and *V. glabra* had growth inhibitory activity (data not shown). DCM and DCM:MeOH extracts of

V. glabra, and DCM extract of *C. gigantea* inhibited all three cancer cell lines. HEX and MeOH extracts of all three species did not show inhibition, with the exception of MeOH extract of *V. glabra*. Extracts of *D. costulata* and *K. fruticosa* did not show any APF activity.

Extracts of the three species were further tested against six human cancer cell lines (MCF-7, MDA-MB-231, HeLa, HT-29, SKOV-3 and HepG2). Results showed that DCM extract of *A. angustiloba* inhibited only MDA-MB-231, HeLa and SKOV-3 cell lines with GI₅₀ values of 20, 20 and 16 µg/ml, respectively (Table 1). DCM and DCM:MeOH extracts of *C. gigantea* inhibited all cancer cell lines except for DCM:MeOH extract against MDA-MB-231. APF activity of DCM extract of *C. gigantea* was the strongest with GI₅₀ values ranging from 1.3 to 3.3 µg/ml. Against MCF-7 and MDA-MB-231, GI₅₀ of DCM extract of *C. gigantea* (1.9 and 1.3 µg/ml) was stronger than that of xanthorrhizol (11 and 8.7 µg/ml), curcumin (4.1 and 8.7 µg/ml) and tamoxifen (8.3 and 4.6 µg/ml), respectively. DCM and DCM:MeOH extracts of *V. glabra* inhibited all cell lines with GI₅₀ values ranging from 7.5-12 µg/ml and 5.8-13 µg/ml, respectively. In addition, MeOH extract of *V. glabra* also inhibited the growth of MCF-7 and HepG2. Against MCF-7, GI₅₀ of DCM and DCM:MeOH extracts of *V. glabra* (7.7 and 7.0 µg/ml) was stronger than xanthorrhizol (11 µg/ml) and comparable to tamoxifen (8.3 µg/ml), respectively.

To the best of our knowledge, this study represents the first report of cytotoxic activity from DCM leaf

extract of *A. angustiloba* and DCM and DCM:MeOH extracts of *V. glabra*. Earlier studies have reported cytotoxic activity from the root bark of *Alstonia macrophylla* Wall. ex G. Don [12] and the stem bark of *Alstonia scholaris* R. Br. [13]. A recent study on *Vallisneria spiralis* (Roth) Kuntz has reported potent cell growth inhibitory activity of cardenolide glycosides isolated from the plant [34]. The finding of strong APF activity from DCM and DCM:MeOH extracts of *C. gigantea* from this study is supported by an earlier report that DCM leaf extracts of *C. gigantea* had strong inhibitory activity against KB, BC and NCI-H187 cancer cell lines [35]. Ethanol root extracts of *C. gigantea* were also reported to be cytotoxic to K562 and SGC-7901 human cell lines [36].

APM activity

Against chloroquine-resistant K1 strain of *P. falciparum*, leaves of *V. glabra* were most effective as all four extracts had APM activity with EC₅₀ less than 10 µg/ml (Table 2). DCM extract was the strongest with EC₅₀ of 0.85 µg/ml. Three extracts of *A. angustiloba*, and two extracts of *C. gigantea*, *D. costulata* and *K. fruticosa* showed APM activity. It should be noted that DCM:MeOH extracts of all five species displayed APM activity with *A. angustiloba* having the strongest activity (EC₅₀ of 0.46 µg/ml).

Against chloroquine-sensitive 3D7 strain of *P. falciparum*, extracts of *A. angustiloba* and *V. glabra* showed

Table 1 Antiproliferative activity of leaf extracts of three Apocynaceae species against six human cancer cell lines^a

Species	Leaf extract	^b GI ₅₀ (µg/ml)					
		MCF-7	MDA-MB-231	HeLa	HT-29	SKOV-3	HepG2
<i>Alstonia angustiloba</i>	HEX	-	-	-	-	-	-
	DCM	-	20 ± 1.7	20 ± 1.1	-	16 ± 1.4	-
	DCM:MeOH	-	-	-	-	-	-
	MeOH	-	-	-	-	-	-
<i>Calotropis gigantea</i>	HEX	-	-	-	-	-	-
	DCM	1.9 ± 0.2	1.3 ± 0.3	2.5 ± 0.5	3.3 ± 0.2	2.5 ± 0.2	1.8 ± 1.7
	DCM:MeOH	13 ± 0.3	-	15 ± 1.0	24 ± 0.7	20 ± 2.3	16 ± 3.5
	MeOH	-	-	-	-	-	-
<i>Vallisneria glabra</i>	HEX	-	-	-	-	-	-
	DCM	7.7 ± 1.3	12 ± 2.0	9.8 ± 1.5	9.3 ± 2.0	7.5 ± 4.5	7.6 ± 0.2
	DCM:MeOH	7.0 ± 2.5	13 ± 6.3	8.5 ± 2.9	12 ± 1.2	7.7 ± 2.4	5.8 ± 1.2
	MeOH	16 ± 2.1	-	-	-	-	19 ± 0.9
^d Standard drug							
	Xanthorrhizol	11 ± 0.7	8.7 ± 0.8				
	Curcumin	4.1 ± 0.9	8.7 ± 0.8				
	Tamoxifen	8.3 ± 0.6	4.6 ± 0.5				

^a Initial screening showed that extracts of *Dyera costulata* and *Kopsia fruticosa* did not show any APF activity.

^b GI₅₀ (µg/ml) is growth inhibition (GI) of cancer cell lines by 50%. Inhibition is not effective (-) with values >20 µg/ml. MCF-7 and MDA-MB-231, HeLa and SKOV-3, HT-29, and HepG2 are human breast, cervical, colon and liver cancer cell lines, respectively.

^c HEX, hexane; DCM, dichloromethane; MeOH, methanol.

^d Data on standard drugs of xanthorrhizol, curcumin and tamoxifen against MCF-7 and MDA-MB-231 cell lines are from an earlier publication [24].

Table 2 Antiplasmodial (APM) activity and selectivity index (SI) of leaf extracts of Apocynaceae species

Species	Leaf extract	^a EC ₅₀ (µg/ml) of APM activity		^b IC ₅₀ (µg/ml) of APF activity		^b SI of APM activity	
		K1	3D7	HepG2	MCF-7	HepG2	MCF-7
<i>Alstonia angustiloba</i>	HEX	7.81	-				
	DCM	-	-				
	DCM:MeOH	0.46	-	>25.0	>25.0	NC	NC
	MeOH	6.46	-				
<i>Calotropis gigantea</i>	HEX	5.82	-				
	DCM	-	-				
	DCM:MeOH	0.97	3.29	>25.0	16.8	NC	17.3
	MeOH	-	-				
<i>Dyera costulata</i>	HEX	-	-				
	DCM	-	8.31				
	DCM:MeOH	7.52	2.13				
	MeOH	7.74	3.56				
<i>Kopsia fruticosa</i>	HEX	-	-				
	DCM	-	7.14				
	DCM:MeOH	4.35	-				
	MeOH	1.01	-				
<i>Vallis glabra</i>	HEX	1.00	-	>25.0	>25.0	NC	NC
	DCM	0.85	-	>25.0	12.0	NC	14.1
	DCM:MeOH	8.45	-				
	MeOH	8.42	-				
^d Standard drug							
	Artemisinin	0.001	0.001				
	Mefloquine	0.008	0.018				

^a EC₅₀ or effective concentration (µg/ml) is the extract concentration that kills 50% of malaria parasites. Activity is not effective (-) if EC₅₀ value >10 µg/ml. K1 and 3D7 are chloroquine-resistant and chloroquine-sensitive strains of *Plasmodium falciparum*, respectively.

^b IC₅₀ of antiproliferative (APF) activity and SI of APM activity against HepG2 and MCF-7 cells were calculated only for the most potent extracts against K1 strain with EC₅₀ ≤1.00 µg/ml. NC, non-cytotoxic with SI values >25.0.

^c HEX, hexane; MeOH, methanol; DCM, dichloromethane.

^d Standard drugs of artemisinin and mefloquine are used as positive controls.

no activity. Extracts of *D. costulata* were the exception in that DCM, DCM:MeOH and MeOH extracts showed positive APM activity with EC₅₀ of 8.31, 2.13 and 3.56 µg/ml, respectively. Generally, extracts were less effective against 3D7 strain.

This finding complements an earlier report that alkaloids from the extracts of *Alstonia* species were effective against chloroquine-resistant K1 strain but not against chloroquine-sensitive T9-96 strain [22]. Previous studies on the APM activity of Apocynaceae were focused on *Alstonia* species [21,22]. This study is the first to report the APM activity of leaf extracts of *D. costulata*, *C. gigantea*, *K. fruticosa* and *V. glabra*. A notable finding is the APM activity of *V. glabra* against K1 strain in all solvent fractions.

Selectivity index

The selectivity index (SI) for APM activity indicates the safety of an extract to be used for antimalarial therapy [37]. The index was calculated based on the ratio of cytotoxicity (IC₅₀) on HepG2 and MCF-7 cells to APM

activity (EC₅₀). The SI was calculated for the most potent extracts against K1 strain of *P. falciparum* with EC₅₀ ≤1.00 µg/ml. They were DCM:MeOH extracts of *A. angustiloba* (0.46 µg/ml) and *C. gigantea* (0.97 µg/ml), and HEX and DCM extracts of *V. glabra* (1.00 and 0.85 µg/ml), respectively (Table 2). Against HepG2 cells, IC₅₀ values were all >25.0 µg/ml, and against MCF-7 cells, IC₅₀ values were >25.0, 16.8, >25.0 and 12.0 µg/ml, respectively.

Against HepG2 cells, HEX and DCM extracts of *V. glabra*, and DCM:MeOH extracts of *A. angustiloba* and *C. gigantea* were non-cytotoxic with SI values >25.0 (Table 2). Against MCF-7 cells, the DCM extract of *V. glabra* and the DCM:MeOH extract of *C. gigantea* had SI values of 14.1 and 17.3, respectively. The HEX extract of *V. glabra* and the DCM:MeOH extract of *A. angustiloba* were non-cytotoxic with SI values >25.0.

Recently, a study on the antimalarial and cytotoxic activity of plants in the Democratic Republic of Congo considered SI values of extracts >10 as high [38]. With SI values relatively higher than 10, the extracts of *A. angustiloba*,

C. gigantea and *V. glabra* are potentially safe for use to treat malaria.

Analysis of TAC, TPC and RSA

Of the five species analysed, MeOH crude and DCM extracts of *K. fruticosa* had the highest TAC (100 and 129 mg BE/g of extract), respectively (Table 3). Other species with high TAC were *D. costulata* and *A. angustiloba* with MeOH crude and DCM:MeOH extracts having values in the range 58-68 and 23-58 mg BE/g of extract, respectively. Based on TAC, the ranking of species was: *K. fruticosa* > *D. costulata* > *A. angustiloba* > *C. gigantea* ≈ *V. glabra*. Extracts of *D. costulata* had the highest TPC and strongest RSA. MeOH crude, DCM:MeOH and MeOH extracts yielded TPC values of 319, 354 and 279 mg GAE/g of extract, and RSA values of 377, 349 and 278 mg AA/g of extract, respectively. Based on TPC and RSA, the ranking of species was: *D. costulata* > *V. glabra* > *K. fruticosa* > *A. angustiloba* > *C. gigantea*. There is a strong correlation between TPC and RSA of extracts ($R^2 = 0.955$) but not with TAC ($R^2 = 0.112$). Correlation of results of phytochemical analysis with APF and APM activities remains unclear. Extracts of *C. gigantea* and *V. glabra* showed strong APF activity. The former had low TAC and TPC, while the latter had low TAC but high TPC. Extracts of *V. glabra* and *D. costulata* were effective against K1 and 3D7 strains of *P. falciparum*, respectively. The former

had low TAC but high TPC, while the latter had high TAC and TPC.

Phytochemical screening

Phytochemical screening showed the presence of terpenoids in all leaf extracts of *V. glabra* except MeOH extract while saponins and tannins were present in DCM and DCM:MeOH extracts, and in DCM:MeOH and MeOH extracts, respectively (Table 4). Cardenolides were not detected.

Terpenoids are used for the treatment of human diseases such as cancer and malaria, and infectious diseases caused by virus and bacterial [39]. Taxol and artemisinin are renowned terpenoid-based anticancer and antimalarial drugs, respectively. Terpenoids inhibit the growth or induce apoptosis of breast cancer cells such as MCF-7, MDA-MB-231 and T47D [40]. They are among the most important natural antimalarial drugs, which also include quinones and alkaloids [41].

Tannins are water-soluble polyphenols that are present in many plant foods [42]. Literature on the effects of tannins on human health is vast and sometimes conflicting. Incidences of esophageal cancer have been attributed to consumption of tannin-rich foods such as herbal teas, suggesting that tannins might be carcinogenic. However, reports have indicated a negative association between consumption of tea and the incidence of cancer. Teas with high tannin content have been suggested to be anticarcinogenic

Table 3 Phytochemical analysis of total alkaloid content, total phenolic content and radical-scavenging activity of leaf extracts of Apocynaceae species

Species	MeOH crude extract	^a Sequential extract			
		HEX	DCM	DCM:MeOH	MeOH
^b Total alkaloid content (mg BE/g)					
<i>Kopsia fruticosa</i>	100 ± 4.2	63 ± 1.1	129 ± 4.0	99 ± 2.5	46 ± 1.6
<i>Dyera costulata</i>	58 ± 2.5	2.4 ± 0.6	11 ± 1.8	68 ± 2.7	41 ± 2.4
<i>Alstonia angustiloba</i>	23 ± 0.3	13 ± 2.3	27 ± 3.1	58 ± 1.2	27 ± 1.0
<i>Calotropis gigantea</i>	2.7 ± 0.3	3.7 ± 1.0	8.6 ± 1.2	9.6 ± 1.9	9.2 ± 2.7
<i>Vallis glabra</i>	2.7 ± 0.4	4.4 ± 0.8	8.9 ± 0.6	9.2 ± 0.2	8.7 ± 2.2
^b Total phenolic content (mg GAE/g)					
<i>Dyera costulata</i>	319 ± 5.5	21 ± 0.2	23 ± 0.1	354 ± 6.0	279 ± 3.0
<i>Vallis glabra</i>	99 ± 3.8	15 ± 0.9	24 ± 0.5	134 ± 1.0	164 ± 13
<i>Kopsia fruticosa</i>	83 ± 1.1	39 ± 0.5	20 ± 0.5	129 ± 1.0	84 ± 1.0
<i>Alstonia angustiloba</i>	68 ± 2.0	17 ± 0.6	24 ± 0.6	96 ± 1.1	94 ± 1.1
<i>Calotropis gigantea</i>	28 ± 0.8	14 ± 0.6	44 ± 1.7	42 ± 0.8	33 ± 0.5
^b Radical-scavenging activity (mg AA/g)					
<i>Dyera costulata</i>	377 ± 25	15 ± 0.3	9.3 ± 0.5	349 ± 21	278 ± 5.4
<i>Vallis glabra</i>	84 ± 0.5	6.0 ± 0.3	8.4 ± 0.4	77 ± 2.2	119 ± 6.8
<i>Kopsia fruticosa</i>	63 ± 3.2	12 ± 1.4	7.5 ± 0.3	70 ± 2.0	48 ± 1.5
<i>Alstonia angustiloba</i>	29 ± 0.9	10 ± 0.2	5.7 ± 0.7	50 ± 1.2	46 ± 2.2
<i>Calotropis gigantea</i>	7.6 ± 0.4	5.6 ± 0.3	6.1 ± 0.4	8.0 ± 0.6	14 ± 1.3

^a HEX, hexane; DCM, dichloromethane; MeOH, methanol.

^b BE, boldine equivalent; GAE, gallic acid equivalent; AA, ascorbic acid. Values are in milligram per gram of extract.

Table 4 Phytochemical screening of leaf extracts of *Vallis glabra*

^a Leaf extract	^b Qualitative test			
	^c Saponin	^c Tannin	^c Cardenolide	^c Terpenoid
HEX	-	-	-	+++
DCM	+	-	-	++
DCM:MeOH	++	++	-	++
MeOH	-	++	-	-

^a HEX, hexane; DCM, dichloromethane; MeOH, methanol.

^b Strong (+++), moderate (++) and weak (+) presence, and absent (-).

Classification was based on observation of colour intensity and amount of precipitate.

^c Saponins, tannins, cardenolides and terpenoids were screened using the frothing, ferric chloride, Kedde and vanillin-H₂SO₄ tests.

and antimutagenic which may be related to their antioxidative property in protecting cellular oxidative damage against lipid peroxidation and superoxide radicals. The antimicrobial activities of tannins are well documented.

Saponins are naturally occurring glycosides with a distinctive foaming characteristic and bitter taste [43,44]. They have a wide range of properties, which include both beneficial and detrimental effects on human health. Saponins affect the immune system in ways that help to protect the human body against cancers, and also lower cholesterol levels. They decrease blood lipids, lower cancer risks and lower blood glucose response.

Conclusions

The DCM extract of *C. gigantea*, and DCM and DCM:MeOH extracts of *V. glabra* inhibited the growth of all six human cancer cell lines. Against MCF-7 and MDA-MB-231 human cell lines, DCM leaf extract of *C. gigantea* had stronger APF activity than standard drugs of xanthorrhizol, curcumin and tamoxifen. With wide spectrum APF activity, leaves of these two species are therefore promising candidates as alternative resources for anticancer drugs. Against K1 strain of *P. falciparum*, all four extracts of *V. glabra* displayed effective APM activity. Selectivity index values suggested that extracts of *V. glabra* are potentially safe for use to treat malaria. The wide spectrum APF and APM activities of *V. glabra* are reported for the first time. This warrants further investigation into its bioactive compounds.

Acknowledgements

The authors are thankful to Monash University Sunway Campus (MUSC) and the Institute of Medical Research (IMR) for funding and supporting the project. The assistance of HT Chan in identifying and locating the plant species is gratefully acknowledged.

Author details

¹School of Science, Monash University Sunway Campus, Bandar Sunway, 46150 Petaling Jaya, Selangor, Malaysia. ²Herbal Medicine Research Centre, Institute for Medical Research, 50588 Kuala Lumpur, Malaysia.

Authors' contributions

The study was conducted by SKW as part of her PhD program in Monash University Sunway Campus in Malaysia which is supervised by YYL. Part of the experiments was done in collaboration with Noor Rain and Fariza Juliana from the Institute of Medical Research. SKW analysed the data and drafted the manuscript which was edited and revised by YYL, with comments from counterparts. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 11 October 2010 Accepted: 14 January 2011

Published: 14 January 2011

References

- Ng FSP: *Tropical Horticulture and Gardening* Clearwater Publications, Kuala Lumpur, Malaysia; 2006.
- Wiat C: *Medicinal Plants of Asia and the Pacific* CRC Press/Taylor & Francis, Boca Raton; 2006.
- Rajakaruna N, Harris CS, Towers GHN: **Antimicrobial activity of plants collected from serpentine outcrops in Sri Lanka.** *Pharm Biol* 2002, **40**:235-244.
- Nalwaya N, Pokharna G, Deb L, Jain NK: **Wound healing activity of latex of *Calotropis gigantea*.** *Inter J Pharm Pharm Sci* 2009, **1**:176-181.
- Johnson T: *CRC Ethnobotany Desk Reference* Cleveland, Ohio: CRC Press LLC; 1999.
- Subhadhirasakul S, Jankeaw B, Malinee A: **Chemical constituents and antioxidative activity of the extract from *Dyera costulata* leaves.** *Songklanakarin J Sci Technol* 2003, **25**:351-357.
- Mulyoutami E, Rismawan R, Joshi L: **Local knowledge and management of simpukng (forest gardens) among the Dayak people in East Kalimantan, Indonesia.** *Forest Ecol Manage* 2009, **257**:2054-2061.
- Lin KW: **Ethnobotanical study of medicinal plants used by the Jah Hut people in Malaysia.** *Indian J Med Sci* 2005, **59**:156-161.
- Wongpornchai S, Sriseadka T, Choonvisase S: **Identification and quantitation of the rice aroma compound, 2-acetyl-1-pyrroline, in bread flowers (*Vallis glabra* Ktze).** *J Agric Food Chem* 2003, **51**:457-462.
- Patel B, Das S, Prakash R, Yasir M: **Natural bioactive compound with anticancer potential.** *Inter J Adv Pharm Sci* 2010, **1**:32-41.
- Schmidt DFN, Yunes RA, Schaab EH, Malheiros A, Filho VC, Franchi GC Jr, Nowill AE, Cardoso AA, Yunes JA: **Evaluation of the anti-proliferative effect of the extracts of *Allamanda blanchetti* and *A. schottii* on the growth of leukemic and endothelial cells.** *J Pharm Pharm Sci* 2006, **9**:200-208.
- Keawpradub N, Eno-Amooquaye E, Burke PJ, Houghton PJ: **Cytotoxic activity of indole alkaloids from *Alstonia macrophylla*.** *Planta Med* 1999, **65**:311-315.
- Jaetia GC, Baliga MS: **Evaluation of anticancer activity of the alkaloid fraction of *Alstonia scholaris* (Sapthaparna) in vitro and in vivo.** *Phytother Res* 2006, **20**:103-109.
- Chang LC, Gills JJ, Bhat KPL, Luyengi L, Farnsworth NR, Pezzuto JM, Kinghorn AD: **Activity-guided isolation of constituents of *Cerbera manghas* with antiproliferative and antiestrogenic activities.** *Bioorg Med Chem Lett* 2000, **10**:2431-2434.
- Nurhanan MY, Asiah O, Mohd Ilham MA, Siti Syarifah MM, Norhayati I, Lili Sahara H: **Anti-proliferative activities of 32 Malaysian plant species in breast cancer cell lines.** *J Trop Forest Sci* 2008, **20**:77-81.
- Siddiqui BS, Begum S, Siddiqui S, Lichter W: **Two cytotoxic pentacyclic triterpenoids from *Nerium oleander*.** *Phytochemistry* 1995, **39**:171-174.
- Pathak S, Multani AS, Narayan S, Kumar V, Newman RA: **Anvirezol™, an extract of *Nerium oleander*, induces cell death in human but not murine cancer cells.** *Anti-Cancer Drugs* 2000, **11**:455-463.
- Kardono LSS, Tsauri S, Padmawinata K, Pezzuto JM, Kinghorn AD: **Cytotoxic constituents of the bark of *Plumeria rubra* collected in Indonesia.** *J Nat Prod* 1990, **53**:1447-1455.
- Lee CC, Houghton P: **Cytotoxicity of plants from Malaysia and Thailand used traditionally to treat cancer.** *J Ethnopharmacol* 2005, **100**:237-243.
- Schwikkard S, van Heerden FR: **Antimalarial activity of plant metabolites.** *Nat Prod Rep* 2002, **19**:675-692.

21. Wright CW, Allen D, Phillipson JD, Kirby GC, Warhurst DC, Massiot G, Le Men-Olivier L: *Alstonia* species: are they effective in malaria treatment? *J Ethnopharmacol* 1993, **40**:41-45.
22. Keawpradub N, Kirby GC, Steele JCP, Houghton PJ: Antiplasmodial activity of extracts and alkaloids of Thai *Alstonia* species from Thailand. *Planta Med* 1999, **65**:690-694.
23. Vichai V, Kirtikara K: Sulforhodamine B colorimetric assay for cytotoxicity screening. *Nat Protoc* 2006, **1**:1112-1116.
24. Cheah YH, Nordin FJ, Tee TT, Azimahtol HLP, Abdullah NR, Ibrahim Z: Antiproliferative property and apoptotic effect of xanthorrhizol on MDA-231 breast cancer cells. *Anticancer Res* 2008, **28**:3677-3690.
25. Gaidhani SN, Lavekar GS, Juvekar AS, Sen S, Singh A, Kumari S: In-vitro anticancer activity of standard extracts used in ayurveda. *Pharmacog Mag* 2009, **5**:425-429.
26. Chan KL, Choo CY, Abdullah NR, Ibrahim Z: Antiplasmodial studies of *Eurycoma longifolia* using the lactate dehydrogenase assay of *Plasmodium falciparum*. *J Ethnopharmacol* 2004, **92**:223-227.
27. Tran QL, Tezuka Y, Ueda JY, Nguyen NT, Maruyama Y, Begum K, Kim HS, Wataya Y, Tran QK, Kadota S: In vitro antiplasmodial activity of antimalarial medicinal plants used in Vietnamese traditional medicine. *J Ethnopharmacol* 2003, **86**:249-252.
28. Ribeiro B, Lopes R, Andrade PB, Seabra RM, Goncalves RF, Baptista P, Quelhas I, Valentao P: Comparative study of phytochemicals and antioxidant potential of wild edible mushroom caps and stipes. *Food Chem* 2008, **110**:47-56.
29. Wong SK, Lim YY, Chan EWC: Antioxidant properties of *Hibiscus*: Species variation, altitudinal change, coastal influence and floral colour change. *J Trop Forest Sci* 2009, **21**:307-315.
30. Rao ASVC, Reddy SG, Babu PP, Reddy AR: The antioxidant and anti-proliferative activities of methanolic extracts from Njavara rice bran. *BMC Complement Altern Med* 2010, **10**:4.
31. Lai HY, Lim YY, Kim KH: *Blechnum orientale* Linn - a fern with potential as antioxidant, anticancer and antibacterial agent. *BMC Complement Altern Med* 2010, **10**:15.
32. Harborne JB: *Phytochemical Method: A Guide to Modern Techniques of Plants Analysis*. Third edition. New York: Chapman and Hall; 1998.
33. Houghton PJ, Raman A: *Laboratory Handbook for the Fractionation of Natural Extracts* New York: Chapman and Hall; 1998.
34. Ahmed F, Sadhu SK, Ohtsuki T, Khatun A, Ishibashi M: Glycosides from *Vallisneria spiralis* with TRAIL resistance overcoming activity. *Heterocycles* 2010, **80**:477-488.
35. Lhinhatrakool T, Sutthivaiyakit S: 19-Nor- and 18,20-epoxy-cardenolides from the leaves of *Calotropis gigantea*. *J Nat Prod* 2006, **69**:1249-1251.
36. Wang ZH, Wang MY, Mei WL, Han Z, Dai HF: A new cytotoxic pregnanone from *Calotropis gigantea*. *Molecules* 2008, **13**:3033-3039.
37. Esmaili S, Naghibi F, Mosaddegh M, Sahranavard S, Ghafari S, Abdullah NR: Screening of antiplasmodial properties among some traditionally used Iranian plants. *J Ethnopharmacol* 2009, **121**:400-404.
38. Lusakibanza M, Mesia G, Tona G, Karemere S, Lukuka A, Tits M, Angenot L, Frédérick M: In vitro and in vivo antimalarial and cytotoxic activity of five plants used in congolese traditional medicine. *J Ethnopharmacol* 2010, **129**:398-402.
39. Wang G, Tang W, Bidigare RR: Terpenoids as therapeutic drugs and pharmaceutical agents. In *Natural Products Drug Discovery and Therapeutic Medicine*. Edited by: Zhang L, Demain AL. Humana Press; 2005:197-227.
40. Rabi T, Bishayee A: Terpenoids and breast cancer chemoprevention. *Breast Cancer Res Treat* 2009, **115**:223-239.
41. Caniato R, Puricelli L: Review: natural antimalarial agents (1995-2001). *Crit Rev Plant Sci* 2003, **22**:79-105.
42. Chung KT, Wong TY, Wei CI, Huang YW, Lin Y: Tannins and human health: a review. *Crit Rev Food Sci Nutr* 1998, **38**:421-464.
43. Desai SD, Desai DG, Kaur H: Saponins and their biological activities. *Pharma Times* 2009, **41**:13-16.
44. Shi J, Arunasalam K, Yeung D, Kakuda Y, Mittal G, Jiang Y: Saponins from edible legumes: chemistry, processing, and health benefits. *J Med Food* 2004, **7**:67-78.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1472-6882/11/3/prepub>

doi:10.1186/1472-6882-11-3

Cite this article as: Wong et al.: Assessment of antiproliferative and antiplasmodial activities of five selected Apocynaceae species. *BMC Complementary and Alternative Medicine* 2011 **11**:3.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

