

Reef sites



Hope for coral reef rehabilitation: massive synchronous spawning by outplanted corals in Okinawa, Japan




Fig. 1 Outplanted coral colonies in Maeganeku, Onna, Okinawa, Japan. **a** Daytime view. **b** Synchronous spawning of outplanted *Acropora tenuis*

Since 1998, the Onna Village Fisheries Cooperative, Okinawa, Japan, has outplanted more than 40 scleractinian coral species, under permits issued by Okinawa Prefecture. The Coop has established a unique and successful rearing method for outplants (Higa and Omori 2014) (Fig. 1a). Seeding outplants at the periphery of existing colonies has been suggested as a means of enhancing larval settlement to increase coral abundance, density, and cover during reef rehabilitation. During sexual reproduction, synchronous spawning maximizes fertilization rate by generating high gamete concentrations (Oliver and Babcock 1992).

Here, we report large-scale synchronous spawning of approximately 2800 colonies of outplanted *Acropora tenuis* (Dana, 1846) in Maeganeku, Onna, Okinawa. On 26 June 2015, 25 nights after the full moon, almost all outplanted colonies released bundles synchronously for approximately 5 min starting at 1947 hrs, 21 min after sunset (Fig. 1b). Average water temperature from the surface to 4 m depth was 29.4 °C.

There have been concerns about reduction in genetic diversity in reef rehabilitation using asexual propagation methods; however, sexual reproduction by synchronized spawning of outplanted colonies should ensure levels of genetic variation equal to those of natural spawning. Along with increasing larval supply, habitat restoration is crucial for maintaining healthy coral reefs, and under unfavorable environmental conditions, coral larval supply cannot be enhanced efficiently by transplantation and outplanting (Ferse et al. 2013). Continuous observation of recruitment, post-settlement survivorship, and growth in recipient habitats will be needed to determine the effects of spawning events such as that reported here.

Acknowledgments The authors thank members of Onna Village Fisheries Cooperative, especially Mr. Masami Yamashiro, Mr. Munekazu Mekar, and Mr. Yoshimi Higa, and members of Okinawa Prefectural Institute of Health and Environment, especially Mr. Tomofumi Nagata, for field observations. We also thank Dr. Steven D. Aird (OIST) for editing the manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Ferse SCA, Nugues MM, Romatzki SBC, Kunzmann A (2013) Examining the use of mass transplantation of brooding and spawning corals to support natural coral recruitment in Sulawesi/Indonesia. *Restoration Ecology* 21:745–754
- Higa Y, Omori M (2014) Production of coral colonies for outplanting using a unique rearing method of donor colonies at Onna Village, Okinawa, Japan. *Galaxea* 16:19–20
- Oliver J, Babcock R (1992) Aspects of the fertilization ecology of broadcast spawning corals: sperm dilution effects and in situ measurements of fertilization. *Biol Bull* 183:409–417

Y. Zayasu (✉) · C. Shinzato

Marine Genomics Unit, Okinawa Institute of Science and Technology Graduate University, 1919-1 Tancha, Onna-son, Okinawa 904-0495, Japan
e-mail: yunazayasu.san@gmail.com

Received: 25 March 2016 / Accepted: 4 May 2016 / Published online: 1 June 2016
© The Author(s). This article is published with open access at Springerlink.com 2016

Coral Reefs (2016) 35:1295
DOI 10.1007/s00338-016-1463-7