

POSTER PRESENTATION


Open Access

Ca²⁺ buffering as a mechanism of short-term synaptic plasticity

Victor Matveev


From Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
 Paris, France. 13-18 July 2013

Spatio-temporal compartmentalization allows Ca²⁺ signals to simultaneously regulate multiple vital cell processes and relies in part on Ca²⁺ buffers that absorb at least 98% of Ca²⁺ ions entering the cytoplasm. Computational modeling has played a central role in the understanding of localized Ca²⁺ signals in neurons and other cell types. Although many models consider only simple, one-to-one Ca²⁺ buffering stoichiometry, practically all buffers have multiple Ca²⁺ binding sites which often display cooperative binding (e.g., calretinin [1-3], calmodulin [4,5]). Given the simplest case of two binding sites, cooperativity manifests itself in an order of magnitude difference in the binding and/or unbinding rates of the two consecutive Ca²⁺ binding steps in the following buffer reaction:



Here we extend recent modeling studies of cooperative buffering [1-5], and find that it can lead to spatio-temporal Ca²⁺ signals that cannot be achieved by any combination of non-cooperative buffers, in particular during a sequence of action potentials or synaptic inputs. Namely, Figure 1B shows that cooperative Ca²⁺ buffering can potentially serve as a mechanism of short-term synaptic depression, in contrast to the case of non-cooperative buffers (Figure 1A), which are believed to underlie short-term synaptic facilitation in certain types of mammalian synapses [6,7].

We explore this phenomenon in detail, demonstrating the dependence of such buffer-induced short-term synaptic plasticity on all relevant buffering parameters. These results may lead to better understanding of *post*-synaptic Ca²⁺ dynamics as well, yielding a deeper insight into synaptic transmission and its dynamic regulation, and may also have relevance for Ca²⁺-dependent processes in other cells such as endocrine cells, myocytes and immune system cells.



Correspondence: matveev@njit.edu
 Department of Mathematical Sciences, New Jersey Institute of Technology,
 Newark, NJ 07102, USA

Acknowledgements

This work was supported in part by NSF grant DMS-0817703.

Published: 8 July 2013

References

1. Saftenku EE: **Effects of Calretinin on Ca^{2+} Signals in Cerebellar Granule Cells: Implications of Cooperative Ca^{2+} Binding.** *Cerebellum* 2011, **11**(1):102-120.
2. Schwaller B: **The continuing disappearance of "pure" Ca^{2+} buffers.** *Cell Mol Life Sci* 2009, **66**(2):275-300.
3. Faas GC, Schwaller B, Vergara JL, Mody I: **Resolving the fast kinetics of cooperative binding: Ca^{2+} buffering by calretinin.** *PLoS Biol* 2007, **5**(11): e311.
4. Faas GC, Raghavachari S, Lisman JE, Mody I: **Calmodulin as a direct detector of Ca^{2+} signals.** *Nat Neurosci* 2011, **14**(3):301-304.
5. Kubota Y, Waxham MN: **Lobe specific Ca^{2+} -calmodulin nano-domain in neuronal spines: a single molecule level analysis.** *PLoS Comput Biol* 2010, **6**(11):e1000987.
6. Burnashev N, Rozov A: **Presynaptic Ca^{2+} dynamics, Ca^{2+} buffers and synaptic efficacy.** *Cell Calcium* 2005, **37**(5):489-495.
7. Matveev V, Zucker RS, Sherman A: **Facilitation through buffer saturation: constraints on endogenous buffering properties.** *Biophys J* 1993, **86**:2691-2701.

doi:10.1186/1471-2202-14-S1-P269

Cite this article as: Matveev: Ca^{2+} buffering as a mechanism of short-term synaptic plasticity. *BMC Neuroscience* 2013 **14**(Suppl 1):P269.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

