



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: March 12, 2015

ACCEPTED: May 11, 2015

PUBLISHED: June 3, 2015

Neutron Majorana mass from exotic instantons in a Pati-Salam model

Andrea Addazi^{a,b} and Massimo Bianchi^c

^a*Dipartimento di Fisica, Università di L'Aquila,
Via Vetoio, 67010 Coppito, L'Aquila, Italy*

^b*LNGS, Laboratori Nazionali del Gran Sasso,
67010 Assergi, L'Aquila, Italy*

^c*Dipartimento di Fisica, Università di Roma Tor Vergata,
I.N.F.N. Sezione di Roma Tor Vergata,
Via della Ricerca Scientifica 1, 00133 Roma, Italy*

E-mail: andrea.addazi@infn.lngs.it, massimo.bianchi@roma2.infn.it

ABSTRACT: We show how exotic stringy instantons can generate an effective interaction between color diquark sextets in a Pati-Salam model, inducing a Majorana mass term for the neutron. In particular, we discuss a simple quiver theory for a Pati-Salam like model, as an example in which the calculations of exotic instanton effects are simple and controllable. We discuss some different possibilities in order to generate $n - \bar{n}$ oscillations testable in the next generation of experiments, Majorana mass matrices for neutrini and a Post-Sphaleron Baryogenesis scenario. Connections with Dark Matter issues and the Higgs mass Hierarchy problem are discussed, in view of implications for LHC and rare processes physics. The model may be viewed as a completion of a Left-Right symmetric extension of the Standard Model, alternative to a GUT-inspired scenario. Combined measures in Neutron-Antineutron physics, FCNC, LHC, Dark Matter could rule out the proposed model or uncover aspects of physics at the Planck scale!

KEYWORDS: Strings and branes phenomenology, Phenomenology of Field Theories in Higher Dimensions

ARXIV EPRINT: [1502.08041](https://arxiv.org/abs/1502.08041)

Contents

1	Introduction	1
2	Exotic instantons and quivers	5
2.1	Instantons	5
2.2	Quivers	5
3	An unoriented quiver for a Pati-Salam model	6
3.1	No-proton decays	9
4	Neutron-Antineutron oscillation through color diquark sextets	9
5	Other comments on phenomenology, dark matter and hierarchy problem	11
6	Conclusions and remarks	12

1 Introduction

How can Matter be generated in our Universe? And how are neutrino masses generated? Has the neutron a Majorana mass?

In principle, these three questions could appear unrelated. However, in Left-Right symmetric models with $SU(2)_L \times SU(2)_R \times SU_c(3) \times U(1)_{B-L}$ gauge group, one can find intriguing and elegant connections between these three issues. A Left-Right model is naturally embedded in a Pati-Salam (P-S) model with $G_{224} = SU(2)_L \times SU(2)_R \times SU(4)_c$ [1], that in turn can be embedded in an $SO(10)$ GUT.

As originally suggested in [2], new Higgses Δ_R in the $(\mathbf{1}, \mathbf{3}, \mathbf{10})$ (and Δ_L in the $(\mathbf{3}, \mathbf{1}, \mathbf{10}^*)$) of G_{224} can be introduced in PS models in order to spontaneously break Left-Right symmetry, through $\langle \Delta_R \rangle = v_R \neq 0$ and $\langle \Delta_L \rangle = 0$. This mechanism also produces Majorana masses for Right-Handed neutrinos, that can trigger a seesaw mechanism as suggested in [3], spontaneously breaking $U(1)_{B-L}$ at the same time.¹ The new Higgs $\Delta_R(\mathbf{1}, \mathbf{3}, \mathbf{10}) \equiv \Delta^c(\mathbf{1}, \mathbf{3}, \mathbf{10}^*)$ decomposes with respect to $SU(2)_L \times SU(2)_R \times SU(3)_C \times U(1)_{B-L}$ as

$$\Delta_R(\mathbf{1}, \mathbf{3}, \mathbf{10}) = \left\{ (\mathbf{1}, \mathbf{3}, \mathbf{1})_{-2} + (\mathbf{1}, \mathbf{3}, \mathbf{3})_{-2/3} + (\mathbf{1}, \mathbf{3}, \mathbf{6})_{2/3} \right\}_R \tag{1.1}$$

with $\Delta_{lc}^c(\mathbf{1}, \mathbf{3}, \mathbf{1})_{-2}$ generating Right-Handed neutrini masses via $\langle \Delta_{\nu^c \nu^c}^c \rangle \nu^c \nu^c$.

In GUT $SO(10)$, the $(\mathbf{1}, \mathbf{3}, \mathbf{10})$ of G_{224} and its conjugate are contained in the **126** representation.² But $\Delta^c(\mathbf{1}, \mathbf{3}, \mathbf{10})$ also contains *color sextet diquark fields* $\Delta_{qq^c}^c(\mathbf{1}, \mathbf{3}, \mathbf{6})_{2/3}$,

¹See also [4, 5] for more about seesaw mechanisms.

²The complete decomposition reads $\mathbf{126} \rightarrow (\mathbf{1}, \mathbf{3}, \mathbf{10}) + (\mathbf{3}, \mathbf{1}, \mathbf{10}^*) + (\mathbf{2}, \mathbf{2}, \mathbf{15}) + (\mathbf{1}, \mathbf{1}, \mathbf{6})$.

leading to possible new effects. In particular, these sextets can induce Baryon number violating processes beyond the Standard Model (BSM). Color sextets can also play an important role in some post-sphaleron baryogenesis mechanism [6–8]. In susy extensions, a quartic superpotential term

$$\mathcal{W}_4 = \Delta^c \Delta^c \Delta^c \Delta^c / \mathcal{M}_0$$

can appear that, among other terms, produces a term coupling three color sextets $\Delta_{q^c q^c}^c$ and one color singlet $\Delta_{\nu^c \nu^c}^c$, as $\Delta_{u^c u^c}^c \Delta_{d^c d^c}^c \Delta_{d^c d^c}^c \Delta_{\nu^c \nu^c}^c$. When the color singlet $\Delta_{\nu^c \nu^c}^c$ takes an expectation value, $U(1)_{B-L}$ is spontaneously broken and Right-handed neutrini get a mass [3]. Moreover a Majorana mass for the neutron is generated through the processes shown in figure 1-(a)-(b). This can be directly tested in Neutron-Antineutron transition experiments, as firstly proposed in [2]. As shown in [9], constraints from post-sphaleron baryogenesis and neutrino oscillations imply a precise prediction about neutron-antineutron transitions: an oscillation time $\tau_{n-\bar{n}} \approx 10^{10} s$ accessible to the next generation of experiments!³

In principle, color sextet scalars could be as light as 1 TeV and they could be directly searched at the LHC, as proposed in [11]: dijet data put constraints on the couplings between colored scalars and quarks. In [12], bounds are shown in comparison with LHC data. On the other hand, processes mediated by FCNC could impose stronger constraints on the sextets with respect to LHC direct searches (see [13] for comparison with experimental limits). For example, the Δ_{dd}^c field couples to two down-type quarks dd, ss, bb and mediates $B_{d,s}^0 - \bar{B}_{d,s}^0, K^0 - \bar{K}^0$ oscillations as well as B mesons decays. On the other hand Δ_{uu}^c mediates $D^0 - \bar{D}^0$ oscillations and D -decays like $D \rightarrow K\pi, \pi\pi$. These analyses show that for coupling strengths of order 10^{-2} , the mass of the color sextets has to exceed the TeV-scale.

In this paper, we propose a (SUSY) PS model that is alternative to the SO(10) GUT inspired model mentioned above. We consider an (un)oriented open string model with intersecting D-branes, producing a susy PS like model. Models of this kind have been previously considered e.g. in [15], where an analysis of the mass spectrum and low-energy phenomenology has been carried out. In oriented string theory, a simple way to generate a $U(N)$ gauge theory is to consider a stack of N D-branes, parallel to each other. In this way the open string excitations stretching between the N D-branes reproduce at low energy the fields in the adjoint of the $U(N)$ gauge symmetry. In type IIA, compactified on a six-dimensional (CY) manifold, one can consider stacks of intersecting D6-branes, filling the 4D ordinary Minkowski spacetime, and wrapping internal 3-cycles. From strings connecting different stacks of branes, we can construct chiral fermions, localised at the four-dimensional intersections of two stacks of D6-branes a and b , in the bi-fundamental representation of $U(N_a) \times U(N_b)$ [16]. The net (positive-negative) number of intersections of two branes a and b is a topological invariant, representing the number of massless fermions. In the case in which D-branes are space-time filling, Ω -planes have to be introduced in order to cancel tadpoles and irreducible anomalies [17–22]. An Ω -plane implements a combination of world-sheet parity and a (non) geometric mirror-like involution in the target space. As a consequence Left- and Right-moving modes of the closed strings are identified; closed and open strings become un-oriented. More choices for the gauge groups and their representations are allowed [17–19]. In this way, one can produce stacks supporting $U(N), SO(N)$ or

³Another string-inspired mechanism for $n - \bar{n}$ was proposed in [10], in heterotic superstrings theories.




Figure 1. a) We show the main diagram for neutron-antineutron oscillation in a non-susy $SU(4)_c \times SU(2)_L \times SU(2)_R$ model [14]. The transition is induced by color sextets $\Delta_{u^c u^c}^c$ and $\Delta_{d^c d^c}^c$. b) We show the main diagram for neutron-antineutron oscillation in a supersymmetric $SU(4)_c \times SU(2)_L \times SU(2)_R$ model [14]. The transition is induced by color sextets $\Delta_{u^c u^c}^c$ and $\Delta_{d^c d^c}^c$ arising from the decomposition of $\Delta^c(1, 3, 10)$. The latter participate in a non-perturbative quartic superpotential term. The diagram involves also gaugini \tilde{g} (gluini, zino or bino), squarks \tilde{d}^c , and susy partners of the color sextets $\tilde{\Delta}_{d^c d^c}^c$. In figures, we use the notation $\Delta_{1,2} = \Delta_{u^c u^c, d^c d^c}$ and s -particle for supersymmetric partners.

$Sp(2N)$ gauge groups. This is interesting in order to construct realistic gauge groups, with chiral matter in a globally consistent model [23, 24]. The closed strings propagate in the entire ten dimensional space-time: some mediate gravitational interactions, some behave as axions or scalar moduli fields.

In principle, one can construct a PS like gauge group $U(4) \times Sp_R(2) \times Sp_L(2)$ or $U(4) \times U_R(2) \times U_L(2)$ in terms of intersecting D-brane stacks and Ω -planes. In [15] the case $U(4) \times U_R(2) \times U_L(2)$ was analysed in some detail. In the present paper, we focus on the $U(4) \times Sp_R(2) \times Sp_L(2)$ case with an Ω^+ -plane that requires a stack of four D-branes and its mirror image under Ω , producing $U(4)$, and two stacks of two D-branes each, identified with their own images under Ω , producing $Sp_L(2)$ and $Sp_R(2)$.

This model has extra anomalous $U(1)$'s that could seem dangerous from a gauge theory point of view. On the other hand, in string theory, Generalized Chern-Simon (GCS) terms appear that cancel anomalies [25, 26], in combination with a generalised Green-Schwarz mechanism [27, 28]. The extra Z' gauge bosons can get a mass through a Stückelberg mechanism [29–32]. We will return onto phenomenological implications of this in the next section. There is however a real problem in this scenario. It is not possible to represent $(1, 3, 10)$ in terms of open strings. Perturbative open strings have two ends and can at most carry fundamental charges with respect to two classical gauge groups. On the other hand, the triplet is the adjoint of $Sp(2)$, i.e. the symmetric product of two doublets, and the decuplet of $SU(4)$ is the symmetric product of two tetraplets. States in the $(1, 3, 10)$ (or its conjugate) would correspond to multi-pronged strings with two ends on the $U(4)$ stack and two ends on the $Sp(2)$ stack⁴ that do not admit a perturbative description.

⁴ $\mathbf{3}$ is also the vector of $SO(3)$ but this would prevent the existence of doublets, which are spinors.

On the other hand, we will show that a spontaneously breaking pattern to the SM, giving masses to the neutrini, can be recovered in this model. In fact, we will see that $\phi_{RR}(1, 3, 1)$ and $\phi_{LL}(1, 1, 3)$ appear as excitations of open strings with both ends attached to $\text{Sp}(2)_R$ or $\text{Sp}(2)_L$, while $\Delta(1, 1, 10)$ and its conjugate $\Delta^c(1, 1, 10^*)$ appear from open strings joining $U(4)$ and $U'(4)$ identified with each other under Ω . As a consequence the breaking $U(4) \times \text{Sp}_R(2) \times \text{Sp}_L(2) \rightarrow \text{SU}(3) \times \text{SU}_L(2) \times U(1)$ is not realized through $(1, 3, 10)$, but through $\phi_{RR}(1, 3, 1)$ and $\Delta(1, 1, 10)$:⁵

- i) Left-Right symmetry breaking through the expectation values $\langle \phi_{RR} \rangle = v_R$ and $\langle \phi_{LL} \rangle = 0$;
- ii) $U(1)_{B-L}$ symmetry breaking through the expectation of value $\langle S \rangle = v_{B-L}$, with S the color singlet contained in $(1, 1, 10)$. Alternatively, $U(1)_{B-L}$ can be broken dynamically by exotic instantons or spontaneously by the compactification.

Similarly to the case of $\Delta(1, 3, 10)$, color sextets are contained in $\Delta(1, 1, 10)$.

Our main suggestion is that the super-potential

$$\mathcal{W}_{\text{eff}} = S^c \Delta_6^c \Delta_6^c \Delta_6^c / \mathcal{M}_0 \tag{1.2}$$

be generated by non-perturbative quantum gravity effects peculiar to string theory, called “exotic instantons”. These are associated to Euclidean branes ($E2$ -branes in our case), wrapping internal 3-cycles, that could directly produce such interactions, in a calculable and controllable way in models like type IIA (un)oriented strings. We would like to stress that this class of instantons exists in string theory only, not in gauge theories. The resulting superpotential term is suppressed by the scale $\mathcal{M}_0 = M_S e^{+S E_2}$, where M_S is the string scale and $e^{+S E_2}$ depends on the ‘size’ of the 3-cycles wrapped by the relevant $E2$ -brane. We would like to remark that the suppression scale is higher (in principle also much higher) than the string scale. This is a peculiarity of the non-renormalizable nature of such a non-perturbative term in the string effective action. As a consequence, the hierarchy depends on the particular model: $e^{+S E_2}$ can be approximately 1 for a ‘small’ 3-cycles or $e^{+S E_2} \gg 1$ for a ‘large’ 3-cycles. So, depending on the string scale M_s , assumed to be larger than some TeV’s at least, and the size of the 3-cycle, it is possible to generate such an operator near the LHC scale or at a much higher scale. This leads to two very different branches for phenomenology. In particular, for $\mathcal{M}_0 \simeq 10^{13}$ GeV, color sextets appear near the TeV scale, with potential implications in meson physics and at LHC, as mentioned above. On the other hand, for $\mathcal{M}_0 \simeq M_S \simeq 10$ TeV, a post-sphaleron scenario is possible and testable at the next generation of experiments on neutron-antineutron oscillations, with heavy color sextets, at a scale $m_6 \gg$ TeV that can be generated by closed-string fluxes, as shown in [33] for quiver theories and reviewed below. In this case, there is no possibility to produce the sextets at LHC, and FCNC’s in the meson sector are strongly suppressed. On the other hand, extra Z' at the TeV scale naturally appears in this scenario. Another relevant and peculiar possibility is a Left-Right breaking scale at TeV, compatible with

⁵In [15] breaking triggered by Higgses in the $(1, 2, 4^*)$ was considered together with mass terms generated by exotic instantons.

neutron-antineutron physics and Post-Sphaleron scenario. This is not possible in a SO(10) scenario, as remarked in [9]. Our string-inspired scenario also naturally provides several candidates of WIMP dark matter as we will see.

We would like to mention that such an operator as (1.2) can emerge from stringy dynamics also in other kinds of models like F-theory [34–40], $E_8 \times E_8$ and SO(32) heterotic strings [41–49], generating an SO(10) GUT. For example in heterotic string theories *world-sheet instantons*, suppressed by $e^{-R^2/\alpha'}$ and thus perturbative in the string coupling g_s , can induce non-vanishing couplings of the desired kind from such amplitudes as $\langle V_{126} V_{126} V_{126} V_{126} \rangle$, for vertex operators V_{126} that can appear in twisted sectors. Unfortunately, in the F-theory case the calculations are more involved [50, 51].

The paper is organized as follows: in section 2, we briefly review what are stringy instantons and quivers. In section 3 we propose a simple and consistent quiver for a Pati-Salam model generating a Majorana mass for the neutron through exotic instantons. In section 4 we discuss some phenomenology resulting from this model. In section 5 we present our conclusions and final remarks.

2 Exotic instantons and quivers

In this section, we briefly review D-brane instantons and unoriented quiver theories.

2.1 Instantons

In 4-dimensional gauge theory, instantons are point-like configurations, that extremize the Euclidean action for a given topological charge. In string theory, instantons admit a simple geometric interpretation: they are special Euclidean branes wrapping some (internal) cycle. In theories with (unoriented) open strings, these are Euclidean D-branes (E-branes) that can intersect the ‘physical’ D-branes.⁶ In (un-)oriented type IIA, gauge instantons can be classified as Euclidean D2 (E2) branes wrapping the same 3-cycle as a stack of “physical” D6-branes. In (un-)oriented IIB, instantons are E(-1) or E3 wrapping the same holomorphic divisor as a stack of “physical” D7-branes. In type I, instantons are E5 branes in the internal space, with the same magnetization as the D9, wrapping the entire CY_3 .

2.2 Quivers

The effective low energy description of the dynamics of D-branes at Calabi-Yau singularities is captured by a quiver field theory. Usually, the (supersymmetric) quiver conventions are the following: the standard D-brane stacks are ‘circle’ nodes, the super-fields in the bi-fundamental representations of two D-brane stacks are oriented lines connecting the nodes, usually termed arrows whence the name ‘quiver’, ‘triangle’ nodes are Euclidean D-branes (instantons), grassmanian moduli or modulini are dashed lines connecting triangles and circles. Square nodes represent flavour branes, i.e. branes wrapping non-compact cycles so much so that $g_{YM,Dp}^2 = g_s(\alpha')^{p+1/2}/V_{p+1} \rightarrow 0$. These simple rules allow one to subsume the system of D-branes and open strings with a simple diagram. In this notation, perturbative

⁶For a pedagogical review see e.g. [67–69].




Figure 2. Unoriented quiver for a Pati-Salam-like model $U(4) \times Sp(2)_L \times Sp(2)_R$. The circles labeled by 4, 4', 2L, 2R represent $U(4)$, $U'(4)$, $Sp(2)_L$, $Sp(2)_R$ stacks respectively. An Ω^+ -plane identifies the $U(4)$ stack with its mirror image, $Sp(2)_{L,R}$ are stacks of two D6-branes laying exactly on top of the Ω -plane. The symmetric representations $\Delta(1, 1, 10)$ and $\Delta^c(1, 1, 10^*)$ appear in between the two stacks 4 and 4'. The triangles represent two possible $E2$ -brane $O(1)$ instantons, laying on top of the Ω -plane.

interaction terms involving the matter super-fields correspond to closed oriented polygons, starting with triangles. On the other hand, interactions between standard super-fields and modulini also correspond to closed oriented polygons involving solid and dashed lines.⁷

3 An unoriented quiver for a Pati-Salam model

In this section, we construct a simple quiver for a Pati-Salam model inducing a Neutron Majorana mass for the neutron. We propose a simple quiver in figure 2, leading to a susy $\mathcal{N} = 1$ Pati-Salam like model, with all the necessary fields and Yukawa's for a spontaneous symmetry breaking pattern to the Standard Model and for the generation of a Neutron Majorana mass from Exotic Instantons.

The gauge group is $U(4) \times Sp(2) \times Sp(2)$: $U(4)$ is generated by a stacks of 4 D-branes and their images $U(4)'$ under an Ω^+ -plane; $Sp(2)_{L,R}$ are generated by two stacks of two D-branes each on top of the Ω^+ -plane. We also consider Exotic $O(1)$ Instantons corresponding to $E2$ -branes on top of the Ω^+ -plane.

In particular, the three generations of Left and Right fundamental representations $F_{L,R}$, containing quarks and leptons, are reproduced as excitations of open strings attached to the $U(4)$ -stack and the Left or Right $Sp(2)_{L,R}$ stacks, respectively. We also get $\Delta = (1, 1, 10)$ and its conjugate from open strings attached to the $U(4)$ -stack and its mirror image $U(4)'$ -stack. $\phi_{RR} = (3, 1, 1)$ and $\phi_{LL} = (1, 3, 1)$ correspond to strings with both end-points attached on the $Sp(2)_{R,L}$ respectively. Higgs fields $h_{LR} = (2, 2, 1)$ are between

⁷One should keep in mind that exotic instantons are brane wrapping empty nodes of the quiver. Their interactions are thus coded in the quiver or dimer.




Figure 3. Amplitude in IIA (un)oriented string theory. Δ_{ab} sextets are excitations of strings attached between two intersecting $D6$ -branes, represented in figure as black lines. The fermionic moduli (or modulini) result from strings localised at the intersection of one $D6$ -brane and an exotic instanton Euclidean $D2$ -brane (or $E2$ brane). They are represented in figure as dashed lines. This system is embedded in some Calabi-Yau compactification CY_3 . In particular $D6$ -branes are wrapping 3-cycles on CY_3 and $E2$ are wrapping a different 3-cycle.

$Sp(2)_L$ and $Sp(2)_R$. It is amusing to observe that a two-generation model of the same kind would result in unoriented Type IIB from ‘fractional’ $D3$ -branes at a \mathbf{C}^3/Z_4 orbifold singularity.

The perturbative super-potential that we obtain from the quiver reads

$$\begin{aligned} \mathcal{W}_{\text{eff}} \sim & y_1 h_{LR} F_L F_R + \frac{1}{M_1} F_L \phi_{LL} F_L \Delta + \frac{1}{M_2} F_R \phi_{RR} F_R \Delta^c + \frac{1}{M_3} h_{LR} \phi_{RR} h_{RL} \phi_{LL} + \mu h_{LR} h_{RL} \\ & + m_\Delta \Delta \Delta^c + \frac{1}{4M_4} (\Delta \Delta^c)^2 + \frac{1}{2} m_L \phi_{LL}^2 + \frac{1}{2} m_R \phi_{RR}^2 + \frac{1}{3!} a_L \phi_{LL}^3 + \frac{1}{3!} a_R \phi_{RR}^3 \end{aligned} \quad (3.1)$$

where $\Delta \equiv \Delta_{qq}$ and the mass scales $M_{1,2,3,4}$ depend on the particular global completion of the model: they could be near M_S or at lower scales. In a T-dual Type IIB context, the mass terms m_Δ and $m_{L,R}$ can be generated by RR-RR or NS-NS three-forms fluxes in the bulk:

$$\begin{aligned} m_{\Delta_{qq}} & \sim \Gamma^{ijk} \langle \tau H_{ijk}^{(qq)} + i F_{ijk}^{(qq)} \rangle \\ m_{L,R} & \sim \Gamma^{ijk} \langle \tau H_{ijk}^{(L,R)} + i F_{ijk}^{(L,R)} \rangle \end{aligned}$$

with H RR-RR and F NS-NS three-forms and in general $H^{dd}, F^{dd} \neq H^{uu}, F^{uu}$ depend on the choice of fluxes through the relevant cycles wrapped by the D-branes.⁸

On the other hand, the non-perturbative superpotential term:

$$\mathcal{W}_{E2} = \frac{1}{\mathcal{M}_0} \epsilon^{ijkl} \epsilon^{i'j'k'l'} \Delta_{ii'}^c \Delta_{jj'}^c \Delta_{kk'}^c \Delta_{ll'}^c \quad (3.2)$$

can be generated by an $E2$ -brane instanton assuming the relevant $E2$ intersect twice the $U(4)$ stack of $D6$ -branes, necessary to produce a four- Δ^c (as well as a four- Δ) interaction.

⁸Mass deformed quivers and dimers have been recently investigated in [33].

The fermionic modulini τ_α^i , with $i = 1, \dots, 4$ and $\alpha, \beta = 1, 2$ interact with the super-fields Δ 's via

$$\mathcal{L}_{E2-D6-D6} \sim \tau_\alpha^i \Delta_{(ij)}^c \tau_\beta^j \epsilon^{\alpha\beta} + \tau_{i\alpha'}^c \Delta^{(ij)} \tau_{j\beta'}^c \epsilon^{\alpha'\beta'}. \quad (3.3)$$

These interactions are induced by mixed disk amplitudes, that emerge at the intersections between two $D6$ -brane stacks and one $E2$ instanton as the one in figure 3.⁹ In our case the two $D6$ -branes are actually the $D6$ -branes of the $U(4)$ stack and their images. Integrating out the fermionic modulini produces two ϵ^{ijkl} so that schematically

$$\mathcal{W}_{E2} = \frac{1}{\mathcal{M}_0} \int d^8 \tau e^{\mathcal{L}_{E2-D6-D6}} = \frac{1}{\mathcal{M}_0} \epsilon^{ijkl} \epsilon^{i'j'k'l'} \Delta_{ii'}^c \Delta_{jj'}^c \Delta_{kk'}^c \Delta_{ll'}^c. \quad (3.4)$$

The suppression scale \mathcal{M}_0 is related to the string scale M_S as $\mathcal{M}_0 \sim M_S e^{+S_{E2}}$, where S_{E2} depends on the closed string moduli that parametrize the (complex) size of the 3-cycles, wrapped by the $E2$ -instantons.¹⁰

After the decomposition $SU(4) \rightarrow SU(3) \times U(1)_{B-L}$, $\mathbf{10}_+ \rightarrow (\mathbf{6}_{+2/3}, \mathbf{3}_{-2/3}, \mathbf{1}_{-2})$, we denote by Δ_6 the ‘diquark’ super-field in the $\mathbf{6}$, T_3 in the triplet $\mathbf{3}$ and S the singlet $\mathbf{1}$ and find

$$\frac{1}{M_2} \phi_{\dot{\alpha}\dot{\beta}}^c F_i^{c\dot{\alpha}} F_j^{c\dot{\beta}} \Delta^{c(ij)} \rightarrow \frac{1}{M_2} \phi_{\dot{\alpha}\dot{\beta}}^c \left[Q_i^{c\dot{\alpha}} Q_j^{c\dot{\beta}} \Delta_6^{cjj} + 2 Q_i^{c\dot{\alpha}} L^{c\dot{\beta}} T_3^{c\dot{\beta}} + L^{c\dot{\alpha}} L^{c\dot{\beta}} S^c \right] \quad (3.5)$$

$$m_\Delta \Delta \Delta^c + \frac{1}{4M_4} (\Delta \Delta^c)^2 \rightarrow m_\Delta (\Delta_6 \Delta_6^c + TT^c + SS^c) + \frac{1}{4M_4} (\Delta_6 \Delta_6^c + TT^c + SS^c)^2 \quad (3.6)$$

$$\begin{aligned} & \frac{1}{\mathcal{M}_0} \epsilon_{ijkl} \epsilon_{i'j'k'l'} \Delta^{cii'} \Delta^{cjj'} \Delta^{ckk'} \Delta^{c ll'} \\ & \rightarrow \frac{1}{\mathcal{M}_0} \left[4 \epsilon_{ijk}^{SU(3)} \epsilon_{i'j'k'}^{SU(3)} \Delta_6^{cii'} \Delta_6^{cjj'} \Delta_6^{ckk'} S + 6 \epsilon_{ijk}^{SU(3)} \epsilon_{i'j'k'}^{SU(3)} \Delta_6^{cii'} \Delta_6^{cjj'} T_3^c T_3^{c k'} \right]. \end{aligned} \quad (3.7)$$

The complete super-potential after the decomposition $SU(4) \rightarrow SU(3) \times U(1)_{B-L}$ reads

$$\begin{aligned} \mathcal{W} & \sim y_1 h_{\alpha\dot{\alpha}} Q^{i\alpha} Q^{c\dot{\alpha}} + y_1 h_{\alpha\dot{\alpha}} L^\alpha L^{c\dot{\alpha}} + \frac{1}{M_1} \phi_{\alpha\beta} \left(Q^{i\alpha} Q^{j\beta} \Delta_{ij} + Q^{i\alpha} L^\beta T_{3i} + L^\alpha L^\beta S \right) \quad (3.8) \\ & + \frac{1}{M_2} \phi_{\alpha\beta}^c (Q^{c\dot{\alpha}} Q^{c\dot{\beta}} \Delta_{ij}^c + Q^{c\dot{\alpha}} L^{c\dot{\beta}} T_{3i}^c + L^{c\dot{\alpha}} L^{c\dot{\beta}} S^c) + \mu h_{\alpha\dot{\alpha}} h^{\alpha\dot{\alpha}} \\ & + \frac{1}{M_3} h^c \phi^c h \phi + m_\Delta (\Delta_6 \Delta_6^c + TT^c + SS^c) + \frac{1}{4M_4} (\Delta_6 \Delta_6^c + TT^c + SS^c)^2 + V(\phi_{L,R}) \\ & + \frac{1}{\mathcal{M}_0} \left[4 \epsilon_{ijk}^{SU(3)} \epsilon_{i'j'k'}^{SU(3)} \Delta_6^{cii'} \Delta_6^{cjj'} \Delta_6^{ckk'} S + 6 \epsilon_{ijk}^{SU(3)} \epsilon_{i'j'k'}^{SU(3)} \Delta_6^{cii'} \Delta_6^{cjj'} T_3^c T_3^{c k'} + c.c \right] \end{aligned}$$

where $V(\phi_{L,R}) = m_{L,R} \phi_{L,R}^2 + a_{L,R} \phi_{L,R}^3 / 3$.

⁹For similar calculations in related contexts, see [70–77].

¹⁰In general, the calculations could be much more complicated, in the presence of bulk fluxes, that can also induce soft susy breaking mass terms for the susy partners. For example gaugino mass terms with $M_\lambda \sim \Omega_{0,3}^{ijk} \langle \tau H_{ijk} + i F_{ijk} \rangle$ can be generated in Type IIB contexts by internal 3-form fluxes. In the presence of fluxes, one has to verify that physical branes and instantons are not lifted, i.e. the cycles they wrap and their intersections are not eliminated. With the introduction of bulk fluxes, one also has to consider the back-reactions on the exotic instantons, that could change then number of zero modes. This could modify our present analysis.

When $\langle \phi_{RR} \rangle = v_R$ and $\langle \phi_{LL} \rangle = 0$, Left-Right symmetry is spontaneously broken. When $\langle S \rangle = v_{B-L}$, $SU(4)$ and its subgroup $U(1)_{B-L}$ are spontaneously broken. MB: contrary to $U(1)_4$ that extends $SU(4)$ to $U(4)$ and is anomalous, both $U(1)_{B-L}$ and $U(1)_Y$ are non-anomalous being associated to traceless generators of the non-abelian groups $SU(4)$ and $Sp(2)_R$. A Majorana mass term for the Neutrino is generated¹¹ as $m_N \sim v_R v_{B-L} / M_2$. For example, $m_N \simeq 10^{12}$ GeV can be obtained if $v_R \simeq M_2$ and $v_{B-L} \simeq 10^{12}$ GeV. In this model the generation of a neutrino Majorana mass is connected to the generation of a Neutron Majorana mass.

In fact, when S takes an expectation value, a cubic interaction term

$$(v_{B-L} / \mathcal{M}_0) \epsilon_{ijk}^{SU(3)} \epsilon_{i'j'k'}^{SU(3)} \Delta_6^{ci'} \Delta_6^{cj'} \Delta_6^{ck'}$$

is generated. In section 4, we will discuss the consequences, for Neutron-Antineutron physics and for LHC phenomenology, in more in details.

3.1 No-proton decays

The quiver in figure 2 is constructed in such a way that, starting from a R-parity preserving susy Pati-Salam model, $E2, E2'$ -instantons non-perturbatively generate only two R-parity violating operators: (3.3) and its complex conjugate $\mathcal{W}_{E2'} = \Delta\Delta\Delta\Delta / \mathcal{M}'_0$, respectively. For example, superpotentials like $\mathcal{W}_{ex1} = F_L F_L F_L F_L / \Lambda_1$ and $\mathcal{W}_{ex2} = F_R F_R F_R F_R / \Lambda_2$ are not generated, simply for construction: exotic instantons that could generate such superpotentials are not introduced in figure 2. As a consequence, there is no proton decay in our model. This is a nice feature of a dynamical R-parity breaking from Exotic Instantons, with respect to an explicit R-violation: we can generate specific superpotentials without other dangerous ones.

4 Neutron-Antineutron oscillation through color diquark sextets

In a susy PS-like model $SU(4) \times SU(2)_R \times SU(2)_L$, we can construct a diagram like the one in figure 1 for Neutron-Antineutron oscillation, through the ‘exotic’ interaction $\Delta_{10}^c \Delta_{10}^c \Delta_{10}^c \Delta_{10}^c / \mathcal{M}_0$, containing

$$\mathcal{W}_{\Delta B=2} = \frac{1}{\mathcal{M}_0} \epsilon^{u^c d^c d^c \nu^c} \epsilon^{u'^c d'^c d'^c \nu'^c} \Delta_{u^c u'^c}^c \Delta_{d^c d'^c}^c \Delta_{d^c d'^c}^c S_{\nu^c \nu'^c}^c \quad (4.1)$$

(with $S_{\nu^c \nu'^c}^c \equiv \Delta_{\nu^c \nu'^c}^c$). The operator (4.1) induces a neutron-antineutron transition depicted in figure 2, as a result of the super-potential term $\tilde{f}_{11} v_R Q^c Q^c \Delta^c / M_2$, whose components include $f_{11} \Delta_{u^c u^c}^c u^c u^c$ and $f_{11} \Delta_{d^c d^c}^c d^c d^c$, with $f_{11} = \tilde{f}_{11} v_R / M_2$, and \tilde{f}_{11} Yukawa couplings.

The process in figure 1-(b) produces an effective operator $G_{n-\bar{n}}(udd)^2$ with

$$G_{n-\bar{n}} \simeq \frac{g_3^2}{16\pi} \frac{f_{11}^2 v_{BL}}{M_{\Delta_{u^c u^c}^c}^2 M_{\Delta_{d^c d^c}^c}^2 M_{SUSY} \mathcal{M}_0}. \quad (4.2)$$

We can now discuss different choices of the parameters leading to very different branches for phenomenology. The motivation of such a variety of possibilities is related

¹¹Dirac masses are generated via Yukawa couplings when h_{LR} gets a VEV.

to the fact that in (4.2) one can produce a scale of 300–1000 TeV, testable in the next generation of experiments, with different choices of the other parameters.

The cases with $\mathcal{M}_0 \simeq 10^{19}$ GeV and $\mathcal{M}_0 \simeq 10^{13}$ GeV are equivalent to the GUT SO(10) inspired scenario, discussed in [14] in figure 1-(b). In these cases $M_{\Delta_{u^c u^c}^c} \sim 1$ TeV. Both cases are well compatible with the mechanism proposed in the previous section. In fact a scenario in which $\mathcal{M}_0 \simeq M_S$ can be envisaged, if $e^{S_{E_2}} \sim 1$ i.e. small 3-cycles wrapped by E_2 in CY . *A priori*, the string scale can be considered as a free parameter, it can be as high as 10^{19} TeV as low as a few TeV's. For instance, if $M_S = 1 \div 10$ TeV, the hierarchy problem of the Higgs mass is automatically solved, and \mathcal{M}_0 can be as high as 10^{13} GeV (or more) if $e^{S_{E_2}} = 10^{10}$ i.e. for an E_2 wrapping a large 3-cycle in the CY .

In TeV-scale gravity scenari, one can also consider an alternative scenario in which $\mathcal{M}_0 \simeq M_S \simeq 10$ TeV, with $e^{S_{E_2}} \sim 1$. Compatibly with $n - \bar{n}$ limits, the four-sextets' interaction (3.2) is suppressed at much lower scales, with $\mathcal{M}_0 \sim 10$ TeV. In this case, the masses of the sextets have to be much higher than 1 TeV, eluding completely a direct observation at LHC and in FCNCs. On the other hand, a post sphaleron baryogenesis scenario as the one proposed in [14] remains possible.

So, combined observations from the next generation of experiments on neutron-antineutron oscillations, LHC, FCNC physics, neutrino physics can provide precious informations not only on a possible PS extension on the SM and its region of the parameters, but also on the dynamical scale generated by the Exotic instantons, and as a consequence on the geometric structure of the Calabi-Yau compactifications, in particular the 3-cycle wrapped by the Exotic Instanton! This is a fundamental information for realistic model building in string phenomenology. Exotic instantons could be *portals* from low energy physics to the quantum gravity scale! In the next section, we will briefly discuss connections with dark matter and the hierarchy problem. Note that in our scenario, $\langle S \rangle \neq \langle \phi_{RR} \rangle$ in general. This is an important difference with respect to Babu-Mohapatra model cited above: $n - \bar{n}$ oscillation time of order 10^{10} s is compatible with a Left-Right symmetry restoration at TeV scale, with intriguing implications for LHC. A recent anomaly with significance near 3σ , compatible with Left-Right symmetry, in $pp \rightarrow l_1 l_2 j j$, was seen by CMS [80]. In a Left-Right model, this is interpreted as sequential W_R and N_R production as [81–84]

$$pp \rightarrow W_R \rightarrow l_1 N_R \rightarrow l_1 l_2 W_R^* \rightarrow l_1 l_2 j j .$$

However, this interpretation requires $g_R(M_{W_R}) \simeq 0.6 g_L(M_{W_R})$, with $1.8 \text{ TeV} < M_{W_R} < 2.4 \text{ TeV}$. Curiously, this situation is not compatible with a D-parity preserving SO(10) GUT scenario. For P-S models emerging from $\text{SO}(10) \rightarrow \text{SU}(4) \times \text{SU}(2)_L \times \text{SU}(2)_R$ D-parity is a symmetry. It is the external automorphisms that exchanges the two SU(2) groups, $\text{SU}(2)_L \leftrightarrow \text{SU}(2)_R$, and at the same time acts by conjugation on SU(4) representations. More explicitly $D_P = \Gamma_7 \gamma_5$ is a symmetry when $g_L = g_R$. At the unification scale $g_L = g_R$, the two coupling constants have the same running if the field content is LR symmetric. For a LR interpretation of CMS anomaly one needs $g_L \neq g_R$, compatibly with a PS model not emerging from SO(10) without D-parity altogether or at least an SO(10) model where D-parity is broken at a high scale. On the other hand, D-parity is not a symmetry in string-

inspired models. The gauge couplings depend on the size of the internal cycle wrapped by the D-branes. In orbifolds or Calabi-Yau singularities, one can tune the blow-up modes so that different cycle have the same size e.g. vanishing, but generically this is not the case and one can start with $g_L \neq g_R$ already at the string scale.

$pp \rightarrow l_1 l_2 j j$ is not the only peculiar channel suggested by our model for LHC, and we will return to other ones in the next section. We conclude this section with another observation. A scenario in which S doesn't take an expectation value at all can be envisaged in our model. In this case, $U(1)_{B-L}$ is not spontaneously broken by S . However, as mentioned above, exotic instantons can dynamically break $U(1)_{B-L}$. For instance, a Majorana mass matrix for RH neutrini can be generated by exotic instantons rather than by S , as cited above. In this case, S could also be a light particle, if a residual discrete symmetry of $U(1)_{B-L}$ stabilizes it. In other words, S can behave as a Majoron, but it is not exactly a Majoron [85]. We can call it an *exoticon*. We suppose that the exoticon interacts with the three color sextets with a coupling μ_S . So, in this case, we have to replace v_{B-L} with μ_S in (4.2). The exoticon carries $B-L = -2$, so $n \rightarrow \bar{n}S$ does not violate $B-L$. We also note another important difference with respect to Majorons: Majoron mass $m_\phi = y_L v_L$, with v_L vev of a global $U(1)_L$ (and y_L coupling), is related to its interaction with neutrini, as $g_{\phi\nu\nu} = m_\nu/v_L$; such a relation, in general, is not satisfied by exoticons. A massive exoticon cannot be emitted in a $n - \bar{n}$ transition, in the vacuum: CPT symmetry protects neutron by transitions $n \rightarrow \bar{n} + S$, i.e. $m_n = m_{\bar{n}}$. However, in a nuclear environment, such a transition is allowed! Such a transition is followed by annihilation of the antineutron with another neutron in the nuclear environment, as $(Z, A) \rightarrow (Z, A - 2) + 3\pi$. We can roughly estimate the corresponding decay width as $\Gamma \simeq (\delta m/\mu_S)^2 \langle \Delta \mathcal{E} \rangle$, where $\langle \Delta \mathcal{E} \rangle \simeq 10 \div 100$ MeV is the average energy in the nuclear environment. Limits on $n - \bar{n}$ oscillation in the nuclei are $\Gamma_{n\bar{n}}^{-1} \sim 10^{-32}$ yr [86], corresponding to $\mu_S > 10^{30} \delta m \simeq$ keV. Another spectacular signature of an exoticon could be a nucleon-nucleon disappearances as $nn \rightarrow S$, $\Delta B = 2$. This could be detected as a nuclear transition $(Z, A) \rightarrow (Z, A - 2) +$ missing energy. We can easily estimate the rate of such a transition as $\Gamma \sim \kappa_{np} (\mathcal{M}_{n\bar{n}})^{-3} m_N^{14} G_{n-\bar{n}}^2$ GeV, where $\kappa_{np} \sim 10^{-6}$ approximately accounts for the hadronic non-perturbative correction. Such an estimate leads us to conclude that such a process is very suppressed, roughly as 10^{40} yr. Finally, an exoticon can be also detected in a neutrinoless double-beta-decay, as a Majoron. However, there are several important differences with respect to the Majoron: a $0\nu 2\beta + S$ does not violate lepton number, it is an *apparent violation*. For a $0\nu 2\beta + S$ process, limits on the exoticon production imply $(m_\nu/\mu_S) < 10^{-5}$ [87, 88] that corresponds to a bound $\mu_S > 10$ keV. Limits from supernovae cooling processes $\nu \rightarrow \nu^c S$ or $\nu\nu \rightarrow SS$ are competitive (for electronic neutrini $m_\nu/\mu_S < 10^{-5}$) [89].

5 Other comments on phenomenology, dark matter and hierarchy problem

In Minimal non-susy Left-Right models, identifying a candidate for cold dark matter is difficult. On the other hand, our model automatically suggests several candidates for cold dark matter. In fact, our model predicts the presence of neutralini and stueckelberg axini

(or stuckelini), mixing with each other, as in [90, 91]. They are good candidate for WIMP dark matter. On the other hand, extra Z' for each anomalous $U(1)$ can get a mass from Stuckelberg mechanism, that is not necessary of the order of string scale M_S . In particular, a scenario in which $M_S \simeq 10 \div 10^3$ TeV can be envisaged, alleviating hierarchy problem of Higgs mass. In this case, $m_{Z'} \simeq 10^{-4} M_S \div M_S \simeq 1 \div 10$ TeV, with Generalized Chern Simon terms inducing peculiar decays as $Z' \rightarrow Z\gamma$ [25–32, 90–92].¹² Curiously, the presence of apparent flavor violations in $B \rightarrow llK$, detected in LHCb [100–103] could be a hint in favor of a new Z' [104–106]. In our model, we also predict extra decays $B \rightarrow l^+l^-l'^+l'^-K$ suppressed by the GCS couplings with respect to $B \rightarrow l^+l^-K$. Extra Z' from anomalous symmetries are different with respect to Z'_R (Z -boson of the $SU(2)_R$), and kinetic mixings $Z - Z'$ or $Z' - Z'_R$ can be envisaged. Z'_R can also interact Z, γ, Z', γ through G.C.S, and a complete study, of resulting cascade processes, is beyond the purpose of this paper. Concerning the hierarchy problem of the Higgs mass, our model is compatible with TeV-scale supersymmetry, but this model has more undetermined parameters than the MSSM, i.e. it could be more elusive and more difficult to constrain.

6 Conclusions and remarks

In this paper, we have shown how to generate a Majorana mass for the neutron, inducing neutron-antineutron transitions with $|\Delta B| = 2$ in the context of a Pati-Salam Left-Right model. Indeed exotic instantons can produce an effective interaction involving color diquark sextets, leading to a Majorana mass term for the neutron. We would like to stress that in the present context no processes with $|\Delta B| = 1$ are allowed that could lead to fast proton decay. We have discussed some possible phenomenological implications, in the main branches of the parameters space, for LHC, FCNC, Dark Matter, $0\nu 2\beta$ -decay. A unifying picture of Dark Matter, Hierarchy problem of the Higgs mass, Baryogenesis and Neutrino mass emerges in a very simple unoriented quiver! In this sense, the model elaborated here could represent a serious alternative to GUT inspired models.

Acknowledgments

We would like to thank R. Mohapatra and A. Pascal for reading the manuscript and suggesting insightful comments. We would also like to thank C. Bachas, F. Quevedo, J. Valle and G. Villadoro for useful discussions. A. A. would like to thank Galileo Galilei Institute for Theoretical Physics for the hospitality, where this paper was prepared.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

¹²We would like to stress that GCS terms generate UV divergent triangles that are cured by considering UV completions with KK states or string excitations. For issues in scattering amplitudes and collider physics see [96], for recent discussion about string theory and causality see [94, 95]. For non-local (string inspired) quantum field theories see [97–99].

References

- [1] J.C. Pati and A. Salam, *Lepton Number as the Fourth Color*, *Phys. Rev. D* **10** (1974) 275 [[INSPIRE](#)].
- [2] R.N. Mohapatra and R.E. Marshak, *Local B-L Symmetry of Electroweak Interactions, Majorana Neutrinos and Neutron Oscillations*, *Phys. Rev. Lett.* **44** (1980) 1316 [[INSPIRE](#)].
- [3] R.N. Mohapatra and G. Senjanović, *Neutrino Mass and Spontaneous Parity Violation*, *Phys. Rev. Lett.* **44** (1980) 912 [[INSPIRE](#)].
- [4] J. Schechter and J.W.F. Valle, *Neutrino Masses in $SU(2) \times U(1)$ Theories*, *Phys. Rev. D* **22** (1980) 2227 [[INSPIRE](#)].
- [5] J. Schechter and J.W.F. Valle, *Neutrino Decay and Spontaneous Violation of Lepton Number*, *Phys. Rev. D* **25** (1982) 774 [[INSPIRE](#)].
- [6] K.S. Babu, R.N. Mohapatra and S. Nasri, *Unified TeV Scale Picture of Baryogenesis and Dark Matter*, *Phys. Rev. Lett.* **98** (2007) 161301 [[hep-ph/0612357](#)] [[INSPIRE](#)].
- [7] K.S. Babu, R.N. Mohapatra and S. Nasri, *Post-Sphaleron Baryogenesis*, *Phys. Rev. Lett.* **97** (2006) 131301 [[hep-ph/0606144](#)] [[INSPIRE](#)].
- [8] K.S. Babu, P.S. Bhupal Dev and R.N. Mohapatra, *Neutrino mass hierarchy, neutron-antineutron oscillation from baryogenesis*, *Phys. Rev. D* **79** (2009) 015017 [[arXiv:0811.3411](#)] [[INSPIRE](#)].
- [9] K.S. Babu, P.S. Bhupal Dev, E.C. F.S. Fortes and R.N. Mohapatra, *Post-Sphaleron Baryogenesis and an Upper Limit on the Neutron-Antineutron Oscillation Time*, *Phys. Rev. D* **87** (2013) 115019 [[arXiv:1303.6918](#)] [[INSPIRE](#)].
- [10] R.N. Mohapatra and J.W.F. Valle, *Neutrino Mass and Baryon Number Nonconservation in Superstring Models*, *Phys. Rev. D* **34** (1986) 1642 [[INSPIRE](#)].
- [11] R.N. Mohapatra, N. Okada and H.-B. Yu, *Diquark Higgs at LHC*, *Phys. Rev. D* **77** (2008) 011701 [[arXiv:0709.1486](#)] [[INSPIRE](#)].
- [12] Y.C. Zhan, Z.L. Liu, S.A. Li, C.S. Li and H.T. Li, *Threshold resummation for the production of a color sextet (antitriplet) scalar at the LHC*, *Eur. Phys. J. C* **74** (2014) 2716 [[arXiv:1305.5152](#)] [[INSPIRE](#)].
- [13] E.C.F.S. Fortes, K.S. Babu and R.N. Mohapatra, *Flavor Physics Constraints on TeV Scale Color Sextet Scalars*, [arXiv:1311.4101](#) [[INSPIRE](#)].
- [14] R.N. Mohapatra, *Neutron-Anti-Neutron Oscillation: Theory and Phenomenology*, *J. Phys. G* **36** (2009) 104006 [[arXiv:0902.0834](#)] [[INSPIRE](#)].
- [15] P. Anastasopoulos, G.K. Leontaris and N.D. Vlachos, *Phenomenological Analysis of D-brane Pati-Salam Vacua*, *JHEP* **05** (2010) 011 [[arXiv:1002.2937](#)] [[INSPIRE](#)].
- [16] M. Berkooz, M.R. Douglas and R.G. Leigh, *Branes intersecting at angles*, *Nucl. Phys. B* **480** (1996) 265 [[hep-th/9606139](#)] [[INSPIRE](#)].
- [17] M. Bianchi and A. Sagnotti, *On the systematics of open string theories*, *Phys. Lett. B* **247** (1990) 517 [[INSPIRE](#)].
- [18] M. Bianchi and A. Sagnotti, *Twist symmetry and open string Wilson lines*, *Nucl. Phys. B* **361** (1991) 519 [[INSPIRE](#)].

- [19] M. Bianchi, G. Pradisi and A. Sagnotti, *Toroidal compactification and symmetry breaking in open string theories*, *Nucl. Phys. B* **376** (1992) 365 [INSPIRE].
- [20] A.M. Uranga, *D-brane probes, RR tadpole cancellation and k-theory charge*, *Nucl. Phys. B* **598** (2001) 225 [hep-th/0011048] [INSPIRE].
- [21] G. Aldazabal, S. Franco, L.E. Ibáñez, R. Rabadán and A.M. Uranga, *D = 4 chiral string compactifications from intersecting branes*, *J. Math. Phys.* **42** (2001) 3103 [hep-th/0011073] [INSPIRE].
- [22] M. Bianchi and J.F. Morales, *Anomalies & tadpoles*, *JHEP* **03** (2000) 030 [hep-th/0002149] [INSPIRE].
- [23] C. Angelantonj, M. Bianchi, G. Pradisi, A. Sagnotti and Y. Stanev, *Chiral asymmetry in four-dimensional open string vacua*, *Phys. Lett. B* **385** (1996) 96 [hep-th/9606169] [INSPIRE].
- [24] C. Angelantonj, M. Bianchi, G. Pradisi, A. Sagnotti and Y.S. Stanev, *Comments on Gepner models and type-I vacua in string theory*, *Phys. Lett. B* **387** (1996) 743 [hep-th/9607229] [INSPIRE].
- [25] P. Anastasopoulos, M. Bianchi, E. Dudas and E. Kiritsis, *Anomalies, anomalous U(1)'s and generalized Chern-Simons terms*, *JHEP* **11** (2006) 057 [hep-th/0605225] [INSPIRE].
- [26] J. De Rydt, J. Rosseel, T.T. Schmidt, A. Van Proeyen and M. Zagermann, *Symplectic structure of N = 1 supergravity with anomalies and Chern-Simons terms*, *Class. Quant. Grav.* **24** (2007) 5201 [arXiv:0705.4216] [INSPIRE].
- [27] D. Feldman, Z. Liu and P. Nath, *The Stueckelberg Z-prime Extension with Kinetic Mixing and Milli-Charged Dark Matter From the Hidden Sector*, *Phys. Rev. D* **75** (2007) 115001 [hep-ph/0702123] [INSPIRE].
- [28] D. Feldman, Z. Liu and P. Nath, *The Stueckelberg extension and milli weak and milli charged dark matter*, *AIP Conf. Proc.* **939** (2007) 50 [arXiv:0705.2924] [INSPIRE].
- [29] B. Körs and P. Nath, *A Stueckelberg extension of the standard model*, *Phys. Lett. B* **586** (2004) 366 [hep-ph/0402047] [INSPIRE].
- [30] B. Körs and P. Nath, *A Supersymmetric Stueckelberg U(1) extension of the MSSM*, *JHEP* **12** (2004) 005 [hep-ph/0406167] [INSPIRE].
- [31] B. Körs and P. Nath, *How Stueckelberg extends the standard model and the MSSM*, [hep-ph/0411406] [INSPIRE].
- [32] B. Körs and P. Nath, *Aspects of the Stueckelberg extension*, *JHEP* **07** (2005) 069 [hep-ph/0503208] [INSPIRE].
- [33] M. Bianchi, S. Cremonesi, A. Hanany, J.F. Morales, D.R. Pacifici and R.K. Seong, *Mass-deformed Brane Tilings*, *JHEP* **1410** (2014) 27 [arXiv:1408.1957] [INSPIRE].
- [34] C. Beasley, J.J. Heckman and C. Vafa, *GUTs and Exceptional Branes in F-theory — I*, *JHEP* **01** (2009) 058 [arXiv:0802.3391] [INSPIRE].
- [35] R. Donagi and M. Wijnholt, *Model Building with F-theory*, *Adv. Theor. Math. Phys.* **15** (2011) 1237 [arXiv:0802.2969] [INSPIRE].
- [36] C. Beasley, J.J. Heckman and C. Vafa, *GUTs and Exceptional Branes in F-theory — II: Experimental Predictions*, *JHEP* **01** (2009) 059 [arXiv:0806.0102] [INSPIRE].

- [37] R. Donagi and M. Wijnholt, *Breaking GUT Groups in F-theory*, *Adv. Theor. Math. Phys.* **15** (2011) 1523 [[arXiv:0808.2223](#)] [[INSPIRE](#)].
- [38] R. Blumenhagen, V. Braun, T.W. Grimm and T. Weigand, *GUTs in Type IIB Orientifold Compactifications*, *Nucl. Phys. B* **815** (2009) 1 [[arXiv:0811.2936](#)] [[INSPIRE](#)].
- [39] C.-M. Chen and Y.-C. Chung, *A Note on Local GUT Models in F-theory*, *Nucl. Phys. B* **824** (2010) 273 [[arXiv:0903.3009](#)] [[INSPIRE](#)].
- [40] J. Marsano, N. Saulina and S. Schäfer-Nameki, *F-theory Compactifications for Supersymmetric GUTs*, *JHEP* **08** (2009) 030 [[arXiv:0904.3932](#)] [[INSPIRE](#)].
- [41] T. Kobayashi, S. Raby and R.-J. Zhang, *Constructing 5D orbifold grand unified theories from heterotic strings*, *Phys. Lett. B* **593** (2004) 262 [[hep-ph/0403065](#)] [[INSPIRE](#)].
- [42] T. Kobayashi, S. Raby and R.-J. Zhang, *Searching for realistic 4d string models with a Pati-Salam symmetry: Orbifold grand unified theories from heterotic string compactification on a Z_6 orbifold*, *Nucl. Phys. B* **704** (2005) 3 [[hep-ph/0409098](#)] [[INSPIRE](#)].
- [43] W. Buchmüller, K. Hamaguchi, O. Lebedev and M. Ratz, *Supersymmetric standard model from the heterotic string*, *Phys. Rev. Lett.* **96** (2006) 121602 [[hep-ph/0511035](#)] [[INSPIRE](#)].
- [44] W. Buchmüller, K. Hamaguchi, O. Lebedev and M. Ratz, *Supersymmetric Standard Model from the Heterotic String (II)*, *Nucl. Phys. B* **785** (2007) 149 [[hep-th/0606187](#)] [[INSPIRE](#)].
- [45] O. Lebedev et al., *A mini-landscape of exact MSSM spectra in heterotic orbifolds*, *Phys. Lett. B* **645** (2007) 88 [[hep-th/0611095](#)] [[INSPIRE](#)].
- [46] O. Lebedev et al., *The Heterotic Road to the MSSM with R parity*, *Phys. Rev. D* **77** (2008) 046013 [[arXiv:0708.2691](#)] [[INSPIRE](#)].
- [47] O. Lebedev, H.P. Nilles, S. Ramos-Sanchez, M. Ratz and P.K.S. Vaudrevange, *Heterotic mini-landscape (II): Completing the search for MSSM vacua in a Z_6 orbifold*, *Phys. Lett. B* **668** (2008) 331 [[arXiv:0807.4384](#)] [[INSPIRE](#)].
- [48] J.E. Kim, J.-H. Kim and B. Kyae, *Superstring standard model from Z_{12-I} orbifold compactification with and without exotics and effective R-parity*, *JHEP* **06** (2007) 034 [[hep-ph/0702278](#)] [[INSPIRE](#)].
- [49] J.E. Kim and B. Kyae, *Kaluza-Klein masses in nonprime orbifolds: Z_{12-I} compactification and threshold correction*, *Phys. Rev. D* **77** (2008) 106008 [[arXiv:0712.1596](#)] [[INSPIRE](#)].
- [50] M. Bianchi, A. Collinucci and L. Martucci, *Magnetized E3-brane instantons in F-theory*, *JHEP* **12** (2011) 045 [[arXiv:1107.3732](#)] [[INSPIRE](#)].
- [51] M. Bianchi, G. Inverso and L. Martucci, *Brane instantons and fluxes in F-theory*, *JHEP* **07** (2013) 037 [[arXiv:1212.0024](#)] [[INSPIRE](#)].
- [52] I.R. Klebanov and E. Witten, *Superconformal field theory on three-branes at a Calabi-Yau singularity*, *Nucl. Phys. B* **536** (1998) 199 [[hep-th/9807080](#)] [[INSPIRE](#)].
- [53] D.R. Morrison and M.R. Plesser, *Nonspherical horizons. 1*, *Adv. Theor. Math. Phys.* **3** (1999) 1 [[hep-th/9810201](#)] [[INSPIRE](#)].
- [54] S. Benvenuti, S. Franco, A. Hanany, D. Martelli and J. Sparks, *An Infinite family of superconformal quiver gauge theories with Sasaki-Einstein duals*, *JHEP* **06** (2005) 064 [[hep-th/0411264](#)] [[INSPIRE](#)].
- [55] S. Franco et al., *Gauge theories from toric geometry and brane tilings*, *JHEP* **01** (2006) 128 [[hep-th/0505211](#)] [[INSPIRE](#)].

- [56] A. Hanany and K.D. Kennaway, *Dimer models and toric diagrams*, [hep-th/0503149](#) [[INSPIRE](#)].
- [57] S. Franco, A. Hanany, K.D. Kennaway, D. Vegh and B. Wecht, *Brane dimers and quiver gauge theories*, *JHEP* **01** (2006) 096 [[hep-th/0504110](#)] [[INSPIRE](#)].
- [58] A. Hanany, V. Jejjala, S. Ramgoolam and R.-K. Seong, *Calabi-Yau Orbifolds and Torus Coverings*, *JHEP* **09** (2011) 116 [[arXiv:1105.3471](#)] [[INSPIRE](#)].
- [59] A. Hanany and R.-K. Seong, *Brane Tilings and Reflexive Polygons*, *Fortsch. Phys.* **60** (2012) 695 [[arXiv:1201.2614](#)] [[INSPIRE](#)].
- [60] A. Hanany and R.-K. Seong, *Brane Tilings and Specular Duality*, *JHEP* **08** (2012) 107 [[arXiv:1206.2386](#)] [[INSPIRE](#)].
- [61] S. Franco et al., *Dimers and orientifolds*, *JHEP* **09** (2007) 075 [[arXiv:0707.0298](#)] [[INSPIRE](#)].
- [62] C. Angelantonj, M. Bianchi, G. Pradisi, A. Sagnotti and Y. Stanev, *Chiral asymmetry in four-dimensional open string vacua*, *Phys. Lett. B* **385** (1996) 96 [[hep-th/9606169](#)] [[INSPIRE](#)].
- [63] G. Aldazabal, L.E. Ibáñez, F. Quevedo and A.M. Uranga, *D-branes at singularities: A bottom up approach to the string embedding of the standard model*, *JHEP* **08** (2000) 002 [[hep-th/0005067](#)] [[INSPIRE](#)].
- [64] M. Cvetič, G. Shiu and A.M. Uranga, *Three family supersymmetric standard-like models from intersecting brane worlds*, *Phys. Rev. Lett.* **87** (2001) 201801 [[hep-th/0107143](#)] [[INSPIRE](#)].
- [65] R. Blumenhagen, B. Körs, D. Lüst and S. Stieberger, *Four-dimensional String Compactifications with D-branes, Orientifolds and Fluxes*, *Phys. Rept.* **445** (2007) 1 [[hep-th/0610327](#)] [[INSPIRE](#)].
- [66] M. Bianchi, G. Inverso, J.F. Morales and D.R. Pacifici, *Unoriented Quivers with Flavour*, *JHEP* **01** (2014) 128 [[arXiv:1307.0466](#)] [[INSPIRE](#)].
- [67] M. Bianchi and M. Samsonyan, *Notes on unoriented D-brane instantons*, *Int. J. Mod. Phys. A* **24** (2009) 5737 [[arXiv:0909.2173](#)] [[INSPIRE](#)].
- [68] M. Bianchi and G. Inverso, *Unoriented D-brane instantons*, *Fortsch. Phys.* **60** (2012) 822 [[arXiv:1202.6508](#)] [[INSPIRE](#)].
- [69] R. Blumenhagen, M. Cvetič, S. Kachru and T. Weigand, *D-Brane Instantons in Type II Orientifolds*, *Ann. Rev. Nucl. Part. Sci.* **59** (2009) 269 [[arXiv:0902.3251](#)] [[INSPIRE](#)].
- [70] R. Blumenhagen, M. Cvetič and T. Weigand, *Spacetime instanton corrections in 4D string vacua: The Seesaw mechanism for D-brane models*, *Nucl. Phys. B* **771** (2007) 113 [[hep-th/0609191](#)] [[INSPIRE](#)].
- [71] L.E. Ibáñez and A.M. Uranga, *Neutrino Majorana Masses from String Theory Instanton Effects*, *JHEP* **03** (2007) 052 [[hep-th/0609213](#)] [[INSPIRE](#)].
- [72] L.E. Ibáñez, A.N. Schellekens and A.M. Uranga, *Instanton Induced Neutrino Majorana Masses in CFT Orientifolds with MSSM-like spectra*, *JHEP* **06** (2007) 011 [[arXiv:0704.1079](#)] [[INSPIRE](#)].
- [73] S. Antusch, L.E. Ibáñez and T. Macri, *Neutrino masses and mixings from string theory instantons*, *JHEP* **09** (2007) 087 [[arXiv:0706.2132](#)] [[INSPIRE](#)].

- [74] M. Bianchi and E. Kiritsis, *Non-perturbative and Flux superpotentials for Type I strings on the Z_3 orbifold*, *Nucl. Phys. B* **782** (2007) 26 [[hep-th/0702015](#)] [[INSPIRE](#)].
- [75] M. Bianchi, F. Fucito and J.F. Morales, *D-brane instantons on the T^6/Z_3 orientifold*, *JHEP* **07** (2007) 038 [[arXiv:0704.0784](#)] [[INSPIRE](#)].
- [76] M. Bianchi, F. Fucito and J.F. Morales, *Dynamical supersymmetry breaking from unoriented D-brane instantons*, *JHEP* **08** (2009) 040 [[arXiv:0904.2156](#)] [[INSPIRE](#)].
- [77] A. Addazi and M. Bianchi, *Neutron Majorana mass from exotic instantons*, *JHEP* **12** (2014) 089 [[arXiv:1407.2897](#)] [[INSPIRE](#)].
- [78] A. Addazi and M. Bianchi, *Un-oriented Quiver Theories for Majorana Neutrons*, [arXiv:1502.01531](#) [[INSPIRE](#)].
- [79] A. Addazi, *‘Exotic vector-like pair’ of color-triplet scalars*, *JHEP* **04** (2015) 153 [[arXiv:1501.04660](#)] [[INSPIRE](#)].
- [80] CMS collaboration, *Search for heavy neutrinos and W bosons with right-handed couplings in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *Eur. Phys. J. C* **74** (2014) 3149 [[arXiv:1407.3683](#)] [[INSPIRE](#)].
- [81] W.-Y. Keung and G. Senjanović, *Majorana Neutrinos and the Production of the Right-handed Charged Gauge Boson*, *Phys. Rev. Lett.* **50** (1983) 1427 [[INSPIRE](#)].
- [82] A. Ferrari et al., *Sensitivity study for new gauge bosons and right-handed Majorana neutrinos in pp collisions at $\sqrt{s} = 14$ TeV*, *Phys. Rev. D* **62** (2000) 013001 [[INSPIRE](#)].
- [83] S.N. Gninenko, M.M. Kirsanov, N.V. Krasnikov and V.A. Matveev, *Detection of heavy Majorana neutrinos and right-handed bosons*, *Phys. Atom. Nucl.* **70** (2007) 441 [[INSPIRE](#)].
- [84] F.F. Deppisch, T.E. Gonzalo, S. Patra, N. Sahu and U. Sarkar, *Signal of Right-Handed Charged Gauge Bosons at the LHC?*, *Phys. Rev. D* **90** (2014) 053014 [[arXiv:1407.5384](#)] [[INSPIRE](#)].
- [85] Y. Chikashige, R.N. Mohapatra and R.D. Peccei, *Are There Real Goldstone Bosons Associated with Broken Lepton Number?*, *Phys. Lett. B* **98** (1981) 265 [[INSPIRE](#)].
- [86] PARTICLE DATA GROUP collaboration, J. Beringer et al., *Review of Particle Physics (RPP)*, *Phys. Rev. D* **86** (2012) 010001 [[INSPIRE](#)].
- [87] M. Agostini et al., *Results on $\beta\beta$ decay with emission of two neutrinos or Majorons in ^{76}Ge from GERDA Phase I*, [arXiv:1501.02345](#) [[INSPIRE](#)].
- [88] EXO-200 collaboration, J. Albert et al., *Search for Majoron-emitting modes of double-beta decay of ^{136}Xe with EXO-200*, *Phys. Rev. D* **90** (2014) 092004 [[arXiv:1409.6829](#)] [[INSPIRE](#)].
- [89] Y. Farzan, *Bounds on the coupling of the Majoron to light neutrinos from supernova cooling*, *Phys. Rev. D* **67** (2003) 073015 [[hep-ph/0211375](#)] [[INSPIRE](#)].
- [90] P. Anastasopoulos, F. Fucito, A. Lionetto, G. Pradisi, A. Racioppi and Y.S. Stanev, *Minimal Anomalous $U(1)'$ Extension of the MSSM*, *Phys. Rev. D* **78** (2008) 085014 [[arXiv:0804.1156](#)] [[INSPIRE](#)].
- [91] C. Corianò, N. Irges and E. Kiritsis, *On the effective theory of low scale orientifold string vacua*, *Nucl. Phys. B* **746** (2006) 77 [[hep-ph/0510332](#)] [[INSPIRE](#)].
- [92] B. de Wit, P.G. Lauwers and A. Van Proeyen, *Lagrangians of $N = 2$ Supergravity-Matter Systems*, *Nucl. Phys. B* **255** (1985) 569 [[INSPIRE](#)].

- [93] A. Addazi and S. Capozziello, *External stability for Spherically Symmetric Solutions in Lorentz Breaking Massive Gravity*, *Int. J. Theor. Phys.* **54** (2015) 1818 [[arXiv:1407.4840](#)] [[INSPIRE](#)].
- [94] X.O. Camanho, J.D. Edelstein, J. Maldacena and A. Zhiboedov, *Causality Constraints on Corrections to the Graviton Three-Point Coupling*, [arXiv:1407.5597](#) [[INSPIRE](#)].
- [95] G. D'Appollonio, P. Di Vecchia, R. Russo and G. Veneziano, *Regge behavior saves String Theory from causality violations*, [arXiv:1502.01254](#) [[INSPIRE](#)].
- [96] M. Bianchi and A.V. Santini, *String predictions for near future colliders from one-loop scattering amplitudes around D-brane worlds*, *JHEP* **12** (2006) 010 [[hep-th/0607224](#)] [[INSPIRE](#)].
- [97] S. Talaganis, T. Biswas and A. Mazumdar, *Towards understanding the ultraviolet behavior of quantum loops in infinite-derivative theories of gravity*, [arXiv:1412.3467](#) [[INSPIRE](#)].
- [98] T. Biswas and N. Okada, *Towards LHC Physics with Non-local Standard Model*, [arXiv:1407.3331](#) [[INSPIRE](#)].
- [99] A. Addazi and G. Esposito, *Nonlocal quantum field theory without acausality and nonunitarity at quantum level: is SUSY the key?*, *Int. J. Mod. Phys. A* **30** (2015) 1550103 [[arXiv:1502.01471](#)] [[INSPIRE](#)].
- [100] S. Jäger and J. Martin Camalich, *Reassessing the discovery potential of the $B \rightarrow K^* \ell^+ \ell^-$ decays in the large-recoil region: SM challenges and BSM opportunities*, [arXiv:1412.3183](#) [[INSPIRE](#)].
- [101] P. Biancofiore, *On the recent anomalies in semileptonic B decays*, *EPJ Web Conf.* **80** (2014) 00048.
- [102] T. Hurth, F. Mahmoudi and S. Neshatpour, *Global fits to $b \rightarrow s \ell \ell$ data and signs for lepton non-universality*, *JHEP* **12** (2014) 053 [[arXiv:1410.4545](#)] [[INSPIRE](#)].
- [103] J. Matias and N. Serra, *Symmetry relations between angular observables in $B^0 \rightarrow K^* \mu^+ \mu^-$ and the LHCb P'_5 anomaly*, *Phys. Rev. D* **90** (2014) 034002 [[arXiv:1402.6855](#)] [[INSPIRE](#)].
- [104] F. Richard, *A Z-prime interpretation of $Bd \rightarrow K^* \mu^+ \mu^-$ data and consequences for high energy colliders*, [arXiv:1312.2467](#) [[INSPIRE](#)].
- [105] A.J. Buras and J. Girrbach, *Left-handed Z' and Z FCNC quark couplings facing new $b \rightarrow s \mu^+ \mu^-$ data*, *JHEP* **12** (2013) 009 [[arXiv:1309.2466](#)] [[INSPIRE](#)].
- [106] R. Gauld, F. Goertz and U. Haisch, *On minimal Z' explanations of the $B \rightarrow K^* \mu^+ \mu^-$ anomaly*, *Phys. Rev. D* **89** (2014) 015005 [[arXiv:1308.1959](#)] [[INSPIRE](#)].