Nandhini and Sujatha *BMC Infectious Diseases* 2014, **14**(Suppl 3):P41 http://www.biomedcentral.com/1471-2334/14/S3/P41



POSTER PRESENTATION

Open Access

Serological response to pandemic influenza A/H1N1 2009 and seasonal influenza A/H3N2 among health care workers (HCWs) in JIPMER, Puducherry

G Nandhini, S Sujatha*

From 2nd International Science Symposium on HIV and Infectious Diseases (HIV SCIENCE 2014) Chennai, India. 30 January - 1 February 2014

Background

Serum antibody titers against influenza viruses are regarded as markers of partial or complete immunoprotection. Antibody titer of ≥40 is associated with at least a 50% reduction in risk for infection or disease.

Methods

Serum was collected from 138 HCWs including laboratory personnel, doctors and nurses working at JIPMER, Puducherry during August - October 2013. Details of influenza vaccination and laboratory confirmed influenza infection were noted. Hemagglutination inhibition assay was performed to determine the serum antibody levels against WHO reference antigens – pandemic influenza A (H1N1) A/California/07/2009, seasonal influenza A (H3N2) A/Victoria/361/2011.

Results

Fifty of the HCWs had received pandemic influenza vaccination at least once in the previous three years. All HCWs had antibody titers ≥40 against seasonal influenza A/H3N2. Antibody titers against pandemic influenza A/H1N1 2009 ranged from ≤20 (16.7%) to 320 with 36.9% showing borderline protective titers (=40). Six of the vaccinated HCWs had non-protective antibody titers while 71 unvaccinated HCWs showed protective titers. Only one HCW developed seasonal influenza A/H3N2 infection despite having borderline protective antibody titer of 40 and his convalescent serum sample after two weeks showed fourfold rise in titers.

* Correspondence: sujathasistla@gmail.com Department of Microbiology, JIPMER, Puducherry, India

Conclusion

This study showed protective antibody levels against pandemic influenza A/H1N1 2009 and seasonal influenza A/H3N2 among a large percentage of HCWs, regardless of vaccination status, which may be the primary reason for the low incidence of influenza cases encountered this year.

Published: 27 May 2014

doi:10.1186/1471-2334-14-S3-P41

Cite this article as: Nandhini and Sujatha: Serological response to pandemic influenza A/H1N1 2009 and seasonal influenza A/H3N2 among health care workers (HCWs) in JIPMER, Puducherry. BMC Infectious Diseases 2014 14(Suppl 3):P41.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



