

REVIEW

Open Access



Moonshine

John FR Duncan¹, Michael J Griffin² and Ken Ono^{2*}

*Correspondence:

ono@mathcs.emory.edu

²Department of Mathematics and
Computer Science, Emory
University, 201 Dowman Drive,
Atlanta, Georgia 30322, USA
Full list of author information is
available at the end of the article**Abstract**

Monstrous moonshine relates distinguished modular functions to the representation theory of the Monster \mathbb{M} . The celebrated observations that

$$1 = 1, \quad 196884 = 1 + 196883, \quad 21493760 = 1 + 196883 + 21296876, \quad \dots \quad (*)$$

illustrate the case of $J(\tau) = j(\tau) - 744$, whose coefficients turn out to be sums of the dimensions of the 194 irreducible representations of \mathbb{M} . Such formulas are dictated by the structure of the graded monstrous moonshine modules. Recent works in moonshine suggest deep relations between number theory and physics. Number theoretic Kloosterman sums have reappeared in quantum gravity, and mock modular forms have emerged as candidates for the computation of black hole degeneracies. This paper is a survey of past and present research on moonshine. We also compute the quantum dimensions of the monster orbifold and obtain exact formulas for the multiplicities of the irreducible components of the moonshine modules. These formulas imply that such multiplicities are asymptotically proportional to dimensions. For example, the proportion of 1's in (*) tends to

$$\frac{\dim(\chi_1)}{\sum_{i=1}^{194} \dim(\chi_i)} = \frac{1}{5844076785304502808013602136} = 1.711 \dots \times 10^{-28}.$$

2010 Mathematics Subject Classification: 11F11; 11F22; 11F37; 11F50; 20C34; 20C35

1 Introduction

This story begins innocently with peculiar numerics, and in its present form exhibits connections to conformal field theory, string theory, quantum gravity, and the arithmetic of mock modular forms. This paper is an introduction to the many facets of this beautiful theory.

We begin with a review of the principal results in the development of monstrous moonshine in Sections 2, 3 and 4. We refer to the introduction of [104], and the more recent survey [116], for more on these topics. After describing these classic works, we discuss the interplay between moonshine and Rademacher sums in Section 5, and related observations which suggest a connection between monstrous moonshine and three-dimensional quantum gravity in Section 6. The remainder of the paper is devoted to more recent works. We describe a generalization of moonshine, the *moonshine tower*, in Section 7, and in Section 8, we compute the quantum dimensions of the monster orbifold and discuss the distributions of irreducible representations of the monster arising

in monstrous moonshine. We present a survey of the recently discovered *umbral moonshine* phenomenon in Section 9, and conclude, in Section 10, with problems for future work.

2 Early days

The classification of finite simple groups [2] distinguishes 26 examples above the others; namely, the *sporadic* simple groups, which are those that belong to none of the naturally occurring infinite families: cyclic groups, alternating groups, or finite groups of Lie type. Distinguished amongst the sporadic simple groups is the *Fischer–Griess monster* \mathbb{M} , on account of its size, which is

$$|\mathbb{M}| = 2^{46} \cdot 3^{20} \cdot 5^9 \cdot 7^6 \cdot 11^2 \cdot 13^3 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71 \tag{2.1}$$

(cf. [119]). Note that the margin is not small, for the order of the monster is

$$2^5 \cdot 3^7 \cdot 5^3 \cdot 7^4 \cdot 11 \cdot 13^2 \cdot 29 \cdot 41 \cdot 59 \cdot 71 \tag{2.2}$$

times that of the next largest sporadic simple group, the baby monster (cf. [169]).

Fischer and Griess independently produced evidence for the monster group in 1973 (cf. [119]). Well before it was proven to exist, Tits gave a lecture on its conjectural properties at the Collège de France in 1975. In particular, he described its order (2.1). Around this time, Ogg had been considering the automorphism groups of certain algebraic curves and had arrived at the set of primes

$$\{2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 41, 47, 59, 71\} \tag{2.3}$$

in a purely geometric way (cf. the Corollaire of [193]). Making what may now be identified as the first observation of monstrous moonshine, Ogg offered a bottle of Jack Daniels^a for an explanation of this coincidence (cf. Remarque 1 of [193]).

Ogg’s observation would ultimately be recognized as reflecting another respect in which the monster is distinguished amongst finite simple groups: as demonstrated by the pioneering construction of Frenkel-Lepowsky-Meurman [102-104], following the astonishing work of Griess [120,121], the ‘most natural’ representation of the monster, is infinite-dimensional.

The explanation of this statement takes us back to McKay’s famous observation, that

$$196884 = 1 + 196883 \tag{2.4}$$

(cf. [58,221]), and the generalizations of this observed by Thompson [221], including^b

$$\begin{aligned} 21493760 &= 1 + 196883 + 21296876, \\ 864299970 &= 2 \times 1 + 2 \times 196883 + 21296876 + 842609326, \\ 20245856256 &= 2 \times 1 + 3 \times 196883 + 2 \times 21296876 + 842609326 + 19360062527. \end{aligned} \tag{2.5}$$

Of course the left-hand sides of (2.4) and (2.5) are familiar to number theorists and algebraic geometers, as coefficients in the Fourier coefficients of the *normalized elliptic modular invariant*

$$\begin{aligned} J(\tau) &:= \frac{1728g_2(\tau)^3}{g_2(\tau)^3 - 27g_3(\tau)^2} - 744 \\ &= q^{-1} + 196884q + 21493760q^2 + 864299970q^3 + 20245856256q^4 + \dots \end{aligned} \tag{2.6}$$

Here $q := e^{2\pi i\tau}$, and we set $g_2(\tau) := 60G_4(\tau)$ and $g_3(\tau) := 140G_6(\tau)$, where $G_{2k}(\tau)$ denotes the Eisenstein series of weight $2k$,

$$G_{2k}(\tau) := \sum_{(m,n) \neq (0,0)} (m + n\tau)^{-2k}, \tag{2.7}$$

for $k \geq 2$. The functions g_2 and g_3 serve to translate between the two most common parameterizations of a complex elliptic curve: as a quotient $\mathbb{C}/(\mathbb{Z} + \mathbb{Z}\tau)$ for τ in the upper-half plane, $\mathbb{H} := \{\tau \in \mathbb{C} \mid \Im(\tau) > 0\}$, and as the locus of a Weierstrass equation, $y^2 = 4x^3 - g_2x - g_3$.

The fundamental property of $J(\tau)$, from both the number theoretic and algebro-geometric points of view, is that it is a modular function for $SL_2(\mathbb{Z})$. In fact, and importantly for the monster’s natural infinite-dimensional representation, $J(\tau)$ is a generator for the field of $SL_2(\mathbb{Z})$ -invariant holomorphic functions on \mathbb{H} that have at most exponential growth as $\Im(\tau) \rightarrow \infty$.

The right-hand sides of (2.4) and (2.5) are familiar to finite group theorists, as simple sums of dimensions of irreducible representations of the monster \mathbb{M} . In fact, the irreducible representations appearing in (2.4) and (2.5) are just the first six, of a total of 194, in the character table of \mathbb{M} (cf. [56]), when ordered by size. We have that

$$\begin{aligned} \chi_1(e) &= 1 \\ \chi_2(e) &= 196883 \\ \chi_3(e) &= 21296876 \\ \chi_4(e) &= 842609326 \\ \chi_5(e) &= 18538750076 \\ \chi_6(e) &= 19360062527 \\ &\vdots \\ \chi_{194}(e) &= 258823477531055064045234375. \end{aligned} \tag{2.8}$$

Here e denotes the identity element of \mathbb{M} , so $\chi_i(e)$ is just the dimension of the irreducible representation of \mathbb{M} with character χ_i .

3 Classical moonshine

The coincidences (2.4) and (2.5) led Thompson to make the following conjecture [221] which realizes the natural representation of the monster alluded to above.

Conjecture 1 (Thompson). There is a naturally defined graded infinite-dimensional monster module, denoted $V^\natural = \bigoplus_{n=-1}^\infty V_n^\natural$, which satisfies

$$\dim(V_n^\natural) = c(n) \tag{3.1}$$

for $n \geq -1$ (cf. (2.6)), such that the decompositions into irreducible representations of the monster satisfy (2.4) and (2.5) for $n = 1, 2, 3$ and 4 (and a similar condition for $n = 5$).

At the time that Thompson’s conjecture was made, the monster had not yet been proven to exist, but Griess [119], and Conway–Norton [58], had independently conjectured

the existence of a faithful representation of dimension 196883, and Fischer-Livingstone-Thorne had constructed the character table of \mathbb{M} , by assuming the validity of this claim (cf. [58]) together with conjectural statements (cf. [119]) about the structure of \mathbb{M} .

Thompson also suggested [220] to investigate the properties of the graded-trace functions

$$T_g(\tau) := \sum_{n=-1}^{\infty} \text{tr}(g|V_n^{\natural})q^n, \tag{3.2}$$

for $g \in \mathbb{M}$, now called the *monstrous McKay-Thompson series*, that would arise from the conjectural monster module V^{\natural} . Using the character table constructed by Fischer-Livingstone-Thorne, it was observed [58,220] that the functions T_g are in many cases directly similar to J in the following respect: the first few coefficients of each T_g coincide with those of a generator for the function field of a discrete group^c $\Gamma_g < \text{SL}_2(\mathbb{R})$, commensurable with $\text{SL}_2(\mathbb{Z})$, containing $-I$, and having *width one at infinity*, meaning that the subgroup of upper-triangular matrices in Γ_g coincides with

$$\Gamma_{\infty} := \left\{ \pm \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \mid n \in \mathbb{Z} \right\}, \tag{3.3}$$

for all $g \in \mathbb{M}$.

This observation was refined and developed by Conway-Norton [58], leading to their famous *monstrous moonshine conjectures*:

Conjecture 2 (Monstrous moonshine: Conway-Norton). For each $g \in \mathbb{M}$ there is a specific group $\Gamma_g < \text{SL}_2(\mathbb{R})$ such that T_g is the unique *normalized principal modulus*^d for Γ_g .

This means that each T_g is the unique Γ_g -invariant holomorphic function on \mathbb{H} which satisfies

$$T_g(\tau) = q^{-1} + O(q), \tag{3.4}$$

as $\Im(\tau) \rightarrow \infty$, and remains bounded as τ approaches any non-infinite cusp of Γ_g . We refer to this feature of the T_g as the *principal modulus property* of monstrous moonshine.

The hypothesis that T_g is Γ_g -invariant, satisfying (3.4) near the infinite cusp of Γ_g but having no other poles, implies that T_g generates the field of Γ_g -invariant holomorphic functions on \mathbb{H} that have at most exponential growth at cusps, in direct analogy with J . In particular, the natural Riemann surface structure on $\Gamma_g \backslash \mathbb{H}$ (cf., e.g., [211]) must be that of the Riemann sphere $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ with finitely many points removed, and for this reason, the groups Γ_g are said to have *genus zero*, and the principal modulus property is often referred to as the *genus zero property* of monstrous moonshine.

The reader will note the astonishing predictive power that the principal modulus property of monstrous moonshine bestows: the fact that a normalized principal modulus for a genus zero group Γ_g is unique, means that we can compute the trace of an element $g \in \mathbb{M}$, on any homogeneous subspace of the monster’s natural infinite-dimensional representation V^{\natural} , without any information about the monster, as soon as we can guess correctly the discrete group Γ_g . The analysis of Conway-Norton in [58] establishes very strong guidelines for the determination of Γ_g , and once Γ_g has been chosen, the ‘theory of replicability’ (cf. [1,58,191]) allows for efficient computation of the coefficients of the

normalized principal modulus T_g , given the knowledge of just a few of them (cf. [99], or (3.10)).

It was verified by Atkin-Fong-Smith [212], using results of Thompson [220] (cf. also [207]), that a graded (possibly virtual) infinite-dimensional monster module V^\natural , such that the functions T_g of (3.2) are exactly those predicted by Conway-Norton in [58], exists.

Theorem 1 (Atkin-Fong-Smith). *There exists a (possibly virtual) graded \mathbb{M} -module $V^\natural = \bigoplus_{n=-1}^{\infty} V_n^\natural$ such that if T_g is defined by (3.2), then T_g is the Fourier expansion of the unique Γ_g -invariant holomorphic function on \mathbb{H} that satisfies $T_g(\tau) = q^{-1} + O(q)$ as τ approaches the infinite cusp, and has no poles at any non-infinite cusps of Γ_g , where Γ_g is the discrete subgroup of $SL_2(\mathbb{R})$ specified by Conway-Norton in [58].*

Thus Thompson’s conjecture was confirmed, albeit indirectly. By this point in time, Griess, in an astonishing tour de force, had constructed the monster explicitly, by hand, by realizing it as the automorphism group of a commutative but non-associative algebra of dimension 196884 [120,121] (See also [55,222]). Inspired by Griess’ construction, and by the representation theory of affine Lie algebras, which also involves graded infinite-dimensional vector spaces whose graded dimensions enjoy good modular properties (cf., e.g., [148-150,153]), Frenkel-Lepowsky-Meurman established Thompson’s conjecture in a strong sense.

Theorem 2 (Frenkel-Lepowsky-Meurman). *Thompson’s Conjecture is true. In particular, the moonshine module V^\natural is constructed in [102,103].*

Frenkel-Lepowsky-Meurman generalized the homogeneous realization of the basic representation of an affine Lie algebra $\hat{\mathfrak{g}}$ due, independently, to Frenkel-Kac [101] and Segal [208], in such a way that *Leech’s lattice* Λ [167,168] - the unique [53] even self-dual positive-definite lattice of rank 24 with no roots - could take on the role played by the root lattice of \mathfrak{g} in the Lie algebra case. In particular, their construction came equipped with rich algebraic structure, furnished by vertex operators, which had appeared first in the physics literature in the late 1960’s.

We refer to [101], and also the introduction to [104] for accounts of the role played by vertex operators in physics (up to 1988) along with a detailed description of their application to the representation theory of affine Lie algebras. The first application of vertex operators to affine Lie algebra representations was obtained by Lepowsky-Wilson in [171].

Borcherds described a powerful axiomatic formalism for vertex operators in [15]. In particular, he introduced the notion of a *vertex algebra*, which can be regarded as similar to a commutative associative algebra, except that multiplications depend upon formal variables z_i , and can be singular, in a certain formal sense, along the canonical divisors $\{z_i = 0\}$, $\{z_i = z_j\}$ (cf. [20,100]).

The appearance of affine Lie algebras above, as a conceptual ingredient for the Frenkel-Lepowsky-Meurman construction of V^\natural hints at an analogy between complex Lie groups and the monster. Borcherds’ vertex algebra theory makes this concrete, for Borcherds showed [15] that both in the case of the basic representation of an affine Lie algebra, and in the case of the moonshine module V^\natural , the vertex operators defined by Frenkel-Kac, Segal, and Frenkel-Lepowsky-Meurmann, extend naturally to vertex algebra structures.

In all of these examples the *Virasoro algebra*, $\mathcal{V} = \bigoplus_n \mathbb{C}L(n) \oplus \mathbb{C}\mathbf{c}$, being the unique universal central extension of the Lie algebra $\mathbb{C}[t, t^{-1}] \frac{d}{dt}$ of polynomial vector fields on the circle,

$$[L(m), L(n)] = (m - n)L(m + n) + \frac{m^3 - m}{12} \delta_{m+n,0} \mathbf{c}, \quad [L(m), \mathbf{c}] = 0, \tag{3.5}$$

acts naturally on the underlying vector space. (See [154] for a detailed analysis of \mathcal{V} . The generator $L(m)$ lies above the vector field $-t^{m+1} \frac{d}{dt}$.) This Virasoro structure, which has powerful applications, was axiomatized in [104], with the introduction of the notion of a *vertex operator algebra*. If V is a vertex operator algebra and the central element \mathbf{c} of the Virasoro algebra acts as c times the identity on V , for some $c \in \mathbb{C}$, then V is said to have *central charge* c .

For the basic representation of an affine Lie algebra $\hat{\mathfrak{g}}$, the group of vertex operator algebra automorphisms - i.e., those vertex algebra automorphisms that commute with the Virasoro action - is the adjoint complex Lie group associated to \mathfrak{g} . For the moonshine module V^\natural , it was shown by Frenkel-Lepowsky-Meurman in [104], that the group of vertex operator algebra automorphisms is precisely the monster.

Theorem 3 (Frenkel-Lepowsky-Meurman). *The moonshine module $V^\natural = \bigoplus_{n=-1}^\infty V_n^\natural$ is a vertex operator algebra of central charge 24 whose graded dimension is given by $J(\tau)$, and whose automorphism group is \mathbb{M} .*

Vertex operator algebras are of relevance to physics, for we now recognize them as ‘chiral halves’ of two-dimensional conformal field theories (cf. [105,106]). From this point of view, the construction of V^\natural by Frenkel-Lepowsky-Meurman constitutes one of the first examples of an orbifold conformal field theory (cf. [72-74]). In the case of V^\natural , the underlying geometric orbifold is the quotient

$$(\mathbb{R}^{24}/\Lambda) / (\mathbb{Z}/2\mathbb{Z}), \tag{3.6}$$

of the 24-dimensional torus $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}/\Lambda \simeq \mathbb{R}^{24}/\Lambda$ by the Kummer involution $x \mapsto -x$, where Λ denotes the Leech lattice. So in a certain sense, V^\natural furnishes a ‘24-dimensional’ construction of \mathbb{M} . We refer to [100,104,152,170] for excellent introductions to vertex algebra, and vertex operator algebra theory.

Affine Lie algebras are special cases of Kac-Moody algebras, first considered by Kac [147] and Moody [184,185], independently. Roughly speaking, a Kac-Moody algebra is ‘built’ from copies of \mathfrak{sl}_2 , in such a way that most examples are infinite-dimensional, but much of the finite-dimensional theory carries through (cf. [151]). Borcherds generalized this further, allowing also copies of the three-dimensional Heisenberg Lie algebra to serve as building blocks, and thus arrived [16] at the notion of *generalized Kac-Moody algebra*, or *Borcherds-Kac-Moody (BKM) algebra*, which has subsequently found many applications in mathematics and mathematical physics (cf. [138,206]).

One of the most powerful such applications occurred in moonshine, when Borcherds introduced a particular example - the *monster Lie algebra* \mathfrak{m} - and used it to prove [18] the moonshine conjectures of Conway-Norton. His method entailed using monster-equivariant versions of the denominator identity for \mathfrak{m} to verify that the coefficients of

the McKay-Thompson series T_g , defined by (3.2) according to the Frenkel-Lepowsky-Meurman construction of V^\natural , satisfy the replication formulas conjectured by Conway-Norton in [58]. This powerful result reduced the proof of the moonshine conjectures to a small, finite number of identities, that he could easily check by hand.

Theorem 4 (Borcherds). *Let V^\natural be the moonshine module vertex operator algebra constructed by Frenkel-Lepowsky-Meurman, whose automorphism group is \mathbb{M} . If T_g is defined by (3.2) for $g \in \mathbb{M}$, and if Γ_g is the discrete subgroup of $SL_2(\mathbb{R})$ specified by Conway-Norton in [58], then T_g is the unique normalized principal modulus for Γ_g .*

Recall that an even self-dual lattice of signature (m, n) exists if and only if $m - n = 0 \pmod{8}$ (cf., e.g., [62]). Such a lattice is unique up to isomorphism if $mn > 0$, and is typically denoted $I_{m,n}$. In the case that $m = n = 1$ we may take

$$I_{1,1} := \mathbb{Z}e + \mathbb{Z}f, \tag{3.7}$$

where e and f are isotropic, $\langle e, e \rangle = \langle f, f \rangle = 0$, and $\langle e, f \rangle = 1$. Then $me + nf \in I_{1,1}$ has square-length $2mn$. Note that $I_{25,1}$ and $\Lambda \oplus I_{1,1}$ are isomorphic, for Λ the Leech lattice, since both lattices are even and self-dual, with signature $(25, 1)$.

In physical terms the monster Lie algebra \mathfrak{m} is ('about half' of) the space of 'physical states' of a bosonic string moving in the quotient

$$(\mathbb{R}^{24}/\Lambda \oplus \mathbb{R}^{1,1}/I_{1,1}) / (\mathbb{Z}/2\mathbb{Z}) \tag{3.8}$$

of the 26-dimensional torus $I_{25,1} \otimes_{\mathbb{Z}} \mathbb{R}/I_{25,1} \simeq \mathbb{R}^{24}/\Lambda \oplus \mathbb{R}^{1,1}/I_{1,1}$ by the involution $(x, y) \mapsto (-x, y)$ (acting as the Kummer involution on the Leech summand, and trivially on the hyperbolic summand). The monster Lie algebra \mathfrak{m} is constructed in a functorial way from V^\natural (cf. [36]), inherits an action by the monster from V^\natural , and admits a monster-invariant grading by $I_{1,1}$.

The denominator identity for a Kac-Moody algebra \mathfrak{g} equates a product indexed by the positive roots of \mathfrak{g} with a sum indexed by the Weyl group of \mathfrak{g} . A BKM algebra also admits a denominator identity, which for the case of the monster Lie algebra \mathfrak{m} is the beautiful *Koike-Norton-Zagier formula*

$$p^{-1} \prod_{\substack{m,n \in \mathbb{Z} \\ m > 0}} (1 - p^m q^n)^{c(mn)} = J(\sigma) - J(\tau), \tag{3.9}$$

where $\sigma \in \mathbb{H}$ and $p = e^{2\pi i \sigma}$ (and $c(n)$ is the coefficient of q^n in $J(\tau)$, cf. (2.6)). Since the right-hand side of (3.9) implies that the left-hand side has no terms $p^m q^n$ with $mn \neq 0$, this identity imposes many non-trivial polynomial relations upon the coefficients of $J(\tau)$. Amongst these is

$$c(4n + 2) = c(2n + 2) + \sum_{k=1}^n c(k)c(2n - k + 1), \tag{3.10}$$

which was first found by Mahler [173] by a different method, along with similar expressions for $c(4n)$, $c(4n + 1)$, and $c(4n + 3)$, which are also entailed in (3.9). Taken together these relations allow us to compute the coefficients of $J(\tau)$ recursively, given just the values

$$\begin{aligned}
 c(1) &= 196884, \\
 c(2) &= 21493760, \\
 c(3) &= 864299970, \\
 c(5) &= 333202640600.
 \end{aligned}
 \tag{3.11}$$

To recover the replication formulas of [58,191], we require to replace J with T_g , and $c(n) = \dim(V_n^\natural)$ with $\text{tr}(g|V_n^\natural)$ in (3.9), and for this we require a categorification of the denominator identity, whereby the positive integers $c(mn)$ are replaced with \mathbb{M} -modules of dimension $c(mn)$.

A categorification of the denominator formula for a finite-dimensional simple complex Lie algebra was obtained by Kostant [165], following an observation of Bott [23]. This was generalized to Kac-Moody algebras by Garland-Lepowsky [118], and generalized further to BKM algebras by Borcherds in [18]. In its most compact form, it is the identity of virtual vector spaces

$$\bigwedge_{-1}(\mathfrak{e}) = H(\mathfrak{e}),
 \tag{3.12}$$

where \mathfrak{e} is the sub Lie algebra of a BKM algebra corresponding to its positive roots (cf. [143,144,151]).

In (3.12), which is a version of the *Euler-Poincaré principle*, we understand $\bigwedge_{-1}(\mathfrak{e})$ to be the specialization of the formal series

$$\bigwedge_t(\mathfrak{e}) := \sum_{k \geq 0} \wedge^k(\mathfrak{e}) t^k
 \tag{3.13}$$

to $t = -1$, where $\wedge^k(\mathfrak{e})$ is the k -th exterior power of \mathfrak{e} , and we write

$$H(\mathfrak{e}) := \sum_{k \geq 0} (-1)^k H_k(\mathfrak{e})
 \tag{3.14}$$

for the alternating sum of the Lie algebra homology groups of \mathfrak{e} .

In the case of the monster Lie algebra \mathfrak{m} , the spaces $\wedge^k(\mathfrak{e})$ and $H_k(\mathfrak{e})$ are graded by $\mathbb{I}_{1,1}$ and acted on naturally by the monster. If we use the variables p and q to keep track of the $\mathbb{I}_{1,1}$ -gradings, then the equality of (3.12) holds in the ring $R(\mathbb{M})[[p, q]][q^{-1}]$ of formal power series in p and q (allowing finitely many negative powers of q), with coefficients in the (integral) representation ring of \mathbb{M} . The so-called no-ghost theorem (cf. Theorem 5.1 in [18]) allows to express the $H_k(\mathfrak{e})$ in terms of the homogenous subspaces of V^\natural , and the identity (3.12) becomes

$$\bigwedge_{-1} \left(\sum_{\substack{m, n \in \mathbb{Z} \\ m > 0}} V_{mn}^\natural p^m q^n \right) = \sum_{m \in \mathbb{Z}} V_m^\natural p^{m+1} - \sum_{n \in \mathbb{Z}} V_n^\natural p q^n.
 \tag{3.15}$$

(It turns out that $H_k(\mathfrak{e}) = 0$ for $k \geq 3$).

The identity (3.15) returns (3.9) once we replace V_k^\natural everywhere with $\dim(V_k^\natural) = c(k)$ and divide both sides by p . More generally, replacing V_k^\natural with $\text{tr}(g|V_k^\natural)$ for $g \in \mathbb{M}$, we deduce from (3.15) that

$$p^{-1} \exp \left(- \sum_{k > 0} \sum_{\substack{m, n \in \mathbb{Z} \\ m > 0}} \frac{1}{k} \text{tr}(g^k | V_{mn}^\natural) p^{mk} q^{nk} \right) = T_g(\sigma) - T_g(\tau)
 \tag{3.16}$$

(cf. [18], and also [146]), which, in turn, implies the replication formulas formulated in [58,191]. Taking $g = e$ in (3.16) we recover (3.9), so (3.16) furnishes a natural, monster-indexed family of analogues of the identity (3.9).

4 Modularity

Despite the power of the BKM algebra theory developed by Borcherds, and despite some conceptual improvements (cf. [64,144,145]) upon Borcherds' original proof of the moonshine conjectures, a conceptual explanation for the principal modulus property of monstrous moonshine is yet to be established. Indeed, there are generalizations and analogues of the notion of replicability which hold for generic modular functions and forms (for example, see [31]), not just those modular functions which are principal moduli.

Zhu explained the modularity of the graded dimension of V^{\natural} in [237], by proving that this is typical for vertex operator algebras satisfying certain natural (but restrictive) hypotheses, and Dong-Li-Mason extended Zhu's work in [79], obtaining modular invariance results for graded trace functions arising from the action of a finite group of automorphisms.

To prepare for a statement of the results of Zhu and Dong-Li-Mason, we mention that the module theory for vertex operators algebras includes so-called *twisted modules*, associated to finite order automorphisms. If g is a finite order automorphism of V , then V is called *g-rational* in case every g -twisted V -module is a direct sum of simple g -twisted V -modules. Dong-Li-Mason proved [78] that a g -rational vertex operator algebra has finitely many simple g -twisted modules up to isomorphism. So in particular, a rational vertex operator algebra has finitely many simple (untwisted) modules.

Given a module M for the vertex operator algebra V , let us write $\dim_* M$ for its graded dimension,

$$\dim_* M := \sum_n \dim(M_n)q^n. \tag{4.1}$$

When the substitution $q = e^{2\pi i\tau}$ in (4.1) yields a locally uniformly convergent series for $\tau \in \mathbb{H}$ write $Z_M(\tau)$ for the resulting holomorphic function on the upper half plane,

$$Z_M(\tau) := \dim_* M|_{q=e^{2\pi i\tau}}. \tag{4.2}$$

Theorem 5 (Zhu, Dong-Li-Mason). *Let V be rational C_2 -cofinite vertex operator algebra. Then the graded dimensions $\dim_* M^i$ of its simple modules M^i define holomorphic functions $Z_{M^i}(\tau)$ which span a finite-dimensional representation of $SL_2(\mathbb{Z})$. More generally, if G is a finite subgroup of $\text{Aut}(V)$ and V is g -rational for every $g \in G$, then the graded trace functions $\sum_n \text{tr}(\tilde{h}|M_n)q^n$, attached to the triples (g, \tilde{h}, M) , where $g, h \in G$ commute, M is a simple h -stable g -twisted module for V , and \tilde{h} is a lift of h to $GL(M)$, span a finite-dimensional representation of $SL_2(\mathbb{Z})$.*

We refer to the Introduction of [79] (see also §2 of [80]) for a discussion of h -stable twisted modules, and the relevant notion of *lift*. Note that any two lifts for h differ only up to multiplication by a non-zero scalar, so $\sum_n \text{tr}(\tilde{h}|M_n)q^n$ is uniquely defined by (g, h, M) , up to a non-zero scalar.

In the case of V^{\natural} , there is a unique simple g -twisted module $V_g^{\natural} = \bigoplus_n \left(V_g^{\natural} \right)_n$ for every $g \in \mathbb{M} = \text{Aut}(V^{\natural})$ (cf. Theorem 1.2 of [78]), and V_g^{\natural} is necessarily h -stable for any $h \in \mathbb{M}$ that commutes with g . Therefore, Theorem 5 suggests that the functions

$$T_{(g,\tilde{h})}(\tau) := \sum_n \text{tr} \left(\tilde{h} \mid \left(V_g^{\natural} \right)_n \right) q^n, \tag{4.3}$$

associated to pairs (g, h) of commuting elements of \mathbb{M} , may be of interest.

Indeed, this was anticipated a decade earlier by Norton, following observations of Conway-Norton [58] and Queen [198], which associated principal moduli to elements of groups that appear as centralizers of cyclic subgroups in the monster. Norton formulated his *generalized moonshine* conjectures in [190] (cf. also [192], and the Appendix to [181]).

Conjecture 3 (Generalized moonshine: Norton). There is an assignment of holomorphic functions $T_{(g,\tilde{h})} : \mathbb{H} \rightarrow \mathbb{C}$ to every pair (g, h) of commuting elements in the monster, such that the following are true:

1. For every $x \in \mathbb{M}$ we have $T_{(x^{-1}gx, x^{-1}\tilde{h}x)} = T_{(g,\tilde{h})}$.
2. For every $\gamma \in \text{SL}_2(\mathbb{Z})$ we have that $T_{(g,\tilde{h})\gamma}(\tau)$ is a scalar multiple of $T_{(g,\tilde{h})}(\gamma\tau)$.
3. The coefficient functions $\tilde{h} \mapsto \text{tr} \left(\tilde{h} \mid \left(V_g^{\natural} \right)_n \right)$, for fixed g and n , define characters of a projective representation of the centralizer of g in \mathbb{M} ,
4. We have that $T_{(g,\tilde{h})}$ is either constant or a generator for the function field of a genus zero group $\Gamma_{(g,h)} < \text{SL}_2(\mathbb{R})$.
5. We have that $T_{(g,\tilde{h})}$ is a scalar multiple of J if and only if $g = h = e$.

Remark. In Conjecture 3 (2) above, the right-action of $\text{SL}_2(\mathbb{Z})$ on commuting pairs of elements of the monster is given by

$$(g, h)\gamma := \left(g^a h^c, g^b h^d \right) \tag{4.4}$$

for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. The (slightly ambiguous) $T_{(g,\tilde{h})\gamma}$ denotes the graded trace of a lift of $g^b h^d$ to $\text{GL} \left(V_{g^a h^c}^{\natural} \right)$. Norton’s generalized moonshine conjectures reduce to the original Conway-Norton moonshine conjectures of [58] when $g = e$.

Conjecture 3 is yet to be proven in full but has been established for a number of special cases. Theorem 5 was used by Dong-Li-Mason in [79], following an observation of Tuite (cf. [76], and [223-225] for broader context), to prove Norton’s conjecture for the case that g and h generate a cyclic subgroup of \mathbb{M} , and this approach, via twisted modules for V^{\natural} , has been extended by Ivanov-Tuite in [141,142]. Höhn obtained a generalization of Borcherds’ method by using a particular twisted module for V^{\natural} to construct a BKM algebra adapted to the case that g is in the class named 2A in [56] - the smaller of the two conjugacy classes of involutions in \mathbb{M} - and in so doing established [135] generalized moonshine for the functions $T_{(g,\tilde{h})}$ with $g \in 2A$. So far the most general results in generalized moonshine have been obtained by Carnahan [34-36] (see [37] for a recent summary).

Theorem 5 explains why the McKay-Thompson series $T_g(\tau)$ of (3.2), and the $T_{(g,\tilde{h})}(\tau)$ of (4.3) more generally, should be invariant under the actions of (finite index) subgroups

of $SL_2(\mathbb{Z})$, but it does not explain the surprising predictive power of monstrous moonshine. That is, it does not explain why the full invariance groups Γ_g of the T_g should be so large that they admit normalized principal moduli, nor does it explain why the T_g should actually be these normalized principal moduli.

A program to establish a conceptual foundation for the principal modulus property of monstrous moonshine, via *Rademacher sums* and *three-dimensional gravity*, was initiated in [84] by the first author and Frenkel.

5 Rademacher sums

To explain the conjectural connection between gravity and moonshine, we first recall some history. The roots of the approach of [84] extend back almost a hundred years, to Einstein’s theory of general relativity, formulated in 1915, and the introduction of the circle method in analytic number theory, by Hardy-Ramanujan [131]. At the same time that pre-war efforts to quantize Einstein’s theory of gravity were gaining steam (see [213] for a review), the circle method was being refined and developed, by Hardy-Littlewood (cf. [130]), and Rademacher [199], amongst others. (See [226] for a detailed account of what is now known as the *Hardy-Littlewood circle method*.) Despite being contemporaneous, these works were unrelated in science until this century: as we will explain presently, Rademacher’s analysis led to a Poincaré series-like expression - the prototypical Rademacher sum - for the elliptic modular invariant $J(\tau)$. It was suggested first in [70] (see also [180]) that this kind of expression might be useful for the computation of partition functions in quantum gravity.

Rademacher ‘perfected’ the circle method introduced by Hardy-Ramanujan, and he obtained an exact convergent series expression for the combinatorial partition function $p(n)$. In 1938 he generalized this work [200] and obtained such exact formulas for the Fourier coefficients of general modular functions. For the elliptic modular invariant $J(\tau) = \sum_n c(n)q^n$ (cf. (2.6)), Rademacher’s formula (which was obtained earlier by Petersson [197], via a different method) may be written as

$$c(n) = 4\pi^2 \sum_{c>0} \sum_{\substack{0<a<c \\ (a,c)=1}} \frac{e^{-2\pi i \frac{a}{c}} e^{2\pi i n \frac{d}{c}}}{c^2} \sum_{k \geq 0} \frac{(4\pi^2)^k}{c^{2k}} \frac{1}{(k+1)!} \frac{n^k}{k!}, \tag{5.1}$$

where d , in each summand, is a multiplicative inverse for a modulo c , and (a, c) is the greatest common divisor of a and c . Having established the formula (5.1), Rademacher sought to reverse the process, and use it to derive the modular invariance of $J(\tau)$. That is, he set out to prove directly that $J_0(\tau + 1) = J_0(-1/\tau) = J_0(\tau)$, when $J_0(\tau)$ is defined by setting $J_0(\tau) = q^{-1} + \sum_{n>0} c(n)q^n$, with $c(n)$ defined by (5.1).

Rademacher achieved this goal in [201], by reorganizing the summation

$$\sum_{n>0} c(n)q^n = 4\pi^2 \sum_{n>0} \sum_{c>0} \sum_{\substack{0<a<c \\ (a,c)=1}} \frac{e^{-2\pi i \frac{a}{c}} e^{2\pi i n(\tau + \frac{d}{c})}}{c^2} \sum_{k \geq 0} \frac{(4\pi^2)^k}{c^{2k}} \frac{1}{(k+1)!} \frac{n^k}{k!} \tag{5.2}$$

into a Poincaré series-like expression for J . More precisely, Rademacher proved that

$$J(\tau) + 12 = e^{-2\pi i \tau} + \lim_{K \rightarrow \infty} \sum_{\substack{0 < c < K \\ -K^2 < d < K^2 \\ (c,d)=1}} e^{-2\pi i \frac{a\tau + b}{c\tau + d}} - e^{-2\pi i \frac{a}{c}}, \tag{5.3}$$

where $a, b \in \mathbb{Z}$ are chosen arbitrarily, in each summand, so that $ad - bc = 1$. We call the right-hand side of (5.3) the first *Rademacher sum*.

Rademacher’s expression (5.3) for the elliptic modular invariant J is to be compared to the formal sum

$$\sum_{\substack{c,d \in \mathbb{Z} \\ (c,d)=1}} e^{-2\pi i m \frac{a\tau+b}{c\tau+d}}, \tag{5.4}$$

for m a positive integer, which we may regard as a (formal) Poincaré series of weight zero for $SL_2(\mathbb{Z})$. In particular, (5.4) is (formally) invariant for the action of $SL_2(\mathbb{Z})$, as we see by recognizing the matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ as representatives for the right coset space $\Gamma_\infty \backslash \Gamma$, where $\Gamma = SL_2(\mathbb{Z})$ and Γ_∞ is defined in (3.3): for a fixed bottom row (c, d) of matrices in $SL_2(\mathbb{Z})$, any two choices for the top row (a, b) are related by left-multiplication by some element of Γ_∞ .

The formal sum (5.4) does not converge for any $\tau \in \mathbb{H}$, so a regularization procedure is required. Rademacher’s sum (5.3) achieves this, for $m = 1$, by constraining the order of summation, and subtracting the limit as $\Im(\tau) \rightarrow \infty$ of each summand $e^{-2\pi i \frac{a\tau+b}{c\tau+d}}$, whenever this limit makes sense. Rademacher’s method has by now been generalized in various ways by a number of authors. The earliest generalizations are due to Knopp [157-160], and a very general negative weight version of the Rademacher construction was given by Niebur in [188]. We refer to [45] for a detailed review and further references. A nice account of the original approach of Rademacher appears in [156].

We note here that one of the main difficulties in establishing formulas like (5.3) is the demonstration of convergence. When the weight w of the Rademacher sum under consideration lies in the range $0 \leq w \leq 2$, then one requires non-trivial estimates on *sums of Kloosterman sums*, like

$$\sum_{c>0} \sum_{\substack{0<a<c \\ (a,c)=1}} \frac{e^{-2\pi i m \frac{a}{c}} e^{2\pi i n \frac{d}{c}}}{c^2} \tag{5.5}$$

(for the case that $w = 0$ or $w = 2$). The demonstration of convergence generally becomes more delicate as w approaches 1.

In [84] the convergence of a weight zero Rademacher sum $R_\Gamma^{(-m)}(\tau)$ is shown, for m a positive integer and Γ an arbitrary subgroup of $SL_2(\mathbb{R})$ that is commensurable with $SL_2(\mathbb{Z})$. Assuming that Γ contains $-I$ and has width one at infinity (cf. (3.3)), we have

$$R_\Gamma^{(-m)}(\tau) = e^{-2\pi i m \tau} + \lim_{K \rightarrow \infty} \sum_{(\Gamma_\infty \backslash \Gamma)_{<K}^\times} e^{-2\pi i m \frac{a\tau+b}{c\tau+d}} - e^{-2\pi i m \frac{a}{c}}, \tag{5.6}$$

where the summation, for fixed K , is over non-trivial right cosets of Γ_∞ in Γ (cf. (3.3)), having representatives $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ such that $0 < c < K$ and $|d| < K^2$.

The modular properties of the $R_\Gamma^{(-m)}$ are also considered in [84], and it is at this point that the significance of Rademacher sums in monstrous moonshine appears. To state the relevant result, we give the natural generalization (cf. §3.2 of [84]) of the Rademacher-Petersson formula (5.1) for $c(n)$, which is

$$c_\Gamma(-m, n) = 4\pi^2 \lim_{K \rightarrow \infty} \sum_{(\Gamma_\infty \backslash \Gamma / \Gamma_\infty)_{<K}^\times} \frac{e^{-2\pi i m \frac{a}{c}} e^{2\pi i n \frac{d}{c}}}{c^2} \sum_{k \geq 0} \frac{(4\pi^2)^k}{c^{2k}} \frac{m^{k+1}}{(k+1)!} \frac{n^k}{k!}, \tag{5.7}$$

where the summation, for fixed K , is over non-trivial double cosets of Γ_∞ in Γ (cf. (3.3)), having representatives $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ such that $0 < c < K$. Note that this formula simplifies for $n = 0$, to

$$c_\Gamma(-m, 0) = 4\pi^2 m \lim_{K \rightarrow \infty} \sum_{(\Gamma_\infty \backslash \Gamma / \Gamma_\infty)_{<K}^\times} \frac{e^{-2\pi i m \frac{a}{c}}}{c^2}. \tag{5.8}$$

The value $c_\Gamma(-1, 0)$ is the *Rademacher constant* attached to Γ (cf. §6 of [191] and §5.1 of [84]).

A *normalized Rademacher sum* $T_\Gamma^{(-m)}(\tau)$ is defined in §4.1 of [84] by introducing an extra complex variable and taking a limit. It is shown in §4.4 of [84] that

$$T_\Gamma^{(-m)}(\tau) = R_\Gamma^{(-m)}(\tau) - \frac{1}{2}c_\Gamma(-m, 0) \tag{5.9}$$

for any group $\Gamma < SL_2(\mathbb{R})$ that is commensurable with $SL_2(\mathbb{Z})$. If Γ has width one at infinity (cf. (3.3)), then also

$$T_\Gamma^{(-m)}(\tau) = q^{-m} + \sum_{n>0} c_\Gamma(-m, n)q^n, \tag{5.10}$$

so in particular, $T_\Gamma^{(-m)}(\tau) = q^{-m} + O(q)$ as $\Im(\tau) \rightarrow \infty$. The following theorem by the first author and Frenkel summarizes the central role of Rademacher sums in the principal modulus property.

Theorem 6 (Duncan-Frenkel [84]). *Let Γ be a subgroup of $SL_2(\mathbb{R})$ that is commensurable with $SL_2(\mathbb{Z})$. Then the normalized Rademacher sum $T_\Gamma^{(-m)}$ is Γ -invariant if and only if Γ has genus zero. Furthermore, if Γ has genus zero then $T_\Gamma^{(-1)}$ is the normalized principal modulus for Γ .*

In the case that the normalized Rademacher sum $T_\Gamma^{(-1)}$ is not Γ -invariant, $T_\Gamma^{(-m)}$ is an *abelian integral of the second kind* for Γ , in the sense that it has at most exponential growth at the cusps of Γ , and satisfies $T_\Gamma^{(-m)}(\gamma\tau) = T_\Gamma^{(-m)}(\tau) + \omega(\gamma)$ for $\gamma \in \Gamma$, for some function $\omega : \Gamma \rightarrow \mathbb{C}$ (depending on m).

Theorem 6 is used as a basis for the formulation of a characterization of the discrete groups Γ_g of monstrous moonshine in terms of Rademacher sums in §6.5 of [84], following earlier work [57] of Conway-McKay-Sebbar. It also facilitates a proof of the following result, which constitutes a uniform construction of the McKay-Thompson series of monstrous moonshine.

Theorem 7 (Duncan-Frenkel [84]). *Let $g \in \mathbb{M}$. Then the McKay-Thompson series T_g coincides with the normalized Rademacher sum $T_{\Gamma_g}^{(-1)}$.*

Proof. Theorem 4 states that T_g is a normalized principal modulus for Γ_g , and in particular, all the Γ_g have genus zero. Given this, it follows from Theorem 6 that $T_{\Gamma_g}^{(-1)}$ is also a normalized principal modulus for Γ_g . A normalized principal modulus is unique if it exists, so we conclude $T_g = T_{\Gamma_g}^{(-1)}$ for all $g \in \mathbb{M}$, as we required to show. \square

Perhaps most importantly, Theorem 6 is an indication of how the principal modulus property of monstrous moonshine can be explained conceptually. For if we can develop a

mathematical theory in which the underlying objects are graded with graded traces that are provably

1. modular invariant, for subgroups of $SL_2(\mathbb{R})$ that are commensurable with $SL_2(\mathbb{Z})$, and
2. given explicitly by Rademacher sums, such as (5.6),

then these graded trace functions are necessarily normalized principal moduli, according to Theorem 6.

We are now led to ask: what kind of mathematical theory can support such results? As we have alluded to above, Rademacher sums have been related to quantum gravity by articles in the physics literature. Also, a possible connection between the monster and three-dimensional quantum gravity was discussed in [232]. This suggests the possibility that three-dimensional quantum gravity and moonshine are related via Rademacher sums, and was a strong motivation for the work [84]. In the next section, we will give a brief review of quantum gravity, since it is an important area of physical inquiry which has played a role in the development of moonshine, but we must first warn the reader: problems have been identified with the existing conjectures that relate the monster to gravity, and the current status of this connection is uncertain.

6 Quantum gravity

Witten was the first to predict a role for the monster in quantum gravity. In [232] Witten considered *pure quantum gravity* in three dimensions with negative cosmological constant and presented evidence that the moonshine module V^{\natural} is a chiral half of the conformal field theory dual to such a quantum gravity theory, at the most negative possible value of the cosmological constant.

To explain some of the content of this statement, note that the action in pure three-dimensional quantum gravity is given explicitly by

$$I_{EG} := \frac{1}{16\pi G} \int d^3x \sqrt{-g} (R - 2\Lambda), \tag{6.1}$$

where G is the *Newton* or *gravitational constant*, R denotes the Ricci scalar, and the *cosmological constant* is the scalar denoted by Λ .

The case that the cosmological constant Λ is negative is distinguished, since then there exist black hole solutions to the action (6.1), as was discovered by Bañados-Teitelboim-Zanelli [9]. These black hole solutions - the *BTZ black holes* - are locally isomorphic to three-dimensional anti-de Sitter space [8], which is a Lorentzian analogue of hyperbolic space, and may be realized explicitly as the universal cover of a hyperboloid

$$-X_{-1}^2 - X_0^2 + X_1^2 + X_2^2 + X_3^2 = -\ell^2 \tag{6.2}$$

in $\mathbb{R}^{2,3}$ (cf., e.g., [174]). The parameter ℓ in (6.2) is called the *radius of curvature*. For a locally anti-de Sitter (AdS) solution to (6.1), the radius of curvature is determined by the cosmological constant, according to

$$\ell^2 = -1/\Lambda. \tag{6.3}$$

In what has become the most cited^e paper in the history of high energy physics, Maldacena opened the door on a new and powerful approach to quantum gravity in [175], by presenting evidence for a gauge/gravity duality, in which gauge theories serve as duals

to gravity theories in one dimension higher. (See [174] for a recent review.) In the simplest examples, the gauge theories are conformal field theories, and the gravity theories involve locally AdS spacetimes. The gauge/gravity duality for these cases is now known as the *AdS/CFT correspondence*.

Maldacena’s duality furnishes a concrete realization of the *holographic principle*, introduced by ‘t Hooft [216], and elaborated on by Susskind [215]. For following refinements to Maldacena’s proposal due to Gubser-Klebanov-Polyakov [126], and Witten [231], it is expected that gravity theories with $(d + 1)$ -dimensional locally AdS spacetimes can be understood through the analysis of d -dimensional conformal field theories defined on the boundaries of these AdS spaces. Thus in the case of AdS solutions to three-dimensional quantum gravity, a governing role may be played by two-dimensional conformal theories, which can be accessed mathematically via vertex operator algebras (as we have mentioned in Section 3).

The conjecture of [232] is that the two-dimensional conformal field theory corresponding to a tensor product of two copies of the moonshine module V^{\natural} (one ‘left-moving,’ the other ‘right-moving’) is the holographic dual to pure three-dimensional quantum gravity with $\ell = 16G$, and therefore

$$\Lambda = -\frac{1}{256G^2}. \tag{6.4}$$

It is also argued that the only physically consistent values of ℓ are $\ell = 16Gm$, for m a positive integer, so that (6.4) is the most negative possible value for Λ , by force of (6.3).

Shortly after this conjecture was formulated, problems with the quantum mechanical interpretation were identified by Maloney-Witten in [178]. Moreover, Gaiotto [114] and Höhn [137] cast doubt on the relevance of the monster to gravity by demonstrating that it cannot act on a holographically dual conformal field theory corresponding to $\ell = 32G$ (i.e., $m = 2$), at least under the hypotheses (namely, an extremality condition, and holomorphic factorization) presented in [232].

Interestingly, the physical problems with the analysis of [232] seem to disappear in the context of *chiral three-dimensional gravity*, which was introduced and discussed in detail by Li-Song-Strominger in [172] (cf. also [177,214]). This is the gravity theory which motivates much of the discussion in §7 of [84].

In order to define chiral three-dimensional gravity, we first describe *topologically massive gravity*, which was introduced in 1982 by Deser-Jackiw-Templeton [68,69] (see also [67]). The action for topologically massive gravity is given by

$$I_{\text{TMG}} := I_{\text{EG}} + I_{\text{CSG}}, \tag{6.5}$$

where I_{EG} is the Einstein-Hilbert action (cf. (6.1)) of pure quantum gravity, and I_{CSG} denotes the *gravitational Chern-Simons term*

$$I_{\text{CSG}} := \frac{1}{32\pi G\mu} \int d^3x \sqrt{-g} \epsilon^{\lambda\mu\nu} \Gamma_{\lambda\sigma}^{\rho} \left(\partial_{\mu} \Gamma_{\rho\nu}^{\sigma} + \frac{2}{3} \Gamma_{\mu\tau}^{\sigma} \Gamma_{\nu\rho}^{\tau} \right). \tag{6.6}$$

The Γ_{**}^* are Christoffel symbols, and the parameter μ is called the *Chern-Simons coupling constant*.

Chiral three-dimensional gravity is the special case of topologically massive gravity in which the Chern-Simons coupling constant is set to $\mu = 1/\ell = \sqrt{-\Lambda}$. It is shown in [172] that at this special value of μ , the left-moving central charges of the boundary conformal field theories vanish, and the right-moving central charges are

$$c = \frac{3\ell}{2G} = 24m, \tag{6.7}$$

for m a positive integer, $\ell = 16Gm$.

Much of the analysis of [232] still applies in this setting, and the natural analogue of the conjecture mentioned above states that V^\natural is holographically dual to chiral three-dimensional quantum gravity at $\ell = 16G$, i.e., $m = 1$. However, as argued in detail in [177], the problem of quantizing chiral three-dimensional gravity may be regarded as equivalent to the problem of constructing a sequence of extremal chiral two-dimensional conformal field theories (i.e., vertex operator algebras), one for each central charge $c = 24m$, for m a positive integer. Here, a vertex operator algebra $V = \bigoplus_n V_n$ with central charge $c = 24m$ is called *extremal*, if its graded^f dimension function satisfies

$$\dim_* V = q^{-m} \frac{1}{\prod_{n>1} (1 - q^n)} + O(q) \tag{6.8}$$

(cf. (4.1)). The moonshine module is the natural candidate for $m = 1$ (indeed, it is the only candidate if we assume the uniqueness conjecture of [104]), as the right-hand side of (6.8) reduces to $q^{-1} + O(q)$ in this case, but the analysis of [114,137] also applies here, indicating that the monster cannot act non-trivially on any candidate^g for $m = 2$. Thus the role of the monster in quantum gravity is still unclear, even in the more physically promising chiral gravity setting.

Nonetheless, the moonshine module V^\natural may still serve as the holographic dual to chiral three-dimensional quantum gravity at $\ell = 16G$, $m = 1$. In this interpretation, the graded dimension, or *genus one partition function* for V^\natural - namely, the elliptic modular invariant J - serves as the exact spectrum of physical states of chiral three-dimensional gravity at $\mu = \sqrt{-\Lambda} = 1/16G$, in spacetime asymptotic to the three-dimensional anti-de Sitter space (cf. (6.2)).

Recall that if V is a representation of the Virasoro algebra \mathcal{V} (cf. (3.5)), then $v \in V$ is called a *Virasoro highest weight vector* with *highest weight* $h \in \mathbb{C}$ if $L(m)v = h\delta_{m,0}v$ whenever $m \geq 0$. A *Virasoro descendant* is a vector of the form

$$L(m_1) \cdots L(m_k)v, \tag{6.9}$$

where v is a Virasoro highest weight vector, and $m_1 \leq \cdots \leq m_k \leq -1$.

Assuming that V^\natural is dual to chiral three-dimensional gravity at $m = 1$, the Virasoro highest weight vectors in V^\natural define operators that create black holes, and the Virasoro descendants of a highest weight vector describe black holes embellished by boundary excitations. In particular, the 196883-dimensional representation of the monster which is contained in the 196884-dimensional homogenous subspace $V_1^\natural < V^\natural$ (cf. (2.4) and (3.1)), represents an 196883-dimensional space of black hole states in the chiral gravity theory.

More generally, the black hole states in the theory are classified, by the monster, into 194 different kinds, according to which monster irreducible representation they belong to.

Question 1. *Assuming that the moonshine module V^\natural serves as the holographic dual to chiral three-dimensional quantum gravity at $m = 1$, how are the 194 different kinds of black hole states distributed amongst the homogeneous subspaces $V_n^\natural < V^\natural$. Are some kinds of black holes more or less common than others?*

This question will be answered precisely in Section 8.

A positive solution to the conjecture that V^{\natural} is dual to chiral three-dimensional gravity at $m = 1$ may furnish a conceptual explanation for why the graded dimension of V^{\natural} is the normalized principal modulus for $SL_2(\mathbb{Z})$. For on the one hand, modular invariance is a consistency requirement of the physical theory - the genus one partition function is really defined on the moduli space $SL_2(\mathbb{Z}) \backslash \mathbb{H}$ of genus one curves, rather than on \mathbb{H} - and on the other hand, the genus one partition function of chiral three-dimensional gravity is given by a Rademacher sum, as explained by Manschot-Moore [180], following earlier work [70] by Dijkgraaf-Maldacena-Moore-Verlinde. (cf. also [177-179].) So, as we discussed in Section 5, the genus one partition function must be the normalized principal modulus $J(\tau)$ for $SL_2(\mathbb{Z})$, according to Theorem 6.

In the analysis of [179,180], the genus one partition function of chiral three-dimensional gravity is a Rademacher sum (5.6), because it is obtained as a sum over three-dimensional hyperbolic structures on a solid torus with genus one boundary, and such structures are naturally parameterized by the coset space $\Gamma_{\infty} \backslash SL_2(\mathbb{Z})$ (cf. (3.3)), as explained in [176] (see also §5.1 of [70]). The terms $e^{-2\pi im \frac{a\tau+b}{c\tau+d}}$ in (5.6) are obtained by evaluating $e^{-I_{\text{TMG}}}$, with $\mu = \sqrt{-\Lambda} = 1/16Gm$, on a solution with boundary curve $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$, and the subtraction of $e^{-2\pi im \frac{a}{c}}$ represents quantum corrections to the classical action.

In [84], the above conjecture is extended so as to encompass the principal modulus property for all elements of the monster, with a view to establishing a conceptual foundation for monstrous moonshine. More specifically, the first main conjecture of [84] states the following.

Conjecture 4 (Duncan-Frenkel). There exists a monster-indexed family of *twisted chiral three-dimensional gravity* theories, whose genus one partition functions at

$$\mu = \sqrt{-\Lambda} = 1/16G \tag{6.10}$$

are given by $T_{\Gamma_g}^{(-1)}(-1/\tau)$, where $T_{\Gamma_g}^{(-1)}(\tau)$ is the normalized Rademacher sum attached to Γ_g , satisfying (5.9).

From the point of view of vertex operator algebra theory, $T_g(-1/\tau)$ - which coincides with $T_{\Gamma_g}^{(-1)}(-1/\tau)$ according to Theorems 4 and 6 - is the graded dimension of the unique simple g -twisted V^{\natural} -module V_g^{\natural} (cf. Section 4). This non-trivial fact about the functions $T_g(-1/\tau)$ is proven by Carnahan in Theorem 5.1.4 of [36].

Geometrically, the twists of the above conjecture are defined by imposing (generalized) spin structure conditions on solutions to the chiral gravity equations, and allowing orbifold solutions of certain kinds. See §7.1 of [84] for a more complete discussion. The corresponding sums over geometries are then indexed by coset spaces $\Gamma_{\infty} \backslash \Gamma$, for various groups $\Gamma < SL_2(\mathbb{R})$, commensurable with $SL_2(\mathbb{Z})$. According to Theorem 6, the genus one partition function corresponding to such a twist, expected to be a Rademacher sum on physical grounds, will only satisfy the basic physical consistency condition of Γ -invariance if Γ is a genus zero group. One may speculate that a finer analysis of physical consistency will lead to the list of conditions given in §6.5 of [84], which characterize the groups Γ_g for $g \in \mathbb{M}$, according to Theorem 6.5.1 of [84]. Thus the discrete groups Γ_g of monstrous

moonshine may ultimately be recovered as those defining physically consistent twists of chiral three-dimensional gravity.

On the other hand, it is reasonable to expect that twisted chiral gravity theories are determined by symmetries of the underlying untwisted theory. Conceptually then, but still conjecturally, the monster group appears as the symmetry group of the chiral three-dimensional gravity for which the corresponding twists exist. The principal modulus property of monstrous moonshine may then be explained: as a consequence of Theorem 6, together with the statement that the genus one partition function of a twisted theory is $T_{\Gamma}^{(-1)}(-1/\tau)$, where $T_{\Gamma}^{(-1)}(\tau)$ is the normalized Rademacher sum attached to the subgroup $\Gamma < \text{SL}_2(\mathbb{R})$ that parameterizes the geometries of the twist.

Before concluding this section, we mention two further variations on three-dimensional gravity which may ultimately prove relevant to moonshine. The first of these is *log gravity* which was initiated by work [123] of Grumiller-Johansson, and discussed also in [177]. We refer to [125] for a detailed review. The second is *flat space chiral gravity* which was introduced in [7], and is also reviewed in [125].

For more background on the mathematics and physics of black holes, we refer the reader to [124]. We refer to [32,33] for reviews that focus on the particular role of conformal field theory in understanding quantum gravity.

7 Moonshine tower

An optimistic view on the relationship between moonshine and gravity is adopted in §7 of [84]. In particular, in §7.2 of [84] the consequences of Conjecture 4 for the *second quantization* of chiral three-dimensional gravity are explored. (We warn the reader that the notion of second quantized gravity is very speculative at this stage.)

Motivated in part by the results on second quantized string theory in [71], the existence of a tower of monster modules

$$V^{(-m)} = \bigoplus_{n=-m}^{\infty} V_n^{(-m)}, \tag{7.1}$$

parameterized by positive integer values of m , is predicted in §7.2 of [84]. Moreover, it is suggested that the graded dimension of $V^{(-m)}$ should be given by

$$J^{(-m)} := m\hat{T}(m)J, \tag{7.2}$$

where $\hat{T}(m)$ denotes the (*order m*) Hecke operator, acting on $\text{SL}_2(\mathbb{Z})$ -invariant holomorphic functions on \mathbb{H} according to the rule

$$(\hat{T}(m)f)(\tau) := \frac{1}{m} \sum_{\substack{ad=m \\ 0 \leq b < d}} f\left(\frac{a\tau + b}{d}\right). \tag{7.3}$$

Standard calculations (cf., e.g., Chp.VII, §5 of [209]) determine that $m\hat{T}(m)J$ is an $\text{SL}_2(\mathbb{Z})$ -invariant holomorphic function on \mathbb{H} , whose Fourier coefficients

$$J^{(-m)}(\tau) = \sum_n c(-m, n)q^n \tag{7.4}$$

are expressed in terms of those of $J(\tau) = \sum_{n=-1}^{\infty} c(n)q^n$, by $c(-m, n) = \delta_{-m,n}$ for $n \leq 0$, and

$$c(-m, n) = \sum_{\substack{k>0 \\ k|(m,n)}} \frac{m}{k} c(mn/k^2), \tag{7.5}$$

for $n > 0$, where (m, n) denotes the greatest common divisor of m and n . In particular, $J^{(-m)}(\tau) = q^{-m} + O(q)$ as $\Im(\tau) \rightarrow \infty$. There is only one such $SL_2(\mathbb{Z})$ -invariant holomorphic function on \mathbb{H} , so we have

$$J^{(-m)}(\tau) = \sum_{n=-m}^{\infty} \dim(V^{(-m)})q^n = T_{\Gamma}^{(-m)}(\tau) \tag{7.6}$$

according to (5.10) and Theorem 6, when $\Gamma = SL_2(\mathbb{Z})$. So the graded dimension of $V^{(-m)}$ is also a normalized Rademacher sum.

We would like to investigate the higher order analogues of the McKay-Thompson series T_g (cf. (3.2)), encoding the graded traces of monster elements on $V^{(-m)}$, but for this we must first determine the \mathbb{M} -module structure on each homogeneous subspace $V_n^{(-m)}$.

A solution to this problem is entailed in Borcherds' proof [18] of the monstrous moonshine conjectures, and the identity (3.15), in particular. To explain this, recall the *Adams operation* ψ^k on virtual G -modules, defined, for $k \geq 0$ and G a finite group, by requiring that

$$\text{tr}(g|\psi^k(V)) = \text{tr}(g^k|V) \tag{7.7}$$

for $g \in G$. (cf. [6,161] for more details on Adams operations.) Using the ψ^k we may equip $V^{(-m)}$ with a virtual \mathbb{M} -module structure (we will see momentarily that it is actually an \mathbb{M} -module, cf. Proposition 1) by defining $V_{-m}^{(-m)} := \mathbb{C}$ to be the one-dimensional trivial \mathbb{M} -module, $V_n^{(-m)} := 0$ for $-m < n \leq 0$, and

$$V_n^{(-m)} := \bigoplus_{\substack{k>0 \\ k|(m,n)}} \mathbb{C}^{m/k} \otimes \psi^k \left(V_{mn/k^2}^{\natural} \right) \tag{7.8}$$

for $n > 0$, where $\mathbb{C}^{m/k}$ denotes the trivial \mathbb{M} -module of dimension m/k . For convenience later on, we also define $V^{(0)} = V_0^{(0)} := \mathbb{C}$ to be the trivial, one-dimensional \mathbb{M} -module, regarded as graded, with grading concentrated in degree $n = 0$.

Evidently ψ^k preserves dimension, so the graded dimension of $V^{(-m)}$ is still given by $J^{(-m)}$, according to (7.5). Define the *order m McKay-Thompson series* $T_g^{(-m)}$, for $m \geq 0$ and $g \in \mathbb{M}$, by setting

$$T_g^{(-m)}(\tau) := q^{-m} + \sum_{n>0} \text{tr} \left(g|V_n^{(-m)} \right) q^n. \tag{7.9}$$

Then $T_g^{(0)} = 1$ for all $g \in \mathbb{M}$, and $T_g^{(-1)}$ is the original McKay-Thompson series T_g . More generally, we have the following result, which constructs the $T_g^{(-m)}$ uniformly and explicitly as Rademacher sums.

Theorem 8. *For $m > 0$ and $g \in \mathbb{M}$ we have $T_g^{(-m)}(\tau) = T_{\Gamma_g}^{(-m)}(\tau)$, where Γ_g is the invariance group of $T_g(\tau)$, and $T_{\Gamma}^{(-m)}$ denotes the normalized Rademacher sum of order m attached to Γ , as in (5.9). In particular, $T_g^{(-m)}(\tau)$ is a monic integral polynomial of degree m in $T_g(\tau)$.*

Proof. We will use Borcherds’ identity (3.15). To begin, note that $T_g^{(-m)}$ is given explicitly in terms of traces on V^\natural by

$$T_g^{(-m)}(\tau) = q^{-m} + \sum_{n>0} \sum_{k|(m,n)} \frac{m}{k} \text{tr} \left(g | \psi^k \left(V_{mn/k^2}^\natural \right) \right) q^n \tag{7.10}$$

according to (7.8) and (7.9). Recall that $R(G)$ denotes the integral representation ring of a finite group G . Extend the ψ^k from $R(G)$ to $R(G)[[p, q]][[q^{-1}]]$, by setting $\psi^k(Mp^mq^n) = \psi^k(M)p^{km}q^{kn}$ for $M \in R(G)$. Then it is a general property of the Adams operations (cf. §5.2 of [146]) that

$$\log \bigwedge_{-1}(X) = - \sum_{k>0} \frac{1}{k} \psi^k(X) \tag{7.11}$$

in $R(G)[[p, q]][[q^{-1}]] \otimes_{\mathbb{Z}} \mathbb{Q}$, for $X \in R(G)[[p, q]][[q^{-1}]]$. So taking $X = \sum_{\substack{m,n \in \mathbb{Z} \\ m>0}} V_{mn}^\natural p^m q^n$ we obtain

$$\begin{aligned} \log \bigwedge_{-1} \left(\sum_{\substack{m,n \in \mathbb{Z} \\ m>0}} V_{mn}^\natural p^m q^n \right) &= - \sum_{k>0} \sum_{\substack{m,n \in \mathbb{Z} \\ m>0}} \frac{1}{k} \psi^k(V_{mn}^\natural) p^{km} q^{kn} \\ &= - \sum_{\substack{m,n \in \mathbb{Z} \\ m>0}} \sum_{k|(m,n)} \frac{1}{k} \psi^k \left(V_{mn/k^2}^\natural \right) p^m q^n \end{aligned} \tag{7.12}$$

for the logarithm of the left-hand side of (3.15). If we now define $V^{(-m)}(q) := \sum_n V_n^{(-m)} q^n$, an element of $R(\mathbb{M})[[q]][[q^{-1}]]$, then the generating series $\sum_{m>0} p^m V^{(-m)}(q)$ is obtained when we apply $-p\partial_p$ to (7.12), according to the definition (7.8) of the $V_n^{(-m)}$ as elements of $R(\mathbb{M})$. So apply $-p\partial_p \log(\cdot)$ to both sides of (3.15) to obtain the identity

$$\sum_{m>0} V^{(-m)}(q) p^m = -1 - (p\partial_p V^\natural(p)) \sum_{k \geq 0} V^\natural(q)^k V^\natural(p)^{-k-1} \tag{7.13}$$

in $R(\mathbb{M})[[p, q]][[q^{-1}]]$, where $V^\natural(q) = V^{(-1)}(q) = q^{-1} + \sum_{n>0} V_n^\natural q^n$. The right-hand side of (7.13) really is a Taylor series in p , for we use $V^\natural(p)^{-1}$ as a short hand for $\sum_{k \geq 0} (-1)^k p^{k+1} V_+^\natural(p)$, where $V_+^\natural(p) := \sum_{n>0} V_n^\natural p^n$ is the regular part of $V^\natural(p)$.

The McKay-Thompson series $T_g^{(-m)}(\tau)$ is just the trace of g on $V^{(-m)}(q)$, so an application of $\text{tr}(g|\cdot)$ to (7.13) replaces $V^{(-m)}(q)$ with $T_g^{(-m)}(\tau)$, and $V^\natural(q)$ with $T_g(\tau)$, etc. and shows that $T_g^{(-m)}$ is indeed a polynomial in T_g , of degree m since the leading term of $T_g^{(-m)}$ is q^{-m} by definition. In particular, $T_g^{(-m)}$ is a modular function for Γ_g , with no poles away from the infinite cusp. Since Γ_g has genus zero, such a function is uniquely determined (up to an additive constant) by the polar terms in its Fourier expansion. The McKay-Thompson series $T_g^{(-m)}$ and the Rademacher sum $T_{\Gamma_g}^{(-m)}$ both satisfy $q^{-m} + O(q)$ as $\Im(\tau) \rightarrow \infty$ (cf. (5.10)), and neither have poles away from the infinite cusp, so they must coincide. This completes the proof. \square

Remark. The identity obtained by taking the trace of $g \in \mathbb{M}$ on (7.13) may be compactly rewritten

$$\sum_{m \geq 0} T_g^{(-m)}(\tau) p^m = \frac{p\partial_p T_g(\sigma)}{T_g(\tau) - T_g(\sigma)}, \tag{7.14}$$

where $p = e^{2\pi i \sigma}$ and $T_g(\sigma) = \sum_m \text{tr} \left(g | V_m^\natural \right) p^m$. This expression (7.14) is proven for some special cases by a different method in [12].

Recall that the monster group has 194 irreducible ordinary representations, up to equivalence. Let us denote these by M_i , for $1 \leq i \leq 194$, where the ordering is as in [56], so that the character of M_i is the function denoted χ_i in [56]. Define $\mathbf{m}_i(-m, n)$ to be the multiplicity of M_i in $V_n^{(-m)}$, so that

$$V_n^{(-m)} \simeq \bigoplus_{i=1}^{194} M_i^{\oplus \mathbf{m}_i(-m, n)} \tag{7.15}$$

as \mathbb{M} -modules, and $c(-m, n) = \sum_{i=1}^{194} \mathbf{m}_i(-m, n) \chi_i(e)$.

A priori, the \mathbb{M} -modules $V_n^{(-m)}$ may be virtual, meaning that some of the integers $\mathbf{m}_i(-m, n)$ are negative.

Proposition 1. *The $V_n^{(-m)}$ are all (non-virtual) modules for the monster. In particular, the integers $\mathbf{m}_i(-m, n)$ are all non-negative.*

Proof. The claim follows from the modification of Borcherds’ proof of Theorem 4 presented by Jurisich-Lepowsky-Wilson in [146]. In [146] a certain free Lie sub algebra u^- of the monster Lie algebra \mathfrak{m} is identified, for which the identity $\Lambda(u^-) = H(u^-)$ (or rather, the logarithm of this) yields

$$\sum_{m, n > 0} \sum_{k | (m, n)} \frac{1}{k} \psi^k \left(V_{mn/k^2}^\natural \right) p^m q^n = \sum_{k > 0} \frac{1}{k} \left(\sum_{m, n > 0} V_{m+n-1}^\natural p^m q^n \right)^k \tag{7.16}$$

in $R(\mathbb{M})[[p, q]][[q^{-1}]] \otimes \mathbb{Q}$. (Notice the different range of summation, compared to (3.15).) We apply $p\partial_p$ to (7.16), and recall the definition (7.8) of $V_n^{(-m)}$ to obtain

$$\sum_{m, n > 0} V_n^{(-m)} p^m q^n = \sum_{k > 0} \left(\sum_{m, n > 0} m V_{m+n-1}^\natural p^m q^n \right) \left(\sum_{m, n > 0} V_{m+n-1}^\natural p^m q^n \right)^{k-1}. \tag{7.17}$$

The coefficient of $p^m q^n$ in the right-hand side of (7.17) is evidently a non-negative integer combination of the \mathbb{M} -modules V_n^\natural , so the proof of the claim is complete. \square

In Section 8, we will determine the behavior of the multiplicity functions $\mathbf{m}_i(-m, n)$ (cf. (7.15)) as $n \rightarrow \infty$. For applications to gravity, a slightly different statistic is more relevant. Recall from Section 6 that it is the Virasoro highest weight vectors - i.e., those $v \in V_n^\natural$ with $L(k)v = 0$ for $k > 0$ - that represent black hole states in chiral three-dimensional gravity at $m = 1$. Such vectors generate *highest weight modules* for \mathcal{V} , the structure of which has been determined by Feigin-Fuchs in [98]. (See [5] for an alternative treatment). Specializing to the case that the central element \mathbf{c} (cf. (3.5)) acts as $c = 24m$ times the identity, for some positive integer m , we obtain from the results of [98] that the isomorphism type of an irreducible highest weight module for \mathcal{V} depends only on the $L(0)$ -eigenvalue of a generating highest weight vector, v , and if $L(0)v = hv$ for h a non-negative integer, then

$$\dim_* L(h, c) = \begin{cases} q^{-m} (1 - q)(q)_\infty^{-1} & \text{if } h = 0, \\ q^{h-m} (q)_\infty^{-1} & \text{if } h > 0, \end{cases} \tag{7.18}$$

(cf. (4.1)) where $L(h, c)$ denotes the irreducible highest weight \mathcal{V} -module generated by ν , and

$$(q)_\infty := \prod_{n>0} (1 - q^n) \tag{7.19}$$

(Note that all the Virasoro modules in this work are graded by $L(0) - \mathbf{c}/24$. See [129] for details of the calculation that returns (7.18) in the case that $m = 1$.)

Remark. We may now recognize the leading terms in (6.8) as exactly those of the graded dimension of the Virasoro module $L(0, 24m)$.

It is known that V^\natural is a direct sum of highest weight modules for the Virasoro algebra (cf., e.g., [129]). Since the Virasoro and monster actions on V^\natural commute, we have an isomorphism

$$V^\natural \simeq L(0, 24) \otimes W_{-1}^\natural \oplus \bigoplus_{n>0} L(n + 1, 24) \otimes W_n^\natural \tag{7.20}$$

of modules for $\mathcal{V} \times \mathbb{M}$, where W_n^\natural denotes the subspace of V_n^\natural spanned by Virasoro highest weight vectors. To investigate how the black hole states in V^\natural are organized by the representation theory of the monster, we define non-negative integers $\mathbf{n}_i(n)$ by requiring that

$$W_n^\natural \simeq \bigoplus_{i=1}^{194} M_i^{\oplus \mathbf{n}_i(n)}, \tag{7.21}$$

for $n \geq -1$.

Evidently $\mathbf{n}_i(n) \leq \mathbf{m}_i(-1, n)$ for all i and n since W_n^\natural is a subspace of V_n^\natural . To determine the precise relationship between the $\mathbf{n}_i(n)$ and $\mathbf{m}_i(-1, n)$, define $U_g(\tau)$ for $g \in \mathbb{M}$ by setting

$$U_g(\tau) := \sum_{n=-1}^{\infty} \text{tr}(g|W_n^\natural)q^n, \tag{7.22}$$

so that $U_g(\tau) = q^{-1} + \sum_{n>0} \sum_{i=1}^{194} \mathbf{n}_i(n)\chi_i(g)q^n$ (cf. (7.21)). Combining (7.18), (7.20) and (7.21), together with the definitions (3.2) of T_g and (7.22) of U_g , we obtain

$$T_g(\tau) = q^{-1} \frac{(1 - q)}{(q)_\infty} + \sum_{n>0} q^n \frac{1}{(q)_\infty} \sum_{i=1}^{194} \mathbf{n}_i(n)\chi_i(g), \tag{7.23}$$

or equivalently,

$$U_g(\tau) = (q)_\infty T_g(\tau) + 1 \tag{7.24}$$

for all $g \in \mathbb{M}$. (This computation also appears in [129]).

In Section 8, we will use (7.24) to determine the asymptotic behavior of the $\mathbf{n}_i(n)$ (cf. Theorem 9), and thus the statistics of black hole states, at $\ell = 16G$, in the conjectural chiral three-dimensional gravity theory dual to V^\natural .

Remark. Note that we may easily construct modules for $\mathcal{V} \times \mathbb{M}$ satisfying the extremal condition (6.8), for each positive integer m , by considering direct sums of the monster modules $V^{(-m)}$ constructed in Proposition 1. A very slight generalization of the argument

just given will then yield formulas for the graded traces of monster elements on the corresponding Virasoro highest weight spaces. Since it has been shown [114,137] that such modules cannot admit vertex operator algebra structure, we do not pursue this here.

8 Monstrous moonshine’s distributions

We now address the problem of determining exact formulas and asymptotic distributions of irreducible components. This work will rely heavily on the modularity of the underlying McKay-Thompson series (i.e., Theorems 1 and 8).

We prove formulas for the multiplicities $\mathbf{m}_i(-m, n)$ and $\mathbf{n}_i(n)$ which in turn imply the following asymptotics.

Theorem 9. *If m is a positive integer and $1 \leq i \leq 194$, then as $n \rightarrow +\infty$ we have*

$$\begin{aligned} \mathbf{m}_i(-m, n) &\sim \frac{\dim(\chi_i)|m|^{1/4}}{\sqrt{2}|n|^{3/4}|\mathbb{M}|} \cdot e^{4\pi\sqrt{|mn|}} \\ \mathbf{n}_i(n) &\sim \frac{\sqrt{12} \dim(\chi_i)}{|24n + 1|^{1/2}|\mathbb{M}|} \cdot e^{\frac{\pi}{6}\sqrt{23|24n+1|}} \end{aligned}$$

These asymptotics immediately imply that the following limits are well-defined

$$\begin{aligned} \delta(\mathbf{m}_i(-m)) &:= \lim_{n \rightarrow +\infty} \frac{\mathbf{m}_i(-m, n)}{\sum_{i=1}^{194} \mathbf{m}_i(-m, n)} \\ \delta(\mathbf{n}_i) &:= \lim_{n \rightarrow +\infty} \frac{\mathbf{n}_i(n)}{\sum_{i=1}^{194} \mathbf{n}_i(n)}. \end{aligned} \tag{8.1}$$

Corollary 1. *In particular, we have that*

$$\delta(\mathbf{m}_i(-m)) = \delta(\mathbf{n}_i) = \frac{\dim(\chi_i)}{\sum_{j=1}^{194} \dim(\chi_j)} = \frac{\dim(\chi_i)}{5844076785304502808013602136}.$$

We illustrate these asymptotics explicitly, for χ_1, χ_2 , and χ_{194} , in the case that $m = 1$, in Table 1, where we let $\delta(\mathbf{m}_i(-m, n))$ denote the proportion of components corresponding to χ_i in $V_n^{(m)}$.

The precise values given in the bottom row of Table 1 admit the following decimal approximations:

$$\begin{aligned} \delta(\mathbf{m}_1(-m)) &= \frac{1}{5844076785304502808013602136} \approx 1.711 \dots \times 10^{-28} \\ \delta(\mathbf{m}_2(-m)) &= \frac{196883}{5844076785304502808013602136} \approx 3.368 \dots \times 10^{-23} \\ \delta(\mathbf{m}_{194}(-m)) &= \frac{258823477531055064045234375}{5844076785304502808013602136} \approx 4.428 \dots \times 10^{-2} \end{aligned} \tag{8.2}$$

Before explaining the proof of Theorem 9, we describe an application to the computation^h of quantum dimensions. Suppose that V is a vertex operator algebra and M is a V -module such that the graded-dimension functions Z_V and Z_M are defined (cf. (4.2)). Then the *quantum dimension* of M relative to V is defined by setting

$$\text{qdim}_V M := \lim_{y \rightarrow 0} \frac{Z_M(iy)}{Z_V(iy)} \tag{8.3}$$

(for y real and positive), assuming the limit exists (cf. §3.1 of [75]).

Table 1 Asymptotic Proportions

n	$\delta(\mathbf{m}_1(-1, n))$	$\delta(\mathbf{m}_2(-1, n))$...	$\delta(\mathbf{m}_{194}(-1, n))$
-1	1	0	...	0
0	0	0	...	0
1	1/2	1/2	...	0
2	1/3	1/3	...	0
⋮	⋮	⋮	⋮	⋮
40	$4.011 \dots \times 10^{-4}$	$2.514 \dots \times 10^{-3}$...	0.00891...
60	$2.699 \dots \times 10^{-9}$	$2.732 \dots \times 10^{-8}$...	0.04419...
80	$4.809 \dots \times 10^{-14}$	$7.537 \dots \times 10^{-13}$...	0.04428...
100	$4.427 \dots \times 10^{-18}$	$1.077 \dots \times 10^{-16}$...	0.04428...
120	$1.377 \dots \times 10^{-21}$	$5.501 \dots \times 10^{-20}$...	0.04428...
140	$1.156 \dots \times 10^{-24}$	$1.260 \dots \times 10^{-22}$...	0.04428...
160	$2.621 \dots \times 10^{-27}$	$3.443 \dots \times 10^{-23}$...	0.04428...
180	$1.877 \dots \times 10^{-28}$	$3.371 \dots \times 10^{-23}$...	0.04428...
200	$1.715 \dots \times 10^{-28}$	$3.369 \dots \times 10^{-23}$...	0.04428...
220	$1.711 \dots \times 10^{-28}$	$3.368 \dots \times 10^{-23}$...	0.04428...
240	$1.711 \dots \times 10^{-28}$	$3.368 \dots \times 10^{-23}$...	0.04428...
⋮	⋮	⋮	⋮	⋮
∞	$\frac{1}{5844076785304502808013602136}$	$\frac{196883}{5844076785304502808013602136}$...	$\frac{258823477531055064045234375}{5844076785304502808013602136}$

Dong-Mason initiated a vertex algebraic *quantum Galois theory* in [81] (cf. also [77, 82,127]). In this theory, inclusions of vertex operator algebras take on the role played by inclusions of fields in the classical setting. It is established in [75] that the quantum dimension $qdim_U V$, for U a sub vertex operator algebra of V , serves as the quantum analogue of the relative dimension $dim_E F$ of a field F over a subfield E .

Write $V^{\mathbb{M}}$ for the fixed points of the action of \mathbb{M} on V^{\natural} . Then $V^{\mathbb{M}}$ is a sub vertex operator algebra of V^{\natural} , called the *monster orbifold*. Using Theorem 9, together with a result from [75], we will compute the quantum dimensions of the monster orbifold. To formulate this precisely, note that, according to the main theorem of [77], we have

$$V^{\natural} \simeq \bigoplus_{i=1}^{194} V_i^{\mathbb{M}} \otimes M_i, \tag{8.4}$$

as $V^{\mathbb{M}} \times \mathbb{M}$ -modules, for some $V^{\mathbb{M}}$ -submodules $V_i^{\mathbb{M}}$ in V^{\natural} , where M_i denotes an irreducible module for \mathbb{M} with character χ_i , as in (7.15).

Corollary 2. *We have $qdim_{V^{\mathbb{M}}} V_i^{\mathbb{M}} = \chi_i(e)$. In particular, the quantum dimension of $V_i^{\mathbb{M}}$ relative to $V^{\mathbb{M}}$ exists, for all $1 \leq i \leq 194$.*

Proof. Recall (cf. (4.2)) that $Z_{V_i^{\mathbb{M}}}(\tau)$ is the function obtained by substituting $q = e^{2\pi i\tau}$ in $dim_* V_i^{\mathbb{M}} = \sum_n (V_i^{\mathbb{M}})_n q^n$, assuming this series converges. Note that $V^{\mathbb{M}} = V_1^{\mathbb{M}}$. According to Proposition 3.6 of [75], if the limit

$$d_i := \lim_{n \rightarrow \infty} \frac{dim(V_i^{\mathbb{M}})_n}{dim(V_1^{\mathbb{M}})_n} \tag{8.5}$$

exists, then the quantum dimension of $V_i^{\mathbb{M}}$ relative to $V_1^{\mathbb{M}}$ also exists, and equals d_i . Comparing with (7.15) we see that $\dim(V_i^{\mathbb{M}})_n = \mathbf{m}_i(-1, n)$. Applying Theorem 9, we obtain $d_i = \chi_i(e)$, and this completes the proof. \square

Note that Corollary 2 confirms a special case of Conjecture 6.7 of [75]. It would be interesting to see how generally this method can be applied, to the computation of quantum dimensions of orbifolds V^G , where V is a vertex operator algebra and G is a group of automorphisms of $\text{Aut}(G)$. (See also Problem 11).

8.1 The modular groups in monstrous moonshine

To obtain exact formulas, we begin by recalling the modular groups which arise in monstrous moonshine. Suppose $\Gamma_* < \text{GL}_2(\mathbb{R})$ is a discrete group which is commensurable with $\text{SL}_2(\mathbb{Z})$. If Γ_* defines a genus zero quotient of \mathbb{H} , then the field of modular functions which are invariant under Γ_* is generated by a single element, the principal modulus (cf. (3.4)). Theorem 4 implies that the T_g (defined by (3.2)) are principal moduli for certain groups Γ_g . We can describe these groups in terms of groups E_g which in turn may be described in terms of the congruence subgroups

$$\Gamma_0(N) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) : c \equiv 0 \pmod{N} \right\}, \tag{8.6}$$

and the Atkin-Lehner involutions W_e for $\Gamma_0(N)$ given by

$$W_e := \begin{pmatrix} ae & b \\ cN & de \end{pmatrix}, \tag{8.7}$$

where e is an exact divisor of N (i.e., $e|N$, and $(e, N/e) = 1$), and a, b, c , and d are integers chosen so that W_e has determinant e .

Following Conway-Norton [58] and Conway-McKay-Sebbar [57], we denote the groups E_g by symbols of the form $\Gamma_0(N|h) + e, f, \dots$ (or simply $N|h + e, f, \dots$), where h divides $(N, 24)$, and each of e, f , etc. exactly divide N/h . This symbol represents the group

$$\Gamma_0(N|h) + e, f, \dots := \begin{pmatrix} 1/h & 0 \\ 0 & 1 \end{pmatrix} \langle \Gamma_0(N/h), W_e, W_f, \dots \rangle \begin{pmatrix} h & 0 \\ 0 & 1 \end{pmatrix},$$

where W_e, W_f , etc. are representative of Atkin-Lehner involutions on $\Gamma_0(N/h)$. We use the notation $\mathcal{W}_g := \{1, e, f, \dots\}$ to denote this list of Atkin-Lehner involutions contained in E_g . We also note that $\Gamma_0(N|h) + e, f, \dots$ contains $\Gamma_0(Nh)$.

The groups E_g are eigengroups for the T_g , so that if $\gamma \in E_g$, then $T_g(\gamma\tau) = \sigma_g(\gamma)T_g$, where σ_g is a multiplicative group homomorphism from E_g to the group of h th roots of unity. Conway and Norton [58] give the following values for σ_g evaluated on generators of $N|h + e, f, \dots$

Lemma 1 (Conway-Norton). *Assuming the notation above, the following are true:*

- (a) $\sigma_g(\gamma) = 1$ if $\gamma \in \Gamma_0(Nh)$
- (b) $\sigma_g(\gamma) = 1$ if γ is an Atkin-Lehner involution of $\Gamma_0(Nh)$ inside E_g
- (c) $\sigma_g(\gamma) = e^{-\frac{2\pi i}{h}}$ if $\gamma = \begin{pmatrix} 1 & 1/h \\ 0 & 1 \end{pmatrix}$

$$(d) \sigma_g(\gamma) = e^{-\lambda_g \frac{2\pi i}{h}} \text{ if } \gamma = \begin{pmatrix} 1 & 0 \\ N & 1 \end{pmatrix},$$

where λ_g in (d) is -1 if $N/h \in \mathcal{W}_g$, and $+1$ otherwise.

This information is sufficient to properly describe the modularity of the series $T_g^{(-m)}(\tau)$ on E_g . In section 8.4, we give an explicit procedure for evaluating σ_g . The invariance group Γ_g , denoted by $\Gamma_0(N|h) + e, f, \dots$ (or by the symbol $N|h + e, f, \dots$), is defined as the kernel of σ_g . A complete list of the groups Γ_g can be found in Appendix 10 of this paper, or in Table two of [58].

Theorems 1 and 8 are summarized by the following uniform statement.

Theorem 10. *Let $g \in \mathbb{M}$ and $m \geq 1$. Then $T_g^{(-m)}$ is the unique weakly holomorphic modular form of weight zero for Γ_g that satisfies $T_g^{(-m)} = q^{-m} + O(q)$ as τ approaches the infinite cusp, and has no poles at any cusps inequivalent to the infinite one.*

Remark. A weakly holomorphic modular form is a meromorphic modular form whose poles (if any) are supported at cusps.

8.2 Harmonic Maass forms

Maass-Poincaré series allow us to obtain formulas for weakly holomorphic modular forms and mock modular forms. We begin by briefly recalling the definition of a *harmonic Maass form* of weight $k \in \frac{1}{2}\mathbb{Z}$ and multiplier ν (a generalization of the notion of a Nebentypus). If $\tau = x + iy$ with x and y real, we define the weight k hyperbolic Laplacian by

$$\Delta_k := -y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + iky \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right), \tag{8.8}$$

and if $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$, define

$$(\gamma : \tau) := (c\tau + d).$$

Suppose Γ is a subgroup of finite index in $\text{SL}_2(\mathbb{Z})$ and $\frac{3}{2} \leq k \in \frac{1}{2}\mathbb{Z}$. Then a real analytic function $F(\tau)$ is a *harmonic Maass form* of weight k on Γ with multiplier ν if:

- (a) The function $F(\tau)$ satisfies the modular transformation with respect to the weight k slash operation,

$$F(\tau)|_k \gamma := (\gamma : \tau)^{-k} F(\gamma\tau) = \nu(\gamma)F(\tau)$$

for every matrix $\gamma \in \Gamma$, where if $k \in \mathbb{Z} + \frac{1}{2}$, the square root is taken to be the principal branch. In particular, if ν is trivial, then F is invariant under the action of the slash operator.

- (b) We have that $\Delta_k F(\tau) = 0$,
- (c) The singularities of F (if any) are supported at the cusps of Γ , and for each cusp ρ there is a polynomial $P_{F,\rho}(q^{-1}) \in \mathbb{C}[q^{-1/t_\rho}]$ and a constant $c > 0$ such that $F(\tau) - P_{F,\rho}(e^{-2\pi i\tau}) = O(e^{-c y})$ as $\tau \rightarrow \rho$ from inside a fundamental domain. Here t_ρ is the width of the cusp ρ .

Remark. The polynomial $P_{F,\rho}$ above is referred to as the *principal part of F at ρ* . In certain applications, condition (c) of the definition may be relaxed to admit larger classes of harmonic Maass forms. However, for our purposes, we will only be interested in those satisfying the given definition, having a holomorphic principal part.

We denote the complex vector space of such functions by $H_k(\Gamma, \nu)$, and note that in order for $H_k(\Gamma, \nu)$ to be nonzero, ν must satisfy $(\gamma : \delta\tau)^k(\delta : \tau)^k\nu(\gamma)\nu(\delta) = (\gamma\delta : \tau)^k\nu(\gamma\delta)$ for every $\gamma, \delta \in \Gamma$.

Let $\mathcal{S}(\Gamma)$ denote some fixed complete set of inequivalent representatives of the cusps of Γ . For each representative $\rho = \frac{\alpha}{\gamma}$ with $(\alpha, \gamma) = 1$, fix a matrix $L_\rho = \begin{pmatrix} -\delta & \beta \\ \gamma & -\alpha \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$ so that $\rho = L_\rho^{-1}\infty$. Following Rankin [205], let t_ρ be the cusp width and let κ_ρ be the cusp parameter, so that $\nu(L_\rho^{-1}T^{t_\rho}L_\rho) = e^{2\pi i\kappa_\rho}$, where $T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. The stabilizer of ρ in Γ is given by $\Gamma_\rho := \langle \pm L_\rho^{-1}T^{t_\rho}L_\rho \rangle$, so for example $\Gamma_\infty = \langle \pm T \rangle$. Given $F(\tau) \in H_{2-k}(\Gamma, \nu)$, we refer to $F_\rho(\tau) := F(\tau)|_{2-k}L_\rho$ as the expansion of F at the cusp ρ . We note that this expansion depends on the choice of L_ρ . These facts imply that the expansion of F_ρ can be given as a Fourier series of the form

$$F_\rho(\tau) = \sum_n a(n, y) e^{2\pi i x(n + \kappa_\rho)/t_\rho}.$$

More precisely, we have the following. The Fourier expansion of harmonic Maass forms F at a cusp ρ (see Proposition 3.2 of [30]) splits into two components. As before, we let $q := e^{2\pi i\tau}$.

Lemma 2. *If $F(\tau)$ is a harmonic Maass form of weight $2 - k$ for Γ where $\frac{3}{2} \leq k \in \frac{1}{2}\mathbb{Z}$, and if ρ is a cusp of Γ , then*

$$F_\rho(\tau) = F_\rho^+(\tau) + F_\rho^-(\tau)$$

where F_ρ^+ is the holomorphic part of F_ρ , given by

$$F_\rho^+(\tau) := \sum_{n \gg -\infty} c_{F,\rho}^+(n) q^{(n + \kappa_\rho)/t_\rho},$$

and F_ρ^- is the nonholomorphic part, given by

$$F_\rho^-(\tau) + \sum_{n < 0} c_{F,\rho}^-(n) \Gamma(k - 1, 4\pi y|(n + \kappa_\rho)/t_\rho|) q^{(n + \kappa_\rho)/t_\rho}.$$

By inspection, we see that weakly holomorphic modular forms are themselves harmonic Maass forms. In fact, under the given definition, all harmonic Maass forms of positive weight are weakly holomorphic. On the other hand, if the weight is non-positive, then the space of harmonic Maass forms may be larger than the space of weakly holomorphic modular forms. However, as with the weakly holomorphic modular forms, a harmonic Maass form is uniquely defined by its principal parts at all cusps. This is clear if the form is weakly holomorphic. If the nonholomorphic part is non-zero, we have the following lemma which follows directly from the work of Bruinier and Funke [30].

Lemma 3. *If $F \in H_{2-k}(\Gamma_0(N))$ has the property that $F^- \neq 0$, then the principal part of F is nonconstant for at least one cusp.*

Proof. Bruinier and Funke define a pairing $\{\bullet, \bullet\}$ on harmonic weak Maass Forms. The particular quantity $\left\{2iy^k \frac{\partial}{\partial \bar{\tau}} F, F\right\}$ can be described in terms of either a Petersson norm, or in terms of products of coefficients of the principal part of F with coefficients of the nonholomorphic part. Since the Petersson norm is non-zero, at least one coefficient of the principal part is also non-zero. \square

Hence, if F and G are harmonic Maass forms of non-positive weight whose principal parts are equal at all cusps, then $F - G$ is holomorphic and vanishes at cusps, and therefore identically 0.

8.3 Maass-Poincaré series

The Maass-Poincaré series define a basis for a space of harmonic Maass forms and provide exact formulas for their coefficients. The following construction of the Maass-Poincaré series follows the method and notation of Bringmann and the third author [26] which builds on the early work of Rademacher, followed by more contemporary work of Fay, Niebur, amongst many others [97,187,188]. The Poincaré series we construct in this section are modular for congruence subgroups $\Gamma_0(N)$ and we will use them to construct the McKay-Thompson series. Although we could follow similar methods to construct Poincaré series for the groups Γ_g directly, we restrict our attention to these groups since the congruence subgroups $\Gamma_0(N)$ are more standard.

For $s \in \mathbb{C}$, $w \in \mathbb{R} \setminus \{0\}$, and $k \geq 3/2$, $k \in \frac{1}{2}\mathbb{Z}$, let

$$\mathcal{M}_s(w) := |y|^{\frac{k}{2}-1} M_{\text{sign}(w)(1-k/2), s-\frac{1}{2}}(|w|), \tag{8.9}$$

where $M_{\nu, \mu}(z)$ is the M -Whittaker function which is a solution to the differential equation

$$\frac{\partial^2 u}{\partial z^2} + \left(-\frac{1}{4} + \frac{\nu}{z} + \frac{\frac{1}{4} - \mu^2}{z^2}\right) u = 0,$$

and (here and throughout this section) $\tau = x + iy$. Using this function, let

$$\phi_s(\tau) := \mathcal{M}_s(4\pi y) e^{2\pi i x}. \tag{8.10}$$

Given a positive integer m and a cusp ρ , Maass-Poincaré series provide a form with principal part equal to $q^{(-m+\kappa_\rho)/t_\rho}$ plus a constant at the cusp ρ , and constant at all other cusps, thereby forming a basis for $H_{2-k}(\Gamma, \nu)$.

Suppose $m > 0$ and $L \in \text{SL}_2(\mathbb{Z})$ with $\rho = L^{-1}\infty$. Then we have the Maass-Poincaré series

$$\mathcal{P}_L(\tau, m, \Gamma, 2-k, s, \nu) := \sum_{M \in \Gamma_\rho \backslash \Gamma} \frac{\phi_s\left(\frac{-m+\kappa_\rho}{t_\rho} \cdot L^{-1}M\tau\right)}{(L^{-1} : M\tau)^{2-k} (M : \tau)^{2-k} \nu(M)}. \tag{8.11}$$

It is easy to check that $\phi_s(\tau)$ is an eigenfunction of Δ_{2-k} with eigenvalue

$$s(1-s) + \frac{k^2 - 2k}{4}.$$

The right-hand side of (8.11) converges absolutely for $\Re(s) > 1$; however, Bringmann and the third author establish conditional convergence when $s \geq 3/4$ [26], giving Theorem 11

below. The theorem is stated for the specific case $\Gamma = \Gamma_0(N)$ for some N and $k \geq 3/2$, in which case we modify the notation slightly and define

$$\mathcal{P}_L(\tau, m, N, 2 - k, \nu) := \frac{1}{\Gamma(k)} \mathcal{P}_L\left(\tau, m, \Gamma_0(N), 2 - k, \frac{k}{2}, \nu\right) \tag{8.12}$$

In the statement of the theorem below, K_c is a modified Kloosterman sum given by

$$K_c(2 - k, L, \nu, m, n) := \sum_{\substack{0 \leq a < ct_\rho \\ a \equiv -\frac{c(\alpha, N)}{\alpha\gamma} \pmod{\frac{N}{(\gamma, N)}} \\ ad \equiv 1 \pmod{c}}} \frac{(S : \tau)^{2-k} \exp\left(2\pi i \left(\frac{a \cdot \frac{(-m+\kappa_\rho)}{t_\rho} + d \cdot \frac{(n+\kappa_\infty)}{t_\infty}}{c}\right)\right)}{(L^{-1} : LS\tau)^{2-k} (LS : \tau)^{2-k} \nu(LS)}, \tag{8.13}$$

where $S = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$. If ν is trivial, we omit it from the notation. We also have that $\delta_{L,S}(m)$ is an indicator function for the cusps $\rho = L^{-1}\infty$ and $\mu = S^{-1}\infty$ given by

$$\delta_{L,S}(m) := \begin{cases} \nu(M)^{-1} e^{2\pi i r \frac{-m+\kappa_\rho}{t_\rho}} & \text{if } M = LT^r S^{-1} \in \Gamma_0(N), \\ 0 & \text{if } \mu \not\sim \rho \text{ under } \Gamma_0(N). \end{cases}$$

Using this notation, we have the following theorem which gives exact formulas for the coefficients and principal part of $\mathcal{P}_L(\tau, m, N, 2 - k, \nu)$, which is a generalization of Theorem 3.2 of [26].

Theorem 11. *Suppose that $\frac{3}{2} \leq k \in \frac{1}{2}\mathbb{Z}$, and suppose $\rho = L^{-1}\infty$ is a cusp of $\Gamma_0(N)$. If m is a positive integer, then $\mathcal{P}_L(\tau, m, N, 2 - k, \nu)$ is in $H_{2-k}(\Gamma_0(N), \nu)$. Moreover, the following are true:*

1. We have

$$\mathcal{P}_L^+(\tau, m, N, 2 - k, \nu) = \delta_{\rho, I}(m) \cdot q^{-m+\kappa_\infty} + \sum_{n \geq 0} a^+(n) q^{n+\kappa_\infty}.$$

Moreover, if $n > 0$, then $a^+(n)$ is given by

$$-t^k 2\pi \left| \frac{-m + \kappa_\rho}{t_\rho(n + \kappa_\infty)} \right|^{\frac{k-1}{2}} \sum_{\substack{c > 0 \\ (c, N) = (\gamma, N)}} \frac{K_c(2 - k, L, \nu, -m, n)}{c} \cdot I_{k-1} \left(\frac{4\pi}{c} \sqrt{\frac{|-m + \kappa_\rho| |n + \kappa_\infty|}{t_\rho}} \right),$$

where I_k is the usual I -Bessel function.

2. If $S \in \text{SL}_2(\mathbb{Z})$, then there is some $c \in \mathbb{C}$ so that the principal part of $\mathcal{P}_L(\tau, m, N, 2 - k)$ at the cusp $\mu = S^{-1}\infty$ is given by

$$\delta_{L,S}(m) q^{\frac{-m+\kappa_\rho}{t_\rho}} + c$$

Remark. We shall use Theorem 11 to prove Theorem 9. We note that one could prove Theorem 9 without referring to the theory of Maass-Poincaré series. One could make use of ordinary weakly holomorphic Poincaré series. However, we have chosen to use Maass-Poincaré series and the theory of harmonic Maass forms because these results are more general, and because of the recent appearance of harmonic Maass forms in the theory of *umbral moonshine* (see Section 9).

Proof. Writing $\rho = \frac{\alpha}{\gamma}$, Bringmann and the third author prove this theorem for the case that $\gamma \mid N$ and $(\alpha, N) = 1$, along with the assumption that μ and ρ are in a fixed complete set of inequivalent cusps, so that $\delta_{\mu, \rho} = 1$ or 0. This general form is useful to us particularly since it works equally well for the cusps ∞ with L taken to be the identity, and for 0 with L taken to be $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

Here and for the remainder of the paper, we let \mathcal{S}_N denote any complete set of inequivalent cusps of $\Gamma_0(N)$, and for each $\rho \in \mathcal{S}_N$, we fix some L_ρ with $\rho = L_\rho^{-1}\infty$. Rankin notes [205] (Proof of Theorem 4.1.1(iii)) that given some choice of \mathcal{S}_N , each right coset of $\Gamma_0(N)\backslash\text{SL}_2(\mathbb{Z})$ is in $\Gamma_0(N) \cdot L_\rho T^r$ for some unique $\rho \in \mathcal{S}_N$. Moreover, the r in the statement is unique modulo t_ρ , so the function $\delta_{L,S}(m)$ given above is well-defined on all matrices in $\text{SL}_q(\mathbb{Z})$.

In the proof given by Bringmann and the third author, the sum of Kloosterman sums $\sum_{\substack{c>0 \\ (c,N)=(\gamma,N)}} \frac{K_c(2-k, L, v, -m, n)}{c} \dots$ is written as a sum over representatives of the double coset $\Gamma_\rho \backslash L^{-1}\Gamma_0(N)/\Gamma_\infty$ (omitting the identity if present). Following similar arguments, but without the assumptions on α and γ , we find the indices of summation given in (8.13) and Theorem 11. As in their case, we find that the principal part of $\mathcal{P}_{L_\rho}(\tau, m, N, 2-k)$ at a cusp μ is constant if $\mu \not\sim \rho$ and is $\delta_{L_\rho, L_\mu} q^{\frac{-m}{t_\rho}} + c$ for some constant if $\mu = \rho$. Therefore, if μ is a cusp with $L_\mu = M^{-1}L_\rho T^r$ for some $M \in \Gamma_0(N, \cdot)$, then the principal part of $\mathcal{P}_{L_\rho}(\tau, m, N, 2-k, v)$ at μ is clearly $v(M)^{-1} e^{2\pi i r \frac{-m+\kappa}{t_\rho}} q^{\frac{-m+\kappa\rho}{t_\rho}} + c$. \square

Since harmonic Maass forms with a nonholomorphic part have a non-constant principal part at some cusp, we have the following theorem.

Theorem 12. ([26], Theorem 1.1) *Assuming the notation above, if $\frac{3}{2} \leq k \in \frac{1}{2}\mathbb{Z}$, and $F(\tau) \in H_{2-k}(\Gamma_0(N), v)$ has principal part $P_\rho(\tau) = \sum_{m \geq 0} a_\rho(-m) q^{\frac{-m+\kappa\rho}{t_\rho}}$ for each cusp $\rho \in \mathcal{S}_N$, then*

$$F(\tau) = \sum_{\rho \in \mathcal{S}_N} \sum_{m > 0} a_\rho(-m) \mathcal{P}_\rho(\tau, m, N, 2-k, v) + g(\tau),$$

where $g(\tau)$ is a holomorphic modular form. Moreover, we have that $c = 0$ whenever $k > 2$, and is a constant when $k = 2$.

8.4 Exact formulas for $Z_g^{(-m)}$

Using Theorems 11 and 12, we can write exact formulas for the coefficients of the $T_g^{(-m)}$ provided we know its principal parts at all cusps of $\Gamma_0(Nh)$. With this in mind, we now regard T_g as a modular function on $\Gamma_0(Nh)$ with trivial multiplier. The location and orders of the poles were determined by Harada and Lang [128].

Lemma 4. ([128], Lemma 7, 9) *Suppose the Γ_g is given by the symbol $N||h + e, f \dots$, and let $L = \begin{pmatrix} -\delta & \beta \\ \gamma & -\alpha \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$. then $T_g|_0L$ has a pole if and only if $\left(\frac{\gamma}{\gamma h}, \frac{N}{h}\right) = \frac{N}{eh}$, for some $e \in \mathcal{W}_g$ (note, here we allow $e = 1$). the order of the pole is given by $\frac{(h, \gamma)^2}{eh^2}$.*

Harada and Lang prove this lemma by showing that if u is an integer chosen such that $\frac{u\gamma - \alpha \cdot (h, \gamma)}{h}$ is integral and divisible by e , and $U = \begin{pmatrix} \frac{e-h}{(h, \gamma)} & \frac{u}{h} \\ 0 & \frac{(h, \gamma)}{h} \end{pmatrix}$, then LU is an Atkin-Lehner involution $W_e \in E_g$. Therefore, we have that

$$T_g|_0L = \sigma_g(LU)T_g \left(\frac{(h, \gamma)^2}{eh^2} \tau - \frac{u \cdot (h, \gamma)}{eh^2} \right). \tag{8.14}$$

Harada and Lang do not compute $\sigma_g(LU)$; however, we will need these values in order to apply Theorem 12. Using Lemma 1, the following procedure allows us to compute $\sigma_g(M)$ for any matrix $M \in E_g$.

Given a matrix $M \in E_g$, we may write M as $M = \begin{pmatrix} ae & \frac{b}{h} \\ cN & de \end{pmatrix}$ with $e \in \mathcal{W}_g$ and $ade - bc\frac{N}{eh} = 1$. We may also write $h = h_e \cdot h_{\bar{e}}$, where $h_{\bar{e}}$ is the largest divisor of h co-prime to e . Since $c\frac{N}{eh}$ is co-prime to both d and e , we may chose integers A, B , and $C \pmod{h}$ such that:

- $c\frac{N}{eh}A + d$ is co-prime to $h_{\bar{e}}$ but is divisible by h_e ,
- $B \equiv -(eaA + b)(c\frac{N}{h}A + ed)^{-1} \pmod{h_{\bar{e}}}$ and $Bc\frac{N}{eh} + b \equiv 0 \pmod{h_e}$,
- $C \equiv -c(c\frac{N}{h}A + ed)^{-1} \pmod{h_{\bar{e}}}$, and $C \equiv 0 \pmod{h_e}$.

A calculation shows that $\widehat{M} := \begin{pmatrix} 1 & \frac{B}{h} \\ 0 & 1 \end{pmatrix} M \begin{pmatrix} 1 & \frac{A}{h} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ cN & 1 \end{pmatrix} \begin{pmatrix} h_e & 0 \\ 0 & h_e \end{pmatrix}$ is an Atkin-Lehner involution W_E for $\Gamma_0(Nh)$ where $E = e \cdot h_e^2$. By Lemma 1, this implies $\sigma_g(\widehat{M}) = 1$, and therefore

$$\sigma_g(M) = \exp \left(\frac{2\pi i}{h} (A + B + \lambda_g C) \right).$$

Combined with Lemma 4, this leads us to the following proposition.

Proposition 2. *Given a matrix $L = \begin{pmatrix} -\delta & \beta \\ \gamma & -\alpha \end{pmatrix} \in SL_2(\mathbb{Z})$, let u and U be chosen as above, and define*

$$\epsilon_g(L) := \sigma_g(LU) \cdot e^{2\pi i \frac{u \cdot (h, \gamma)}{eh^2}}.$$

Then by (8.14), we have that

$$T_g|_0L = \epsilon_g(L)q^{-\frac{(h, \gamma)^2}{eh^2}} + O(q).$$

Using this notation, we are equipped to find exact formulas for the $T_g^{(-m)}$.

Theorem 13. *Let $g \in \mathbb{M}$, with $\Gamma_g = N|h + e, f, \dots$, and let S_{Nh} and \mathcal{W}_g be as above. if m and n are positive integers, then there is a constant c for which*

$$T_g^{(-m)}(\tau) = c + \sum_{e \in \mathcal{W}_g} \sum_{\substack{\frac{\alpha}{\gamma} \in S_{Nh} \\ \left(\frac{\gamma}{(h, \gamma)}, \frac{N}{h}\right) = \frac{N}{eh}}} \epsilon_g(L_\rho)^m \mathcal{P}_{\alpha/\gamma}^+(\tau, m, Nh, 0).$$

The n -th coefficient of $T_g^{(-m)}(\tau)$ is given by

$$\sum_{e \in \mathcal{W}_g} \sum_{\substack{\rho = \frac{\alpha}{\gamma} \in \mathcal{S}_{Nh} \\ \left(\frac{\gamma}{(\gamma, h)}, \frac{N}{h}\right) = \frac{N}{eh}}} \epsilon_g(L_\rho)^m 2\pi \left| \frac{-m}{n} \cdot \frac{(h, \gamma)^2}{eh^2} \right|^{\frac{1}{2}} \times \sum_{\substack{c > 0 \\ (c, Nh) = (\gamma, Nh)}} \frac{K_c(0, L, -m, n)}{c} \cdot I_1 \left(\frac{4\pi}{c} \sqrt{\left| \frac{-mn \cdot (h, \gamma)^2}{eh^2} \right|} \right).$$

Proof. Every modular function is a harmonic Maass form. Therefore, the idea is to exhibit a linear combination of Maass-Poincaré series with exactly the same principal parts at all cusps as $T_g^{(-m)}$. By Lemma 3 and Theorem 12, this form equals $T_g^{(-m)}$ up to an additive constant. Lemma 1 (c) implies that the coefficients $c_g(n)$ of T_g are supported on the arithmetic progression $n \equiv -1 \pmod{h}$. The function $T_g^{(-m)}$ is a polynomial in T_g , and as such must be the sum of powers of T_g each of which is congruent to $m \pmod{h}$. Therefore, if $M \in \Gamma_g$, then $T^{(-m)}|_0 M = \sigma_g(M)^m T^{(-m)}$. Given $L \in \text{SL}_2(\mathbb{Z})$, let U be a matrix as in (8.14) so that $LU \in \Gamma_g$. By applying Proposition 2, we find

$$T_g^{(-m)}|_0 L = \sigma_g(LU)^m T_g^{(-m)}|_0 U^{-1} = \epsilon_g(L)^m q^{-m \frac{(h, \gamma)^2}{eh^2}} + O(q).$$

Theorem 12, along with the observations that $t_\rho = \frac{(h, \gamma)^2}{eh^2}$ and $\kappa_\rho = 0$ for every $\rho = \frac{\alpha}{\gamma} = L_\rho^{-1}\infty$, implies the first part of the theorem. The formula for the coefficients follows by Theorem 11. □

8.5 Exact formulas for U_g up to a theta function

Following a similar process to that in the previous section, we may construct a series $\widehat{U}_g(\tau) = q^{-\frac{23}{24}} + O\left(q^{\frac{1}{24}}\right)$ with principal parts matching those of $\eta(\tau)T_g(\tau)$ at all cusps. Then according to (7.24), the difference $q^{\frac{1}{24}}(U_g - 1) - \widehat{U}_g$ is a weight $\frac{1}{2}$ holomorphic modular form, which by a celebrated result [210] of Serre-Stark, is a finite linear combination of unary theta functions. This will not affect the asymptotics in Theorem 9. The functions T_g and \widehat{U}_g differ primarily in their weight, and in that \widehat{U}_g has a non-trivial multiplier $\nu_\eta : M \rightarrow \frac{\eta(M\tau)}{(M:\tau)^{1/2}\eta(\tau)}$. They also have slightly different orders of poles, which is accounted for by the fact that the multiplier ν_η implies that $\kappa_\rho = t_\rho/24$ at every cusp ρ for the \widehat{U}_g , rather than 0 for the T_g . The proof of the following theorem is the same as that of Theorem 13, *mutatis mutandis*.

Theorem 14. *Let $g \in \mathbb{M}$, with $\Gamma_g = N|h + e, f, \dots$, and let \mathcal{S}_{Nh} and \mathcal{W}_g be as above. If m is a positive integer then*

$$\widehat{U}_g = \sum_{e \in \mathcal{W}_g} \sum_{\substack{\rho = \frac{\alpha}{\gamma} \in \mathcal{S}_{Nh} \\ \left(\frac{\gamma}{(\gamma, h)}, \frac{N}{h}\right) = \frac{N}{eh}}} \epsilon_g(L_\rho) \mathcal{P}_{L_\rho}^+(\tau, 1, Nh, 1/2, \nu_\eta).$$

For n a non-negative integer, the coefficient of $q^{n+\frac{1}{24}}$ in \widehat{U}_g is given by

$$\sum_{e \in \mathcal{W}_g} \sum_{\substack{\rho = \frac{\alpha}{\gamma} \in \mathcal{S}_{Nh} \\ \left(\frac{\gamma}{(\gamma, h)}, \frac{N}{h}\right) = \frac{N}{eh}}} \epsilon_g(L_\rho) \frac{1-i}{\sqrt{2}} 2\pi \left| \frac{-(h, \gamma)^2}{eh^2} + \frac{1}{24} \right|^{\frac{1}{4}} \times$$

$$\sum_{\substack{c > 0 \\ (c, Nh) = (\gamma, Nh)}} \frac{K_c\left(\frac{1}{2}, L, v_\eta, -1, n\right)}{c} \cdot I_{\frac{1}{2}} \left(\frac{4\pi}{c} \sqrt{\left| -\frac{(h, \gamma)^2}{eh^2} + \frac{1}{24} \right| \left| n + \frac{1}{24} \right|} \right).$$

This immediately admits the following corollary.

Corollary 3. *Given the notation above, there is a weight $\frac{1}{2}$ linear combination of theta functions $h_g(\tau)$ for which the coefficient q^n in $U_g(\tau) - q^{-\frac{1}{24}}h_g(\tau)$ coincides with the coefficient of $q^{n+\frac{1}{24}}$ in \widehat{U}_g , given explicitly in Theorem 14.*

8.6 Proof of Theorem 9

Proof. Following Harada and Lang [128], we begin by defining the functions

$$T_{\chi_i}^{(-m)}(\tau) := \frac{1}{|\mathbb{M}|} \sum_{g \in \mathbb{M}} \chi_i(g) T_g^{(-m)}(\tau). \tag{8.15}$$

The orthogonality of characters imply that for g and $h \in \mathbb{M}$,

$$\sum_{i=1}^{194} \overline{\chi_i(g)} \chi_i(h) = \begin{cases} |C_{\mathbb{M}}(g)| & \text{if } g \text{ and } h \text{ are conjugate,} \\ 0 & \text{otherwise.} \end{cases} \tag{8.16}$$

Here $|C_{\mathbb{M}}(g)|$ is the order of the centralizer of g in \mathbb{M} . Since the order of the centralizer times the order of the conjugacy class of an element is the order of the group, (8.16) and (8.15) together imply the inverse relation

$$T_g^{(-m)}(\tau) = \sum_{i=1}^{194} \overline{\chi_i(g)} T_{\chi_i}^{(-m)}(\tau).$$

In particular we have that $T_e^{(-m)}(\tau) = \sum_{i=1}^{194} \dim(\chi_i) T_{\chi_i}^{(-m)}(\tau)$, and therefore we can identify

the $\mathbf{m}_i(-m, n)$ as the Fourier coefficients of the $T_{\chi_i}^{(-m)}(\tau) = \sum_{n=-m}^{\infty} \mathbf{m}_i(-m, n) q^n$.

Using Theorem 13, we obtain exact formulas for the coefficients of $T_{\chi_i}^{(-m)}(\tau)$. Let $g \in \mathbb{M}$ with $\Gamma_g = N_g || h_g + e_g, f_g, \dots$. If m and n are positive integers, then the n th coefficient is given exactly by

$$\frac{1}{|\mathbb{M}|} \sum_{g \in \mathbb{M}} \chi_i(g) \sum_{e \in \mathcal{W}_g} \sum_{\substack{\alpha \in \mathcal{S}_{N_g h_g} \\ \left(\frac{\gamma}{(\gamma, h_g)}, \frac{N_g}{h_g}\right) = \frac{N_g}{eh_g}}} \epsilon_g(L_\rho)^m 2\pi \left| \frac{-m}{n} \cdot \frac{(h_g, \gamma)^2}{eh_g^2} \right|^{\frac{1}{2}}$$

$$\sum_{\substack{c > 0 \\ (c, N_g h_g) = (\gamma, N_g h_g)}} \frac{K_c(2-k, L, v, -m, n)}{c} \cdot I_1 \left(\frac{4\pi}{c} \sqrt{\frac{-mn \cdot (h_g, \gamma)^2}{eh_g^2}} \right),$$

where $\mathcal{S}_{N_g h_g}$ and \mathcal{W}_g are given as above.

Using the well-known asymptotics for the I -Bessel function

$$I_k(x) \sim \frac{e^x}{\sqrt{2\pi x}} \left(1 - \frac{4k^2 - 1}{8x} + \dots \right),$$

we see that the formula for $\mathbf{m}_i(-m, n)$ is dominated by the $c = 1$ term which appears only for $g = e$ (so that $N_e = h_e = 1$). This term yields the asymptotic

$$\mathbf{m}_i(-m, n) \sim \frac{\chi_i(e) \cdot |m|^{1/4}}{\sqrt{2}n^{3/4}|\mathbb{M}|} \cdot e^{4\pi\sqrt{|mn|}}$$

as in the statement of the theorem.

The asymptotics for $\mathbf{n}_i(n)$ follows similarly, using the formula

$$U_{\chi_i}(\tau) := \frac{1}{|\mathbb{M}|} \sum_{g \in \mathbb{M}} \chi_i(g) U_g^{(-m)}(\tau).$$

We note that the coefficients of the theta functions $h_g(\tau)$ in Corollary 3 are bounded by constants and so do not affect the asymptotics. This yields

$$\mathbf{n}_i(n) \sim \frac{\sqrt{12} \chi_i(e)}{|24n + 1|^{1/2}|\mathbb{M}|} \cdot e^{\frac{\pi}{6}\sqrt{23|24n+1|}}$$

as in the theorem. □

8.7 Examples of the exact formulas

We conclude with a few examples illustrating the exact formulas for the McKay-Thompson series. These formulas for the coefficients generally converge rapidly. However, the rate of convergence is not uniform and often requires many more terms to converge to a given precision.

Example. We first consider the case that g is the identity element. Then we have $\Gamma_g = \text{SL}_2(\mathbb{Z})$, which has only the cusp at infinity. In this case Theorem 13 reduces to the well-known expansion

$$T_g = J(\tau) - 744 = q^{-1} + \sum_{n \geq 1} \frac{2\pi}{\sqrt{n}} \cdot \sum_{c > 0} \frac{K_c(\infty, -m, n)}{c} \cdot I_1\left(\frac{4\pi\sqrt{n}}{c}\right) q^n.$$

Table 2 below contains several approximations made by bounding the size of the c term in the summation.

Example. The second example we consider is g in the conjugacy class 4B. In this case we have $\Gamma_g = 4||2 + 2 \supset \Gamma_0(8)$. The function T_g has a pole at each of the four cusps of $\Gamma_0(8)$:

1. The cusp ∞ has $e = 1$, width $t = 1$, and coefficient $\epsilon(L_\infty) = 1$.
2. The cusp 0 has $e = 2$, width $t = 8$, and coefficient $\epsilon(L_0) = 1$.

Table 2 Convergence to Coefficients 1A

	$n = 1$	$n = 5$	$n = 10$
$c \leq 25$	196883.661 ...	333202640598.254 ...	22567393309593598.047 ...
≤ 50	196883.881 ...	333202640599.429 ...	22567393309593598.660 ...
≤ 75	196883.840 ...	333202640599.828 ...	22567393309593599.369 ...
≤ 100	196883.958 ...	333202640599.827 ...	22567393309593599.681 ...
∞	196884	333202640600	22567393309593600

3. The cusp $1/2$ has $e = 2$, width $t = 2$, and coefficient $\epsilon(L_{1/2}) = i$.
4. The cusp $1/4$ has $e = 1$, width $t = 1$, and $\epsilon(L_{1/4}) = -1$.

Table 3 below contains several approximations as in Table 2.

9 Umbral moonshine

In this penultimate section, we review the recently discovered and rapidly developing field of umbral moonshine.

9.1 K3 surfaces

Eguchi-Ooguri-Tachikawa reignited the field of moonshine with their 2010 observation [93] that dimensions of representations of the largest Mathieu group, M_{24} , occur as multiplicities of superconformal algebra characters in the K3 elliptic genus.

To formulate their observation more precisely, recall that a *complex K3 surface* is a compact connected complex manifold M of dimension 2, with $\Omega_M^2 \simeq \mathcal{O}_M$ and $H^1(M, \mathcal{O}_M) = 0$ (see [10,11] for introductory accounts). Following Witten [166,230], the *elliptic genus* of a complex manifold M of dimension d is defined to be

$$Z_M := \int_M \text{ch}(\mathbb{E}_{q,y}) \text{td}(M), \tag{9.1}$$

where $\text{td}(M)$ is the Todd class of M , and $\text{ch}(\mathbb{E}_{q,y})$ is the Chern character of the formal power series

$$\mathbb{E}_{q,y} = y^{\frac{d}{2}} \bigotimes_{n=1}^{\infty} \left(\bigwedge_{-y^{-1}q^{n-1}} T_M \otimes \bigwedge_{-yq^n} T_M^* \otimes \bigvee_{q^n} T_M \otimes \bigvee_{q^n} T_M^* \right), \tag{9.2}$$

whose coefficients are virtual vector bundles, obtained as sums of tensor products of the exterior and symmetric powers of the holomorphic tangent bundle T_M , and its dual bundle T_M^* (cf. also §1 of [122] or Appendix A of [71]). In (9.2) we interpret $\bigvee_t E$ in direct analogy with $\bigwedge_t E$ (cf. (3.13)), replacing exterior powers $\bigwedge^k E$ with symmetric powers $\bigvee^k E$.

Since a complex K3 surface M has trivial canonical bundle, and hence vanishing first Chern class, its elliptic genus is a *weak Jacobi form* $Z_M(\tau, z)$ of weight zero and index $\dim(M)/2 = 1$ - see [122] or [136] for proofs of this fact - once we set $q = e(\tau)$ and $y = e(z)$. This meansⁱ that $Z_M(\tau, z)$ is a holomorphic function on $\mathbb{H} \times \mathbb{C}$, satisfying

$$\begin{aligned} Z_M(\tau, z) &= e\left(\frac{-cz^2}{c\tau + d}\right) Z_M\left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}\right) \\ &= e(\lambda^2\tau + 2\lambda z) Z_M(\tau, z + \lambda\tau + \mu) \end{aligned} \tag{9.3}$$

for $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$ and $\lambda, \mu \in \mathbb{Z}$, with a Fourier expansion $Z_M(\tau, z) = \sum_{n,r} c(n, r) q^n y^r$ such that $c(n, r) = 0$ whenever $n < 0$.

Table 3 Convergence to Coefficients 4B

	$n = 1$	$n = 5$	$n = 10$
$c \leq 25$	51.975...	4760.372...	0.107...
≤ 50	52.003...	4759.860...	0.117...
≤ 75	52.041...	4760.066...	0.092...
≤ 100	51.894...	4760.049...	0.040...
∞	52	4760	0

It is known that $Z_M(\tau, z)$ specializes to the Euler characteristic $\chi(M) = 24$ when $z = 0$ (cf., e.g., [122]). The space of weak Jacobi forms of weight zero and index one is one-dimensional (cf. [96]), so Z_M is independent of the choice of M . In fact, we have

$$Z_M(\tau, z) = 8 \left(\frac{\theta_2(\tau, z)^2}{\theta_2(\tau, 0)^2} + \frac{\theta_3(\tau, z)^2}{\theta_3(\tau, 0)^2} + \frac{\theta_4(\tau, z)^2}{\theta_4(\tau, 0)^2} \right), \tag{9.4}$$

where the $\theta_i(\tau, z)$ are the usual Jacobi theta functions:

$$\begin{aligned} \theta_2(\tau, z) &:= \sum_{n \in \mathbb{Z}} y^{n+1/2} q^{(n+1/2)^2/2} = y^{1/2} q^{1/8} \prod_{n>0} (1 + y^{-1} q^{n-1})(1 + yq^n)(1 - q^n) \\ \theta_3(\tau, z) &:= \sum_{n \in \mathbb{Z}} y^n q^{n^2/2} = \prod_{n>0} (1 + y^{-1} q^{n-1/2})(1 + yq^{n-1/2})(1 - q^n) \\ \theta_4(\tau, z) &:= \sum_{n \in \mathbb{Z}} (-1)^n y^n q^{n^2/2} = \prod_{n>0} (1 - y^{-1} q^{n-1/2})(1 - yq^{n-1/2})(1 - q^n). \end{aligned} \tag{9.5}$$

In Witten’s original analysis [166,230], the elliptic genus Z_M is the graded trace of an integer-valued operator on a Hilbert space arising from a *supersymmetric nonlinear sigma model* on M . In the case that M is a K3 surface - see [3,4] for analyses of the sigma models associated to K3 surfaces - it is expected that the corresponding Hilbert space admits a unitary action by the (small) $N = 4$ superconformal algebra (cf. [95]). At least this is known for some special cases, so (9.4) can be written as an integer combination of the irreducible unitary $N = 4$ algebra characters. This leads (cf. [91,94]) to an expression

$$Z_M(\tau, z) = 24\mu(\tau, z) \cdot \frac{\theta_1(\tau, z)^2}{\eta(\tau)^3} + H(\tau) \cdot \frac{\theta_1(\tau, z)^2}{\eta(\tau)^3}, \tag{9.6}$$

where $\theta_1(\tau, z)$ is the Jacobi theta function

$$\theta_1(\tau, z) := i \sum_{n \in \mathbb{Z}} (-1)^n y^{n+1/2} q^{(n+1/2)^2/2} = iy^{1/2} q^{1/8} \prod_{n>0} (1 - y^{-1} q^{n-1})(1 - yq^n)(1 - q^n), \tag{9.7}$$

$\mu(\tau, z)$ denotes the Appell-Lerch sum defined by

$$\mu(\tau, z) := \frac{iy^{1/2}}{\theta_1(\tau, z)} \sum_{n \in \mathbb{Z}} (-1)^n \frac{y^n q^{n(n+1)/2}}{1 - yq^n}, \tag{9.8}$$

and $q^{1/8}H(\tau)$ is a power series in q with integer coefficients,

$$H(\tau) = -2q^{-1/8} + 90q^{7/8} + 462q^{15/8} + 1540q^{23/8} + 4554q^{31/8} + 11592q^{39/8} + \dots \tag{9.9}$$

The surprising observation of [93] is that each coefficient of a non-polar term appearing in (9.9) is twice the dimension of an irreducible representation (cf. [56]) of the sporadic simple group M_{24} , discovered by Émile Mathieu [182,183] more than 150 years ago (Generally, the coefficient of a positive power of q in (9.9) is some non-negative integer combination of dimensions of representations of M_{24} .) Thus $H(\tau)$ serves as an analogue of $J(\tau)$ (cf. (2.6)) for M_{24} , and

$$90 = 45 + 45 \tag{9.10}$$

is the Mathieu analogue of McKay’s monstrous observation (2.4).

The analogy with monstrous moonshine was quickly taken up, with the determination by Cheng [38], Eguchi-Hikami [92], and Gaberdiel-Hohenegger-Volpato [107,108], of *Mathieu McKay-Thompson series*

$$H_g(\tau) = -2q^{-1/8} + \sum_{n>0} \text{tr}(g|K_{n-1/8})q^{n-1/8} \tag{9.11}$$

associated to a graded M_{24} -module $K = \bigoplus_{n>0} K_{n-1/8}$, such that $K_{7/8}$ is the sum of the two 45-dimensional irreducible representations of M_{24} , and $K_{15/8}$ the sum of the two 231-dimensional irreducible representations, etc. (See [42] for a detailed review of Mathieu moonshine, and explicit descriptions of the H_g in particular.) We have the following beautiful recent result of Gannon [42].

Theorem 15 (Gannon). *There is a graded M_{24} -module $K = \bigoplus_{n>0} K_{n-1/8}$ for which (9.11) is true (given that the H_g are as described in [42]).*

Remark. A concrete construction of K remains unknown.

The observer may ask: how were the H_g determined, if the module K is as yet unknown? To explain this, note that the subscript in M_{24} is a reference to the fact that M_{24} is distinguished amongst permutation groups: it may be characterized as the unique proper subgroup of the alternating group A_{24} that acts quintuply transitively on 24 points (cf. [65]). Write χ_g for the number of fixed points of an element $g \in M_{24}$, in this defining permutation representation.

The first few terms of H_g are determined by the Eguchi-Ooguri-Tachikawa observation on (9.9), for it indicates that the coefficient of $q^{7/8}$ in H_g should be the trace of g on the sum of the two 45-dimensional irreducible representations, and the coefficient of $q^{15/8}$ should be the trace of g on the sum of the two 231-dimensional irreducible representations, etc. To determine the remaining infinitely many terms, modularity may be used: the series H_g , determined in [38,92,107,108], have the property that

$$Z_g(\tau, z) := \chi_g \mu(\tau, z) \cdot \frac{\theta_1(\tau, z)^2}{\eta(\tau)^3} + H_g(\tau) \cdot \frac{\theta_1(\tau, z)^2}{\eta(\tau)^3} \tag{9.12}$$

is a weak Jacobi form of weight zero and index one for $\Gamma_0^J(N) := \Gamma_0(N) \times \mathbb{Z}^2$ (with non-trivial multiplier when $\chi_g \neq 0$), where $N = o(g)$ is the order of g , and $\Gamma_0(N)$ is as in (8.6).

Thus Mathieu moonshine entails twisted or *twined* versions (9.12) of the K3 elliptic genus (9.4), but the single variable series $H_g(\tau)$ may also be studied in their own right, as automorphic objects of a particular kind: it turns out that they are *mock modular forms*¹ of weight 1/2, for various groups $\Gamma_0(N)$, with *shadows* $\chi_g \eta(\tau)^3$. This means that the *completed* functions

$$\widehat{H}_g(\tau) := H_g(\tau) + \chi_g \frac{1}{2\sqrt{i}} \int_{-\bar{\tau}}^{\infty} \frac{\overline{\eta(-\bar{w})^3}}{\sqrt{w+\tau}} dw \tag{9.13}$$

are harmonic Maass forms of weight 1/2, with the same multiplier system as $\eta(\tau)^{-3}$ when $\chi_g \neq 0$. (In case $\chi_g = 0$, i.e., when H_g is already a modular form, the multiplier is slightly different. See, e.g., [42]. The groups $\Gamma_0(o(g))$ for which $\chi_g \neq 0$ are characterized in [44].) The function $H_g(\tau)$, being the holomorphic part of $\widehat{H}_g(\tau)$, is the *mock modular form*.

In contrast to the twined K3 elliptic genera Z_g , the mock modular forms H_g are distinguished, in a manner directly analogous to the McKay-Thompson series T_g of monstrous moonshine: it is shown in [43] that the H_g admit a uniform description in terms of Rademacher sums, in direct analogy with Theorem 7. (We refer to [43] or the review [42] for a precise statement of this result.) Since the coincidence between the monstrous McKay-Thompson series and (normalized) Rademacher sums depends in a crucial way upon the genus zero property of monstrous moonshine, as evidenced by Theorem 6, it is natural to identify the Rademacher sum realization of the H_g as the Mathieu moonshine counterpart to the genus zero property of monstrous moonshine.

As we have hinted above, the Rademacher sum property that distinguishes the T_g and H_g does not hold for the weight zero Jacobi forms Z_g (cf. (9.12)). A Poincaré series approach to Jacobi forms is described in [24], using the foundations established in [27,28], and it is verified there that the Z_g are not all realized in this way. On the other hand, the main result of [24] is the Poincaré series construction of certain Maass-Jacobi forms of weight one, naturally associated to elements of M_{24} . Thus we can expect that Jacobi forms of weight one, rather than the Z_g of (9.12), will play an important role in a comprehensive conceptual explanation of the Mathieu moonshine phenomenon.

Note that some of the functions Z_g admit a geometric interpretation in terms of K3 surfaces. Namely, it has been established in [63] that if \bar{g} is a symplectic automorphism of a K3 surface M then the natural \bar{g} -equivariant modification of (9.1) coincides with Z_g , for a suitable element $g \in M_{24}$. However, not all Z_g arise in this way. Please see Section 9.4 for a fuller discussion of this.

9.2 Niemeier lattices

Vector-valued versions of the Rademacher sums that characterize the H_g were used in [47] to identify Mathieu moonshine as a special case of six directly similar correspondences, between conjugacy classes in certain finite groups and distinguished (vector-valued) mock modular forms of weight $1/2$. Since the mock modular forms arising seemed to be characterized by their shadows, this was dubbed *umbral moonshine* in [47].

The conjectures of [47] were greatly expanded in [48], following an observation of Glauberman (cf. the Acknowledgement in [48]), that the finite groups identified in [47] also appear as automorphism groups of codes associated to deep holes in the Leech lattice (cf. [59] or [56]).

To explain the significance of this, recall that an *integral lattice* is a free abelian group L together with a symmetric bilinear form $\langle \cdot, \cdot \rangle : L \times L \rightarrow \mathbb{Z}$. A lattice L is called *positive-definite* if $\langle \lambda, \lambda \rangle \geq 0$ for all $\lambda \in L$, with equality only when $\lambda = 0$. It is called *even* if $\langle \lambda, \lambda \rangle \in 2\mathbb{Z}$ for all $\lambda \in L$, and *self-dual* if $L = L^*$, for L^* the *dual* of L ,

$$L^* := \{ \mu \in L \otimes_{\mathbb{Z}} \mathbb{Q} \mid \langle \lambda, \mu \rangle \in \mathbb{Z} \iff \lambda \in L \}. \tag{9.14}$$

The even self-dual positive-definite lattices of rank 24 have been classified [189] (see also [60,227]) by Niemeier: there are 24 in total, up to isomorphism. They are characterized by their root systems - i.e., the configurations of their vectors with square length equal to 2 - and the Leech lattice Λ (cf. Section 3) is the unique such lattice whose root

system is empty. We refer to the remaining 23 as the *Niemeyer lattices*. The *Niemeyer root systems* are the root systems of the Niemeier lattices, and they are described explicitly as

$$A_1^{24}, A_2^{12}, A_3^8, A_4^6, A_6^4, A_{12}^2, \tag{9.15}$$

$$A_5^4 D_4, A_7^2 D_5^2, A_8^3, A_9^2 D_6, A_{11} D_7 E_6, A_{15} D_9, A_{17} E_7, A_{24}, \\ D_4^6, D_6^4, D_8^3, D_{10} E_7^2, D_{12}^2, D_{16} E_8, D_{24}, E_6^4, E_8^3, \tag{9.16}$$

in terms of the irreducible, simply-laced (i.e., ADE type) root systems. (See [62] or [139] for more on root systems.)

In (9.15) and (9.16) we use juxtaposition as a shorthand for direct sum, so that A_1^{24} denotes 24 copies of the A_1 root system, and $A_{11} D_7 E_6$ is shorthand for $A_{11} \oplus D_7 \oplus E_6$, etc. The subscripts indicate ranks. The Coxeter numbers of the ADE root systems are given by

$$m(A_n) = n + 1, m(D_n) = 2n - 2, m(E_6) = 12, m(E_7) = 18, m(E_8) = 30, \tag{9.17}$$

and one can check that the Niemeier root systems (9.15), (9.16) are exactly those unions of ADE type root systems for which the total rank is 24, and the Coxeter number is constant across irreducible components.

For X a Niemeier root system and N^X the corresponding Niemeier lattice, define the *outer automorphism group* of N^X by setting

$$\text{Out}(N^X) := \text{Aut}(N^X)/W^X, \tag{9.18}$$

where W^X denotes the subgroup of $\text{Aut}(N^X)$ generated by reflections in root vectors. Applying this construction to the Leech lattice, corresponding to $X = \emptyset$, we obtain the *Conway group*,

$$Co_0 := \text{Aut}(\Lambda), \tag{9.19}$$

so named in light of Conway’s detailed description [52,54] of its structure. A number of the 26 sporadic simple groups appear as subgroups, or quotients of subgroups of Co_0 , including the three sporadic simple Conway groups, Co_1 , Co_2 , and Co_3 . The Conway group Co_0 is a double cover of the first, and largest of these,

$$Co_1 \simeq \text{Aut}(\Lambda)/\{\pm \text{Id}\}. \tag{9.20}$$

Note that M_{24} is naturally a subgroup of Co_0 , and also Co_1 , for if $\{\lambda_i\} \subset \Lambda$ is a set of 24 vectors such that $\langle \lambda_i, \lambda_j \rangle = 8\delta_{ij}$, then the subgroup of Co_0 that stabilizes this set $\{\lambda_i\}$ is a copy of M_{24} .

$$M_{24} \simeq \{g \in Co_0 \mid \{g(\lambda_i)\} \subset \{\lambda_i\}\}. \tag{9.21}$$

According to Conway-Parker-Sloane [59], the Niemeier root systems classify the *deepest holes* in the Leech lattice, being the points in $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ at maximal distance from vectors in Λ . Moreover, this correspondence is strong enough that the Niemeier outer automorphism groups $\text{Out}(N^X)$ are also visible inside the Conway group, Co_0 . More precisely, if $x \in \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ is a deep hole, with corresponding Niemeier root system X according to [59], then the stabilizer $\text{Aut}(\Lambda, x)$ of x in $\text{Aut}(\Lambda)$ has a normal subgroup C^x such that

$$\text{Aut}(\Lambda, x)/C^x \simeq \text{Out}(N^X). \tag{9.22}$$

The subgroup C^x even encodes a method for constructing N^X , as is explained in detail in [61], for if L^X denotes the sub lattice of N^X generated by roots, then N^X is determined by its image in $(L^X)^*/L^X$ (cf. (9.14)) under the natural map $N^X \rightarrow (L^X)^*/L^X$. Write C^x for this subgroup of $(L^X)^*/L^X$, called the *glue code* of X in [61] (see also [60]). Then C^x is isomorphic to C^X , according to [61].

$$1 \rightarrow C^X \simeq C^x \rightarrow \text{Aut}(\Lambda, x) \rightarrow \text{Out}(N^X) \rightarrow 1 \tag{9.23}$$

Thus $\text{Out}(N^X)$ acts as automorphisms on the glue code C^X , and Glauber’s observation suggests an extension of the results of [47], whereby distinguished vector-valued mock modular forms $H_g^X = (H_{g,r}^X)$ are associated to elements g in the *umbral groups*

$$G^X := \text{Out}(N^X), \tag{9.24}$$

for each Niemeier root system X . The realization of this suggestion is described in detail in [48]. For $X = A_1^{24}$, the glue code C^X is a copy of the *extended binary Golay code* (cf. [62] or [202]), and G^X is its full automorphism group, M_{24} . Thus, from the Niemeier root system perspective, Mathieu moonshine is the special case of umbral moonshine corresponding to the root system A_1^{24} .

In (9.15) we have separated out the Niemeier root systems of the form A_n^d with $d = 24/n$ even. It is exactly these cases of umbral moonshine that are discussed in [47]. The original umbral moonshine observation of Eguchi-Ooguri-Tachikawa stemmed from consideration of the weight zero, index one weak Jacobi form Z_M (cf. (9.4)), realized as the K3 elliptic genus. The analysis of [47] is, to some extent, similarly motivated, including the attachment of a weight zero, index n weak Jacobi form $Z_g^{(n+1)}(\tau, z)$ to each $g \in G^X$, for each Niemeier root system $X = A_n^d$ with $d = 24/n$ even.

A notion of *extremal Jacobi form* is formulated in [47], motivated by the representation theory of the $N = 4$ superconformal algebra, and it is proven^k there that the six functions $Z^{(n+1)} := Z_e^{(n+1)}$, for $n \in \{1, 2, 3, 4, 6, 12\}$, exhaust all examples. Thus the cases (9.15) of umbral moonshine considered in [47] are distinguished from the point of view of Jacobi forms of weight zero.

By contrast, there seems to be no natural way to associate weight zero Jacobi forms to the Niemeier root systems not^l of the *pure A-type*, A_n^d . Rather, the mock modular forms H_g^X described in [48] naturally appear as the *theta-coefficients of finite parts* of certain meromorphic Jacobi forms ψ_g^X of weight 1 and index m ,

$$\psi_g^X(\tau, z) = \psi_g^{X,P}(\tau, z) + \sum_{r \pmod{2m}} H_{g,r}^X(\tau)\theta_{m,r}(\tau, z), \tag{9.25}$$

where $m = m(X)$ is the Coxeter number of any irreducible component of X (cf. (9.17)).

Here, meromorphic means that we allow poles in the functions $z \mapsto \psi_g^X(\tau, z)$, at torsion points $z \in \mathbb{Q}\tau + \mathbb{Q}$. The Weierstrass \wp function

$$\wp(\tau, z) := \frac{1}{z^2} + \sum_{\substack{\omega \in \mathbb{Z}\tau + \mathbb{Z} \\ \omega \neq 0}} \frac{1}{(z + \omega)^2} - \frac{1}{\omega^2} \tag{9.26}$$

is a natural example (with weight two and index zero).

The decomposition (9.25) is described in detail in [48], following the general structural results on meromorphic Jacobi forms established in [66,238]. For now let us just mention that the first summand on the right-hand side is the *polar part* of ψ_g^X , defined as in §8.2 of [66], and

$$\theta_{m,r}(\tau, z) := \sum_{k \in \mathbb{Z}} y^{2km+r} q^{(2km+r)^2/4m} \tag{9.27}$$

evidently depends only on r modulo $2m$.

A number of the meromorphic Jacobi forms attached to Niemeier root systems by umbral moonshine also appear amongst the specific examples of [66], where the main application is the computation of quantum degeneracies of black holes in certain string theories. However, whilst some speculations are offered in §5.5 of [47], no direct relationship between umbral moonshine and string theory has been formulated as yet.

We have seen in Section 9.1 that the mock modular forms attached to M_{24} by Mathieu moonshine (i.e., umbral moonshine for $X = A_1^{24}$) may be characterized as Rademacher sums, and this serves as an umbral analogue of the principal modulus/genus zero property of monstrous moonshine, on the strength of Theorem 6. It is natural to ask for an extension of this result to all cases of umbral moonshine.

Conjecture 5.4 of [47] amounts to the prediction that vector-valued generalizations of the Rademacher sums of [43] will recover the H_g^X for $X = A_n^d$ with d even (cf. (9.15)), and Conjecture 3.2 of [49] is an extension of this to all Niemeier root systems X . Thus a positive solution to Conjecture 3.2 of [49] will verify the umbral analogue of the principal modulus property of monstrous moonshine. So far, the Rademacher sum conjecture for umbral moonshine is known to be true only in the case that $X = A_1^{24}$, but a program to analyze the Rademacher sum conjecture for more general cases of umbral moonshine, via the theory of Maass-Jacobi forms (cf. [27,28]), has been initiated in [24].

A notion of optimal growth was formulated in §6.3 of [48], following the work [66], with a view to extending Conjecture 5.4 of [47]. It is now known that this condition does not uniquely determine the H_g^X for general X (see [49] for a full discussion of this), but all the H_g^X serve as examples. With this in mind, it is interesting to note that many of Ramanujan’s mock theta functions [203,204] appear as components of the umbral McKay-Thompson series H_g^X (cf. §4.7 of [47] and §5.4 of [48]).

9.3 Modules

As we have explained above, the Rademacher sum property of umbral moonshine is a natural counterpart to the principal modulus, or genus zero property of monstrous moonshine (cf. Section 3), formulated in a detailed way by Conway-Norton [58].

The natural counterpart to Thompson’s conjecture, Conjecture 1, verified by the Frenkel-Lepowsky-Meurman construction [102-104] of the moonshine module V^\natural , together with Borcherds’ work [18], is the following (cf. §6.1 of [48], and §2 of [49]).

Conjecture 5 (Cheng-Duncan-Harvey). For each Niemeier root system X , there is a bi-graded G^X -module

$$\check{K}^X = \bigoplus_{\substack{r \in I^X \\ D=r^2}} \bigoplus_{\substack{D \in \mathbb{Z} \\ (\text{mod } 4m)}} \check{K}_{r,-D/4m}^X, \tag{9.28}$$

such that the vector-valued umbral McKay-Thompson series $H_g^X = (H_{g,r}^X)$ is recovered^m from the graded trace of g on \check{K}^X via

$$H_{g,r}^X(\tau) = -2q^{-1/4m}\delta_{r,1} + \sum_{\substack{D \in \mathbb{Z} \\ D \equiv r^2 \pmod{4m}}} \text{tr} \left(g | \check{K}_{r,-D/4m}^X \right) q^{-D/4m} \tag{9.29}$$

for $r \in I^X$.

In (9.28) and (9.29), $m = m(X)$ is the Coxeter number of any irreducible component of X , as in (9.25). The $H_{g,r}^X$ satisfy $H_{g,-r}^X = -H_{g,r}^X$, so the umbral McKay-Thompson series H_g^X is determined by its components $H_{g,r}^X$ with $0 < r < m$. If the highest rank irreducible component of X is of type D or E then there are more symmetries amongst the $H_{g,r}^X$, and the definition of the set $I^X \subset \mathbb{Z}/2m\mathbb{Z}$ reflects this: if X has an A-type component then $I^X := \{1, 2, 3, \dots, m - 1\}$. If the highest rank component of X is of type D then $m = 2 \pmod{4}$, and $I^X := \{1, 3, 5, \dots, m/2\}$. The remaining cases are $X = E_6^4$, in which case $I^X := \{1, 4, 5\}$, and $X = E_8^3$, for which $I^X := \{1, 7\}$.

As mentioned in Section 9.1, the existence of the module \check{K}^X for $X = A_1^{24}$ has been proven by Gannon [117]. More specifically, Gannon has shown that the coefficients of the non-negative powers of q in H_g^X for $X = A_1^{24}$ are traces of elements of M_{24} on direct sums of irreducible M_{24} -modules. A priori, we might have needed \mathbb{C} -linear combinations of such traces in order to recover the $H_{g,r}^X$.

In forthcoming work [85], the authors confirm the validity of Conjecture 5.

Theorem 16 (Duncan-Griffin-Ono). *Conjecture 5 is true.*

Theorem 16 serves, to a certain extent, as the umbral counterpart to Borcherds' result, Theorem 4. Indeed, the method of [85] may be used to give an alternative proof of the existence of the \mathbb{M} -module V^\natural , for which the associated graded trace functions are the normalized principal moduli of the genus zero groups Γ_g , identified by Conway-Norton in [58].

Nonetheless, there is still work to be done; for in order to have a direct counterpart to Theorem 4, we require concrete constructions of the \check{K}^X . In the case of monstrous moonshine, the construction of V^\natural due to Frenkel-Lepowsky-Meurman came equipped with rich algebraic structure, ultimately leading to the notion of vertex operator algebra, and powerful connections to physics. We can expect that a full explanation of the umbral moonshine phenomena will require analogues of this for all the \check{K}^X .

Just such an analogue for $X = E_8^3$ has recently been obtained in [86], where a super vertex operator algebra V^X is constructed, together with an action of $G^X \simeq S_3$, such that the components of the vector-valued mock modular forms $H_g^X = (H_{g,r}^X)$ are recovered from traces of elements of G^X on canonically twisted modules for V^X . The main ingredient in the construction of [86] is an adaptation of the familiar (to specialists) lattice vertex algebra construction (cf. [15,104]), to cones in indefinite lattices. The choice of cone is in turn inspired by Zwegers' work [239] on a particular pair of the fifth order mock theta functions of Ramanujan.

In [46,89] a different approach to the module problem is considered, whereby meromorphic Jacobi forms associated to the H_g^X are recovered as graded traces on canonically twisted modules for certain super vertex algebras. In [89] constructions are given for the

ψ_g^X of (9.25), for $X \in \{A_3^8, A_4^6, A_6^4, A_{12}^2\}$. In [46] certain half-integral index analogues of the ψ_g^X are recovered, for $X \in \{D_6^4, D_8^3, D_{12}^2, D_{24}\}$.

As we will explain in more detail in the next section, recent work [40,51] constructs modules underlying assignments of vector-valued mock modular forms to the sporadic simple groups M_{24} , M_{23} , and M_{22} . Here, M_{23} denotes the maximal subgroup of M_{24} composed of elements fixing any given point in the defining permutation representation (cf. Section 9.1), and M_{22} is obtained similarly from M_{23} , as the subgroup stabilizing a point in its natural permutation representation of degree 23. Although the mock modular forms realized in [40,51] are not directly related to the H_g^X , it seems likely that the construction used therein holds important hints for future developments in the module problem for umbral moonshine.

9.4 Sigma models

Recall from the Section 9.1 that $\Omega_M^2 \simeq \mathcal{O}_M$ for M a complex K3 surface. An automorphism of M that induces the trivial action on $H^0(M, \Omega_M^2)$ is called *symplectic*. It is a celebrated result of Mukai [186] (cf. also [162]), that the finite groups of symplectic automorphisms of complex K3 surfaces are, up to isomorphism, precisely the subgroups of M_{24} that have at least five orbits in the unique non-trivial permutation representation on 24 points, including at least one fixed point.

Since a symplectic automorphism of a complex K3 surface M induces a supersymmetry preserving automorphism of a sigma model attached to M (cf. Section 9.1), and since it is the supersymmetry preserving automorphisms of a K3 sigma model that can be used to twine the K3 elliptic genus (9.4), the problem of classifying the supersymmetry preserving automorphism groups of nonlinear K3 sigma models was considered by Gaberdiel-Hohenegger-Volpato in [109].

One might have anticipated that all supersymmetry preserving K3 sigma model automorphism groups would be contained in M_{24} , but this is not the case. Rather, the main result of [109], being a quantum analogue of Mukai’s classification of finite symplectic automorphism groups of K3 surfaces, is that the groups of supersymmetry preserving automorphisms of K3 sigma models are, up to isomorphism, precisely the subgroups of $Co_0 = \text{Aut}(\Lambda)$ (cf. (9.19)) that fix a sublattice of Λ with rank at least four.

Note that the results of [109] are obtained subject to certain conjectural assumptions about the structure of the moduli space of K3 sigma models. Nonetheless, it seems fair to conclude that the K3 sigma models do not furnish quite the right theoretical setting for solving the mysteries of umbral moonshine. For not all of the M_{24} -twinnings (9.12) of the K3 elliptic genus (9.4) arise as twinings defined by K3 sigma model automorphisms, since, for example, there are elements of M_{24} (cf. (9.21)) that do not fix a rank four lattice in the Leech lattice, Λ (cf. also, the last sentence of Section 9.1).

That notwithstanding, we can expect to learn useful information about umbral moonshine from further investigation of K3 sigma models. The history of monstrous moonshine provides a useful point of comparison: in advance of his proof of the Conway-Norton conjectures, Borcherds considered a certain BKM algebra (cf. Section 3) in [17], which was, at the time, called the monster Lie algebra, although it turned out to be only indirectly connected to the monster. The Lie algebra constructed in [17] is now known as the *fake monster Lie algebra* (cf. §2 of [18]) and has found a number of applications outside of moonshine. For example, the denominator function of the fake monster Lie

algebra (cf. Example 2 in §10 of [19]) is used to prove facts about families of K3 surfaces in [21,163,235].

At the level of vertex operator algebras, the fake monster Lie algebra corresponds to the lattice vertex algebra V_Λ attached to the Leech lattice. This may be regarded as a ‘fake’ moonshine module, for it has exactly the same graded dimension as V^\natural , up to the constant term,

$$\dim_* V_\Lambda = J(\tau) + 24. \tag{9.30}$$

(Cf. (4.1).) There is no action of the monster on V_Λ , although there is an action by a groupⁿ of the shape $2^{24}.Co_0 = 2^{24}.(2.Co_1)$ (cf. (9.20)), whereas the monster contains a maximal subgroup with the shape $(2.2^{24}).Co_1$. The Frenkel-Lepowsky-Meurman construction of V^\natural takes V_Λ as a main ingredient (cf. (3.6)).

It is striking that the Conway group $Co_0 = 2.Co_1$ plays a prominent role in so many of the objects we have discussed: it is visible within the monster, and within the automorphism group of V_Λ . It serves for K3 sigma models as M_{24} does for K3 surfaces, as discussed above, and all of the umbral moonshine groups (9.24) are visible within Co_0 , according to (9.22).

Moreover, there is moonshine for the Conway group, in direct analogy with that for the monster, in the sense that there is an assignment of normalized principal moduli T_g^s to elements $g \in Co_0$ which are realized as trace functions on a graded infinite-dimensional Co_0 -module. A proof of this statement has recently appeared in [87].

To explain this, take $g \in Co_0$, let $\{\varepsilon_i\}$ be the eigenvalues associated to the action of g on $\Lambda \otimes_{\mathbb{Z}} \mathbb{C}$, and define

$$T_g^s(\tau) := q^{-1/2} \prod_{n>0} \prod_{i=1}^{24} (1 - \varepsilon_i q^{n-1/2}) + \chi_g, \tag{9.31}$$

where

$$\chi_g := \sum_i \varepsilon_i \tag{9.32}$$

is the character value associated to the action of g on $\Lambda \otimes_{\mathbb{Z}} \mathbb{C}$. Then $T_g^s(2\tau) = q^{-1} + O(q)$ is the normalized principal modulus for a genus zero group, according to Conway-Norton [58] and Queen [198] (see also [164]).

It has been demonstrated in [87] that the functions T_g^s are the graded traces attached to the action of Co_0 on a distinguished^o super vertex operator algebra, $V^{sn} = \bigoplus_{n=-1}^{\infty} V_{n/2}^{sn}$.

$$T_g^s(\tau) = \sum_{n=-1}^{\infty} \text{tr} \left(\mathfrak{z}g|V_{n/2}^{sn} \right) q^{n/2} \tag{9.33}$$

(In (9.33) we write \mathfrak{z} for the super space involution, acting as $(-1)^n \text{Id}$ on $V_{n/2}^{sn}$.) Thus the super vertex operator algebra V^{sn} solves the Conway moonshine analogue of Thompson’s Conjecture 1, and V^{sn} is the natural analogue of the moonshine module V^\natural for the Conway group, Co_0 .

The Conway module V^{sn} is closely related to monstrous moonshine, for, in addition to being directly analogous to V^\natural , many of the discrete groups $\Gamma_g < \text{SL}_2(\mathbb{R})$, for $g \in \mathbb{M}$, also arise as invariance groups of principal moduli attached to Co_0 via its action on V^{sn} (cf. [87]). On the other hand, V^{sn} enjoys a close connection to K3 sigma models, for it

is shown in [88] that the data defining a K3 sigma model gives rise to a bi-grading on a distinguished, canonically-twisted^P V^{sq} -module

$$V_{tw}^{sq} = \bigoplus_{n,r} (V_{tw}^{sq})_{n,r}, \tag{9.34}$$

such that the associated graded traces of compatible elements of Co_0 are weak Jacobi forms.

More specifically, following §2.1 of [109], we may regard the data of a K3 sigma model as equivalent^Q to a choice of positive-definite 4-space $\Pi < \mathbb{I}_{4,20} \otimes_{\mathbb{Z}} \mathbb{R}$ (cf. (3.7)), such that

$$\delta \in \Pi^\perp \cap \mathbb{I}_{4,20} \implies \langle \delta, \delta \rangle \neq -2. \tag{9.35}$$

Then the supersymmetry preserving automorphism group of the nonlinear sigma model defined by Π is the group

$$G_\Pi := \text{Aut}(\mathbb{I}_{4,20}, \Pi), \tag{9.36}$$

composed of orthogonal transformations of $\mathbb{I}_{4,20}$ that extend to the identity on Π , according to §2.2 of [109]. One of the main results of [109] is that G_Π may be identified with a subgroup of Co_0 .

The construction of [88] uses V_{tw}^{sq} to attach a graded trace function

$$\phi_g(\tau, z) := - \sum_{n,r} \text{tr}(\mathfrak{z}g|(V_{tw}^{sq})_{n,r})q^n y^r \tag{9.37}$$

to each pair (g, Π) , where $\Pi < \mathbb{I}_{4,20} \otimes_{\mathbb{Z}} \mathbb{R}$ satisfies (9.35), and $g \in G_\Pi$, and \mathfrak{z} is a certain naturally defined involution on V_{tw}^{sq} (analogous to the \mathfrak{z} in (9.33)). It is shown in [88] that ϕ_g is a weak Jacobi form of weight zero and index one for $\Gamma_0^J(N)$ (cf. (9.12)), for some N , for all choices of Π and $g \in G_\Pi$. Moreover, ϕ_g is found to coincide with the g -twined K3 elliptic genus associated to the sigma model defined by Π , for all the examples computed in [109,113,228]. (These examples account for about half of the conjugacy classes of Co_0 that fix a 4-space in $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$.) In particular, taking $g = e$ in (9.37) recovers the K3 elliptic genus (9.4), but in the form

$$\phi_e(\tau, z) = -2^{11} \frac{\theta_2(\tau, z)^2}{\theta_2(\tau, 0)^2} \frac{\Delta(2\tau)}{\Delta(\tau)} + \frac{1}{2} \frac{\theta_3(\tau, z)^2}{\theta_3(\tau, 0)^2} \frac{\Delta(\tau)^2}{\Delta(2\tau)\Delta(\tau/2)} - \frac{1}{2} \frac{\theta_4(\tau, z)^2}{\theta_4(\tau, 0)^2} \frac{\Delta(\tau/2)}{\Delta(\tau)}, \tag{9.38}$$

where $\Delta(\tau) := \eta(\tau)^{24} = q \prod_{n=1}^\infty (1 - q^n)^{24}$.

Thus V_{tw}^{sq} serves as a kind of universal object for K3 sigma models. This is interesting, for generally it is difficult to construct the Hilbert spaces underlying a K3 sigma model, and therefore difficult to compute the associated twined K3 elliptic genera, for instance, for all but a few special examples.

In [140], Huybrechts has related the positive-definite 4-spaces $\Pi < \mathbb{I}_{4,20} \otimes_{\mathbb{Z}} \mathbb{R}$ satisfying (9.35) to pairs (X, σ) , where X is a projective complex K3 surface, and σ is a stability condition on the bounded derived category of coherent sheaves on X . In this way he has obtained an alternative analogue of Mukai’s result [186], whereby symplectic automorphisms of K3 surfaces are replaced by symplectic derived autoequivalences. (The results of [88] are formulated in this language.)

A number of the functions Z_g (cf. (9.12)) arising in Mathieu moonshine are realized as ϕ_g for some $g \in Co_0$. So the construction of [88] relates V^{sq} to umbral moonshine,

but the connection goes deeper, for it is shown in [88] that a natural generalization of the definition (9.37) recovers a number of the Jacobi forms attached to other root systems of the form $X = A_n^d$ (cf. Section 9.2), beyond the special case $X = A_1^{24}$. It is interesting to compare this to the results of [50] (see also the precursor [133]), which demonstrate a role for K3 surface geometry in all cases of umbral moonshine (i.e., for all the Niemeier root systems), by considering sigma models attached to K3 surfaces admitting du Val singularities. Indeed, many of the Jacobi forms computed in [50] appear also in [88].

From the discussion above, we see that the Conway module V^{st} provides evidence for a deep connection between monstrous and umbral moonshine. Further support for the notion that monstrous and umbral moonshine share a common origin is obtained in [195], where the generalized Borcherds products of [29] are used to relate the trace functions of monstrous and umbral moonshine directly.

We elaborate now upon the results of [40,51], which, as mentioned at the end of Section 9.3, fall outside of umbral moonshine as formulated in [48], but are nonetheless related. Actually, these works are further applications of the Conway moonshine module V^{st} , for in [40] the canonically twisted V^{st} -module $V_{\text{tw}}^{\text{st}}$ (cf. (9.34)) is equipped with module structures for the $N = 2$ and $N = 4$ superconformal algebras, which in turn give rise to an assignment of distinguished vector-valued mock modular forms to elements of the sporadic simple Mathieu groups M_{23} and M_{22} , respectively. This work furnishes the first examples of concretely constructed modules for sporadic simple groups, such that the associated graded trace functions define mock modular forms. The methods of [40] are extended in [51], to the case of the $Spin(7)$ algebra (a certain extension of the $N = 1$ superconformal algebra, cf. [13] and references therein) and vector-valued mock modular forms for M_{24} are obtained. Interestingly, Conway's sporadic groups Co_2 and Co_3 , and the sporadic groups of McLaughlin and Higman-Sims (all rather larger than M_{24} , or any of the other umbral groups G^X) also appear in the analysis of [40,51].

As a further indication of the important role that K3 sigma models will play in illuminating umbral moonshine, we mention the interesting work [112,217-219], which seeks to explain the Mathieu moonshine observation by formulating a precise mechanism for combining symmetries of distinct K3 sigma models into a single group. We note in particular, that a fixed-point-free maximal subgroup of M_{24} is constructed in this way in [219].

We conclude this section with references [41,132-134,155,196,233] to a number of other occurrences of umbral groups in geometry and physics, all promising connections to Mathieu moonshine, or umbral moonshine more generally. We also note the recent work [39], which analyzes all cases of generalized umbral moonshine, thereby extending the investigation of generalized Mathieu moonshine that was initiated in [111].

10 Open problems

We conclude the article by identifying some open problems for future research that are suggested by our results in Sections 7 and 8, and the developments described in Section 9.

Problem 1. We have seen that the known connections between monstrous moonshine and physics owe much to the Frenkel-Lepowsky-Meurman construction of the moonshine module V^{\natural} , and its associated vertex operator algebra structure. Just as the vertex operator algebra structure on V^{\natural} gives a strong solution to Thompson's conjecture, Conjecture 1, we can expect concrete constructions of the \check{K}^X - whose existence is now guaranteed thanks to Theorem 16 - to be necessary for the elucidation of the physical origins of umbral moonshine. As we have described in Section 9.3, progress on this problem has been obtained recently in [46,86,89], and the related work [40] may also be useful, in the determination of a general, algebraic solution to the module problem for umbral moonshine.

Problem 2. Norton's generalized moonshine conjectures were discussed in Section 4, and generalized umbral moonshine has been investigated in [39,111]. A special case of generalized moonshine for the Conway group is established in [87], but the full formulation and proof of generalized Conway moonshine remains open. Given the close connections between $V^{s^{\natural}}$ and umbral moonshine discussed in Section 9.4, it will be very interesting to determine the precise relationship between the corresponding generalized moonshine theories. We can expect that the elucidation of these structures will be necessary for a full understanding of the role that umbral moonshine plays in physics.

Problem 3. As discussed in Section 9.2, the fact (cf. Theorem 7) that the McKay-Thompson series of monstrous moonshine are realized as Rademacher sums admits conjectural analogues for umbral moonshine. (See Conjecture 3.2 of [49] for a precise formulation.) So far this has been established only for $X = A_1^{24}$, corresponding to Mathieu moonshine (cf. [43]), and the general case remains open. As explained in Section 9.2, a positive solution to Conjecture 3.2 of [49] will establish an umbral moonshine counterpart to the principal modulus/genus zero property of monstrous moonshine.

Problem 4. As discussed above, in Sections 5 and 9.2, Rademacher sums play a crucial role in both monstrous and umbral moonshine, by serving to demonstrate the distinguished nature of the automorphic functions arising in each setting. In the case of monstrous moonshine, the Rademacher sum property also indicates a potentially powerful connection to physics, via three-dimensional gravity, as explained in Section 6. Thus it is an interesting problem to formulate umbral moonshine analogues of the conjectures of [84], discussed here in Sections 6 and 7.

Problem 5. Relatedly, it follows from the results of [84] that the McKay-Thompson series T_g^s (cf. (9.33)), attached to elements g in the Conway group Co_0 via its action on $V^{s^{\natural}}$ (cf. Section 9.4), are also realized as Rademacher sums. That is, Theorem 7 generalizes naturally to Conway moonshine. Thus it is natural to investigate the higher order analogues $V^{s^{(-m)}}$ of the super vertex operator algebra $V^{s^{\natural}}$, and the Conway group analogues of the three-dimensional gravity conjectures of [84]. Some perspectives on this are available in [137,178,232].

Problem 6. The notion of extremal vertex operator algebra is defined by (6.8). So far the only known example is the moonshine module V^{\natural} . As explained in Section 6, the construction of a series of extremal vertex operator algebras, with central charges the positive integer multiples of 24, would go a long way towards the construction of a chiral three-dimensional quantum gravity theory. This problem also has a super analogue, cf. [137].

Problem 7. The monster modules $V^{(-m)}$, defined in Section 7, cannot be vertex operator algebras for $m > 1$, for an action of the Virasoro algebra would generate a non-zero vector with non-positive eigenvalue for $L(0) - c/24$, and this would violate the condition $\sum_n \dim(V_n^{(-m)}) q^n = q^{-m} + O(q)$. Nonetheless, we may ask: do the $V^{(-m)}$ admit vertex algebra structure? Or, is there another natural algebraic structure, which characterizes the monster group actions on the $V^{(-m)}$?

Problem 8. Relatedly, $\mathcal{V} \times \mathbb{M}$ -modules satisfying the extremal condition (6.8) may be easily constructed from the monster modules $V^{(-m)}$, as is mentioned at the conclusion of Section 7. What is the algebraic significance of these spaces? We know from [114,137] that they cannot admit vertex operator algebra structure compatible with the given $\mathcal{V} \times \mathbb{M}$ actions. Is there some other kind of algebraic structure which is compatible with this symmetry?

Problem 9. The result of Corollary 1 implies that the $V_n^{(-m)}$ and W_n^{\natural} tend to direct sums of copies of the regular representation of \mathbb{M} , as $n \rightarrow \infty$. This means that if we write each homogeneous subspace of each module, particularly the moonshine module V^{\natural} , as the sum of a free part (free over the group ring of \mathbb{M}) and a non-free part, then the non-free part tends to 0 (relative to the free part) as $n \rightarrow \infty$. Is there something to be learnt from an analysis of the non-free parts of $V^{(-m)}$, W^{\natural} ? As one can see from Table 1, some irreducible representations of the monster feature more often in the non-free part than others. We thank Bob Griess for posing this question.

Problem 10. It is a natural problem to generalize the methods employed in Section 8, to determine the distributions of irreducible representations of the umbral groups G^X (cf. (9.24)) in the umbral moonshine modules \check{K}^X (cf. (9.28)). Similarly, one may also consider the distributions of the Co_0 -modules in $V^{s^{\natural}} = V^{s^{(-1)}}$, and in the $V^{s^{(-m)}}$ more generally. In all of these cases, questions analogous to Problem 9 may reward investigation.

Problem 11. In Corollary 2 we have used our asymptotic results (Theorem 9) on multiplicities of monster modules inside V^{\natural} to compute the quantum dimensions of the monster orbifold, and in so doing confirmed a special case of Conjecture 6.7 of [75]. How generally can this method be applied, to orbifolds V^G , where V is a vertex operator algebra and G is a compact group of automorphisms of V ? Note the following strengths of the asymptotic approach: thanks to Proposition 3.6 of [75], we did not need to verify that the monster orbifold of V^{\natural} is a rational vertex operator algebra, nor did we need to assume the positivity condition of Theorem 6.3 in [75] (which in any case does not hold for V^{\natural}).

Endnotes

^aWe refer the reader to [90] for a recent analysis of the Jack Daniels problem.

^bAs was pointed out to us by J.-P. Serre, the decomposition for 20245856256 that appears in [221] differs from that which develops from the monster's natural, infinite-dimensional representation. We reproduce Serre's decomposition in (2.5).

^cThe relevant groups Γ_g are discussed in detail in Section 8.1.

^dA principal modulus is also referred to as a *Hauptmodul*.

^eMaldacena's groundbreaking paper [175] on the gauge/gravity duality has over 10,000 citations at the time of writing, according to inspirehep.net.

^fWe regard all vertex operator algebras as graded by $L(0) - c/24$. Cf. (3.5).

^gThe existence of extremal vertex operator algebras with central charge $c = 24m$ for $m > 1$ remains an open question. We refer to [110,115,137,234] for analyses of this problem.

^hWe are grateful to the referee for suggesting this.

ⁱSee [66] or [96] for more on Jacobi forms, including the general transformation formula in case of weight different from zero or index different from one.

^jThe notion of mock modular form has arisen recently, from Zwegers' foundational work [238] on Ramanujan's mock theta functions [203,204], and the subsequent contributions [25] and [236]. We refer to [66,194,236] for nice introductions to the theory.

^kThe main step in the classification given in [47] is a demonstration that the existence of an extremal Jacobi form of index $m - 1$ implies the vanishing of $L(f, 1)$ for all new forms f of weight 2 and level m , where $L(f, s)$ is the Dirichlet series naturally attached to f (cf. e.g. §3.6 of [211]). At this point one expects extremal Jacobi forms to be very few in number, on the strength of the Birch-Swinnerton-Dyer conjecture (cf. [14,229]), for example. This machinery is evidently quite powerful, and we may anticipate further applications to umbral moonshine in the future.

^lThe cases A_8^3 and A_{24} do come with weight zero Jacobi forms attached, which are obtained via a slight weakening of the notion of extremal Jacobi form formulated in [47]. Cf. §4.3 of [48].

^mIn the original formulation, Conjecture 6.1 of [48], the function $H_{g,r}^X$ in (9.29) is replaced with $3H_{g,r}^X$ in the case that $X = A_8^3$. It also predicted that $\check{K}_{r,-D/4m}^X$ is a virtual G^X -module in case $X = A_8^3$ and $D = 0$. Recently, a modification of the specification of the H_g^X for $X = A_8^3$ has been discovered, which leads to the simpler, more uniform formulation appearing here. We refer to [49] for a full discussion of this.

ⁿGiven groups A and B , say that a group G has the *shape* $A.B$, and write $G = A.B$, if G contains a normal subgroup isomorphic to A such that $G/A \simeq B$. In this setting it is typical to write p^n as a shorthand for an elementary abelian p -group with p^n elements.

^oThe super vertex operator algebra V^{st} admits actions by both Co_0 (cf. (9.19)) and the simple group Co_1 (cf. (9.20)). Its construction as a Co_1 -module was sketched first in §15 of [103], described later in §5 of [22], and subsequently studied in detail in [83]. The Co_0 -module structure on V^{st} is mentioned in [83], following [22], but it seems that the modular properties of the trace functions associated to the Co_0 -action were not considered until [87].

^pTwisted modules for vertex algebras are discussed in Section 4. In the case of a super vertex algebra there is a canonically defined involution coming from the superspace structure, called the *parity involution*. A module for a super vertex algebra that is twisted with respect to the parity involution is called *canonically-twisted*.

^qThis convention excludes some interesting K3 sigma models, such as those considered in [50]. We refer to [3,4] for detailed discussions of K3 sigma model moduli.

Appendix A Monstrous groups

The table below contains the symbols $\Gamma_g = N||h + e, f, \dots$, for each conjugacy class of the monster. Following [57], if $h = 1$, we omit the ‘||1’ from the symbol. If $\mathcal{W}_g = \{1\}$, then we write $N||h$, whereas if it contains every exact divisor of N/h , we write $N||h+$.

The naming of the conjugacy classes is as in [56]. We follow the convention of writing $23AB$ as a shorthand for $23A \cup 23B$, since these conjugacy classes are related by inversion in the monster. (There are 22 such pairs. The Monster group has 196 conjugacy classes in total.) Since the monstrous McKay-Thompson series have real coefficients, $T_g = T_{g^{-1}}$ and $\Gamma_g = \Gamma_{g^{-1}}$ for all g in the monster. Note however that $27A$ and $27B$ are not related by inversion, even though $\Gamma_{27A} = \Gamma_{27B}$. To the authors best knowledge, this coincidence has not yet been explained.

1A	1	11A	11+	20D	20 2 + 5
2A	2+	12A	12+	20E	20 2 + 10
2B	2	12B	12 + 4	20F	20 + 20
3A	3+	12C	12 2+	21A	21+
3B	3	12D	12 3+	21B	21 + 3
3C	3 3	12E	12 + 3	21C	21 3+
4A	4+	12F	12 2 + 6	21D	21 + 21
4B	4 2+	12G	12 2 + 2	22A	22+
4C	4	12H	12 + 12	22B	22 + 11
4D	4 2	12I	12	23AB	23+
5A	5+	12J	12 6	24A	24 2+
5B	5	13A	13+	24B	24+
6A	6+	13B	13	24C	24 + 8
6B	6 + 6	14A	14+	24D	24 2 + 3
6C	6 + 3	14B	14 + 7	24E	24 6+
6D	6 + 2	14C	14 + 14	24F	24 4 + 6
6E	6	15A	15+	24G	24 4 + 2
6F	6 3	15B	15 + 5	24H	24 2 + 12
7A	7+	15C	15 + 15	24I	24 + 24
7B	7	15D	15 3	24J	24 12
8A	8+	16A	16 2+	25A	25+
8B	8 2+	16B	16	26A	26+
8C	8 4+	16C	16+	26B	26 + 26
8D	8 2	17A	17+	27A	27+
8E	8	18A	18 + 2	27B	27+
8F	8 4	18B	18+	28A	28 2+
9A	9+	18C	18 + 9	28B	28+
9B	9	18D	18	28C	28 + 7
10A	10+	18E	18 + 18	28D	28 2 + 14
10B	10 + 5	19A	19+	29A	29+
10C	10 + 2	20A	20+	30A	30 + 6, 10, 15
10D	10 + 10	20B	20 2+	30B	30+
10E	10	20C	20 + 4	30C	30 + 3, 5, 15

30D	30 + 5, 6, 30	42B	42 + 6, 14, 21	62AB	62+
30E	30 3 + 10	42C	42 3 + 7	66A	66+
30F	30 + 2, 15, 30	42D	42 + 3, 14, 42	66B	66 + 6, 11, 66
30G	30 + 15	44AB	44+	68A	68 2+
31AB	31+	45A	45+	69AB	69+
32A	32+	46AB	46 + 23	70A	70+
32B	32 2+	46CD	46+	70B	70 + 10, 14, 35
33A	33 + 11	47AB	47+	71AB	71+
33B	33+	48A	48 2+	78A	78+
34A	34+	50A	50+	78BC	78 + 6, 26, 39
35A	35+	51A	51+	84A	84 2+
35B	35 + 35	52A	52 2+	84B	84 2 + 6, 14, 21
36A	36+	52B	52 2 + 26	84C	84 3+
36B	36 + 4	54A	54+	87AB	87+
36C	36 2+	55A	55+	88AB	88 2+
36D	36 + 36	56A	56+	92AB	92+
38A	38+	56BC	56 4 + 14	93AB	93 3+
39A	39+	57A	57 3+	94AB	94+
39B	39 3+	59AB	59+	95AB	95+
39CD	39 + 39	60A	60 2+	104AB	104 4+
40A	40 4+	60B	60+	105A	105+
40B	40 2+	60C	60 + 4, 15, 60	110A	110+
40CD	40 2 + 20	60D	60 + 12, 15, 20	119AB	119+
41A	41+	60E	60 2 + 5, 6, 30		
42A	42+	60F	60 6 + 10		

Acknowledgements

The authors are supported by the NSF. The first author also thanks the Simons Foundation (#316779), and the third author also thanks the support of the Asa Griggs Candler Fund. The authors thank John McKay and John Thompson for answering questions about the history related to the study of distributions within the moonshine module. Thanks are due also to Miranda Cheng, Igor Frenkel, Bob Griess, Daniel Grumiller, Jeff Harvey, Shamit Kachru, Barry Mazur, John McKay, Jean-Pierre Serre and the anonymous referee, for comments, corrections and excellent suggestions.

Author details

¹Department of Mathematics, Applied Mathematics and Statistics, Case Western Reserve University, 10900 Euclid Avenue, Cleveland, Ohio 44106, USA. ²Department of Mathematics and Computer Science, Emory University, 201 Dowman Drive, Atlanta, Georgia 30322, USA.

Received: 11 November 2014 Accepted: 28 April 2015

Published online: 10 July 2015

References

- Alexander, D, Cummins, C, McKay, J, Simons, C: Completely replicable functions, Groups, combinatorics & geometry (Durham, 1990), London Math. Soc. Lecture Note Ser., Vol. 165. Cambridge Univ. Press, Cambridge (1992). pp. 87–98. MR 1200252 (94g:11029)
- Aschbacher, M: The status of the classification of the finite simple groups. *Notices Amer. Math. Soc.* **51**(7), 736–740 (2004). MR 2072045
- Aspinwall, PS: $K3$ surfaces and string duality. *Fields, strings and duality* (Boulder, CO, 1996). World Sci. Publ., River Edge, NJ (1997). pp. 421–540. MR 1479699 (98j:81243)
- Aspinwall, PS, Morrison, DR: String theory on $K3$ surfaces, Mirror symmetry, II. *AMS/IP Stud. Adv. Math.*, vol. 1, Amer. Math. Soc., Providence, RI, 703–716 (1997). MR 1416354 (97i:81128)
- Astashkevich, A: On the structure of Verma modules over Virasoro and Neveu-Schwarz algebras. *Comm. Math. Phys.* **186**(3), 531–562 (1997). MR 1463812 (98h:17028)
- Atiyah, MF: K -theory. second ed. *Advanced Book Classics*, Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA (1989). Notes by D. W. Anderson. MR 1043170 (90m:18011)
- Bagchi, A, Detournay, S, Grumiller, D: Flat-space chiral gravity. *Phys. Rev. Lett.* **109**, 151301 (2012)

8. Bañados, M, Henneaux, M, Teitelboim, C, Zanelli, J: Geometry of the $2+1$ black hole. *Phys. Rev. D.* **48**(3 no. 4), 1506–1525 (1993). MR 1236812 (94g:83079)
9. Bañados, M, Teitelboim, C, Zanelli, J: Black hole in three-dimensional spacetime. *Phys. Rev. Lett.* **69**(3), 1849–1851 (1992). MR MR1181663 (93e:83042)
10. Barth, WP, Hulek, K, Peters, CAM, Van de Ven, A: Compact complex surfaces. second ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, Vol. 4. Springer-Verlag, Berlin (2004). MR 2030225 (2004m:14070)
11. Beauville, A, Bourguignon, JP, Demazure, M, Gauduchon, P, Méridol, JY, Potier, JL, Risler, JJ: Géométrie des surfaces $K3$: modules et périodes. Société Mathématique de France, Paris, 1985, Papers from the seminar held in Palaiseau, October 1981–January 1982, Astérisque No. 126 (1985). MR785216 (87h:32052)
12. Beneish, L, Larson, H: Traces of singular values of Hauptmoduln. *Int. J. Number Th.* **11**(3), 1027–1048 (2015). <http://dx.doi.org/10.1142/S1793042115500542> accepted for publication
13. Benjamin, N, Harrison, SM, Kachru, S, Paquette, NM, Whalen, D: On the elliptic genera of manifolds of Spin (7) holonomy (2014). ArXiv e-prints
14. Birch, BJ: HPF. Swinnerton-Dyer, Notes on elliptic curves. II. *J. Reine Angew. Math.* **218**, 79–108 (1965). MR 0179168 (31 #3419)
15. Borcherds, R: Vertex algebras, Kac-Moody algebras, and the Monster. *Proc. Nat. Acad. Sci. USA.* **83**(10), 3068–3071 (1986)
16. Borcherds, R: Generalized Kac-Moody algebras. *J. Algebra.* **115**(2), 501–512 (1988). MR 943273 (89g:17004)
17. Borcherds, R: The monster Lie algebra. *Adv. Math.* **83**(1), 30–47 (1990). MR 1069386 (91k:17027)
18. Borcherds, R: Monstrous moonshine and monstrous Lie superalgebras. *Invent Math.* **109**(2), 405–444 (1992)
19. Borcherds, R: Automorphic forms with singularities on Grassmannians. *Invent. Math.* **132**(3), 491–562 (1998). MR 1625724 (99c:11049)
20. Borcherds, R: Quantum vertex algebras. In: Taniguchi Conference on Mathematics Nara '98, *Adv. Stud. Pure Math.*, vol. 31, Math. Soc., pp. 51–74, Japan, Tokyo, (2001). MR 1865087 (2002k:17054)
21. Borcherds, R, Katzarkov, L, Pantev, T, Shepherd-Barron, NI: Families of $K3$ surfaces. *J. Algebraic Geom.* **7**(1), 183–193 (1998). MR 1620702 (99d:14029)
22. Borcherds, RA, Ryba, JE: Modular Moonshine. II. *Duke Math. J.* **83**(2), 435–459 (1996). MR 1390654 (98b:17030)
23. Bott, R: Homogeneous vector bundles. *Ann. of Math.* **66**(2), 203–248 (1957). MR 0089473 (19,681d)
24. Bringmann, K, Duncan, J, Rolen, L: Maass-Jacobi Poincaré series and Mathieu Moonshine. **281**, 248–278 (2014). ArXiv e-prints <http://www.sciencedirect.com/science/article/pii/S0001870815001796>
25. Bringmann, K, Ono, K: The $f(q)$ mock theta function conjecture and partition ranks. *Invent. Math.* **165**(2), 243–266 (2006). MR 2231957 (2007e:11127)
26. Bringmann, K, Ono, K: Coefficients of harmonic maass forms. Partitions, q-Series, and Modular Forms (Krishnaswami Alladi and Frank Garvan, eds.), *Developments in Mathematics*, Vol. 23. Springer, New York (2012). pp. 23–38
27. Bringmann, K, Richter, O: Zagier-type dualities and lifting maps for harmonic Maass-Jacobi forms. *Adv. Math.* **225**(4), 2298–2315 (2010). MR 2680205 (2011h:11052)
28. Bringmann, K, Richter, O: Exact formulas for coefficients of Jacobi forms. *Int. J. Number Theory.* **7**(3), 825–833 (2011). MR 2805582 (2012f:11095)
29. Bruinier, JH, Ono, K: Heegner divisors, L -functions and harmonic weak Maass forms. *Ann. Math.* **172**(2 no.3), 2135–2181 (2010). MR 2726107 (2012c:11101)
30. Bruinier, JH, Funke, J: On two geometric theta lifts. *Duke Math. J.* **125**(1), 45–90 (2004). MR 2097357 (2005m:11089)
31. Bruinier, JH, Kohnen, W, Ono, K: The arithmetic of the values of modular functions and the divisors of modular forms. *Compos. Math.* **140**(3), 552–566 (2004). MR 2041768 (2005h:11083)
32. Carlip, S: Conformal field theory, $(2+1)$ -dimensional gravity, and the BTZ black hole. *Class. Quant. Grav.* **22**, R85–R124 (2005)
33. Carlip, S: Effective conformal descriptions of black hole entropy. *Entropy.* **13**, 1355–1379 (2011)
34. Carnahan, S: Generalized moonshine I: genus-zero functions. *Algebra Number Theory.* **4**(6), 649–679 (2010). MR 2728485 (2012b:11065)
35. Carnahan, S: Generalized moonshine, II: Borcherds products. *Duke Math. J.* **161**(5), 893–950 (2012). MR 2904095
36. Carnahan, S: Generalized Moonshine IV: Monstrous Lie algebras (2012). arXiv:1208.6254 [math.RT]
37. Carnahan, S: Monstrous Lie Algebras. *RIMS Kôkyûroku.* **1872** (2014)
38. Cheng, M: $K3$ surfaces, $N = 4$ dyons and the Mathieu group M_{24} . *Commun. Number Theory Phys.* **4**(4), 623–657 (2010). MR 2793423 (2012e:11076)
39. Cheng, M, de Lange, P, Whalen, D: Generalised Umbral Moonshine (2015)
40. Cheng, M, Dong, X, Duncan, J, Harrison, S, Kachru, S, Wrase, T: Mock Modular Mathieu Moonshine Modules (2014). ArXiv e-prints, arXiv:1406.5502 [hep-th]
41. Cheng, M, Dong, X, Duncan, J, Harvey, J, Kachru, S, Wrase, T: Mathieu Moonshine and $N=2$ String Compactifications. *JHEP.* **1309**, 030 (2013)
42. Cheng, M, Duncan, J: The largest Mathieu group and (mock) automorphic forms. *String-Math 2011, Proc. Sympos. Pure Math.*, vol. 85, Amer. Math. Soc., Providence, RI, 53–82 (2012). MR 2985326
43. Cheng, M, Duncan, J: On Rademacher sums, the largest Mathieu group, and the holographic modularity of moonshine. *Commun. Number Theory Phys.* **6**(3), 697–758 (2012)
44. Cheng, M, Duncan, J: On the discrete groups of Mathieu moonshine. *Perspect. Representation Theory, Contemp. Math.* **610**, 65–78 (2014)
45. Cheng, M, Duncan, J: Rademacher sums and rademacher series, Conformal Field Theory, Automorphic Forms and Related Topics. In: Winfried Kohnen, Rainer Weissauer (eds.) *Contributions in Mathematical and Computational Sciences*, vol. 8, pp. 143–182. Springer, Berlin Heidelberg, (2014). English
46. Cheng, M, Duncan, J: Meromorphic Jacobi Forms of Half-Integral Index and Umbral Moonshine Modules (2015)
47. Cheng, M, Harvey, J, Duncan, J: Umbral moonshine. *Commun. Number Theory Phys.* **8**(2), 101–242 (2014)
48. Cheng, M, Harvey, J, Duncan, J: Umbral moonshine and the Niemeier lattices. *Res. Math. Sci.* **1**(3), 1–81 (2014)

49. Cheng, M, Harvey, J, Duncan, J: Weight One Jacobi Forms and Umbral Moonshine (2015)
50. Cheng, M, Harrison, S: Umbral Moonshine and K3 Surfaces (2014). ArXiv e-prints, arXiv:1406.0619 [hep-th]
51. Cheng, MCN, Harrison, SM, Kachru, S, Whalen, D: Exceptional Algebra and Sporadic Groups at $c=12$ (2015). ArXiv e-prints, arXiv:1503.07219 [hep-th]
52. Conway, JH: A perfect group of order **8, 315, 553, 613, 086, 720, 000** and the sporadic simple groups. Proc. Nat. Acad. Sci. U.S.A. **61**, 398–400 (1968). MR 0237634 (38 #5915)
53. Conway, JH: A characterisation of Leech's lattice. Invent. Math. **7**, 137–142 (1969). MR 0245518 (39 #6824)
54. Conway, JH: A group of order **8, 315, 553, 613, 086, 720, 000**. Bull. London Math. Soc. **1**, 79–88 (1969). MR 0248216 (40 #1470)
55. Conway, JH: A simple construction for the Fischer-Griess monster group. Invent. Math. **79**(3), 513–540 (1985). MR MR782233 (86h:20019)
56. Conway, JH, Curtis, RT, Norton, SP, Parker, RA, Wilson, RA: Atlas of finite groups. Maximal subgroups and ordinary characters for simple groups. With comput. assist. from J. G. Thackray. Clarendon Press, Oxford (1985)
57. Conway, J, McKay, J, Sebbar, A: On the discrete groups of moonshine. Proc. Amer. Math. Soc. **132**, 2233–2240 (2004)
58. Conway, JH, Norton, SP: Monstrous moonshine. Bull. London Math Soc. **11**(3), 308–339 (1979). MR 554399 (81j:20028)
59. Conway, JH, Parker, RA, Sloane, NJA: The covering radius of the Leech lattice. Proc. Roy. Soc. London Ser. A. **380**(1779), 261–290 (1982). MR 660415 (84m:10022b)
60. Conway, JH, Sloane, NJA: On the enumeration of lattices of determinant one. J. Number Theory. **15**(1), 83–94 (1982). MR 666350 (84b:10047)
61. Conway, JH, Sloane, NJA: Twenty-three constructions for the Leech lattice. Proc. Roy. Soc. London Ser. A. **381**(1781), 275–283 (1982). MR 661720 (84m:10020)
62. Conway, JH, Sloane, NJA: Sphere packings, lattices and groups. third ed. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 290, Springer-Verlag, New York, (1999) With additional contributions by E. Bannai, R. E. Borcherds, J. Leech, S. P. Norton, A. M. Odlyzko, R. A. Parker, L. Queen and B. B. Venkov. MR 1662447 (2000b:11077)
63. Creutzig, T, Höhn, G: Mathieu moonshine and the geometry of K3 surfaces. Commun. Number Theory Phys. **8**(2), 295–328 (2014). MR 3271177
64. Cummins, CJ, Gannon, T: Modular equations and the genus zero property of moonshine functions. Invent. Math. **129**(3), 413–443 (1997). MR 1465329 (98k:11046)
65. Curtis, RT: Natural constructions of the Mathieu groups. Math. Proc Cambridge Philos. Soc. **106**(3), 423–429 (1989). MR 1010366 (90h:20020)
66. Dabholkar, A, Murthy, S, Zagier, D: Quantum Black Holes, Wall Crossing, and Mock Modular Forms (2012)
67. Deser, S, Jackiw, R: Erratum: "Topologically massive gauge theories" [Ann. Physics **140**, (1982), no. 2, 372–411; MR0665601 (84j:81128)] by Deser, Jackiw and S. Templeton. Ann. Physics. **185**(2), 406 (1988). MR 965585 (90a:81077b)
68. Deser, S, Jackiw, R, Templeton, S: Topologically massive gauge theories. Ann. Phys. **140**(2), 372–411 (1982). MR 665601 (84j:81128)
69. Deser, S, Jackiw, R, Templeton, S: Three-Dimensional Massive Gauge Theories. Phys. Rev. Lett. **48**, 975–978 (1982)
70. Dijkgraaf, R, Maldacena, J, Moore, G, Verlinde, E: A black hole farey tail (2007)
71. Dijkgraaf, R, Moore, G, Verlinde, E, Verlinde, H: Elliptic genera of symmetric products and second quantized strings. Commun. Math. Phys. **185**, 197–209 (1997)
72. Dixon, L, Ginsparg, P, Harvey, J: Beauty and the beast: superconformal symmetry in a Monster module. Comm. Math. Phys. **119**(2), 221–241 (1988). MR 968697 (90b:81119)
73. Dixon, L, Harvey, JA, Vafa, C, Witten, E: Strings on orbifolds. Nuclear Phys. B. **261**(4), 678–686 (1985). MR 818423 (87k:81104a)
74. Dixon, L, Harvey, JA, Vafa, C, Witten, E: Strings on orbifolds. II. Nuclear Phys. B. **274**(2), 285–314 (1986). MR 851703 (87k:81104b)
75. Dong, C, Jiao, X, Xu, F: Quantum dimensions and quantum Galois theory. Trans. Amer. Math. Soc. **365**(2), 6441–6469 (2013). MR 3105758
76. Dong, C, Li, H, Mason, G: Some twisted sectors for the Moonshine module, Moonshine, the Monster, and related topics (South Hadley, MA, 1994), Contemp. Math., vol. 193, Amer. Math. Soc., Providence, RI, 25–43 (1996). MR 1372716 (97b:17021)
77. Dong, C, Li, H, Mason, G: Compact automorphism groups of vertex operator algebras. Int. Math. Res. Notices. **18**, 913–921 (1996). MR 1420556 (98a:17044)
78. Dong, C, Li, H, Mason, G: Twisted representations of vertex operator algebras. Math. Ann. **310**(3), 571–600 (1998). MR 1615132 (99d:17030)
79. Dong, C, Li, H, Mason, G: Modular invariance of trace functions in orbifold theory and generalized Moonshine. Commun. Math. Phys. **214**, 1–56 (2000)
80. Dong, C, Mason, G: Nonabelian orbifolds and the boson-fermion correspondence. Comm. Math. Phys. **163**(3), 523–559 (1994). MR 1284796 (95i:17031)
81. Dong, C, Mason, G: On quantum Galois theory. Duke Math. J. **86**(2), 305–321 (1997). MR 1430435 (97k:17042)
82. Dong, C, Mason, G: Quantum Galois theory for compact Lie groups. J. Algebra. **214**(1), 92–102 (1999). MR 1684904 (2000g:17043b)
83. Duncan, JF: Super-Moonshine for Conway's largest sporadic group. Duke Math. J. **139**(2), 255–315 (2007)
84. Duncan, JFR, Frenkel, IB: Rademacher sums, moonshine and gravity. Commun. Number Theory Phys. **5**(4), 1–128 (2011)
85. Duncan, JFR, Griffin, M, Ono, K: Proof of the Umbral Moonshine Conjecture (2015). ArXiv e-prints <http://adsabs.harvard.edu/abs/2015arXiv150301472D>
86. Duncan, JFR, Harvey, JA: The Umbral Moonshine Module for the Unique Unimodular Niemeier Root System. ArXiv e-prints (2014). <http://adsabs.harvard.edu/abs/2014arXiv1412.8191D>

87. Duncan, JFR, Mack-Crane, S: The Moonshine Module for Conway's Group. ArXiv e-prints (2014). <http://adsabs.harvard.edu/abs/2014arXiv1409.3829D>
88. Duncan, JFR, Mack-Crane, S: Derived Equivalences of K3 Surfaces and Twined Elliptic Genera (2015)
89. Duncan, J, O'Desky, A: Super Vertex Algebras, Meromorphic Jacobi Forms, and Umbral Moonshine (2015)
90. Duncan, JFR, Ono, K: The Jack Daniels Problem. ArXiv e-prints (2014). <http://adsabs.harvard.edu/abs/2014arXiv1411.5354D>
91. Eguchi, T, Hikami, K: Superconformal Algebras and Mock Theta Functions 2. Rademacher Expansion for K3 Surface. *Commun. Number Theory Phys.* **3**, 531–554 (2009)
92. Eguchi, T, Hikami, K: Note on Twisted Elliptic Genus of K3 Surface. *Phys. Lett.* **B694**, 446–455 (2011)
93. Eguchi, T, Ooguri, H, Tachikawa, Y: Notes on the K3 Surface and the Mathieu group M_{24} . *Exper. Math.* **20**, 91–96 (2011)
94. Eguchi, T, Ooguri, H, Taormina, A, Yang, S-K: Superconformal algebras and string compactification on manifolds with $SU(N)$ Holonomy. *Nucl. Phys.* **B315**, 193 (1989)
95. Eguchi, T, Taormina, A: Unitary representations of the $N = 4$, superconformal algebra. *Phys. Lett. B.* **196**(1), 75–81 (1987). MR 910253 (88j:17022)
96. Eichler, M, Zagier, D: *The theory of Jacobi forms*. Birkhäuser (1985)
97. Fay, J: Fourier coefficients of the resolvent for a Fuchsian group. *J. Reine Angew. Math.* **293/294**, 143–203 (1977). MR 0506038 (58 #21944)
98. Feigin, BL, Fuchs, DB: Verma modules over the Virasoro algebra, *Topology (Leningrad, 1982)*, Lecture Notes in Math., vol. 1060. Springer, Berlin (1984). pp. 230–245. MR 770243 (86g:17004)
99. Ford, D, McKay, J, Norton, S: More on replicable functions. *Comm. Algebra.* **22**(13), 5175–5193 (1994). MR 1291027 (95i:11036)
100. Frenkel, E, Ben-Zvi, D: *Vertex algebras and algebraic curves*. second ed. *Mathematical Surveys and Monographs*, vol. 88, American Mathematical Society, Providence, RI (2004). MR 2082709 (2005d:17035)
101. Frenkel, IB, Kac, VG: Basic representations of affine Lie algebras and dual resonance models. *Invent. Math.* **62**(1), 23–66. (1980/81) MR 84f:17004
102. Frenkel, I, Lepowsky, J, Meurman, A: A natural representation of the Fischer-Griess Monster with the modular function J as character. *Proc. Nat. Acad. Sci. U.S.A.* **81**(10), 3256–3260 (1984). MR 747596 (85e:20018)
103. Frenkel, I, Lepowsky, J, Meurman, A: *A Moonshine Module for the Monster*. *Vertex operators in mathematics and physics (Berkeley, Calif., 1983)*, *Math. Sci. Res. Inst. Publ.*, vol. 3. Springer, New York (1985). pp. 231–273. MR 86m:20024
104. Frenkel, I, Lepowsky, J, Meurman, A: *Vertex Operator Algebras and the Monster*. *Pure and Applied Mathematics*, vol. 134. Academic Press Inc., Boston, MA (1988). MR 90h:17026
105. Gaberdiel, M: An Introduction to conformal field theory. *Rept. Prog. Phys.* **63**, 607–667 (2000)
106. Gaberdiel, M: *2D conformal field theory and vertex operator algebras* (2005)
107. Gaberdiel, M, Hohenegger, S, Volpato, R: Mathieu moonshine in the elliptic genus of K3. *JHEP.* **1010**, 062 (2010)
108. Gaberdiel, M, Hohenegger, S, Volpato, R: Mathieu twining characters for K3. *JHEP.* **1009**(058), 19 (2010)
109. Gaberdiel, M, Hohenegger, S, Volpato, R: Symmetries of K3 sigma models. *Commun. Number Theory Phys.* **6**(1), 1–50 (2012). MR 2955931
110. Gaberdiel, M, Keller, C, Volpato, R: Genus two partition functions of chiral conformal field theories. *Commun. Number Theory Phys.* **4**(2), 295–363 (2010). MR 2725054 (2011h:81208)
111. Gaberdiel, M, Persson, D, Ronellenfitsch, HR: Volpato: Generalized Mathieu Moonshine. *Commun. Number Theory Phys.* **7**(1), 145–223 (2013). MR3108775
112. Gaberdiel, M, Taormina, A, Volpato, R, Wendland, K: A K3 sigma model with $Z_2^8 : M_{20}$ symmetry. **1402** (2014). doi: 10.1007/JHEP02(2014)022
113. Gaberdiel, M, Volpato, R: Mathieu Moonshine and Orbifold K3s. **8**, 109–141 (2014). doi: 10.1007/978-3-662-43831-2_5
114. Gaiotto, D: *Monster symmetry and Extremal CFTs* (2008)
115. Gaiotto, D, Yin, X: Genus two partition functions of extremal conformal field theories. *JHEP.* **0708**, 029 (2007)
116. Gannon, T: Monstrous moonshine: the first twenty-five years. *Bull. London Math. Soc.* **38**(1), 1–33 (2006). MR 2201600 (2006k:11071)
117. Gannon, T: Much ado about Mathieu. ArXiv e-prints (2012). <http://adsabs.harvard.edu/abs/2012arXiv1211.5531G>
118. Garland, H, Lepowsky, J: Lie algebra homology and the Macdonald-Kac formulas. *Invent. Math.* **34**(1), 37–76 (1976). MR 0414645 (54 #2744)
119. Griess, RL Jr: The structure of the "monster" simple group. In: *Proceedings of the Conference on Finite Groups (Univ. Utah, Park City, Utah, 1975)*, pp. 113–118. Academic Press, New York, (1976). MR 0399248 (53 #3099)
120. Griess, RL Jr: A construction of F_1 as automorphisms of a **196, 883**-dimensional algebra. *Proc. Nat. Acad. Sci. U.S.A.* **78**(2, part 1), 686–691 (1981). MR 605419 (82d:20022)
121. Griess, RL Jr: The friendly giant. *Invent. Math.* **69**(1), 1–102 (1982). MR 671653 (84m:20024)
122. Gritsenko, V: Elliptic genus of Calabi-Yau manifolds and Jacobi and Siegel modular forms. *Algebra i Analiz.* **11**(5), 100–125 (1999). MR 1734348 (2001i:11051)
123. Grumiller, D, Johansson, N: Instability in cosmological topologically massive gravity at the chiral point. *J High Energy Phys.* **7**, 1–16 (2008)
124. Grumiller, D, McNees, R, Salzer, J: *Black holes and thermodynamics - The first half century* (2014)
125. Grumiller, D, Riedler, W, Rosseel, J, Zojer, T: *Holographic applications of logarithmic conformal field theories*, Vol. A46 (2013)
126. Gubser, SS, Klebanov, IR, Polyakov, AM: Gauge theory correlators from noncritical string theory. *Phys. Lett.* **B428**, 105–114 (1998)
127. Hanaki, A, Miyamoto, M, Tambara, D: Quantum Galois theory for finite groups. *Duke Math. J.* **97**(3), 541–544 (1999). MR 1682988 (2000g:17043a)

128. Harada, K, Lang, M-L: The McKay-Thompson series associated with the irreducible characters of the Monster, Moonshine, the Monster, and related topics (South Hadley, MA, 1994). *Contemp. Math.*, vol. 193, Amer. Math. Soc., Providence, RI (1996). MR 1372718 (97b:11059)
129. Harada, K, Lang, M-L: Modular forms associated with the Monster module. *The Monster and Lie algebras* (Columbus, OH, 1996), Ohio State Univ. Math. Res. Inst. Publ., vol. 7, de Gruyter, Berlin (1998). pp. 59–83. MR 1650637 (2000f:11046)
130. Hardy, GH: *Collected papers of G. H. Hardy* (Including Joint papers with J. E. Littlewood and others). Vol. I. Edited by a committee appointed by the London Mathematical Society, Clarendon Press, Oxford (1966). MR 0201267 (34 #1151)
131. Hardy, GH, Ramanujan, S: Asymptotic formulæ in combinatory analysis [*Proc. London Math. Soc.* (2) **17** (1918), 75–115] *Collected papers of Srinivasa Ramanujan*, AMS Chelsea Publ., Providence, RI, 276–309 (2000). MR 2280879
132. Harrison, S, Kachru, S, Paquette, N: Twining Genera of (0,4) Supersymmetric Sigma Models on K3. *JHEP.* **1404**, 048 (2014)
133. Harvey JA, Murthy, S: Moonshine in Fivebrane Spacetimes. *ArXiv e-prints.* **1401**, 146 (2013). doi: 10.1007/JHEP01(2014)146
134. Harvey, JA, Murthy, S, Nazarov, C: ADE Double Scaled Little String Theories, Mock Modular Forms and Umbral Moonshine. *ArXiv e-prints.* **1505**, 126 (2015). doi: 10.1007/JHEP05(2015)126
135. Hoehn, G: Generalized Moonshine for the Baby Monster (2003). <http://www.math.ksu.edu/~gerald/papers/>
136. Hoehn, G: Komplexe elliptische Geschlechter und S^1 -equivariante Kobordismustheorie (Complex elliptic genera and S^1 -equivariant cobordism theory). *ArXiv Mathematics e-prints* (2004). <http://adsabs.harvard.edu/abs/2004math.....5232H>
137. Hoehn, G: Self-dual vertex operator superalgebras of large minimal weight. *arXiv:0801.1822v1* (2008)
138. Hohenegger, S: Automorphic products, generalized Kac-Moody algebras and string amplitudes. *J. Phys. Conf. Ser.* **346**, 012008 (2012)
139. Humphreys, J: *Introduction to Lie algebras and representation theory*. Springer-Verlag, New York (1972). Graduate Texts in Mathematics, Vol. 9. MR 0323842 (48 #2197)
140. Huybrechts, D: On derived categories of K3 surfaces, symplectic automorphisms and the Conway group. *ArXiv e-prints* (2013). <http://adsabs.harvard.edu/abs/2013arXiv1309.6528H>
141. Ivanov, R, Tuite, M: Rational generalised moonshine from abelian orbifoldings of the moonshine module. *Nuclear Phys. B.* **635**(3), 435–472 (2002). MR 1915258 (2003g:11040)
142. Ivanov, R, Tuite, M: Some irrational generalised moonshine from orbifolds. *Nuclear Phys. B.* **635**(3), 473–491 (2002). MR 1915259 (2003f:11057)
143. Jurisch, E: An exposition of generalized Kac-Moody algebras. *Lie algebras and their representations* (Seoul, 1995), *Contemp. Math.*, vol. 194, Amer. Math. Soc., Providence, RI, 121–159 (1996). MR 1395597 (97e:17035)
144. Jurisch, E: Generalized Kac-Moody Lie algebras, free Lie algebras and the structure of the Monster Lie algebra. *J. Pure Appl. Algebra.* **126**, 233–266 (1998). MR 1600542 (99b:17032)
145. Jurisch, E: Borcherd's proof of the Conway-Norton conjecture. *Moonshine: the first quarter century and beyond*, *London Math. Soc. Lecture Note Ser.*, vol. 372, Cambridge Univ. Press, Cambridge (2010). pp. 219–235. MR 2681781 (2012a:20026)
146. Jurisch, E, Lepowsky, J, Wilson, RL: Realizations of the monster Lie algebra. *Selecta Math. (N.S.)* **1**(1), 129–161 (1995). MR 1327230 (96e:17059)
147. Kac, VG: Simple irreducible graded Lie algebras of finite growth. *Izv. Akad. Nauk SSSR Ser. Mat.* **32**, 1323–1367 (1968). MR 0259961 (41 #4590)
148. Kac, VG: Infinite-dimensional Lie algebras, and the Dedekind η -function. *Funkcional. Anal. i Priložen.* **8**(1), 77–78 (1974). MR 0374210 (51 #10410)
149. Kac, VG: Infinite-dimensional algebras, Dedekind's η -function, classical Möbius function and the very strange formula. *Adv. Math.* **30**(2), 85–136 (1978). MR 513845 (83a:17014a)
150. Kac, VG: An elucidation of: "Infinite-dimensional algebras, Dedekind's η -function, classical Möbius function and the very strange formula". $E_8^{(1)}$ and the cube root of the modular invariant j . *Adv. Math.* **35**(3), 264–273 (1980). MR 563927 (83a:17014b)
151. Kac, VG: *Infinite-dimensional Lie algebras*. third ed. Cambridge University Press, Cambridge (1990). MR 1104219 (92k:17038)
152. Kac, VG: *Vertex algebras for beginners*. second ed. University Lecture Series, vol. 10, American Mathematical Society, Providence, RI (1998). MR 1651389 (99f:17033)
153. Kac, VG, Peterson, DH: Infinite-dimensional Lie algebras, theta functions and modular forms. *Adv. Math.* **53**(2), 125–264 (1984). MR750341 (86a:17007)
154. Kac, VG, Raina, AK: *Bombay lectures on highest weight representations of infinite-dimensional Lie algebras*. *Advanced Series in Mathematical Physics*, vol. 2, World Scientific Publishing Co., Inc., Teaneck, NJ (1987). MR 1021978 (90k:17013)
155. Katz, S, Klemm, A, Pandharipande, R: On the motivic stable pairs invariants of K3 surfaces. *ArXiv e-prints* (2014). <http://adsabs.harvard.edu/abs/2014arXiv1407.3181K>
156. Knopp, MI: Rademacher on $J(\tau)$, Poincaré series of nonpositive weights and the Eichler cohomology. *Notices Amer. Math. Soc.* **37**(4), 385–393 (1990). MR MR1050700 (91e:11047)
157. Knopp, MI: Construction of a class of modular functions and forms. *Pacific J. Math.* **11**, 275–293 (1961). MR MR0122994 (23#A325)
158. Knopp, MI: Construction of a class of modular functions and forms. II. *Pacific J. Math.* **11**, 661–678 (1961). MR MR0130229 (24 #A96)
159. Knopp, MI: Construction of automorphic forms on H -groups and supplementary Fourier series. *Trans. Am. Math. Soc.* **103**, 168–188 (1962). MR MR0136735 (25 #197)
160. Knopp, MI: On abelian integrals of the second kind and modular functions. *Amer. J. Math.* **84**, 615–628 (1962). MR MR0146379 (26 #3901)

161. Knutson, D: λ -rings and the representation theory of the symmetric group. Lecture Notes in Mathematics, Vol. 308, Springer-Verlag, Berlin-New York (1973). MR 0364425 (51 #679)
162. Kondó, S: Niemeier lattices, Mathieu groups, and finite groups of symplectic automorphisms of $K3$ surfaces. *Duke Math. J.* **92**(3), 593–603 (1998). With an appendix by Shigeru Mukai. MR 1620514 (99i:14042)
163. Kondó, S: On the Kodaira dimension of the moduli space of $K3$ surfaces. II. *Compositio Math.* **116**(2), 111–117 (1999). MR 1686793 (2000j:14055)
164. Kondo, T: The automorphism group of Leech lattice and elliptic modular functions. *J. Math. Soc. Japan.* **37**(2), 337–362 (1985). MR 780666 (86k:11033)
165. Kostant, B: Lie algebra cohomology and the generalized Borel-Weil theorem. *Ann. Math.* **74**(2), 329–387 (1961). MR 0142696 (26 #265)
166. Landweber, PS: Elliptic curves and modular forms in algebraic topology, Lecture Notes in Mathematics, Vol. 1326. Springer-Verlag, Berlin (1988). MR 970278 (91a:57021)
167. Leech, J: Some sphere packings in higher space. *Canad. J. Math.* **16**, 657–682 (1964). MR 0167901 (29 #5166)
168. Leech, J: Notes on sphere packings. *Canad. J. Math.* **19**, 251–267 (1967). MR 0209983 (35 #878)
169. Leon, J, Sims, C: The existence and uniqueness of a simple group generated by $\{3, 4\}$ -transpositions. *Bull. Amer. Math. Soc.* **83**(5), 1039–1040 (1977). MR 0444765 (56 #3113)
170. Lepowsky, J, Li, H: Introduction to vertex operator algebras and their representations, Progress in Mathematics, Vol. 227. Birkhäuser Boston Inc., Boston, MA (2004). MR 2023933 (2004k:17050)
171. Lepowsky, J, Wilson, RL: Construction of the affine Lie algebra $A_1^{(1)}$. *Comm. Math. Phys.* **62**(1), 43–53 (1978). MR 0573075 (58 #28089)
172. Li, W, Song, W, Strominger, A: Chiral gravity in three dimensions. *J. High Energy Phys.* **4**, 082, 15 (2008). MR 2425221 (2010a:83101)
173. Mahler, K: On a class of non-linear functional equations connected with modular functions. *J. Austral. Math. Soc. Ser. A.* **22**(1), 65–118 (1976). MR 0441867 (56 #258)
174. Maldacena, J: The gauge gravity duality. In: Proceedings of Xth Quark Confinement and the Hadron Spectrum (Confinement X), (2012). TUM Camous Garching, Munich, Germany
175. Maldacena, J: The large N limit of superconformal field theories and supergravity. *Adv. Theor. Math. Phys.* **2**, 231–252 (1998)
176. Maldacena, J, Strominger, A: AdS (3), black holes and a stringy exclusion principle. *JHEP.* **9812**, 005 (1998)
177. Maloney, A, Song, W, Strominger, A: Chiral Gravity, Log Gravity and Extremal CFT. *Phys. Rev.* **D81**, 064007 (2010)
178. Maloney, A, Witten, E: Quantum gravity partition functions in three dimensions. *J. High Energy Phys.* **029**(2), 58 (2010). MR 2672754 (2011m:83022)
179. Manschot, J: AdS_3 partition functions reconstructed. *J. High Energy Phys.* **103**(10), 7 (2007). MR MR2357875
180. Manschot, J, Moore, GW: A Modern Fareytail. *Commun. Num. Theor. Phys.* **4**, 103–159 (2007)
181. Mason, G: Finite groups and modular functions. The Arcata Conference on Representations of Finite Groups (Arcata, Calif., 1986) (1987). MR 933359 (89c:11066)
182. Mathieu, É: Mémoire sur l'étude des fonctions de plusieurs quantités, sur la manière de les former et sur les substitutions qui les laissent invariables. *J. de Mathématiques Pures et Appliquées.* **6**, 241–323 (1861)
183. Mathieu, É: Sur la fonction cinq fois transitive de 24 quantités. *Journal de Mathématiques Pures et Appliquées.* **18**, 25–46 (1873)
184. Moody, RV: Lie algebras associated with generalized Cartan matrices. *Bull. Amer. Math. Soc.* **73**, 217–221 (1967). MR 0207783 (34 #7598)
185. Moody, RV: A new class of Lie algebras. *J. Algebra.* **10**, 211–230 (1968). MR 0229687 (37 #5261)
186. Mukai, S: Finite groups of automorphisms of $K3$ surfaces and the Mathieu group. *Invent. Math.* **94**(1), 183–221 (1988). MR 958597 (90b:32053)
187. Niebur, D: A class of nonanalytic automorphic functions. *Nagoya Math. J.* **52**, 133–145 (1973). MR MR0337788 (49 #2557)
188. Niebur, D: Construction of automorphic forms and integrals. *Trans. Amer. Math. Soc.* **191**, 373–385 (1974). MR MR0344196 (49 #8936)
189. Niemeier, HV: Definite quadratische Formen der Dimension 24 und Diskriminante 1. *J. Number Theory.* **5**, 142–178 (1973). MR 0316384 (47 #4931)
190. Norton, SP: Generalized Moonshine. *Proc. Symp. Pure Math.* **47**, 208–209 (1987)
191. Norton, SP: More on moonshine. Computational group theory (Durham, 1982). Academic Press, London (1984). MR 760657 (86h:20020)
192. Norton, SP: From Moonshine to the Monster. In: Proceedings on Moonshine and related topics (Montréal, QC, 1999), CRM Proc. Lecture Notes, vol. 30, Amer. Math. Soc., Providence, RI, pp. 163–171, (2001). MR 1877765 (2002k:20025)
193. Ogg, AP: Automorphismes de courbes modulaires. Séminaire Delange-PisotPoitou (16e année: 1974/75), Théorie des nombres, Fasc. 1, Exp. No. 7. Secrétariat Mathématique, Paris (1975). MR 0417184 (54 #5242)
194. Ono, K: Unearthing the visions of a master: harmonic Maass forms and number theory. Current developments in mathematics, 2008. Int. Press, Somerville, MA (2009). MR 2555930 (2010m:11060)
195. Ono, K, Rolin, L, Trebat-Leder, S: Classical and Umbral Moonshine: Connections and p -adic Properties. *J. Ramanujan Math. Soc.* (2014). <http://adsabs.harvard.edu/abs/2014arXiv1403.37120>
196. Paquette, N, Wrase, T: Comments on M_{24} representations and CY_3 geometries. **1411**, 155 (2014). doi: 10.1007/JHEP11(2014)155
197. Petersson, H: Über die Entwicklungskoeffizienten der automorphen Formen. *Acta Math.* **58**(1), 169–215 (1932). MR 1555346
198. Queen, L: Modular functions arising from some finite groups. *Math. Comp.* **37**(156), 547–580 (1981). MR 628715 (83d:20008)
199. Rademacher, H: On the Partition Function $p(n)$. *Proc. London Math. Soc.* **43**(2), 241–254 (1937)
200. Rademacher, H: The Fourier Coefficients of the Modular Invariant $J(\tau)$. *Amer. J. Math.* **60**(2), 501–512 (1938). MR MR1507331

201. Rademacher, H: The Fourier Series and the Functional Equation of the Absolute Modular Invariant $J(\tau)$. *Amer. J. Math.* **61**(1), 237–248 (1939). MR MR1507375
202. Rains, E, Sloane, NJA: Self-dual codes. *Handbook of coding theory*, Vol. I, II. North-Holland, Amsterdam (1998). MR 1667939
203. Ramanujan, S: The lost notebook and other unpublished papers. Springer-Verlag, Berlin (1988). With an introduction by George E. Andrews, MR 947735 (89j:01078)
204. Ramanujan, S: Collected papers of Srinivasa Ramanujan. AMS Chelsea Publishing, Providence, RI (2000). Edited by G. H. Hardy, P. V. Seshu Aiyar and B. M. Wilson, Third printing of the 1927 original, With a new preface and commentary by Bruce C. Berndt, MR 2280843 (2008b:11002)
205. Rankin, RA: Modular forms and functions. Cambridge University Press, Cambridge-New York-Melbourne (1977). MR 0498390 (58 #16518)
206. Ray, U: Generalized Kac-Moody algebras and some related topics. *Bull. Amer. Math. Soc. (N.S.)* **38**(1), 1–42 (electronic 2001). MR 1803076 (2001j:17044)
207. Reynolds, W: Thompson’s characterization of characters and sets of primes. *J. Algebra.* **156**(1), 237–243 (1993). MR 1213794 (94b:20012)
208. Segal, G: Unitary representations of some infinite-dimensional groups. *Comm. Math. Phys.* **80**(3), 301–342 (1981). MR 626704 (82k:22004)
209. Serre, J-P: A course in arithmetic. Springer-Verlag, New York (1973). Translated from the French, Graduate Texts in Mathematics, No. 7. MR 0344216 (49 #8956)
210. Serre, J-P, Stark, HM: Modular forms of weight $1/2$. In: *Modular functions of one variable, VI* (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976) *Lecture Notes in Math.*, pp. 27–67. Springer, Berlin, (1977)
211. Shimura, G: Introduction to the arithmetic theory of automorphic functions, Publications of the Mathematical Society of Japan, No. 11. Iwanami Shoten, Publishers, Tokyo (1971). Kanô Memorial Lectures, No. 1. MR0314766 (47 #3318)
212. Smith, SD: On the head characters of the Monster simple group, Finite groups—coming of age (Montreal, Que., 1982), *Contemp. Math.*, vol. 45. Amer. Math. Soc., Providence, RI (1985). MR 822245 (87h:20037)
213. Stachel, J: The early history of quantum gravity (1916–1940). Black holes, gravitational radiation and the universe, *Fund. Theories Phys.*, vol. 100. Kluwer Acad. Publ., Dordrecht (1999). MR 1663704
214. Strominger, A: A Simple Proof of the Chiral Gravity Conjecture (2008)
215. Susskind, L: The world as a hologram. *J. Math. Phys.* **36**, 6377–6396 (1995)
216. ’t Hooft, G: Dimensional Reduction in Quantum Gravity. *ArXiv General Relativity and Quantum Cosmology e-prints*, 284–296 (1993)
217. Taormina, A, Wendland, K: The overarching finite symmetry group of Kummer surfaces in the Mathieu group M_{24} . *J. High Energy Phys.* **8**, 125 (2013). front matter+62. MR 3106313
218. Taormina, A, Wendland, K: A twist in the M_{24} moonshine story (2013)
219. Taormina, A, Wendland, K: Symmetry-surfing the moduli space of Kummer $K3$ s (2013)
220. Thompson, JG: Finite groups and modular functions. *Bull. London Math. Soc.* **11**(3), 347–351 (1979). MR MR554401 (81j:20029)
221. Thompson, JG: Some numerology between the Fischer-Griess Monster and the elliptic modular function. *Bull. London Math. Soc.* **11**(3), 352–353 (1979). MR MR554402 (81j:20030)
222. Tits, J: On R. Griess’ “friendly giant”. *Invent. Math.* **78**(3), 491–499 (1984). MR 768989 (86f:20019)
223. Tuite, MP: Monstrous Moonshine from orbifolds. *Commun. Math. Phys.* **146**, 277–309 (1992)
224. Tuite, MP: On the relationship between monstrous moonshine and the uniqueness of the moonshine module. *Commun. Math. Phys.* **166**, 495–532 (1995)
225. Tuite, MP: Generalized moonshine from abelian orbifold constructions. *Contemp. Maths.* **193**, 353–368 (1995)
226. Vaughan, RC: The Hardy-Littlewood method. second ed. Cambridge, Tracts in Mathematics, vol. 125, Cambridge University Press, Cambridge (1997). MR 1435742 (98a:11133)
227. Venkov, BB: On the classification of integral even unimodular 24 -dimensional quadratic forms. *Trudy Mat. Inst. Steklov.* **148**, 65–76 (1978). 273, *Algebra, number theory and their applications.* MR 558941 (81d:10024)
228. Volpato, R: On symmetries of $N=(4,4)$ sigma models on T^4 . *ArXiv e-prints.* **1408**, 094 (2014). doi: 10.1007/JHEP08(2014)094
229. Wiles, A: The Birch and Swinnerton-Dyer conjecture. The millennium prize problems, Clay Math. Inst., Cambridge, MA (2006). pp. 31–41. MR2238272
230. Witten, E: Elliptic genera and quantum field theory. *Comm. Math. Phys.* **109**(4), 525–536 (1987). MR 885560 (89i:57017)
231. Witten, E: Anti-de Sitter space and holography. *Adv. Theor. Math. Phys.* **2**, 253–291 (1998)
232. Witten, E: Three-Dimensional Gravity Revisited (2007)
233. Wrase, T: Mathieu moonshine in four dimensional $\mathcal{N} = 1$ theories. *JHEP.* **1404**, 069 (2014)
234. Yin, X: Partition functions of three-dimensional pure gravity. *Commun. Number Theory Phys.* **2**(2), 285–324 (2008). MR 2442775 (2009m:83105)
235. Yoshikawa, K-I: Nikulin’s $K3$ surfaces, adiabatic limit of equivariant analytic torsion, and the Borcherds Φ -function, *Complex analysis in several variables—Memorial Conference of Kiyoshi Oka’s Centennial Birthday.* *Adv. Stud. Pure Math.*, vol. 42, Math. Soc., Japan Tokyo (2004). pp. 339–345. MR 2087068 (2005h:14093)
236. Zagier, D: Ramanujan’s mock theta functions and their applications (after Zwegers and Ono-Bringmann). *Astérisque* (2009), no. 326, Exp. No. 986, vii–viii, 143–164 (2010), *Séminaire Bourbaki.* Vol. 2007/2008. MR 2605321 (2011h:11049)
237. Zhu, Y: Modular invariance of characters of vertex operator algebras. *J. Am. Math. Soc.* **9**(1), 237–302 (1996)
238. Zwegers, S: Mock Theta Functions. Ph.D. thesis, Utrecht University (2002)
239. Zwegers, S: On two fifth order mock theta functions. *Ramanujan J.* **20**(2), 207–214 (2009). MR 2558702 (2011a:11046)