

Short Report

## Sero-prevalence of latent *Toxoplasma gondii* infection among HIV-infected and HIV-uninfected people in Addis Ababa, Ethiopia: A comparative cross-sectional study

Techalew Shimelis\*<sup>1</sup>, Mekashaw Tebeje<sup>2</sup>, Endale Tadesse<sup>1</sup>, Belete Tegbaru<sup>2</sup> and Ashenafi Terefe<sup>3</sup>

Address: <sup>1</sup>Hawassa University, Department of Medical Laboratory Science, Hawassa, Ethiopia, <sup>2</sup>Ethiopian Health and Nutrition Research Institute, National HIV Laboratory, Addis Ababa, Ethiopia and <sup>3</sup>Jimma University, School of Medical Laboratory Technology, Jimma, Ethiopia

Email: Techalew Shimelis\* - [techalew03@yahoo.com](mailto:techalew03@yahoo.com); Mekashaw Tebeje - [mekashawtebeje@yahoo.com](mailto:mekashawtebeje@yahoo.com); Endale Tadesse - [endaletd@yahoo.com](mailto:endaletd@yahoo.com); Belete Tegbaru - [beletegbaru@gmail.com](mailto:beletegbaru@gmail.com); Ashenafi Terefe - [ashenafidejene@gmail.com](mailto:ashenafidejene@gmail.com)

\* Corresponding author

Published: 23 October 2009

Received: 20 February 2009

BMC Research Notes 2009, 2:213 doi:10.1186/1756-0500-2-213

Accepted: 23 October 2009

This article is available from: <http://www.biomedcentral.com/1756-0500/2/213>

© 2009 Shimelis et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Toxoplasmosis in immuno-compromised hosts manifests primarily as a life threatening condition, toxoplasmic encephalitis. However, there is scarce information about the magnitude of *Toxoplasma gondii* infection among HIV-infected people in Ethiopia. This study was, therefore, conducted to determine the sero-prevalence of *T. gondii* infection among HIV-infected and HIV-uninfected subjects.

**Findings:** Sera were collected from people with and without HIV infection for the purpose of studying hepatitis B virus (HBV) at St. Paul Hospital, Addis Ababa, Ethiopia from 24 January 2007 to 15 February 2007. Among these sera, the first 330 consecutive sera, 165 from each HIV serogroup, were selected and tested for anti-*T. gondii* IgG antibodies using Enzyme Linked Immunosorbent Assay. The seroprevalence of *Toxoplasma* infection was assessed against socio-demographic characteristics, HIV and HBV serostatus and HBV-related risk factors. The overall sero-prevalence of latent *T. gondii* infection among the study subjects was 90.0%. *Toxoplasma* infection was observed with respective prevalence of 93.3% and 86.7% among HIV-infected and HIV-uninfected people. Though *Toxoplasma* infection seems to be influenced by age, gender and HIV serostatus, only HBV serostatus was significantly associated (OR 2.71, CI 1.12 to 6.57) in multivariate logistic regression analysis.

**Conclusion:** The seroprevalence of latent *T. gondii* infection is high and similar by HIV status. Educating people to prevent acquisition of new *Toxoplasma* infection and minimizing the risk of disease manifestations among HIV-*Toxoplasma* co-infected individuals is important.

### Background

*Toxoplasma gondii* is one of the most prevalent protozoan parasites of man and livestock [1]. It has been estimated that up to one third of the world's population is infected

by *T. gondii* [2]. Most infections among humans occur by eating undercooked or raw meat containing tissue cysts or by exposure to oocysts through ingestion of contaminated foods and drinks with cat's faeces [3-5]. Other modes of

transmission include the transplacental route, blood product transfusion and tissue transplantation [6,7].

In vast majority of immunocompetent human host, *T. gondii* ensue a latent infection characterized by the persistence of the organism in tissues (primarily brain, skeletal muscle, and heart) without causing disease [8]. However, in chronically infected individuals who develop defects in cell-mediated immunity a symptomatic disease more likely occurs as a result of reactivation of latent infection [9,10]. Toxoplasmosis among Acquired Immunodeficiency Syndrome (AIDS) patients manifests primarily as a life threatening condition, toxoplasmic encephalitis (TE) [9-11]. Early diagnosis and appropriate management of toxoplasmosis decreases the incidence rates of TE; subsequently reduce morbidity and mortality among HIV infected individuals [7].

In Ethiopia, up to 80% prevalence of *Toxoplasma* infection has been reported in different risk groups [12-15]. Although latent *Toxoplasma* infection has great importance among HIV infected people, it has been poorly studied. Therefore, this study was conducted to determine the sero-prevalence of latent *T. gondii* infection among HIV-infected and HIV-uninfected subjects in order to get some baseline information, from which clinical implication may be drawn.

## Methods

From 24 January 2007 to 15 February 2007, blood samples were collected from a total of 305 HIV-positive and 315 HIV-negative clients seeking either HIV or immunological testing at St. Paul's Hospital, Addis Ababa, Ethiopia. Parts of separated sera were originally used for studying seroprevalence of hepatitis B virus (HBV) infection and leftover samples were stored at -70°C for further investigations. Detailed description of methods used to study HBV infection was published elsewhere [16].

HIV and HBV sero-status of all samples were, therefore, known and selected sera from each HIV serogroup were used for the purpose of studying seroprevalence of *Toxoplasma* infection. In the present study, because of limited reagent kits, we only included the first 330 consecutive serum samples, 165 from each HIV sero-group.

Laboratory investigation of *Toxoplasma* infection was carried out with in six months after initial blood collection. Frozen sera were thawed at room temperature and those having unsuitable appearance in terms of turbidity and hemolysis were not included. Sera were tested in duplicate for anti-*Toxoplasma* IgG antibody using the Enzyme Linked Immunosorbent Assay (BioCheck, Inc, CA, USA). Positive and negative controls were included per each batch of test run to ensure kits were working properly and

technical procedures were carried out correctly. As per the instruction of the manufacturer, the mean absorbance value of each sample was divided by the cut - off calibrator mean value to obtain a Toxo G Index. A sample was considered positive for anti-*Toxoplasma* IgG antibody whenever a Toxo G Index value is equal or greater than 1.0 (> 32 IU/ml).

*Toxoplasma* infection was assessed against socio-demographic characteristics, HIV and hepatitis B core antibody (anti-HBc) serostatus. HBV-related risk factors including history of blood transfusion, tattooing, blood letting, multiple sexual partners, and surgery was also analyzed if they have association with *Toxoplasma* infection.

Data entry and analysis was performed using SPSS Version-12 software. Summaries were presented in terms of mean, range and percentage as appropriate. Differences in proportions were evaluated by Pearson's Chi-square test. In cases in which more than 20% of the expected values were less than 5, Fisher's exact test was used in lieu of chi-square test. Bivariate and Multivariate logistic regression analyses were used to assess the crude and adjusted effect of HIV and other correlates on *Toxoplasma* infection. Odds ratio was used to measure the strength of association between the *Toxoplasma* infection and its correlates. A statistical test result was considered significant whenever a p-value was < 0.05.

Institutional ethical clearance and appropriate written informed consent was obtained from each participant to use surplus sera for the present study.

## Results

A total of 330 sera were included for studying sero-prevalence of latent *Toxoplasma* infection among people with and without HIV infection. Half of the samples (50%) were HIV positives and 44.8% (148/330) had anti-HBc marker. The mean age of HIV infected participants was 35 years (range 20-66 years; SD 9.3) compared with mean age of 25 years (range 17-64 years; SD 7.3) in HIV uninfected clients. The male to female ratio was 0.8:1 in HIV positive and 1:1 in HIV negative individuals.

The cumulative sero-prevalence of anti-*Toxoplasma* IgG antibody among the study subjects was 90.0% (297/330). In bivariate analysis, rate of *Toxoplasma* infection was significantly higher among males than females (93.6% versus 86.6%;  $p = 0.04$ ). In addition, rate of *Toxoplasma* infection was shown to increase significantly with age ( $p = 0.003$ ) (Table 1).

The prevalence of latent *Toxoplasma* infection was 93.3% (154/165) among HIV positive and 86.7% (143/165) among HIV negative participants. This difference was also

**Table 1: Sero-prevalence of anti-Toxoplasma IgG antibody in relation to socio-demographic characteristics of the study subjects, Addis Ababa, 2007.**

Characteristics	Total (%)	Number (%) positive for anti-Toxoplasma IgG antibody	P-value
Residence			
Rural	24 (7.3)	24 (100)	0.15
Urban	306 (92.7)	273 (89.2)	
Sex			
Male	156 (47.3)	146 (93.6)	0.04
Female	174 (52.7)	151 (86.8)	
Age (years)			
17-24	106 (32.1)	87 (82.1)	0.003
25-34	132 (40.0)	122 (92.4)	
≥ 35	92 (27.9)	88 (95.7)	
Marital status			
Married	112 (33.9)	105 (93.8)	0.07
Single	167 (50.6)	144 (86.2)	
Divorced/Widowed	51 (15.5)	48 (94.1)	
Religion			
Christian	286 (86.7)	254 (88.8)	0.10
Muslim	44 (13.3)	43 (97.7)	
Ethnicity			
Amhara	148 (44.8)	131 (88.5)	0.70
Oromo	94 (28.5)	84 (89.4)	
Gurage	55 (16.7)	51 (92.7)	
Other	33 (10.0)	31 (93.9)	
Occupation			
Gov't/private employee	82 (24.8)	75 (91.5)	0.17
Housewife/house maid	56 (17.0)	51 (91.1)	
Student	59 (17.9)	48 (81.4)	
Merchant	40 (12.1)	38 (95.0)	
No work	93 (28.2)	85 (91.4)	
Educational status			
Illiterate	37 (11.2)	36 (97.3)	0.09
Primary school	117 (35.5)	108 (92.3)	
Secondary school and above	176 (53.3)	153 (86.9)	

statistically significant in bivariate analysis ( $p = 0.04$ ). Regardless of their HIV status, anti-Toxoplasma IgG antibody was more frequently detected in sera with anti-HBc

antibody compared with those with no anti-HBc serological marker (95.3% versus 85.7%,  $p = 0.004$ ). However, similar rate of anti-Toxoplasma IgG antibodies was

**Table 2: Multivariate analysis on seemingly significant predictors of *T. gondii* infection in bivariate analysis, Addis Ababa, 2007**

Characteristics	Crude odds ratio (95% confidence interval)	Adjusted odds ratio (95% confidence interval)	P-value
Sex			
Female	1	1	0.13
Male	2.22 (1.02-4.83)	1.9 (0.84-4.32)	
Age (years)			
17-24	1	1	0.07
25-34	2.66 (1.18-6.01)	2.33 (0.93-5.81)	
≥ 35	4.81 (1.57-14.70)	2.96 (0.81-10.81)	0.10
HIV status			
Negative	1	1	0.56
positive	2.15 (1.01-4.60)	1.32 (0.52-3.33)	
Anti-HBc status			
Negative	1	1	0.03
Positive	3.56 (1.41-7.97)	2.71 (1.12-6.57)	

detected in sera positive for both HBV and HIV compared to those with HBV alone (48.7% versus 46.2%,  $p > 0.05$ ). Any of HBV-related risk factors analyzed to have association with *Toxoplasma* infection were not found to be significant.

In further analysis, after adjustment for those significantly associated variables using the multivariate logistic regression, age, gender and HIV status were not found to affect rate of *Toxoplasma* infection. However, sera with anti-HBc antibody were shown more likely to have anti-*Toxoplasma* IgG antibody (OR 2.71, CI 1.12 to 6.57) (Table 2).

## Discussion

The sero-prevalence of anti-*Toxoplasma* IgG antibody among HIV-infected and HIV-uninfected participants was determined in the present study. The overall prevalence rate of latent *T. gondii* infection was found to be 90%. The high prevalence of anti-*Toxoplasma* IgG antibody observed in this hospital based study was in agreement with the result of a previous survey carried out in the general population of Ethiopia (75%) [13]. Similar sero-prevalence of *T. gondii* infection (80%) was also reported in a study, which involved HIV-infected and HIV-uninfected people in Addis Ababa [15]. The high sero-prevalence of latent *Toxoplasma* infection among the study population seems reasonable as raw or insufficiently cooked meat prepared in a various favorite cultural food is consumed. In addition, cats are abundant to cause environmental contamination and the climate is favorable to favor survival of the parasite.

In this study, sero-prevalence of latent *T. gondii* infection was shown to increase with age though the difference was not statistically significant. Similarly, no association between prevalence of latent *Toxoplasma* infection and age was reported elsewhere [17]. In contrast, significantly higher risk of having latent *Toxoplasma* infection was observed with increasing age [18,19]. The non-significant effect of age on *T. gondii* infection in our study subjects may suggest that exposure was irrespective of the risk behaviors pertinent to specific age group.

Moreover, the present study showed that HIV-positive and HIV-negative subjects had similar exposure to *T. gondii* infection, as reported by others [15,17,18]. The high prevalence of latent *T. gondii* infection in these risk groups, particularly among HIV-infected people, is of great concern. Evidences have shown that high prevalence of latent *T. gondii* infection and level of immunodeficiency in a given population are directly related to the incidence of TE [7,20]. It has been estimated that approximately one-third of HIV positive people with latent *T. gondii* infection will develop toxoplasmosis [11]. Accordingly, high incidence

rate of TE in our study population is expected unless preventive measures are undertaken.

However, our result indicating the more frequent occurrence of latent *T. gondii* infection among people with anti-HBc antibody needs further study, as a similar observation is not yet reported, in which additional results may confirm or challenge our finding. The lack of association between *Toxoplasma* serostatus and HBV-related risk-behaviors including blood transfusion may be the role of these routes in the transmission of *Toxoplasma* infection was non-significant among the study population.

Finally, because this study was not specifically designed to study toxoplasmosis, we missed data on basic risk factors such as cat ownership, dietary habits, soil exposure and other important risk factors for disease acquisition. Moreover, because of limited resources, sufficient number of sera was not included so that the study has lower statistical power. Our findings, therefore, should be interpreted in light of the study limitations.

In conclusion, the sero-prevalence of latent *T. gondii* infection was high and similar among HIV-infected and HIV-uninfected people. Therefore, burden of *T. gondii* infection should not be overlooked and educating people to prevent acquisition of new infection has great importance. Efforts should also be made to minimize the risk of subsequent reactivation and disease manifestation among HIV-positive people with latent *Toxoplasma* infection.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

TS, AT and BT designed the study protocol. TS and ET performed data analysis. MT carried out the laboratory work. All authors wrote the manuscript.

## Acknowledgements

The authors would like to thank Mr. Oscar Kay and Mrs. Anna Pao (Bio-check, Inc, USA) for their kind supply of reagent kits. We would also like to acknowledge Ethiopian Health and Nutrition Research Institute for laboratory facilities.

## References

1. Kloos H, Berhane Y: **Zoonotic diseases of public health importance. In Epidemiology and Ecology of Health and Diseases in Ethiopia.** 1st edition. Edited by: Berhane Y, Hailemariam D, Kloos H. Addis Ababa: Shama books; 2006:692-700.
2. Motoya JG, Liesenfeld O: **Toxoplasmosis.** *Lancet* 2004, **363**:1965-74.
3. Dubey JP: **Toxoplasmosis - a waterborne zoonosis.** *Vet Parasitol* 2004, **126**:57-72.
4. Dawson D: **Foodborne protozoan parasites.** *Int J Food Microbiol* 2005, **103**:207-227.
5. Weigel RM, Dubey JP, Dyer D, Siegel AM: **Risk factors for infection with *Toxoplasma gondii* for residents and workers on swine farms in Illinois.** *Am J Trop Med Hyg* 1999, **60**:793-798.

6. Alvarado-Esquivel C, Sifuentes-Alvarez A, Narro-Duarte SG, Estrada-Martínez S, Díaz-García JH, Liesenfeld O, Martínez-García SA, Canales-Molina A: **Seroepidemiology of *Toxoplasma gondii* infection in pregnant women in a public hospital in northern Mexico.** *BMC Infect Dis* 2006, **6**:1-7.
7. Trikha I, Wig N: **Management of Toxoplasmosis in AIDS.** *Indian J Med Sci* 2001, **55**:87-98.
8. Cheesbrough M: **District Laboratory Practice in Tropical Countries.** In Part 1 2nd edition. UK: Cambridge University press; 2005.
9. Luft BJ, Remington JS: **Toxoplasmic encephalitis in AIDS.** *Clin Infect Dis* 1992, **15**:211-222.
10. Porter SB, Sande MA: **Toxoplasmosis of the central nervous system in the acquired immune deficiency syndrome.** *N Engl J Med* 1992, **327**:1643-1648.
11. Grant IH, Gold WM, Rosenblum M, Niedzwieki D, Armstrong D: ***Toxoplasma gondii* serology in HIV infected patients: the development of central nervous system toxoplasmosis in AIDS.** *AIDS* 1990, **4**:519-521.
12. Tsega E, Behailu A: **Toxoplasmosis in hospitalized Ethiopians with lymphadenopathy.** *East Afr Med J* 1980, **57**:35-38.
13. Guebre-xabier M, Nurilign A, Gebre-Hiwot A, Hailu A, Sissay Y, Getachew E, Frommel D: **Seroepidemiological survey of *Toxoplasma gondii* infection in Ethiopia.** *Ethiopian Med J* 1993, **31**:201-208.
14. Eshete H, Tessema S, Abebe A, Seitz HM: **Some notes on toxoplasmosis in pregnant women in Ethiopia.** *Ethiopian Med J* 1994, **32**:135-136.
15. Woldemichael T, Fontanet AL, Sahlu T, Gilis H, Messele T, Rinke de Wit TF, Yeneneh H, Coutinho RA, Gool TV: **Evaluation of the Eiken latex agglutination test for anti-*Toxoplasma* antibodies and seroprevalence of *Toxoplasma* infection among factory workers in Addis Ababa, Ethiopia.** *Trans R Soc Trop Med Hyg* 1998, **92**:401-403.
16. Shimelis T, Torben W, Medhin G, Tebeje M, Andualm A, Demessie F, Mulu A, Tegbaru B, Gebre-Selassie S: **Hepatitis B virus infection among people attending the voluntary counseling and testing center and antiretroviral therapy clinic at St Paul's general specialized hospital Addis Ababa Ethiopia.** *Sex Transm Infect* 2007, **84**:37-41.
17. Nissapatorn V, Kamarulzaman A, Init I, Tan LH, Rohela M, Norliza A, Chan LL, Latt HM, Anuar AK, Quek KF: **Seroepidemiology of toxoplasmosis among HIV-infected patients and healthy blood donors.** *Med J Malaysia* 2002, **57**:304-310.
18. Falusi O, French AL, Seaberg EC, Tien PC, Watts DH, Minkoff H, Piesens E, Kovacs A, Anastos K, Cohen MH: **Prevalence and Predictors of *Toxoplasma* Seropositivity in Women with and at Risk for Human Immunodeficiency Virus Infection.** *Clin Infect Dis* 2002, **35**:1414-1417.
19. Jones JL, Kruszon-Moran D, Wilson M, McQuillan G, Navin T, McAuley JB: ***Toxoplasma gondii* Infection in the United States: Seroprevalence and Risk Factors.** *Am J Epidemiol* 2001, **154**(4):351-364.
20. Davarpanah MA, Mehrabani D, Neirami R, Ghahremanpoori M, Darvishi M: **Toxoplasmosis in HIV/AIDS patients in Shiraz, southern Iran.** *Iranian Red Crescent Med J* 2007, **9**(1):22-27.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

