



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: November 13, 2013

REVISED: March 23, 2014

ACCEPTED: April 3, 2014

PUBLISHED: April 30, 2014

Gravitational dynamics from entanglement “thermodynamics”

Nima Lashkari, Michael B. McDermott and Mark Van Raamsdonk

*Department of Physics and Astronomy, University of British Columbia,
6224 Agricultural Road, Vancouver, B.C., V6T 1W9, Canada*

E-mail: nlashkari@gmail.com, mbmcdermott@gmail.com, mav@phas.ubc.ca

ABSTRACT: In a general conformal field theory, perturbations to the vacuum state obey the relation $\delta S = \delta E$, where δS is the change in entanglement entropy of an arbitrary ball-shaped region, and δE is the change in “hyperbolic” energy of this region. In this note, we show that for holographic conformal field theories, this relation, together with the holographic connection between entanglement entropies and areas of extremal surfaces and the standard connection between the field theory stress tensor and the boundary behavior of the metric, implies that geometry dual to the perturbed state satisfies Einstein’s equations expanded to linear order about pure AdS.

KEYWORDS: Gauge-gravity correspondence, AdS-CFT Correspondence

ARXIV EPRINT: [1308.3716](https://arxiv.org/abs/1308.3716)

JHEP04(2014)195

Contents

1	Introduction	1
2	Entropy-energy relation	2
3	Gravitational implications of $dS = dE$ in holographic theories	4
3.1	Gravitational calculation of dS	4
3.2	Gravitational calculation of dE	5
4	Derivation of linearized Einstein's equations from $dE = dS$	5
4.1	Proof that $\delta S = \delta E$ for solutions of Einstein's equations	6
4.2	Proof that $\delta S = \delta E$ implies the linearized Einstein's equations	8
5	Discussion	11
A	Alternative derivation of linearized Einstein's equations from $\delta E = \delta S$	12
A.1	Expansion of $\delta E = \delta S$ in powers of R	12
A.2	Checking that solutions of Einstein's equations satisfy $\delta S = \delta E$	13

1 Introduction

Since the first connections between gravity and thermodynamics were realized in the study of black hole physics [1–3], various attempts have been made to derive Einstein's equations from the thermodynamics of some underlying degrees of freedom, starting with Jacobson's intriguing paper [4] (see also [5, 6]). With the AdS/CFT correspondence [7, 8], the underlying degrees of freedom for certain theories of gravity with AdS asymptotics have been explicitly identified as the degrees of freedom of a conformal field theory. It is thus interesting to ask whether the Einstein's equations in the gravitational theory can be derived from some thermodynamic relations for the CFT degrees of freedom.

In this note, following [9–13] we demonstrate that at least to linear order in perturbations around pure AdS, Einstein's equations do follow from a relation $dE = dS$ closely related to the First Law of Thermodynamics, but where the entropy S is the *entanglement entropy* of a spatial region in the field theory, and E is a certain energy associated with this region. A key point is that dS and dE can be defined and the relation $dS = dE$ shown to hold for *arbitrary* perturbations around the vacuum state; thus, the relation is more general than the ordinary first law which applies only in situations of thermodynamic equilibrium.

The specific relation we employ, which we write as

$$\delta S_A = \delta E_A^{\text{hyp}} \tag{1.1}$$

was derived recently by Blanco, Casini, Hung, and Myers in [13]. Here A represents a ball-shaped spatial region, δS_A represents the change in entanglement entropy of the region A

relative to the vacuum state, and δE_A^{hyp} represents the “hyperbolic” energy of the perturbed state in the region A , the expectation value of an operator which maps to the Hamiltonian of the CFT on hyperbolic space times time under a conformal transformation that takes the domain of dependence of the region A to $H^d \times \text{time}$. We review the derivation of this relation in section 2 below.

For holographic conformal field theories, each side of (1.1) has an interpretation in the dual gravity theory. Assuming that the perturbed state $|\Psi\rangle$ corresponds to some weakly-curved classical spacetime, the entanglement entropy S_A may be calculated (at the leading order in the $1/N$ to which we work) via the Ryu-Takayanagi proposal [9] and its covariant generalization [10] as the area of an extremal surface in the bulk, as we review in section 3.1. In section 3.2, we recall that the energy δE_A can be calculated from the asymptotic behavior of the metric. Thus, the field theory relation $\delta S_A = \delta E_A^{\text{hyp}}$ translates to a constraint on the dual geometry.

In section 4, we show that this constraint is precisely that the bulk metric corresponding to $|\Psi\rangle$ must satisfy Einstein’s equations to linear order in the perturbation around pure AdS (the geometry corresponding to the CFT vacuum state). That solutions of Einstein’s equations satisfy $\delta S_A = \delta E_A^{\text{hyp}}$ has already been shown in [13] (see also the related earlier work [12, 14, 15]). For completeness, we provide an alternate demonstration of this in section 4.1. In section 4.2, we go the other direction, showing that any perturbation to pure AdS satisfying $\delta S_A = \delta E_A^{\text{hyp}}$ must satisfy Einstein’s equations. This requires more than simply reversing the arguments of section 4.2 (or of [13]). In particular, demanding that $\delta S_A = \delta E_A^{\text{hyp}}$ for all ball-shaped spatial regions A in a particular Lorentz frame only places mild constraints on the metric, determining the combination $H_{xx} + H_{yy}$ in terms of the other components. It is only when we demand that $\delta S_A = \delta E_A^{\text{hyp}}$ in an arbitrary Lorentz frame (i.e. for ball-shaped regions on arbitrary spatial slices) that the full set of linearized Einstein’s equations is implied.

In appendix A, we give an alternative proof that Einstein’s equations imply $\delta S_A = \delta E_A^{\text{hyp}}$ that is perhaps more straightforward, but assumes that the metric is analytic.

We conclude in section 5 with a discussion.

2 Entropy-energy relation

In this section, we review the relation $\delta S_A = \delta E_A^{\text{hyp}}$, derived by Blanco, Casini, Hung, and Myers in [13] as a special case of an inequality that follows from the positivity of relative entropy.

General expression for variation of the entanglement entropy. Consider a CFT on $R^{d,1}$ in some state $|\Psi\rangle$. Choosing a spatial region A , define ρ_A to be the reduced density matrix associated with this region for the state $|\Psi\rangle$,

$$\rho_A = \text{tr}_{\bar{A}} |\Psi\rangle\langle\Psi| .$$

From this, we can define the modular Hamiltonian H_A by

$$\rho_A = e^{-H_A} .$$

For general states, this modular Hamiltonian is not related to the usual Hamiltonian, and cannot be written as the integral of a local density. We now consider an arbitrary variation of the state $|\Psi\rangle$. The change in entanglement entropy S_A for the region A is given by

$$\begin{aligned} \delta S_A &= \delta(-\text{tr}(\rho_A \log \rho_A)) \\ &= -\text{tr}(\delta\rho_A \log \rho_A) \\ &= \text{tr}(\delta\rho_A H_A) \\ &= \delta\langle H_A \rangle \end{aligned}$$

where we have used the fact that $\text{tr}(\delta\rho_A) = 0$, a consequence of assuming that the density matrix has a fixed normalization. In the last line, H_A is the original modular Hamiltonian associated with the density matrix ρ_A for the original state. Thus, we have the general relation

$$\delta S_A = \delta\langle H_A \rangle, \tag{2.1}$$

valid in any spatial region A for arbitrary perturbations of an arbitrary state.

“Thermodynamic” relation for perturbations around the vacuum state. We now specialize to the case where $|\Psi\rangle$ is the vacuum state, and the region A is a ball of radius R . In this case, the domain of dependence of the ball-shaped region¹ can be mapped by a conformal transformation to hyperbolic space times time. As shown in [11], such a transformation maps the vacuum density matrix for the region A to the thermal density matrix $e^{-\beta H_{\text{hyp}}}$ for the hyperbolic space theory, where the temperature is related to the hyperbolic space curvature radius R_H by $\beta = 2\pi R_{\text{hyp}}$. In this case H_{hyp} is the integral of the local operator T_{hyp}^{00} over hyperbolic space. Mapping back to the ball-shaped region of Minkowski space, it follows [11] that the modular Hamiltonian can be written as

$$H_A^{\text{vac}} = 2\pi \int_A d^d x \frac{R^2 - r^2}{2R} T^{00}$$

where T^{00} is the energy density operator for the CFT and r is a radial coordinate centered at the center of the ball.

In this case, we have

$$\delta\langle H_A \rangle = 2\pi \int_A d^d x \frac{R^2 - r^2}{2R} \delta T^{00} \equiv \delta E_A^{\text{hyp}}, \tag{2.2}$$

i.e. the variation in the expectation value of the vacuum modular Hamiltonian H_A^{vac} under a small perturbation away from the vacuum state is equal to the change in the “hyperbolic” energy of the region. Thus, the general relation (2.1) gives

$$\delta S_A = \delta E_A^{\text{hyp}}, \tag{2.3}$$

reminiscent of the First Law of Thermodynamics. We emphasize however that the entanglement entropy S_A can be defined for any state, in contrast to the usual thermodynamic entropy which applies to equilibrium states. Thus, (2.3) represents a much more general result.

¹The domain of dependence of A is the set of points p for which all inextendible causal curves passing through p also pass through A .

3 Gravitational implications of $dS = dE$ in holographic theories

Let us now consider the case of a holographic conformal field theory on Minkowski space, whose states correspond to asymptotically AdS spacetimes in some quantum theory of gravity. In this case, each side of the relation $\delta S_A = \delta E_A^{\text{hyp}}$ has a straightforward gravitational interpretation. As we review below, the left side may be calculated using the Ryu-Takayanagi proposal [9, 10], while the right side can be calculated from the asymptotic form of the metric. The equality of these quantities represents a constraint on the gravitational dynamics implied by the dual field theory. In the next section, we show that this constraint is precisely equivalent to Einstein's equations linearized about AdS.

3.1 Gravitational calculation of dS

According to the Ryu-Takayanagi proposal [9] and its covariant generalization [10], the entanglement entropy S_A for a state with a geometrical gravity dual is proportional² to the area of the extremal co-dimension two surface \tilde{A} in the bulk whose boundary coincides with the boundary of the region A on the AdS boundary,

$$S_A = \frac{\text{Area}(\tilde{A})}{4G_N} .$$

The surface \tilde{A} is an extremum of the area functional

$$A(G, X_{\text{ext}}) = \int d^d \sigma \sqrt{g}$$

where

$$g = \det(g_{ab}) = \det(G_{\mu\nu}) \frac{dX^\mu}{d\sigma^a} \frac{dX^\nu}{d\sigma^b} .$$

Starting from pure AdS, with metric³

$$ds^2 = G_{\mu\nu}^0 dx^\mu dx^\nu = \frac{1}{z^2} (-dt^2 + dz^2 + d\vec{x}^2) \tag{3.1}$$

the extremal surface ending on the spatial boundary sphere of radius R is described by the spacetime surface

$$\vec{x}^2 + z^2 = R^2 . \tag{3.2}$$

We now consider a small variation

$$G_{\mu\nu} = G_{\mu\nu}^0 + \delta G_{\mu\nu} . \tag{3.3}$$

In this case, the extremal surface changes, and the new area is

$$A(G_0 + \delta G, X_{\text{ext}}^0 + \delta X)$$

²Here, we are working to leading order in $1/N$. See the discussion section for comments on $1/N$ corrections.

³Throughout this paper, we set the AdS radius to one.

where the variation δX will be of order δG . Since the original surface was extremal, we have

$$A(G_0, X_{\text{ext}}^0 + \delta X) = A(G_0, X_{\text{ext}}^0) + \mathcal{O}(\delta X^2) .$$

Thus, the variation of the surface gives rise to changes in the area that start at order δG^2 . To find the order δG variation of the area, we need only evaluate

$$A(G_0 + \delta G, X_{\text{ext}}^0) - A(G, X_{\text{ext}}^0)$$

expanded to linear order in δG . We find that

$$\delta A = \int d^d \sigma \frac{1}{2} \sqrt{g_0} g_0^{ab} \delta g_{ab} , \tag{3.4}$$

where we have used lower-case letters to represent pullbacks to the extremal surface. Thus, for field theory state $|\Psi\rangle$ close to the vacuum state with dual geometry described by (3.3), the change in the entanglement entropy for region A relative to the vacuum state is given by an integral of the metric perturbation over the original extremal surface \tilde{A} . Using the explicit metric (3.1) and parameterizing the extremal surface (3.2) by the boundary coordinates x^i , we have finally that

$$\delta S = \frac{R}{8G_N} \int d^d x \left(\delta_{ij} - \frac{1}{R^2} x_i x_j \right) H_{ij} . \tag{3.5}$$

3.2 Gravitational calculation of dE

General asymptotically AdS spacetimes with a Minkowski space boundary geometry may be described using Fefferman-Graham coordinates by a metric

$$ds^2 = \frac{1}{z^2} (dz^2 + dx_\mu dx^\mu + z^d H_{\mu\nu}(x, z) dx^\mu dx^\nu) . \tag{3.6}$$

where pure AdS, dual to the CFT vacuum, corresponds to $H_{\mu\nu} = 0$. With this parametrization, the expectation value $t_{\mu\nu}$ of the field theory stress-energy tensor is simply related to the asymptotic metric by [16, 17]

$$t_{\mu\nu}(x) = \frac{d+1}{16\pi G_N} H_{\mu\nu}(z=0, x) .$$

Thus, we may write the change in the hyperbolic energy (2.2) relative to the vacuum state as

$$\delta E_A^{\text{hyp}} = \frac{d+1}{16G_N} \int_A d^d x \frac{R^2 - r^2}{R} \delta H_{00}(0, x) . \tag{3.7}$$

This is an integral of the boundary value of H over the region A .

4 Derivation of linearized Einstein's equations from $dE = dS$

We are now ready to demonstrate that using the holographic dictionary reviewed in the previous section, the CFT relation $\delta S_A = \delta E_A^{\text{hyp}}$ is equivalent to the constraint that metric corresponding to the perturbed CFT satisfies Einstein's equations to linear order. For

clarity, we focus on the case of 2+1 dimensional conformal field theories, corresponding to gravitational theories with four non-compact dimensions. However, the result can also be proven for general higher-dimensional theories.

Using the results (3.5) and (3.7), the CFT relation $\delta S_A = \delta E_A^{\text{hyp}}$ implies that a disk of any radius R centered at any point (x_0, y_0) on the boundary, the integral

$$\delta \hat{S} = \int_{D_R} dx dy \left\{ H_{xx}(\sqrt{R^2 - x^2 - y^2}, t, x + x_0, y + y_0)(R^2 - x^2) \right. \\ \left. + H_{yy}(\sqrt{R^2 - x^2 - y^2}, t, x + x_0, y + y_0)(R^2 - y^2) \right. \\ \left. - 2H_{xy}(\sqrt{R^2 - x^2 - y^2}, t, x + x_0, y + y_0)xy \right\} \quad (4.1)$$

over the bulk extremal surface must equal the integral

$$\delta \hat{E} = \frac{3}{2} \int_{D_R} dx dy (R^2 - x^2 - y^2) H_{tt}(0, t, x + x_0, y + y_0) \quad (4.2)$$

over the $z = 0$ surface, where we have absorbed a factor of $1/8G_N R$ to define $\delta \hat{S}(R, x_0, y_0)$ and $\delta \hat{E}(R, x_0, y_0)$ (we drop the hats from now on). We will now show that this equality is true for all disks *in all Lorentz frames* if and only if the bulk metric satisfies Einstein's equations to linear order in H . As shown in [13], these are equivalent to the set of equations

$$H_\alpha{}^\alpha = 0 \quad \partial_\mu H^{\mu\nu} = 0 \quad \frac{1}{z^4} \partial_z \{ z^4 \partial_z H_{\mu\nu} \} + \partial^2 H_{\mu\nu} = 0 \quad (4.3)$$

that arise by plugging the Fefferman-Graham form of the metric (3.6) into the zz , $z\mu$, and $\mu\nu$ components of Einstein's equations

$$W_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R - 3g_{\mu\nu} = 0,$$

respectively and using the fact that H is regular at $z = 0$. In (4.3), the last equation is equivalent to saying that each component of $z^3 H$ must satisfy the Laplace equation on the AdS background.

4.1 Proof that $\delta S = \delta E$ for solutions of Einstein's equations

We begin by showing that solutions of the linearized Einstein's equations obey the equality $\delta S = \delta E$. This has already been checked in section 3.1 of [13] by demonstrating the result for a complete basis of solutions to the equations (4.3). In this section, we offer an alternative proof that does not require using an explicit basis of solutions. A third proof that is perhaps more straightforward but assumes a series expansion of H is given in appendix A.

Using the equations (4.3), we have:

$$\begin{aligned} \partial_t^2 H_{tt} &= \partial_t^2 (H_{xx} + H_{yy}) \\ \Rightarrow \partial_t (\partial_x H_{xt} + \partial_y H_{yt}) &= \partial_t^2 (H_{xx} + H_{yy}) \\ \Rightarrow \partial_x^2 H_{xx} + \partial_y^2 H_{yy} + 2\partial_x \partial_y H_{xy} &= \partial_t^2 (H_{xx} + H_{yy}) \\ \Rightarrow \partial_x^2 H_{xx} + \partial_y^2 H_{yy} + 2\partial_x \partial_y H_{xy} &= (\partial_x^2 + \partial_y^2)(H_{xx} + H_{yy}) + \frac{1}{z^4} \partial_z (z^4 \partial_z (H_{xx} + H_{yy})) \\ \Rightarrow 2\partial_x \partial_y H_{xy} &= \partial_x^2 H_{xx} + \partial_y^2 H_{yy} + \frac{1}{z^4} \partial_z (z^4 \partial_z (H_{xx} + H_{yy})). \end{aligned} \quad (4.4)$$

We would like to use the last equation to eliminate H_{xy} from (4.1). However, we have H_{xy} rather than $\partial_x \partial_y H_{xy}$ in (4.1). To make progress, we begin by differentiating δS by x_0 and y_0 (the coordinates of the center of the boundary disk). This gives

$$\begin{aligned} \partial_{x_0} \partial_{y_0} \delta S = \int_{D_R} dx dy \left\{ \partial_x \partial_y H_{xx}(\sqrt{R^2 - x^2 - y^2}, t, x + x_0, y + y_0)(R^2 - x^2) \right. \\ \left. + \partial_x \partial_y H_{yy}(\sqrt{R^2 - x^2 - y^2}, t, x + x_0, y + y_0)(R^2 - y^2) \right. \\ \left. - 2 \partial_x \partial_y H_{xy}(\sqrt{R^2 - x^2 - y^2}, t, x + x_0, y + y_0)xy \right\}. \end{aligned} \quad (4.5)$$

Now, using (4.4), we have

$$\begin{aligned} \partial_{x_0} \partial_{y_0} \delta S = \int_{D_R} dx dy \left\{ \partial_x \partial_y H_{xx}(R^2 - x^2) + \partial_x \partial_y H_{yy}(R^2 - y^2) \right. \\ \left. - xy \left(\partial_y^2 H_{xx} + \partial_x^2 H_{yy} + \frac{1}{z^4} \partial_z (z^4 \partial_z (H_{xx} + H_{yy})) \right) \right\}. \end{aligned} \quad (4.6)$$

It is straightforward to check that this expression is equal to the integral over the extremal surface of an exact form dA , where A is defined for all (x, y, z, t) as

$$\begin{aligned} A = & (-xz \partial_z H_{xx} - 3x H_{xx} + z^2 \partial_x H_{yy}) dx \\ & + (z^2 \partial_y H_{xx} - yz \partial_z H_{yy} - 3y H_{yy}) dy \\ & + (-yz \partial_y H_{xx} - xz \partial_x H_{yy}) dz. \end{aligned} \quad (4.7)$$

By Stokes theorem, this equals the integral of A over the boundary of the extremal surface, so we have

$$\begin{aligned} \partial_{x_0} \partial_{y_0} \delta S &= \int_{\partial D_R} A \\ &= -3 \int_{\partial D_R} (x H_{xx} dx + y H_{yy} dy) \\ &= 3 \int d\theta (H_{xx} + H_{yy}) \cos(\theta) \sin(\theta). \end{aligned}$$

In the second step, we have used the fact that all other terms in A vanish for $z = 0$.

Similarly, we find that $\partial_{x_0} \partial_{y_0} \delta E$ may be written as

$$\begin{aligned} \partial_{x_0} \partial_{y_0} \delta E &= \frac{3}{2} \int_{D_R} dx dy \partial_{x_0} \partial_{y_0} H_{tt}(0, t, x + x_0, y + y_0)(R^2 - x^2 - y^2) \\ &= \frac{3}{2} \int_{D_R} dx dy \partial_x \partial_y (H_{xx}(0, t, x + x_0, y + y_0) \\ &\quad + H_{yy}(0, t, x + x_0, y + y_0))(R^2 - x^2 - y^2) \\ &= \frac{3}{2} \int_{D_R} d\hat{A}, \end{aligned}$$

where we can choose

$$\hat{A} = (-2x H_{xx} + (R^2 - x^2 - y^2) \partial_x H_{yy}) dx + (-2y H_{yy} + (R^2 - x^2 - y^2) \partial_y H_{xx}) dy.$$

Again, using Stokes theorem, this reduces to the integral of $(3/2)\hat{A}$ over the boundary, so

$$\begin{aligned} \partial_{x_0}\partial_{y_0}\delta E &= \frac{3}{2} \int_{\partial D_R} \hat{A} \\ &= -3 \int_{\partial D_R} (xH_{xx}dx + yH_{yy}dy) \\ &= \partial_{x_0}\partial_{y_0}\delta S. \end{aligned}$$

We conclude that for any H satisfying Einstein's equations,

$$\delta S(x_0, y_0, R; H) - \delta E(x_0, y_0, R; H) = C_x(x_0, R; H) + C_y(y_0, R; H),$$

where C_x and C_y are some functionals linear in H that do not depend on y_0 or x_0 respectively. Now, consider the class of functions H that vanish for sufficiently large $x_0^2 + y_0^2$ at the time $t = 0$ where we evaluate δS and δE . In this case, fixing any x_0 and taking $y_0 \rightarrow \infty$ or fixing any y_0 and taking $x_0 \rightarrow \infty$, the left side must vanish. For this to be true on the right side, both C_x and C_y must be constant (as functions of x_0 and y_0), with $C_x + C_y = 0$. Thus, the right side vanishes for any H that vanishes as $x_0^2 + y_0^2 \rightarrow \infty$. But more general H can be written as linear combinations of such functions, and since the right side is a linear functional in H , it must vanish for all H . This completes the argument that $\delta S_A = \delta E_A^{\text{hyp}}$ for solutions of Einstein's equations.

4.2 Proof that $\delta S = \delta E$ implies the linearized Einstein's equations

In this section, we go the other direction to show that the relation $\delta S = \delta E$ implies that the metric satisfies Einstein's equations to linear order, i.e. that the equivalence of (4.1) and (4.2) implies the relations (4.3).

Given the boundary stress tensor $t_{\mu\nu}$, let $H_{\mu\nu}^{EE}$ be the corresponding metric perturbation that follows from Einstein's equations, i.e. the solution of (4.3) satisfying $H_{\mu\nu}^{EE}(0, t, x, y) = (16\pi G_N/3)t_{\mu\nu}$. We will show that there is no other H with these boundary conditions for which $\delta S = \delta E$ in all frames of reference.

Suppose there were another H for which $\delta S = \delta E$ for all disk shaped regions in all Lorentz frames. Then the difference $\Delta = H - H^{EE}$ must satisfy

$$\Delta_{\mu\nu}(z = 0, t, x, y) = 0, \tag{4.8}$$

and

$$\begin{aligned} 0 = \int_{D_R} dx dy \left\{ \Delta_{xx} \left(\sqrt{R^2 - x^2 - y^2}, x + x_0, y + y_0 \right) \left(1 - \frac{x^2}{R^2} \right) \right. \\ \left. + \Delta_{yy} \left(\sqrt{R^2 - x^2 - y^2}, x + x_0, y + y_0 \right) \left(1 - \frac{y^2}{R^2} \right) \right. \\ \left. - 2\Delta_{xy} \left(\sqrt{R^2 - x^2 - y^2}, x + x_0, y + y_0 \right) \frac{xy}{R^2} \right\} \end{aligned} \tag{4.9}$$

for arbitrary R , x_0 , and y_0 , and in an arbitrary Lorentz frame.

Let us first see the consequences of demanding this result in a fixed frame. To begin, we note that (4.9) may be expanded in powers of R using the basic integral

$$\int_{D_R} dx dy (R^2 - x^2 - y^2)^{\frac{n}{2}} x^{2m_x} y^{2m_y} = R^{n+2m_x+2m_y+2} I_{n,m_x,m_y},$$

where

$$I_{n,m_x,m_y} = \frac{\Gamma(m_x + \frac{1}{2})\Gamma(m_y + \frac{1}{2})\Gamma(\frac{n}{2} + 1)}{\Gamma(\frac{n}{2} + m_x + m_y + 2)}. \quad (4.10)$$

Defining

$$\Delta_{\mu\nu}(z, x, y) = \sum_{n=0}^{\infty} z^n \Delta_{\mu\nu}^{(n)}(x, y) \quad (4.11)$$

we find that (4.9) becomes⁴

$$0 = \sum R^{n+2m_x+2m_y+2} \left\{ \frac{1}{(2m_x)!(2m_y)!} \partial_x^{2m_x} \partial_y^{2m_y} \Delta_{xx}^{(n)}(t, x_0, y_0) (I_{n,m_x,m_y} - I_{n,m_x+1,m_y}) \right. \\ + \frac{1}{(2m_x)!(2m_y)!} \partial_x^{2m_x} \partial_y^{2m_y} \Delta_{yy}^{(n)}(t, x_0, y_0) (I_{n,m_x,m_y} - I_{n,m_x,m_y+1}) \\ \left. - 2R^2 \frac{1}{(2m_x+1)!(2m_y+1)!} \partial_x^{2m_x+1} \partial_y^{2m_y+1} \Delta_{xy}^{(n)}(t, x_0, y_0) I_{n,m_x+1,m_y+1} \right\}. \quad (4.12)$$

The vanishing of the terms at order R^{N+2} implies that

$$\Delta_{xx}^{(N)}(t, x_0, y_0) + \Delta_{yy}^{(N)}(t, x_0, y_0) = \sum_{(m_x,m_y) \neq (0,0)} C_{xx}^{N,m_x,m_y} \partial_x^{2m_x} \partial_y^{2m_y} \Delta_{xx}^{(N-2m_x-2m_y)} \\ + C_{yy}^{N,m_x,m_y} \partial_x^{2m_x} \partial_y^{2m_y} \Delta_{yy}^{(N-2m_x-2m_y)} \\ + C_{xy}^{N,m_x,m_y} \partial_x^{2m_x-1} \partial_y^{2m_y-1} \Delta_{xy}^{(N-2m_x-2m_y)},$$

where the C coefficients can be read off from (4.12). As examples, the first few equations give

$$\begin{aligned} \Delta_{xx}^{(0)}(t, x_0, y_0) + \Delta_{yy}^{(0)}(t, x_0, y_0) &= 0 \\ \Delta_{xx}^{(1)}(t, x_0, y_0) + \Delta_{yy}^{(1)}(t, x_0, y_0) &= 0 \\ \Delta_{xx}^{(2)}(t, x_0, y_0) + \Delta_{yy}^{(2)}(t, x_0, y_0) &= -\frac{1}{4}(\partial_y^2 \Delta_{xx}^{(0)}(t, x_0, y_0) + \partial_x^2 \Delta_{yy}^{(0)}(t, x_0, y_0)) \\ &\quad - \frac{3}{20}(\partial_x^2 \Delta_{xx}^{(0)}(t, x_0, y_0) + \partial_y^2 \Delta_{yy}^{(0)}(t, x_0, y_0)) \\ &\quad + \frac{1}{5} \partial_x \partial_y \Delta_{xy}^{(0)}(t, x_0, y_0) \\ \Delta_{xx}^{(3)}(t, x_0, y_0) + \Delta_{yy}^{(3)}(t, x_0, y_0) &= -\frac{1}{6}(\partial_y^2 \Delta_{xx}^{(1)}(t, x_0, y_0) + \partial_x^2 \Delta_{yy}^{(1)}(t, x_0, y_0)) \\ &\quad - \frac{1}{6}(\partial_x^2 \Delta_{xx}^{(1)}(t, x_0, y_0) + \partial_y^2 \Delta_{yy}^{(1)}(t, x_0, y_0)) \\ &\quad + \frac{1}{9} \partial_x \partial_y \Delta_{xy}^{(1)}(t, x_0, y_0). \end{aligned} \quad (4.13)$$

⁴Here, we are assuming that the function Δ is analytic. It would be useful to find a derivation of our result that holds more generally.

We see that this set of equations completely determines the combination $\Delta_{xx} + \Delta_{yy}$ at each order in z in terms of the lower order terms in the expansion of Δ . However, apart from the constraint (4.8) on the boundary behavior (equivalent to $\Delta_{\mu\nu}^{(0)} = 0$), the remaining elements of $\Delta_{\mu\nu}$ are completely unconstrained.

To constrain $\Delta_{\mu\nu}$ further, we need to use the requirement that the relation (4.9) should hold in an arbitrary Lorentz frame. Thus, for each choice of reference frame, we will have equations analogous to (4.13). Specifically, consider a general boost

$$\Lambda = \begin{pmatrix} \gamma & \gamma\beta_x & \gamma\beta_y \\ \gamma\beta_x & 1 + \beta_x^2 \frac{\gamma^2}{\gamma+1} & \beta_x\beta_y \frac{\gamma^2}{\gamma+1} \\ \gamma\beta_y & \beta_x\beta_y \frac{\gamma^2}{\gamma+1} & 1 + \beta_y^2 \frac{\gamma^2}{\gamma+1} \end{pmatrix}.$$

In the equations for a general frame of reference obtained by such a boost, the left sides in (4.13) will be replaced by

$$\Lambda_x^\mu \Lambda_x^\nu \Delta_{\mu\nu} + \Lambda_y^\mu \Lambda_y^\nu \Delta_{\mu\nu}.$$

Up to an overall constant factor, this gives

$$\Delta_{ii} + 2\beta_i \Delta_{it} + \beta^2 \left(\Delta_{tt} - \frac{1}{2} \Delta_{ii} \right) + \left(\beta_i \beta_j - \frac{1}{2} \delta_{ij} \beta^2 \right) \Delta_{ij}.$$

Now, consider the general version of the second equation in (4.13) (the first equation already holds by (4.8)). This requires the vanishing of

$$\Delta_{ii}^{(1)} + 2\beta_i \Delta_{it}^{(1)} + \beta^2 \left(\Delta_{tt}^{(1)} - \frac{1}{2} \Delta_{ii}^{(1)} \right) + \left(\beta_i \beta_j - \frac{1}{2} \delta_{ij} \beta^2 \right) \Delta_{ij}^{(1)}.$$

For a fixed x_0 and y_0 , this is a polynomial in β_i that must vanish for all values of β_i . Thus, the polynomial must be identically zero. At order β^0 , this gives

$$\Delta_{ii}^{(1)}(t, x_0, y_0) = 0$$

as we had before. At order β , we get

$$\Delta_{it}^{(1)}(t, x_0, y_0) = 0.$$

At order β^2 , this gives

$$\Delta_{tt}^{(1)}(t, x_0, y_0) = \frac{1}{2} \Delta_{ii}^{(1)}(t, x_0, y_0) = 0$$

and

$$\Delta_{ij}^{(1)}(t, x_0, y_0) - \frac{1}{2} \Delta_{ij} \Delta_{kk}^{(1)}(t, x_0, y_0) = 0.$$

Thus, we have $\Delta_{\mu\nu}^{(1)} = 0$. We can now continue to analyze the remaining equations in (4.13) in turn. Supposing that we have shown $\Delta_{\mu\nu}^{(k)} = 0$ for $k < n$, the general version of the n th equation in (4.13) requires the vanishing of

$$\Delta_{ii}^{(n)} + 2\beta_i \Delta_{it}^{(n)} + \beta^2 \left(\Delta_{tt}^{(n)} - \frac{1}{2} \Delta_{ii}^{(n)} \right) + \left(\beta_i \beta_j - \frac{1}{2} \delta_{ij} \beta^2 \right) \Delta_{ij}^{(n)},$$

since the right hand side in (4.13) will be zero. Repeating the analysis above, we conclude that $\Delta_{\mu\nu}^{(n)} = 0$. By induction, this holds for all n , so we have shown that $\Delta_{\mu\nu} = 0$, completing the proof.

5 Discussion

In this paper, we have seen that to linear order in perturbations about the vacuum state, the emergence of gravitational dynamics in the theory dual to a holographic CFT is directly related to a general relation satisfied by CFT entanglement entropies on ball-shaped regions. This relation is closely related to the First Law of Thermodynamics, but is more general since it applies to arbitrary perturbations of the state rather than perturbations for which the system remains in thermal equilibrium.

While the CFT relation (1.1) is an exact equivalence, we have made use of this relation only at the leading order in $1/N$ where the entanglement entropy maps over to the extremal surface area. This corresponds to working in the classical limit in the bulk. According to [18], $1/N$ corrections to the CFT entanglement entropy correspond to bulk quantum corrections including the entropy of entanglement of bulk quantum fields across the extremal surface. It will be interesting to understand the implications of the CFT relation (1.1) beyond the classical level in the bulk, but we leave this for future work.

The derivations in section 4 were written specifically for the case of four-dimensional gravity. However, the proof given in [13] that Einstein's equations imply $\delta S = \delta E$, and our method of proof in section 4.2 that $\delta S = \delta E$ implies the linearized Einstein's equations work for general dimensions.⁵

The linearized Einstein's equations we derived are for the metric components in the field theory directions and radial direction of the bulk. Any additional fields in the gravitational theory, including metric components in any compactified directions, are not constrained by the CFT relation we have considered. At linear order, the equations for these fields decouple from the linearized Einstein's equations for the metric in the non-compact directions. Thus, we can say that the universal relation $\delta S = \delta E$ is equivalent to the universal sector of the linearized bulk equations.

Our results do not imply that all holographic theories are dual to gravitational theories whose metric perturbations satisfy Einstein's equations. In this paper, we assumed that entanglement entropies are related to areas via the usual Ryu-Takayanagi formula, and that the stress-energy tensor in the dual field theory is related to the asymptotic form of the metric. In more general theories, the entanglement entropy may correspond to a more complicated functional of the bulk geometry and the relation between the stress tensor and asymptotic metric may be modified. In these cases, we expect that the bulk equations will be different, for example involving α' corrections with higher-derivative terms. However, it may be possible following the methods in this paper to derive the linearized version of these more general equations given a particular choice for the holographic entanglement entropy formula and the holographic formula for the stress tensor.

It will be interesting to see whether the first non-linear corrections to Einstein's equations in the bulk are equivalent to some simple property of entanglement entropies.

Finally, we comment on the relation to the work of Jacobson [4], which partly motivated our investigations. Jacobson realized that Einstein's equations could be derived from the

⁵Specifically, eq. (4.9) becomes $0 = \int_{D_R} d^d x (\Delta_{ii} - x_i x_j / R^2 \Delta_{ij})$; expanding this in powers of R using the generalization of eq. (4.10) yields at each order in R an equation that relates $\Delta_{ii}(n)$ to quantities calculated from Δ at lower orders in n . The steps in the proof are as before.

assumption that the energy flux through a part of any bulk Rindler horizon gives rise to a proportional local change in area of this horizon. Interpreting the area as an entropy, such a relation looks like the first law of thermodynamics. However, in Jacobson’s work, it was not clear why areas of segments of an arbitrary bulk Rindler horizon (not necessarily associated with any black hole) should correspond to an entropy, so the origin of the thermodynamic relation remained mysterious.

In our case, the “thermodynamic relation” $dS = dE$ is an exact quantum relation (i.e. not really thermodynamics) derived to hold for the underlying fundamental degrees of freedom associated with our gravitational system. Thus, while our final result (in contrast to Jacobson’s work) applies so far only at the linearized level, the starting point is well understood. In detail, the bulk interpretation of our $dS = dE$ relation is somewhat different than Jacobson’s starting point (the bulk surfaces/horizons we deal with are global rather than local and the energy has a different interpretation), but the two relations were similar enough to motivate the question of whether Einstein’s equations could be derived from the first law of [13].

Acknowledgments

This research was supported in part by the Natural Sciences and Engineering Research Council of Canada.

A Alternative derivation of linearized Einstein’s equations from $\delta E = \delta S$

In this appendix, we offer an alternative proof that solutions of Einstein’s equations satisfy $\delta S_A = \delta E_A^{\text{hyp}}$. This proof replaces $\delta S_A = \delta E_A^{\text{hyp}}$ with the infinite set of relations obtained by matching the terms in the power series expansion of this relation in R , the radius of the disk A , as we did in section 4.2.

A.1 Expansion of $\delta E = \delta S$ in powers of R

To begin, we expand both (4.1) and (4.2) in powers of R . Defining

$$H_{\mu\nu}(z, x, y) = \sum_{n=0}^{\infty} z^n H_{\mu\nu}^{(n)}(x, y) \tag{A.1}$$

we have

$$\delta E = \frac{3}{2} \sum_{m_x, m_y=0} R^{2+2m_x+2m_y} I_{2, m_x, m_y} \partial_x^{2m_x} \partial_y^{2m_y} H_{tt}^{(0)}(t, x_0, y_0) \tag{A.2}$$

while

$$\delta S = \sum R^{n+2m_x+2m_y+2} \left\{ \frac{1}{(2m_x)!(2m_y)!} \partial_x^{2m_x} \partial_y^{2m_y} H_{xx}^{(n)}(t, x_0, y_0) (I_{n, m_x, m_y} - I_{n, m_x+1, m_y}) \right. \tag{A.3}$$

$$+ \frac{1}{(2m_x)!(2m_y)!} \partial_x^{2m_x} \partial_y^{2m_y} H_{yy}^{(n)}(t, x_0, y_0) (I_{n, m_x, m_y} - I_{n, m_x, m_y+1})$$

$$\left. - 2R^2 \frac{1}{(2m_x+1)!(2m_y+1)!} \partial_x^{2m_x+1} \partial_y^{2m_y+1} H_{xy}^{(n)}(t, x_0, y_0) I_{n, m_x+1, m_y+1} \right\}$$

where I was defined in (4.10).

A.2 Checking that solutions of Einstein's equations satisfy $\delta S = \delta E$

Using these expansions, it is straightforward to verify that any solution of the linearized Einstein's equations (4.3) satisfies $\delta E = \delta S$, as was done originally in [13] and by another alternative approach in section 4.

Using the expansion (A.1), the equations (4.3) become

$$H_{tt}^{(n)} = H_{xx}^{(n)} + H_{yy}^{(n)} \quad (\text{A.4})$$

$$\partial_t H_{tt}^{(n)} = \partial_x H_{tx}^{(n)} + \partial_y H_{ty}^{(n)} \quad (\text{A.5})$$

$$\partial_t H_{tx}^{(n)} = \partial_x H_{xx}^{(n)} + \partial_y H_{xy}^{(n)} \quad (\text{A.6})$$

$$\partial_t H_{ty}^{(n)} = \partial_x H_{xy}^{(n)} + \partial_y H_{yy}^{(n)} \quad (\text{A.7})$$

$$H_{\mu\nu}^{(n)} = \frac{1}{n(n+3)} (\partial_t^2 - \partial_x^2 - \partial_y^2) H_{\mu\nu}^{(n-2)} \quad n \geq 2 \quad (\text{A.8})$$

$$H_{\mu\nu}^{(1)} = 0. \quad (\text{A.9})$$

Starting with (A.4) and then using (A.5), (A.6), (A.7), and finally (A.8), we find:

$$\begin{aligned} \partial_t^2 H_{tt}^{(n)} &= \partial_t^2 (H_{xx}^{(n)} + H_{yy}^{(n)}) \\ \Rightarrow \partial_t (\partial_x H_{xt}^{(n)} + \partial_y H_{yt}^{(n)}) &= \partial_t^2 (H_{xx}^{(n)} + H_{yy}^{(n)}) \\ \Rightarrow \partial_x^2 H_{xx}^{(n)} + \partial_y^2 H_{yy}^{(n)} + 2\partial_x \partial_y H_{xy}^{(n)} &= \partial_t^2 (H_{xx}^{(n)} + H_{yy}^{(n)}) \\ \Rightarrow \partial_x^2 H_{xx}^{(n)} + \partial_y^2 H_{yy}^{(n)} + 2\partial_x \partial_y H_{xy}^{(n)} &= \partial_t^2 (H_{xx}^{(n)} + H_{yy}^{(n)}) \\ \Rightarrow \partial_x^2 H_{xx}^{(n)} + \partial_y^2 H_{yy}^{(n)} + 2\partial_x \partial_y H_{xy}^{(n)} &= (\partial_x^2 + \partial_y^2) (H_{xx}^{(n)} + H_{yy}^{(n)}) \\ &\quad + (n+2)(n+5) (H_{xx}^{(n+2)} + H_{yy}^{(n+2)}) \\ \Rightarrow 2\partial_x \partial_y H_{xy}^{(n)} &= \partial_y^2 H_{xx}^{(n)} + \partial_x^2 H_{yy}^{(n)} + (n+2)(n+5) (H_{xx}^{(n+2)} + H_{yy}^{(n+2)}). \end{aligned}$$

Using this last equation, we can eliminate $H_{xy}^{(n)}$ from (A.3). This gives

$$\delta S = \sum R^{n+2m_x+2m_y+2} \left\{ \frac{1}{(2m_x)!(2m_y)!} \partial_x^{2m_x} \partial_y^{2m_y} H_{xx}^{(n)}(t, x_0, y_0) C_{n,m_x,m_y}^{xx} \right. \\ \left. + \frac{1}{(2m_x)!(2m_y)!} \partial_x^{2m_x} \partial_y^{2m_y} H_{yy}^{(n)}(t, x_0, y_0) C_{n,m_x,m_y}^{yy} \right\} \quad (\text{A.10})$$

where for $n \geq 2$ we have

$$\begin{aligned} C_{n,m_x,m_y}^{xx} &= I_{n,m_x,m_y} - I_{n,m_x+1,m_y} - \frac{2m_y}{2m_x+1} I_{n,m_x+1,m_y} \\ &\quad - \frac{n(n+3)}{(2m_x+1)(2m_y+1)} I_{n-2,m_x+1,m_y+1} \\ &= 0 \end{aligned}$$

$$\begin{aligned} C_{n,m_x,m_y}^{yy} &= I_{n,m_x,m_y} - I_{n,m_x,m_y+1} - \frac{2m_x}{2m_y+1} I_{n,m_x,m_y+1} \\ &\quad - \frac{n(n+3)}{(2m_x+1)(2m_y+1)} I_{n-2,m_x+1,m_y+1} \\ &= 0 \end{aligned}$$

while for $n = 1$ and $n = 0$, we have

$$C_{1,m_x,m_y}^{xx} = I_{1,m_x,m_y} - I_{1,m_x+1,m_y} - \frac{2m_y}{2m_x+1} I_{1,m_x+1,m_y} = \frac{4}{3} I_{3,m_x,m_y}$$

$$C_{1,m_x,m_y}^{yy} = I_{1,m_x,m_y} - I_{1,m_x,m_y+1} - \frac{2m_x}{2m_y+1} I_{1,m_x,m_y+1} = \frac{4}{3} I_{3,m_x,m_y}$$

and

$$C_{0,m_x,m_y}^{xx} = I_{0,m_x,m_y} - I_{0,m_x+1,m_y} - \frac{2m_y}{2m_x+1} I_{0,m_x+1,m_y} = \frac{3}{2} I_{2,m_x,m_y}$$

$$C_{0,m_x,m_y}^{yy} = I_{0,m_x,m_y} - I_{0,m_x,m_y+1} - \frac{2m_x}{2m_y+1} I_{0,m_x,m_y+1} = \frac{3}{2} I_{2,m_x,m_y}.$$

In each case, we have made simplifications using the definition (4.10) of I . Using these results together with (A.9), we find that (A.10) simplifies to

$$\begin{aligned} \delta S &= \sum R^{2m_x+2m_y+2} \frac{1}{(2m_x)!(2m_y)!} \partial_x^{2m_x} \partial_y^{2m_y} (H_{xx}^{(0)}(t, x_0, y_0) + H_{yy}^{(0)}(t, x_0, y_0)) \left(\frac{3}{2} I_{2,m_x,m_y} \right) \\ &= \frac{3}{2} \sum R^{2m_x+2m_y+2} \frac{1}{(2m_x)!(2m_y)!} \partial_x^{2m_x} \partial_y^{2m_y} H_{tt}^{(0)}(t, x_0, y_0) I_{2,m_x,m_y} \\ &= \delta E. \end{aligned}$$

Thus, we have verified that $\delta S = \delta E$ for linearized solutions of Einstein's equations, providing an alternate argument to the one in [13].

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J.D. Bekenstein, *Black holes and entropy*, *Phys. Rev. D* **7** (1973) 2333 [[INSPIRE](#)].
- [2] J.M. Bardeen, B. Carter and S.W. Hawking, *The four laws of black hole mechanics*, *Commun. Math. Phys.* **31** (1973) 161 [[INSPIRE](#)].
- [3] S.W. Hawking, *Particle creation by black holes*, *Commun. Math. Phys.* **43** (1975) 199 [*Erratum ibid.* **46** (1976) 206-206] [[INSPIRE](#)].
- [4] T. Jacobson, *Thermodynamics of space-time: The Einstein equation of state*, *Phys. Rev. Lett.* **75** (1995) 1260 [[gr-qc/9504004](#)] [[INSPIRE](#)].
- [5] T. Padmanabhan, *Thermodynamical aspects of gravity: new insights*, *Rept. Prog. Phys.* **73** (2010) 046901 [[arXiv:0911.5004](#)] [[INSPIRE](#)].
- [6] E.P. Verlinde, *On the origin of gravity and the laws of Newton*, *JHEP* **04** (2011) 029 [[arXiv:1001.0785](#)] [[INSPIRE](#)].
- [7] J.M. Maldacena, *The large- N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [*Int. J. Theor. Phys.* **38** (1999) 1113] [[hep-th/9711200](#)] [[INSPIRE](#)].

- [8] O. Aharony, S.S. Gubser, J.M. Maldacena, H. Ooguri and Y. Oz, *Large- N field theories, string theory and gravity*, *Phys. Rept.* **323** (2000) 183 [[hep-th/9905111](#)] [[INSPIRE](#)].
- [9] S. Ryu and T. Takayanagi, *Holographic derivation of entanglement entropy from AdS/CFT*, *Phys. Rev. Lett.* **96** (2006) 181602 [[hep-th/0603001](#)] [[INSPIRE](#)].
- [10] V.E. Hubeny, M. Rangamani and T. Takayanagi, *A covariant holographic entanglement entropy proposal*, *JHEP* **07** (2007) 062 [[arXiv:0705.0016](#)] [[INSPIRE](#)].
- [11] H. Casini, M. Huerta and R.C. Myers, *Towards a derivation of holographic entanglement entropy*, *JHEP* **05** (2011) 036 [[arXiv:1102.0440](#)] [[INSPIRE](#)].
- [12] M. Nozaki, T. Numasawa, A. Prudenziati and T. Takayanagi, *Dynamics of entanglement entropy from Einstein equation*, *Phys. Rev. D* **88** (2013) 026012 [[arXiv:1304.7100](#)] [[INSPIRE](#)].
- [13] D.D. Blanco, H. Casini, L.-Y. Hung and R.C. Myers, *Relative entropy and holography*, *JHEP* **08** (2013) 060 [[arXiv:1305.3182](#)] [[INSPIRE](#)].
- [14] D. Allahbakhshi, M. Alishahiha and A. Naseh, *Entanglement thermodynamics*, *JHEP* **08** (2013) 102 [[arXiv:1305.2728](#)] [[INSPIRE](#)].
- [15] G. Wong, I. Klich, L.A. Pando Zayas and D. Vaman, *Entanglement temperature and entanglement entropy of excited states*, *JHEP* **12** (2013) 020 [[arXiv:1305.3291](#)] [[INSPIRE](#)].
- [16] R.C. Myers, *Stress tensors and Casimir energies in the AdS/CFT correspondence*, *Phys. Rev. D* **60** (1999) 046002 [[hep-th/9903203](#)] [[INSPIRE](#)].
- [17] S. de Haro, S.N. Solodukhin and K. Skenderis, *Holographic reconstruction of space-time and renormalization in the AdS/CFT correspondence*, *Commun. Math. Phys.* **217** (2001) 595 [[hep-th/0002230](#)] [[INSPIRE](#)].
- [18] T. Faulkner, A. Lewkowycz and J. Maldacena, *Quantum corrections to holographic entanglement entropy*, *JHEP* **11** (2013) 074 [[arXiv:1307.2892](#)] [[INSPIRE](#)].