

## POSTER PRESENTATION


## Open Access

# Looking at the role of direct and indirect pathways in basal ganglia networks at different levels

Rahmi Elibol\*, Neslihan Serap Şengör

*From* 24th Annual Computational Neuroscience Meeting: CNS\*2015  
Prague, Czech Republic. 18-23 July 2015

The role of basal ganglia in motor action initiation and selection has been well studied and now it is evident that impairment in this structure causes not only movement disorders as Parkinson's disease, Huntington's disease but also behavioral dysfunctions as addiction, obsessive-compulsive disorder [1-4]. In order to understand the mechanisms giving rise to motor actions, cognitive processes related to these actions as decision making and the diseases occurring due to malfunctioning of these structures, various computational models of direct and indirect pathways have been proposed [5-9]. Here, in order to set a simple relation between models of basal ganglia at different levels, a simple mass model indicating the controversial role of direct and indirect pathways will be introduced first. While dopamine (DA) in direct pathway enhances the activity in Thalamus giving rise to inhibition of action, arise of DA in indirect pathway disinhibits Thalamus activity promoting the action to take place. This activity can be followed from Figure 1 for different DA levels. Based on the results of this simple mass model, spiking neural network (SNN) is built by point neurons and the relation between the local field potential of this SNN and simple mass model will be discussed. The aim is to build a simple relation between different levels of computational models which would help investigating the mechanisms behind the cognitive processes without engaging in detailed models initially. Thus, the simple mass model proposed would be primary model giving a chance to test the initial interpretation of the concepts formed and lead to setting up more detailed, realistic models.



**Figure 1** DA level is normal, high and low at top, middle and bottom, respectively. THL activity is normal, high and low similar to STRD1 activity and contrary to STRD2 activity.

\* Correspondence: [rahmielibol@itu.edu.tr](mailto:rahmielibol@itu.edu.tr)  
Electronics and Communication Engineering, Istanbul Technical University,  
Istanbul, Turkey

Published: 18 December 2015

## References

1. Kropotov J, Etlinger S: **Selection of actions in the basal ganglia-thalamocortical circuits: Review and model.** *International Journal of Psychophysiology* 1999, **31**(3):197-217.
2. DeLong MR, Wichmann T: **Circuits and circuit disorders of the basal ganglia.** *Archives of Neurology* 2007, **64**(1):20-24.
3. Nicola S: **The nucleus accumbens as part of a basal ganglia action selection circuit.** *Psychopharmacology* 2007, **191**(3):521-550.
4. Graybiel AM, Rauch SL: **Toward a Neurobiology of Obsessive-Compulsive Disorder.** *Neuron* 2000, **28**(2):343-347.
5. Terman D, Rubin JE, Yew AC, Wilson CJ: **Activity patterns in a model for the subthalamopallidal network of the basal ganglia.** *The Journal of Neuroscience* 2002, **22**(7):2963-2976.
6. Chersi F, Mirolii M, Pezzulo G, Baldassarre G: **A spiking neuron model of the cortico-basal ganglia circuits for goal-directed and habitual action learning.** *Neural Networks* 2013, **41**:212-224.
7. McCarthy MM, Moore-Kochlacs C, Gu X, Boyden ES, Han X, Kopell N: **Striatal origin of the pathologic beta oscillations in Parkinson's disease.** *Proceedings of the National Academy of Sciences* 2011, **108**(28):11620-11625.
8. Marreiros AC, Cagnan H, Moran RJ, Friston KJ, Brown P: **Basal ganglia cortical interactions in Parkinsonian patients.** *NeuroImage* 2013, **66**:301-310.
9. Yucelgen C, Denizdurduran B, Metin S, Elibol R, Sengor NS: **A biophysical network model displaying the role of basal ganglia pathways in action selection.** *Artificial Neural Networks and Machine Learning ICANN 2012*, **7552**:177-184.

doi:10.1186/1471-2202-16-S1-P225

**Cite this article as:** Elibol and Şengör: Looking at the role of direct and indirect pathways in basal ganglia networks at different levels. *BMC Neuroscience* 2015 **16**(Suppl 1):P225.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

