

AR-AR AGE AND HALOGEN CHARACTERISTICS OF NAKHLITE MIL 03346: RECORDS OF CRUSTAL PROCESSES ON MARS

M. Anand^{1,2}, R. Burgess³, V. Fernandes⁴ and M.M. Grady¹
¹Centre for Earth, Planetary, Space and Astronomical Research (CEPSAR), The Open University, Milton Keynes, UK. E-mail: M.Anand@open.ac.uk ²Department of Mineralogy, The Natural History Museum, London, UK. ³Department of Earth Sciences, University of Manchester, Manchester, UK. ⁴Instituto Geofísico, Universidade de Coimbra, Portugal

Introduction: MIL 03346 is one of the 7 members of the nakhlite group of Martian meteorites. The texture and mineralogy of this rock distinguishes it from other nakhlites [1]. Almost complete absence of crystalline plagioclase and ferrohedenbergitic rim on pyroxene phenocrysts are some of the unique characteristics of this sample. Previous Ar-Ar age study of MIL 03346 reported a total Ar-Ar age of 1.37 Ga [2]. Recent reports have also highlighted presence of exotic minerals such as K-Cl-rich amphibole in melt inclusions in pyroxenes and olivines in MIL 03346 [3]. The non-chondritic halogen ratios in nakhlites have been interpreted as a result of fluid activity on Mars [4]. In the present study we have combined measurements of step-heating Ar-Ar ages with that of halogen contents. The work was conducted on ~12 mg whole-rock sample, sliced out from 1g allocation of MIL 03346,37 from the MWG to the senior author.

Results: The total Ar-Ar age is 1360 ± 2 Ma, similar to that reported by [2]. The main release of K is between 600-900 °C and there is a clear decrease in age with temperature, most likely recoil-related. The major Cl release occurs at around 1000 °C, is therefore separate from K release, and accompanies the release of Ca from pyroxene. The measured halogen contents in MIL 03346 are high relative to other SNC meteorites, only lower than Nakhla [4]. The halogen data for MIL 03346 are: Cl = 156 ppm; Br = 0.41 ppm; I = 0.014 ppm. The halogen ratios of Nakhla are slightly higher than MIL (Br/Cl = 0.002; I/Cl = 6×10^{-5}) but this may be explained by high Br/Cl and I/Cl ratios of martian weathering components in alteration veins in Nakhla olivines [5]. For comparison the martian regolith Br/Cl = 0.007 and I/Cl = 16×10^{-5} [6] while shergottites are more similar at Br/Cl = 0.005; I/Cl = 3×10^{-5} .

The presence of K-Cl-rich amphibole in melt inclusions in pyroxene could explain the major Cl release at high temperature and some of the recoil effect if $^{39}\text{Ar}_K$ moves from the melt inclusions into the pyroxene. Alternatively, $^{39}\text{Ar}_K$ may also be released from the mesostasis glass but then we would also expect to see major Cl release at the same time. Between 600-850 °C 73% of total K and 26% of total Cl are released; >900 °C the values are 21% and 67%; so the majority of halogen release presumably comes from the melt inclusions. This indicates that the Br/Cl and I/Cl are representative of the melt (although they may originate from a soil contaminated melt).

References: [1] Anand M. et al., 2005. 36th Lunar and Planetary Science Conference. Abs # 1639. [2] Bogard, D.D. and Garrison, D.H. 2006. 37th Lunar and Planetary Science Conference. Abs # 1108. [3] Sautter V. et al. 2006. 37th Lunar and Planetary Science Conference. Abs# 1318. [4] Dreibus et al. 2006. 37th Lunar and Planetary Science Conference. Abs# 1180. [5] Sutton S.R. et al. 2002. 33rd Lunar and Planetary Science Conference. Abs # 1278. [6] Rao M.N. et al. 2002. *Icarus* 156, pp. 352-372.