



Dmitri Finkelshtein · Yuri Kondratiev · Oleksandr Kutoviy

Statistical dynamics of continuous systems: perturbative and approximative approaches

Received: 6 February 2014 / Accepted: 2 July 2014 / Published online: 30 July 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract We discuss general concept of Markov statistical dynamics in the continuum. For a class of spatial birth-and-death models, we develop a perturbative technique for the construction of statistical dynamics. Particular examples of such systems are considered. For the case of Glauber type dynamics in the continuum we describe a Markov chain approximation approach that gives more detailed information about statistical evolution in this model.

Mathematics Subject Classification 46E30 · 82C21 · 47D06

المخلص

نناقش مفهوماً عاماً لديناميكية ماركوف الإحصائية في المُتَّصِل. لصف خاص من نماذج الولادة والموت الفضائية، نطور تقنية تشويشية لتركيب ديناميكيات إحصائية. يتم اعتبار أمثلة محددة من هذه الأنظمة. في حالة الديناميكيات من نوع جلاوبر في المُتَّصِل، نصف مقارنة تقريب سلسلة ماركوف تعطينا معلومات مفصلة بشكل أكبر عن التجذير الإحصائي في هذا النموذج.

1 Introduction

Dynamics of interacting particle systems appear in several areas of the complex systems theory. In particular, we observe a growing activity in the study of Markov dynamics for continuous systems. The latter fact is motivated, in particular, by modern problems of mathematical physics, ecology, mathematical biology, and genetics, see, e.g. [27, 28, 31–34, 36–39, 51–53, 68] and literature cited therein. Moreover, Markov dynamics are used for the construction of social, economic, and demographic models. Note that Markov processes for continuous systems are considered in the stochastic analysis as dynamical point processes [43, 44, 46] and they appear even in the representation theory of big groups [10–14].

A mathematical formalization of the problem may be described as follows. As a phase space of the system we use the space $\Gamma(\mathbb{R}^d)$ of locally finite configurations in the Euclidean space \mathbb{R}^d . An heuristic Markov generator which describes the considered model is given by its expression on a proper set of functions (observables) over

D. Finkelshtein
Department of Mathematics, Swansea University, Singleton Park, Swansea SA2 8PP, UK
E-mail: d.i.finkelshtein@swansea.ac.uk

Y. Kondratiev (✉) · O. Kutoviy
Fakultät für Mathematik, Universität Bielefeld, 33615 Bielefeld, Germany
E-mail: kondrat@math.uni-bielefeld.de

O. Kutoviy
Department of Mathematics, Massachusetts Institute of Technology, 77 Massachusetts Avenue E18-420, Cambridge, MA, USA
E-mail: kutoviy@mit.edu; kutoviy@math.uni-bielefeld.de



$\Gamma(\mathbb{R}^d)$. With this operator we can relate two evolution equations, namely Kolmogorov backward equation for observables and Kolmogorov forward equation on probability measures on the phase space $\Gamma(\mathbb{R}^d)$ (macroscopic states of the system). The latter equation is also known as Fokker–Planck equation in the mathematical physics terminology. Compared with the usual situation in stochastic analysis, there is an essential technical difficulty: the corresponding Markov process in the configuration space may be constructed only in very special cases. As a result, a description of Markov dynamics in terms of random trajectories is absent for most models under considerations.

As an alternative approach we use a concept of statistical dynamics that substitutes the notion of a Markov stochastic process. A central object now is an evolution of states of the system that will be defined by mean of the Fokker–Planck equation. This evolution equation w.r.t. probability measures on $\Gamma(\mathbb{R}^d)$ may be reformulated as a hierarchical chain of equations for correlation functions of considered measures. Such kind of evolution equations are well known in the study of Hamiltonian dynamics for classical gases as BBGKY chains but now they appear as a tool for construction and analysis of Markov dynamics. As an essential technical step, we consider related pre-dual evolution chains of equations on the so-called quasi-observables. As it will be shown in the paper, such hierarchical equations may be analyzed in the framework of semigroup theory with the use of powerful techniques of perturbation theory for the semigroup generators, etc. Considering the dual evolution for the constructed semigroup on quasi-observables we then introduce the dynamics on correlation functions. The described scheme of the dynamics construction looks quite surprising because any perturbation techniques for initial Kolmogorov evolution equations one cannot expect. The point is that states of infinite interacting particle systems are given by measures which are, in general, orthogonal to each other. As a result, we cannot compare their evolutions or apply a perturbative approach. But under quite general assumptions they have correlation functions and corresponding dynamics may be considered in a common Banach space of correlation functions. A proper choice of this Banach space means, in fact, that we find an admissible class of initial states for which the statistical dynamics may be constructed. There we see again a crucial difference with the framework of Markov stochastic processes where the initial distribution evolution is defined for any initial data.

The structure of the paper is as follows. In Sect. 2 we discuss general concept of statistical dynamics for Markov evolutions in the continuum and introduce necessary mathematical structures. Then, in Sect. 3, this concept is applied to an important class of Markov dynamics of continuous systems, namely to birth-and-death models. Here general conditions for the existence of a semigroup evolution in a space of quasi-observables are obtained. Then we construct evolutions of correlation functions as dual objects. It is shown how to apply general results to the study of particular models of statistical dynamics coming from mathematical physics and ecology.

Finally, in Sect. 4 we describe an alternative technique for the construction of solutions to hierarchical chains evolution equations by means of an approximative approach. For concreteness, this approach is discussed in the case of the so-called Glauber-type dynamics in the continuum. We construct a family of Markov chains on configuration space in finite volumes with concrete transition kernels adopted to the Glauber dynamics. Then the solution to the hierarchical equation for correlation functions may be obtained as the limit of the corresponding object for the Markov chain dynamics. This limiting evolution generates the state dynamics. Moreover, in the uniqueness regime for the corresponding equilibrium measure of Glauber dynamics which is, in fact, Gibbs, dynamics of correlation functions is exponentially ergodic.

This paper is based on a series of our previous works [26, 28–30, 34, 53], but certain results and constructions are detailed and generalized, in particular, in more complete analysis of the dual dynamics on correlation functions.

2 Statistical description for stochastic dynamics of complex systems in the continuum

2.1 Complex systems in the continuum

In recent decades, different branches of natural and life sciences have been addressing to a unifying point of view on a number of phenomena occurring in systems composed of interacting subunits. This leads to formation of an interdisciplinary science which is referred to as the theory of complex systems. It provides reciprocation of concepts and tools involving wide spectrum of applications as well as various mathematical theories such that statistical mechanics, probability, nonlinear dynamics, chaos theory, numerical simulation, and many others.



Nowadays complex systems theory is a quickly growing interdisciplinary area with a very broad spectrum of motivations and applications. For instance, having in mind biological applications, Levin [61] characterized complex adaptive systems by such properties as diversity and individuality of components, localized interactions among components, and the outcomes of interactions used for replication or enhancement of components. We will use a more general informal description of a complex system as a specific collection of interacting elements which have so-called collective behavior. This means appearance of properties of the system which are not peculiar to inner nature of each element itself. The significant physical example of such properties is thermodynamical effects which were a basis for creation by Boltzmann of statistical physics as a mathematical language for studying complex systems of molecules.

We assume that all elements of a complex system are identical by properties and possibilities. Thus, one can model these elements as points in a proper space whereas the complex system will be modeled as a discrete set in this space. Mathematically this means that for study of complex systems a proper language and techniques are delivered by the interacting particle models which form a rich and powerful direction in modern stochastic and infinite dimensional analysis. Interacting particle systems have a wide use as models in condensed matter physics, chemical kinetics, population biology, ecology (individual based models), sociology, and economics (agent based models). For instance, a population in biology or ecology may be represented by a configuration of organisms located in a proper habitat.

In spite of completely different orders of numbers of elements in real physical, biological, social, and other systems (typical numbers start from 10^{23} for molecules and, say, 10^5 for plants) their complexities have analogous phenomena and need similar mathematical methods. One of them consists in mathematical approximation of a huge but finite real-world system by an infinite system realized in an infinite space. This approach was successfully applied to the thermodynamic limit for models of statistical physics and appeared quite useful for the ecological modeling in the infinite habitat to avoid boundary effects in a population evolution.

Therefore, our phase space for the mathematical description should consist of countable sets from an underlying space. This space itself may have discrete or continuous nature that leads to segregation of the world of complex systems on two big classes. Discrete models correspond to systems whose elements can occupy some prescribing countable set of positions, for example, vertices of the lattice \mathbb{Z}^d or, more generally, of some graph embedded to \mathbb{R}^d . These models are widely studied and the corresponding theories were realized in numerous publications, see, e.g. [62,63] and the references therein. Continuous models, or models in the continuum, were studied not so intensively and broadly. We concentrate our attention exactly on continuous models of systems whose elements may occupy any points in Euclidean space \mathbb{R}^d . (Note that most part of our results may be easily transferred to much more general underlying spaces). Having in mind that real elements have physical sizes we will consider only the so-called locally finite subsets of the underlying space \mathbb{R}^d that means that in any bounded region we assume to have finite number of elements. Another restriction will be prohibition of multiple elements at the same position of the space.

We will consider systems of elements of the same type only. The mathematical realization of considered approaches may be successfully extended to multi-type systems; meanwhile such systems will have richer qualitative properties and will be an object of interest for applications. Some particular results can be found, e.g. in [21,22,39].

2.2 Mathematical description for a complex systems

We proceed to the mathematical realization of complex systems.

Let $\mathcal{B}(\mathbb{R}^d)$ be the family of all Borel sets in \mathbb{R}^d , $d \geq 1$; $\mathcal{B}_b(\mathbb{R}^d)$ denotes the system of all bounded sets from $\mathcal{B}(\mathbb{R}^d)$.

The configuration space over space \mathbb{R}^d consists of all locally finite subsets (configurations) of \mathbb{R}^d , namely

$$\Gamma = \Gamma(\mathbb{R}^d) := \left\{ \gamma \subset \mathbb{R}^d \mid |\gamma_\Lambda| < \infty, \text{ for all } \Lambda \in \mathcal{B}_b(\mathbb{R}^d) \right\}.$$

Here $|\cdot|$ means the cardinality of a set, and $\gamma_\Lambda := \gamma \cap \Lambda$. We may identify each $\gamma \in \Gamma$ with the non-negative Radon measure $\sum_{x \in \gamma} \delta_x \in \mathcal{M}(\mathbb{R}^d)$, where δ_x is the Dirac measure with unit mass at x , $\sum_{x \in \emptyset} \delta_x$ is, by definition, the zero measure, and $\mathcal{M}(\mathbb{R}^d)$ denotes the space of all non-negative Radon measures on $\mathcal{B}(\mathbb{R}^d)$. This identification allows to endow Γ with the topology induced by the vague topology on $\mathcal{M}(\mathbb{R}^d)$, i.e. the weakest topology on Γ with respect to which all mappings

$$\Gamma \ni \gamma \mapsto \sum_{x \in \gamma} f(x) \in \mathbb{R} \tag{2.1}$$

are continuous for any $f \in C_0(\mathbb{R}^d)$ that is the set of all continuous functions on \mathbb{R}^d with compact supports. It is worth noting the vague topology may be metrizable in such a way that Γ becomes a Polish space (see, e.g. [50] and references therein).

Corresponding to the vague topology the Borel σ -algebra $\mathcal{B}(\Gamma)$ appears the smallest σ -algebra for which all mappings

$$\Gamma \ni \gamma \mapsto N_\Lambda(\gamma) := |\gamma_\Lambda| \in \mathbb{N}_0 := \mathbb{N} \cup \{0\} \tag{2.2}$$

are measurable for any $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$, see, e.g. [1]. This σ -algebra may be generated by the sets

$$Q(\Lambda, n) := \{ \gamma \in \Gamma \mid N_\Lambda(\gamma) = |\gamma_\Lambda| = n \}, \quad \Lambda \in \mathcal{B}_b(\mathbb{R}^d), n \in \mathbb{N}_0. \tag{2.3}$$

Clearly, for any $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$,

$$\Gamma = \bigsqcup_{n \in \mathbb{N}_0} Q(\Lambda, n).$$

Among all measurable functions $F : \Gamma \rightarrow \bar{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$ we mark out the set $\mathcal{F}_0(\Gamma)$ consisting of such of them for which $|F(\gamma)| < \infty$ at least for all $|\gamma| < \infty$. The important subset of $\mathcal{F}_0(\Gamma)$ formed by cylindric functions on Γ . Any such a function is characterized by a set $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ such that $F(\gamma) = F(\gamma_\Lambda)$ for all $\gamma \in \Gamma$. The class of cylindric functions we denote by $\mathcal{F}_{cyl}(\Gamma) \subset \mathcal{F}_0(\Gamma)$.

Functions on Γ are usually called *observables*. This notion is borrowed from statistical physics and means that typically in course of empirical investigation we may estimate, check, and see only some quantities of a whole system rather than look on the system itself.

Example 2.1 Let $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$ and consider the so-called *linear function* on Γ , cf. (2.1),

$$\langle \varphi, \gamma \rangle := \begin{cases} \sum_{x \in \gamma} \varphi(x), & \text{if } \sum_{x \in \gamma} |\varphi(x)| < \infty, \quad \gamma \in \Gamma, \\ +\infty, & \text{otherwise.} \end{cases}$$

Then, evidently, $\langle \varphi, \cdot \rangle \in \mathcal{F}_0(\Gamma)$. If, additionally, $\varphi \in C_0(\mathbb{R}^d)$, then $\langle \varphi, \cdot \rangle \in \mathcal{F}_{cyl}(\Gamma)$. Not that for, e.g. $\varphi(x) = \|x\|_{\mathbb{R}^d}$ (the Euclidean norm in \mathbb{R}^d) we have that $\langle \varphi, \gamma \rangle = \infty$ for any infinite $\gamma \in \Gamma$.

Example 2.2 Let $\phi : \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{R}$ be an even function, namely $\phi(-x) = \phi(x)$, $x \in \mathbb{R}^d$. Then one can consider the so-called *energy function*

$$E^\phi(\gamma) := \begin{cases} \sum_{\{x,y\} \subset \gamma} \phi(x-y), & \text{if } \sum_{\{x,y\} \subset \gamma} |\phi(x-y)| < \infty, \quad \gamma \in \Gamma, \\ +\infty, & \text{otherwise.} \end{cases} \tag{2.4}$$

Clearly, $E^\phi \in \mathcal{F}_0(\Gamma)$. However, even for ϕ with a compact support, E^ϕ will not be a cylindric function.

As we discussed before, any configuration γ represents some system of elements in a real-world application. Typically, investigators are not able to take into account exact positions of all elements due to huge number of them. For quantitative and qualitative analysis of a system researchers mostly need some its statistical characteristics such as density, correlations, spatial structures, and so on. This leads to the so-called statistical description of complex systems when people study distributions of countable sets in an underlying space instead of sets themselves. Moreover, the main idea in Boltzmann’s approach to thermodynamics based on giving up the description in terms of evolution for groups of molecules and using statistical interpretation of molecules motion laws. Therefore, the crucial role for studying of complex systems plays distributions (probability measures) on the space of configurations. In statistical physics these measures usually called *states* that accentuates their role for description of considered systems.

We denote the class of all probability measures on $(\Gamma, \mathcal{B}(\Gamma))$ by $\mathcal{M}^1(\Gamma)$. Given a distribution $\mu \in \mathcal{M}^1(\Gamma)$ one can consider a collection of random variables $N_\Lambda(\cdot)$, $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ defined in (2.2). They describe random

numbers of elements inside bounded regions. The natural assumption is that these random variables should have finite moments. Thus, we consider the class $\mathcal{M}_{\text{fin}}^1(\Gamma)$ of all measures from $\mathcal{M}^1(\Gamma)$ such that

$$\int_{\Gamma} |\gamma_{\Lambda}|^n d\mu(\gamma) < \infty, \quad \Lambda \in \mathcal{B}_b(\mathbb{R}^d), n \in \mathbb{N}. \tag{2.5}$$

Example 2.3 Let σ be a non-atomic Radon measure on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$. Then the *Poisson measure* π_{σ} with intensity measure σ is defined on $\mathcal{B}(\Gamma)$ by

$$\pi_{\sigma}(Q(\Lambda, n)) = \frac{(\sigma(\Lambda))^n}{n!} \exp\{-\sigma(\Lambda)\}, \quad \Lambda \in \mathcal{B}_b(\mathbb{R}^d), n \in \mathbb{N}_0. \tag{2.6}$$

This formula is nothing but the statement that the random variables N_{Λ} have Poissonian distribution with mean value $\sigma(\Lambda)$, $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$. Note that by the Rényi theorem [47, 74] a measure π_{σ} will be Poissonian if (2.6) holds for $n = 0$ only. In the case then $d\sigma(x) = \rho(x) dx$ one can say about nonhomogeneous Poisson measure π_{ρ} with density (or intensity) ρ . This notion goes back to the famous Campbell formula [15, 16] which states that

$$\int_{\Gamma} \langle \varphi, \gamma \rangle d\pi_{\rho}(\gamma) = \int_{\mathbb{R}^d} \varphi(x) \rho(x) dx, \tag{2.7}$$

if only the right-hand side of (2.7) is well defined. The generalization of (2.7) is the Mecke identity [65]

$$\int_{\Gamma} \sum_{x \in \gamma} h(x, \gamma) d\pi_{\sigma}(\gamma) = \int_{\Gamma} \int_{\mathbb{R}^d} h(x, \gamma \cup x) d\sigma(x) d\pi_{\sigma}(\gamma), \tag{2.8}$$

which holds for all measurable nonnegative functions $h : \mathbb{R}^d \times \Gamma \rightarrow \mathbb{R}$. Here and in the sequel we will omit brackets for the one-point set $\{x\}$. In [65], it was shown that the Mecke identity is a characterization identity for the Poisson measure. In the case $\rho(x) = z > 0$, $x \in \mathbb{R}^d$ one can say about the homogeneous Poisson distribution (measure) π_z with constant intensity z . We will omit sub-index for the case $z = 1$, namely $\pi := \pi_1 = \pi_{dx}$. Note that the property (2.5) is followed from (2.8) easily.

Example 2.4 Let ϕ be as in Example 2.2 and suppose that the energy given by (2.4) is *stable*: there exists $B \geq 0$ such that, for any $|\gamma| < \infty$, $E^{\phi}(\gamma) \geq -B|\gamma|$. An example of such ϕ may be given by the expansion

$$\phi(x) = \phi^+(x) + \phi^p(x), \quad x \in \mathbb{R}^d, \tag{2.9}$$

where $\phi^+ \geq 0$, whereas ϕ^p is a positive defined function on \mathbb{R}^d (the Fourier transform of a measure on \mathbb{R}^d), see, e.g. [40, 75]. Fix any $z > 0$ and define the *Gibbs measure* $\mu \in \mathcal{M}^1(\Gamma)$ with potential ϕ and activity parameter z as a measure which satisfies the following generalization of the Mecke identity:

$$\int_{\Gamma} \sum_{x \in \gamma} h(x, \gamma) d\mu(\gamma) = \int_{\Gamma} \int_{\mathbb{R}^d} h(x, \gamma \cup x) \exp\{-E^{\phi}(x, \gamma)\} z dx d\mu(\gamma), \tag{2.10}$$

where

$$E^{\phi}(x, \gamma) := \langle \phi(x - \cdot), \gamma \rangle = \sum_{y \in \gamma} \phi(x - y), \quad \gamma \in \Gamma, x \in \mathbb{R}^d \setminus \gamma. \tag{2.11}$$

The identity (2.10) is called the Georgii–Nguyen–Zessin identity, see [45, 67]. If potential ϕ is additionally satisfied the so-called integrability condition

$$\beta := \int_{\mathbb{R}^d} |e^{-\phi(x)} - 1| dx < \infty, \tag{2.12}$$

then it can be checked that the condition (2.5) for the Gibbs measure holds. Note that under conditions $z\beta \leq (2e)^{-1}$ there exists a unique measure on $(\Gamma, \mathcal{B}(\Gamma))$ which satisfies (2.10). Heuristically, the measure μ may be given by the formula

$$d\mu(\gamma) = \frac{1}{Z} e^{-E^{\phi}(\gamma)} d\pi_z(\gamma), \tag{2.13}$$

where Z is a normalizing factor. To give rigorous meaning for (2.13) it is possible to use the so-called DLR-approach (named after R. L. Dobrushin, O. Lanford, D. Ruelle), see, e.g. [2] and references therein. As was shown in [67], this approach gives the equivalent definition of the Gibbs measures which satisfies (2.10).

Note that (2.13) could have a rigorous sense if we restrict our attention on the space of configuration which belong to a bounded domain $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$. The space of such (finite) configurations will be denoted by $\Gamma(\Lambda)$. The σ -algebra $\mathcal{B}(\Gamma(\Lambda))$ may be generated by family of mappings $\Gamma(\Lambda) \ni \gamma \mapsto N_{\Lambda'}(\gamma) \in \mathbb{N}_0$, $\Lambda' \in \mathcal{B}_b(\mathbb{R}^d)$, $\Lambda' \subset \Lambda$. A measure $\mu \in \mathcal{M}_{\text{fm}}^1(\Gamma)$ is called locally absolutely continuous with respect to the Poisson measure π if for any $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ the projection of μ onto $\Gamma(\Lambda)$ is absolutely continuous with respect to (w.r.t.) the projection of π onto $\Gamma(\Lambda)$. More precisely, if we consider the projection mapping $p_\Lambda : \Gamma \rightarrow \Gamma(\Lambda)$, $p_\Lambda(\gamma) := \gamma_\Lambda$ then $\mu^\Lambda := \mu \circ p_\Lambda^{-1}$ is absolutely continuous w.r.t. $\pi_\Lambda := \pi \circ p_\Lambda^{-1}$.

Remark 2.5 Having in mind (2.13), it is possible to derive from (2.10) that the Gibbs measure from Example 2.4 is locally absolutely continuous w.r.t. the Poisson measure, see, e.g. [24] for the more general case.

By, e.g. [48], for any $\mu \in \mathcal{M}_{\text{fm}}^1(\Gamma)$ which is locally absolutely continuous w.r.t the Poisson measure there exists the family of (symmetric) correlation functions $k_\mu^{(n)} : (\mathbb{R}^d)^n \rightarrow \mathbb{R}_+ := [0, \infty)$ which is defined as follows. For any symmetric function $f^{(n)} : (\mathbb{R}^d)^n \rightarrow \mathbb{R}$ with a finite support the following equality holds

$$\begin{aligned} & \int_\Gamma \sum_{\{x_1, \dots, x_n\} \subset \gamma} f^{(n)}(x_1, \dots, x_n) d\mu(\gamma) \\ &= \frac{1}{n!} \int_{(\mathbb{R}^d)^n} f^{(n)}(x_1, \dots, x_n) k_\mu^{(n)}(x_1, \dots, x_n) dx_1 \dots dx_n \end{aligned} \tag{2.14}$$

for $n \in \mathbb{N}$, and $k_\mu^{(0)} := 1$.

The meaning of the notion of correlation functions is the following: the correlation function $k_\mu^{(n)}(x_1, \dots, x_n)$ describes the non-normalized density of probability to have points of our systems in the positions x_1, \dots, x_n .

Remark 2.6 Iterating the Mecke identity (2.8), it can be easily shown that

$$k_{\pi_\rho}^{(n)}(x_1, \dots, x_n) = \prod_{i=1}^n \rho(x_i), \tag{2.15}$$

in particular,

$$k_{\pi_z}^{(n)}(x_1, \dots, x_n) \equiv z^n. \tag{2.16}$$

Remark 2.7 Note that if potential ϕ from Example 2.4 satisfies to (2.9), (2.12), then, by [76], there exists $C = C(z, \phi) > 0$ such that for μ defined by (2.10)

$$k_\mu^{(n)}(x_1, \dots, x_n) \leq C^n, \quad x_1, \dots, x_n \in \mathbb{R}^d. \tag{2.17}$$

The inequality (2.17) is referred to as the Ruelle bound.

We dealt with symmetric function of n variables from \mathbb{R}^d ; hence, they can be considered as functions on n -point subsets from \mathbb{R}^d . We proceed now to the exact constructions.

The space of n -point configurations in $Y \in \mathcal{B}(\mathbb{R}^d)$ is defined by

$$\Gamma^{(n)}(Y) := \{\eta \subset Y \mid |\eta| = n\}, \quad n \in \mathbb{N}.$$

We put $\Gamma^{(0)}(Y) := \{\emptyset\}$. As a set, $\Gamma^{(n)}(Y)$ may be identified with the symmetrization of

$$\widetilde{Y}^n = \{(x_1, \dots, x_n) \in Y^n \mid x_k \neq x_l \text{ if } k \neq l\}.$$

Hence, one can introduce the corresponding Borel σ -algebra, which we denote by $\mathcal{B}(\Gamma^{(n)}(Y))$. The space of finite configurations in $Y \in \mathcal{B}(\mathbb{R}^d)$ is defined as

$$\Gamma_0(Y) := \bigsqcup_{n \in \mathbb{N}_0} \Gamma^{(n)}(Y). \tag{2.18}$$

This space is equipped with the topology of the disjoint union. Let $\mathcal{B}(\Gamma_0(Y))$ denote the corresponding Borel σ -algebra. In the case of $Y = \mathbb{R}^d$ we will omit the index Y in the previously defined notations, namely

$$\Gamma_0 := \Gamma_0(\mathbb{R}^d), \quad \Gamma^{(n)} := \Gamma^{(n)}(\mathbb{R}^d), \quad n \in \mathbb{N}_0. \tag{2.19}$$

The restriction of the Lebesgue product measure $(dx)^n$ to $(\Gamma^{(n)}, \mathcal{B}(\Gamma^{(n)}))$ we denote by $m^{(n)}$. We set $m^{(0)} := \delta_{\{\emptyset\}}$. The Lebesgue–Poisson measure λ on Γ_0 is defined by

$$\lambda := \sum_{n=0}^{\infty} \frac{1}{n!} m^{(n)}. \tag{2.20}$$

For any $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ the restriction of λ to $\Gamma_0(\Lambda) = \Gamma(\Lambda)$ will be also denoted by λ .

Remark 2.8 The space $(\Gamma, \mathcal{B}(\Gamma))$ is the projective limit of the family of measurable spaces $\{(\Gamma(\Lambda), \mathcal{B}(\Gamma(\Lambda)))\}_{\Lambda \in \mathcal{B}_b(\mathbb{R}^d)}$. The Poisson measure π on $(\Gamma, \mathcal{B}(\Gamma))$ from Example 2.3 may be defined as the projective limit of the family of measures $\{\pi^\Lambda\}_{\Lambda \in \mathcal{B}_b(\mathbb{R}^d)}$, where $\pi^\Lambda := e^{-m(\Lambda)}\lambda$ is the probability measure on $(\Gamma(\Lambda), \mathcal{B}(\Gamma(\Lambda)))$ and $m(\Lambda)$ is the Lebesgue measure of $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ (see, e.g. [1] for details).

Functions on Γ_0 will be called *quasi-observables*. Any $\mathcal{B}(\Gamma_0)$ -measurable function G on Γ_0 , in fact, is defined by a sequence of functions $\{G^{(n)}\}_{n \in \mathbb{N}_0}$ where $G^{(n)}$ is a $\mathcal{B}(\Gamma^{(n)})$ -measurable function on $\Gamma^{(n)}$. We preserve the same notation for the function $G^{(n)}$ considered as a symmetric function on $(\mathbb{R}^d)^n$. Note that $G^{(0)} \in \mathbb{R}$.

A set $M \in \mathcal{B}(\Gamma_0)$ is called bounded if there exists $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ and $N \in \mathbb{N}$ such that

$$M \subset \bigsqcup_{n=0}^N \Gamma^{(n)}(\Lambda).$$

The set of bounded measurable functions on Γ_0 with bounded support we denote by $B_{bs}(\Gamma_0)$, i.e. $G \in B_{bs}(\Gamma_0)$ iff $G \upharpoonright_{\Gamma_0 \setminus M} = 0$ for some bounded $M \in \mathcal{B}(\Gamma_0)$. For any $G \in B_{bs}(\Gamma_0)$ the functions $G^{(n)}$ have finite supports in $(\mathbb{R}^d)^n$ and may be substituted into (2.14). But, additionally, the sequence of $G^{(n)}$ vanishes for big n . Therefore, one can summarize equalities (2.14) by $n \in \mathbb{N}_0$. This leads to the following definition.

Let $G \in B_{bs}(\Gamma_0)$; then we define the function $KG : \Gamma \rightarrow \mathbb{R}$ such that

$$\begin{aligned} (KG)(\gamma) &:= \sum_{\eta \in \gamma} G(\eta) \\ &= G^{(0)} + \sum_{n=1}^{\infty} \sum_{\{x_1, \dots, x_n\} \subset \gamma} G^{(n)}(x_1, \dots, x_n), \quad \gamma \in \Gamma, \end{aligned} \tag{2.21}$$

see, e.g. [48, 59, 60]. The summation in (2.21) is taken over all finite subconfigurations $\eta \in \Gamma_0$ of the (infinite) configuration $\gamma \in \Gamma$; we denote this by the symbol, $\eta \Subset \gamma$. The mapping K is linear, positivity preserving, and invertible, with

$$(K^{-1}F)(\eta) := \sum_{\xi \subset \eta} (-1)^{|\eta \setminus \xi|} F(\xi), \quad \eta \in \Gamma_0. \tag{2.22}$$

By [48], for any $G \in B_{bs}(\Gamma_0)$, $KG \in \mathcal{F}_{cyl}(\Gamma)$, moreover, there exists $C = C(G) > 0$, $\Lambda = \Lambda(G) \in \mathcal{B}_b(\mathbb{R}^d)$, and $N = N(G) \in \mathbb{N}$ such that

$$|KG(\gamma)| \leq C(1 + |\gamma_\Lambda|)^N, \quad \gamma \in \Gamma. \tag{2.23}$$

The expression (2.21) can be extended to the class of all nonnegative measurable $G : \Gamma_0 \rightarrow \mathbb{R}_+$, in this case, evidently, $KG \in \mathcal{F}_0(\Gamma)$. Stress that the left-hand side (l.h.s.) of (2.22) has a meaning for any $F \in \mathcal{F}_0(\Gamma)$, moreover, in this case $(KK^{-1}F)(\gamma) = F(\gamma)$ for any $\gamma \in \Gamma_0$.

For G as above we may summarize (2.14) by n and rewrite the result in a compact form:

$$\int_{\Gamma} (KG)(\gamma) d\mu(\gamma) = \int_{\Gamma_0} G(\eta) k_{\mu}(\eta) d\lambda(\eta). \tag{2.24}$$

As was shown in [48], the equality (2.21) may be extended on all functions G such that the l.h.s. of (2.24) is finite. In this case (2.21) holds for μ -a.a. $\gamma \in \Gamma$ and (2.24) holds too.

Remark 2.9 The equality (2.24) may be considered as definition of the correlation function k_{μ} . In fact, the definition of correlation functions in statistical physics, given by Bogolyubov in [7], based on a similar relation. More precisely, consider for a $\mathcal{B}(\mathbb{R}^d)$ -measurable function f the so-called coherent state, given as a function on Γ_0 by

$$e_{\lambda}(f, \eta) := \prod_{x \in \eta} f(x), \quad \eta \in \Gamma_0 \setminus \{\emptyset\}, \quad e_{\lambda}(f, \emptyset) := 1. \tag{2.25}$$

Then for any $f \in C_0(\mathbb{R}^d)$ we have the point-wise equality

$$(Ke_{\lambda}(f))(\gamma) = \prod_{x \in \gamma} (1 + f(x)), \quad \eta \in \Gamma_0. \tag{2.26}$$

As a result, the correlation functions of different orders may be considered as kernels of a Taylor-type expansion

$$\begin{aligned} \int_{\Gamma} \prod_{x \in \gamma} (1 + f(x)) d\mu(\gamma) &= 1 + \sum_{n=1}^{\infty} \frac{1}{n!} \int_{(\mathbb{R}^d)^n} \prod_{i=1}^n f(x_i) k_{\mu}^{(n)}(x_1, \dots, x_n) dx_1 \dots dx_n \\ &= \int_{\Gamma_0} e_{\lambda}(f, \eta) k_{\mu}(\eta) d\lambda(\eta). \end{aligned} \tag{2.27}$$

Remark 2.10 By (2.18)–(2.20), we have that for any $f \in L^1(\mathbb{R}^d, dx)$

$$\int_{\Gamma_0} e_{\lambda}(f, \eta) d\lambda(\eta) = \exp\left\{ \int_{\mathbb{R}^d} f(x) dx \right\}. \tag{2.28}$$

As a result, taking into account (2.15), we obtain from (2.27) the expression for the Laplace transform of the Poisson measure

$$\begin{aligned} \int_{\Gamma} e^{-\langle \varphi, \gamma \rangle} d\pi_{\rho}(\gamma) &= \int_{\Gamma_0} e_{\lambda}(e^{-\varphi(x)} - 1, \eta) e_{\lambda}(\rho, \eta) d\lambda(\eta) \\ &= \exp\left\{ - \int_{\mathbb{R}^d} (1 - e^{-\varphi(x)}) \rho(x) dx \right\}, \quad \varphi \in C_0(\mathbb{R}^d). \end{aligned}$$

Remark 2.11 Of course, to obtain convergence of the expansion (2.27) for, say, $f \in L^1(\mathbb{R}^d, dx)$ we need some bounds for the correlation functions $k_{\mu}^{(n)}$. For example, if the generalized Ruelle bound holds, that is, cf. (2.17),

$$k_{\mu}^{(n)}(x_1, \dots, x_n) \leq AC^n (n!)^{1-\delta}, \quad x_1, \dots, x_n \in \mathbb{R}^d \tag{2.29}$$

for some $A, C > 0, \delta \in (0, 1]$ independent on n , then the l.h.s. of (2.27) may be estimated by the expression

$$1 + A \sum_{n=1}^{\infty} \frac{(C \|f\|_{L^1(\mathbb{R}^d)})^n}{(n!)^{\delta}} < \infty.$$

For a given system of functions $k^{(n)}$ on $(\mathbb{R}^d)^n$ the question about existence and uniqueness of a probability measure μ on Γ which has correlation functions $k_{\mu}^{(n)} = k^{(n)}$ is an analog of the moment problem in classical analysis. Significant results in this area were obtained by Lenard.

Proposition 2.12 ([58,60]) *Let $k : \Gamma_0 \rightarrow \mathbb{R}$.*

1. Suppose that k is a positive definite function that means that for any $G \in B_{bs}(\Gamma_0)$ such that $(KG)(\gamma) \geq 0$ for all $\gamma \in \Gamma$ the following inequality holds:

$$\int_{\Gamma_0} G(\eta)k(\eta) d\lambda(\eta) \geq 0. \tag{2.30}$$

Suppose also that $k(\emptyset) = 1$. Then there exists at least one measure $\mu \in \mathcal{M}_{fm}^1(\Gamma)$ such that $k = k_\mu$.

2. For any $n \in \mathbb{N}$, $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$, we set

$$s_n^\Lambda := \frac{1}{n!} \int_{\Lambda^n} k^{(n)}(x_1, \dots, x_n) dx_1 \cdots dx_n.$$

Suppose that for all $m \in \mathbb{N}$, $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$

$$\sum_{n \in \mathbb{N}} (s_{n+m}^\Lambda)^{-\frac{1}{n}} = \infty. \tag{2.31}$$

Then there exists at most one measure $\mu \in \mathcal{M}_{fm}^1(\Gamma)$ such that $k = k_\mu$.

- Remark 2.13*
1. In [58,60], the wider space of multiple configurations was considered. The adaptation for the space Γ was realized in [57].
 2. It is worth noting also that the growth of correlation functions $k^{(n)}$ up to $(n!)^2$ is admissible to have (2.31).
 3. Other conditions for existence and uniqueness for the moment problem on Γ were studied in [4,48].

2.3 Statistical descriptions of Markov evolutions

Spatial Markov processes in \mathbb{R}^d may be described as stochastic evolutions of configurations $\gamma \subset \mathbb{R}^d$. In course of such evolutions points of configurations may disappear (die), move (continuously or with jumps from one position to another), or new particles may appear in a configuration (that is birth). The rates of these random events may depend on whole configuration that reflect an interaction between elements of the our system.

The construction of a spatial Markov process in the continuum is highly difficult question which is not solved in a full generality at present, see, e.g. a review [71] and more detail references about birth-and-death processes in Sect. 3. Meanwhile, for the discrete systems the corresponding processes are constructed under quite general assumptions, see, e.g. [62]. One of the main difficulties for continuous systems includes the necessity to control number of elements in a bounded region. Note that the construction of spatial processes on bounded sets from \mathbb{R}^d are typically well solved, see, e.g. [41].

The existing Markov process $\Gamma \ni \gamma \mapsto X_t^\gamma \in \Gamma$, $t > 0$ provides solution to the backward Kolmogorov equation for bounded continuous functions:

$$\frac{\partial}{\partial t} F_t = L F_t, \tag{2.32}$$

where L is the Markov generator of the process X_t . The question about existence and properties of solutions to (2.32) in proper spaces itself is also highly nontrivial problem of infinite-dimensional analysis. The Markov generator L should satisfy the following two (informal) properties: (1) to be conservative, that is $L1 = 0$, (2) maximum principle, namely if there exists $\gamma_0 \in \Gamma$ such that $F(\gamma) \leq F(\gamma_0)$ for all $\gamma \in \Gamma$, then $(LF)(\gamma_0) \leq 0$. These properties might yield that the semigroup, related to (2.32) (provided it exists), will preserve constants and positive functions, correspondingly.

To consider an example of such L let us consider a general Markov evolution with appearing and disappearing of groups of points (giving up the case of continuous moving of particles). Namely, let $F \in \mathcal{F}_{cyl}(\Gamma)$ and set

$$(LF)(\gamma) = \sum_{\eta \in \gamma} \int_{\Gamma_0} c(\eta, \xi, \gamma \setminus \eta) [F((\gamma \setminus \eta) \cup \xi) - F(\gamma)] d\lambda(\xi). \tag{2.33}$$

Heuristically, it means that any finite group η of points from the existing configuration γ may disappear and simultaneously a new group ξ of points may appear somewhere in the space \mathbb{R}^d . The rate of this random event is equal to $c(\eta, \xi, \gamma \setminus \eta) \geq 0$. We need some minimal conditions on the rate c to guarantee that at least

$$LF \in \mathcal{F}_0(\Gamma) \quad \text{for all } F \in \mathcal{F}_{\text{cyl}}(\Gamma) \tag{2.34}$$

(see Sect. 3 for a particular case). The term in the sum in (2.33) with $\eta = \emptyset$ corresponds to a pure birth of a finite group ξ of points whereas the part of integral corresponding to $\xi = \emptyset$ (recall that $\lambda(\{\emptyset\}) = 1$) is related to pure death of a finite sub-configuration $\eta \subset \gamma$. The parts with $|\eta| = |\xi| \neq 0$ corresponds to jumps of one group of points into another positions in \mathbb{R}^d . The rest parts present splitting and merging effects. In the present paper the technical realization of the ideas below is given for one-point birth-and-death parts only, i.e. for the cases $|\eta| = 0, |\xi| = 1$ and $|\eta| = 1, |\xi| = 0$, correspondingly.

As we noted before, for most cases appearing in applications, the existence problem for a corresponding Markov process with a generator L is still open. On the other hand, the evolution of a state in the course of a stochastic dynamics is an important question in its own right. A mathematical formulation of this question may be realized through the forward Kolmogorov equation for probability measures (states) on the configuration space Γ , namely we consider the pairing between functions and measures on Γ given by

$$\langle F, \mu \rangle := \int_{\Gamma} F(\gamma) d\mu(\gamma). \tag{2.35}$$

Then we consider the initial value problem

$$\frac{d}{dt} \langle F, \mu_t \rangle = \langle LF, \mu_t \rangle, \quad t > 0, \quad \mu_t|_{t=0} = \mu_0, \tag{2.36}$$

where F is an arbitrary function from a proper set, e.g. $F \in K(B_{\text{bs}}(\Gamma_0)) \subset \mathcal{F}_{\text{cyl}}(\Gamma)$. In fact, the solution to (2.36) describes the time evolution of distributions instead of the evolution of initial points in the Markov process. We rewrite (2.36) in the following heuristic form:

$$\frac{\partial}{\partial t} \mu_t = L^* \mu_t, \tag{2.37}$$

where L^* is the (informally) adjoint operator of L with respect to the pairing (2.35).

In the physical literature, (2.37) is referred to the *Fokker–Planck equation*. The Markovian property of L yields that (2.37) might have a solution in the class of probability measures. However, the mere existence of the corresponding Markov process will not give us much information about properties of the solution to (2.37), in particular, about its moments or correlation functions. To do this, we suppose now that a solution $\mu_t \in \mathcal{M}_{\text{fm}}^1(\Gamma)$ to (2.36) exists and remains locally absolutely continuous with respect to the Poisson measure π for all $t > 0$ provided μ_0 has such a property. Then one can consider the correlation function $k_t := k_{\mu_t}$, $t \geq 0$.

Recall that we suppose (2.34). Then, one can calculate $K^{-1}LF$ using (2.22), and, by (2.24), we may rewrite (2.36) in the following way:

$$\frac{d}{dt} \langle K^{-1}F, k_t \rangle = \langle K^{-1}LF, k_t \rangle, \quad t > 0, \quad k_t|_{t=0} = k_0, \tag{2.38}$$

for all $F \in K(B_{\text{bs}}(\Gamma_0)) \subset \mathcal{F}_{\text{cyl}}(\Gamma)$. Here the pairing between functions on Γ_0 is given by

$$\langle\langle G, k \rangle\rangle := \int_{\Gamma_0} G(\eta)k(\eta) d\lambda(\eta). \tag{2.39}$$

Let us recall that then, by (2.20),

$$\langle\langle G, k \rangle\rangle = \sum_{n=0}^{\infty} \frac{1}{n!} \int_{(\mathbb{R}^d)^n} G^{(n)}(x_1, \dots, x_n)k^{(n)}(x_1, \dots, x_n) dx_1 \dots dx_n,$$

Next, if we substitute $F = KG$, $G \in B_{\text{bs}}(\Gamma_0)$ in (2.38), we derive

$$\frac{d}{dt} \langle\langle G, k_t \rangle\rangle = \langle\langle \widehat{L}G, k_t \rangle\rangle, \quad t > 0, \quad k_t|_{t=0} = k_0, \tag{2.40}$$

for all $G \in B_{bs}(\Gamma_0)$. Here the operator

$$(\widehat{L}G)(\eta) := (K^{-1}LKG)(\eta), \quad \eta \in \Gamma_0$$

is defined point-wise for all $G \in B_{bs}(\Gamma_0)$ under conditions (2.34). As a result, we are interested in a weak solution to the equation

$$\frac{\partial}{\partial t}k_t = \widehat{L}^*k_t, \quad t > 0, \quad k_t|_{t=0} = k_0, \tag{2.41}$$

where \widehat{L}^* is dual operator to \widehat{L} with respect to the duality (2.39), namely

$$\int_{\Gamma_0} (\widehat{L}G)(\eta)k(\eta) d\lambda(\eta) = \int_{\Gamma_0} G(\eta)(\widehat{L}^*k)(\eta) d\lambda(\eta). \tag{2.42}$$

The procedure of deriving the operator \widehat{L} for a given L is fully combinatorial; meanwhile, to obtain the expression for the operator \widehat{L}^* we need an analog of integration by parts formula. For a difference operator L considered in (2.33) this discrete integration by parts rule is presented in Lemma 3.4 below.

We recall that any function on Γ_0 may be identified with an infinite vector of symmetric functions of the growing number of variables. In this approach, the operator \widehat{L}^* in (2.41) will be realized as an infinite matrix $(\widehat{L}_{n,m}^*)_{n,m \in \mathbb{N}_0}$, where $\widehat{L}_{n,m}^*$ is a mapping from the space of symmetric functions of n variables into the space of symmetric functions of m variables. As a result, instead of Eq. (2.36) for infinite-dimensional objects we obtain an infinite system of equations for functions $k_t^{(n)}$; each of them is a function of a finite number of variables, namely

$$\begin{aligned} \frac{\partial}{\partial t}k_t^{(n)}(x_1, \dots, x_n) &= (\widehat{L}_{n,m}^*k_t^{(n)})(x_1, \dots, x_n), \quad t > 0, \quad n \in \mathbb{N}_0, \\ k_t^{(n)}(x_1, \dots, x_n)|_{t=0} &= k_0^{(n)}(x_1, \dots, x_n). \end{aligned} \tag{2.43}$$

Of course, in general, for a fixed n , any equation from (2.43) itself is not closed and includes functions $k_t^{(m)}$ of other orders $m \neq n$; nevertheless, the system (2.43) is a closed linear system. The chain evolution equations for $k_t^{(n)}$ consists the so-called *hierarchy* which is an analog of the BBGKY hierarchy for Hamiltonian systems, see, e.g. [18].

One of the main aims of the present paper was to study the classical solution to (2.41) in a proper functional space. The choice of such a space might be based on estimates (2.17), or more generally, (2.29). However, even the correlation functions (2.16) of the Poisson measures show that it is rather natural to study the solutions to the Eq. (2.41) in weighted L^∞ -type space of functions with the Ruelle-type bounds. Integrable correlation functions are not natural for the dynamics on the spaces of locally finite configurations. For example, it is well known that the Poisson measure π_ρ with integrable density $\rho(x)$ is concentrated on the space Γ_0 of finite configurations [since in this case one can consider \mathbb{R}^d instead of Λ in (2.6)]. Therefore, typically, the case of integrable correlation functions yields that effectively our stochastic dynamics evolves through finite configurations only. Note that the case of an integrable first-order correlation function is referred to *zero density* case in statistical physics.

In the present paper we restrict our attention to the so-called *sub-Poissonian* correlation functions. Namely, for a given $C > 0$ we consider the following Banach space:

$$\mathcal{K}_C := \{k : \Gamma_0 \rightarrow \mathbb{R} \mid k \cdot C^{-|\cdot|} \in L^\infty(\Gamma_0, d\lambda)\} \tag{2.44}$$

with the norm

$$\|k\|_{\mathcal{K}_C} := \|C^{-|\cdot|}k(\cdot)\|_{L^\infty(\Gamma_0, \lambda)}.$$

It is clear that $k \in \mathcal{K}_C$ implies, cf. (2.17),

$$|k(\eta)| \leq \|k\|_{\mathcal{K}_C} C^{|\eta|} \quad \text{for } \lambda\text{-a.a. } \eta \in \Gamma_0. \tag{2.45}$$

In the following, we distinguish two possibilities for a study of the initial value problem (2.41). We may try to solve this equation in one space \mathcal{K}_C . The well-posedness of the initial value problem in this case is

equivalent with an existence of the strongly continuous semigroup (C_0 -semigroup in the sequel) in the space \mathcal{K}_C with a generator \widehat{L}^* . However, the space \mathcal{K}_C is isometrically isomorphic to the space $L^\infty(\Gamma_0, C^{|\cdot|}d\lambda)$, whereas by the Lotz theorem [3,64], in a L^∞ space any C_0 -semigroup is uniformly continuous, that is it has a bounded generator. Typically, for the difference operator L given in (2.33), any operator $\widehat{L}_{n,m}^*$, cf. (2.43), might be bounded as an operator between two spaces of bounded symmetric functions of n and m variables, whereas the whole operator \widehat{L}^* is unbounded in \mathcal{K}_C .

To avoid these difficulties we use a trick which goes back to Phillips [72]. The main idea is to consider the semigroup in L^∞ space not itself but as a dual semigroup $T^*(t)$ to a C_0 -semigroup $T(t)$ with a generator A in the pre-dual L^1 space. In this case $T^*(t)$ appears strongly continuous semigroup not on the whole L^∞ but on the closure of the domain of A^* only.

In our case this leads to the following scheme. We consider the pre-dual Banach space to \mathcal{K}_C , namely for $C > 0$,

$$\mathcal{L}_C := L^1(\Gamma_0, C^{|\cdot|}d\lambda). \tag{2.46}$$

The norm in \mathcal{L}_C is given by

$$\|G\|_C := \int_{\Gamma_0} |G(\eta)| C^{|\eta|} d\lambda(\eta) = \sum_{n=0}^{\infty} \frac{C^n}{n!} \int_{(\mathbb{R}^d)^n} |G^{(n)}(x_1, \dots, x_n)| dx_1 \dots dx_n.$$

Consider the initial value problem, cf. (2.40), (2.41),

$$\frac{\partial}{\partial t} G_t = \widehat{L}G_t, \quad t > 0, \quad G_t|_{t=0} = G_0 \in \mathcal{L}_C. \tag{2.47}$$

Let (2.47) be well-posed in \mathcal{L}_C ; then there exists a C_0 -semigroup $\widehat{T}(t)$ in \mathcal{L}_C . Then using Philips' result we obtain that the restriction of the dual semigroup $\widehat{T}^*(t)$ onto $\text{Dom}(\widehat{L}^*)$ will be C_0 -semigroup with generator which is a part of \widehat{L}^* (the details see in Sect. 3 below). This provides a solution to (2.41) which continuously depends on an initial data from $\text{Dom}(\widehat{L}^*)$. And after we would like to find a more useful universal subspace of \mathcal{K}_C which is not dependent on the operator \widehat{L}^* . The realization of this scheme for a birth-and-death operator L is presented in Sect. 3 below. As a result, we obtain the classical solution to (2.41) for $t > 0$ in a class of sub-Poissonian functions which satisfy the Ruelle-type bound (2.45). Of course, after this we need to verify existence and uniqueness of measures whose correlation functions are solutions to (2.41), cf. Proposition 2.12 above. This usually can be done using proper approximation schemes, see, e.g. Sect. 4.

There is another possibility for a study of the initial value problem (2.41) which we will not touch below, namely one can consider this evolutionary equation in a proper scale of spaces $\{\mathcal{K}_C\}_{C_* \leq C \leq C^*}$. In this case we will have typically that the solution is local in time only. More precisely, there exists $T > 0$ such that for any $t \in [0, T)$ there exists a unique solution to (2.41) and $k_t \in \mathcal{K}_{C_t}$ for some $C_t \in [C_*, C^*]$. We realized this approach in series of papers [5,25,37,38] using the so-called Ovsyannikov method [69,77,78]. This method provides less restrictions on systems parameters; however, the price for this is a finite time interval. And, of course, the question about possibility to recover measures via solutions to (2.41) should be also solved separately in this case.

3 Birth-and-death evolutions in the continuum

3.1 Microscopic description

One of the most important classes of Markov evolution in the continuum is given by the birth-and-death Markov processes in the space Γ of all configurations from \mathbb{R}^d . These are processes in which an infinite number of individuals exist at each instant, and the rates at which new individuals appear and some old ones disappear depend on the instantaneous configuration of existing individuals [46]. The corresponding Markov generators have a natural heuristic representation in terms of birth and death intensities. The birth intensity $b(x, \gamma) \geq 0$ characterizes the appearance of a new point at $x \in \mathbb{R}^d$ in the presence of a given configuration $\gamma \in \Gamma$. The death intensity $d(x, \gamma) \geq 0$ characterizes the probability of the event that the point x of the configuration

γ disappears, depending on the location of the remaining points of the configuration, $\gamma \setminus x$. Heuristically, the corresponding Markov generator is described by the following expression, cf. (2.33):

$$(LF)(\gamma) := \sum_{x \in \gamma} d(x, \gamma \setminus x) [F(\gamma \setminus x) - F(\gamma)] + \int_{\mathbb{R}^d} b(x, \gamma) [F(\gamma \cup x) - F(\gamma)] dx, \tag{3.1}$$

for proper functions $F : \Gamma \rightarrow \mathbb{R}$.

The study of spatial birth-and-death processes was initiated by Preston [73]. This paper dealt with a solution of the backward Kolmogorov equation (2.32) under the restriction that only a finite number of individuals are alive at each moment of time. Under certain conditions, corresponding processes exist and are temporally ergodic, that is, there exists a unique stationary distribution. Note that a more general setting for birth-and-death processes only requires that the number of points in any compact set remains finite at all times. A further progress in the study of these processes was achieved by Holley and Stroock in [46]. They described in detail an analytic framework for birth-and-death dynamics. In particular, they analyzed the case of a birth-and-death process in a bounded region.

Stochastic equations for spatial birth-and-death processes were formulated in [42], through a spatial version of the time-change approach. Further, in [43], these processes were represented as solutions to a system of stochastic equations, and conditions for the existence and uniqueness of solutions to these equations, as well as for the corresponding martingale problems, were given. Unfortunately, quite restrictive assumptions on the birth and death rates in [43] do not allow an application of these results to several particular models that are interesting for applications (see, e.g. some of examples below).

A growing interest to the study of spatial birth-and-death processes, which we have recently observed, is stimulated by (among others) an important role which these processes play in several applications. For example, in spatial plant ecology, a general approach to the so-called individual based models was developed in a series of works, see, e.g. [8, 9, 17, 66] and the references therein. These models are described as birth-and-death Markov processes in the configuration space Γ with specific rates b and d which reflect biological notions such as competition, establishment, fecundity, etc. Other examples of birth-and-death processes may be found in mathematical physics. In particular, the Glauber-type stochastic dynamics in Γ is properly associated with the grand canonical Gibbs measures for classical gases. This gives a possibility to study these Gibbs measures as equilibrium states for specific birth-and-death Markov evolutions [6]. Starting with a Dirichlet form for a given Gibbs measure, one can consider an equilibrium stochastic dynamics [54]. However, these dynamics give the time evolution of initial distributions from a quite narrow class, namely the class of admissible initial distributions is essentially reduced to the states which are absolutely continuous with respect to the invariant measure. Below we construct non-equilibrium stochastic dynamics which may have a much wider class of initial states.

This approach was successfully applied to the construction and analysis of state evolutions for different versions of the Glauber dynamics [28, 34, 53] and for some spatial ecology models [26]. Each of the considered models required its own specific version of the construction of a semigroup, which takes into account particular properties of corresponding birth and death rates.

In this section, we realize a general approach considered in Sect. 2 to the construction of the state evolution corresponding to the birth-and-death Markov generators. We present conditions on the birth-and-death intensities which are sufficient for the existence of corresponding evolutions as strongly continuous semigroups in proper Banach spaces of correlation functions satisfying the Ruelle-type bounds. Also we consider weaker assumptions on these intensities which provide the corresponding evolutions for finite time intervals in scales of Banach spaces as above.

3.2 Expressions for \widehat{L} and \widehat{L}^* . Examples of rates b and d

We always suppose that rates $d, b : \mathbb{R}^d \times \Gamma \rightarrow [0; +\infty]$ from (3.1) satisfy the following assumptions:

$$d(x, \eta), b(x, \eta) > 0, \quad \eta \in \Gamma_0 \setminus \{\emptyset\}, x \in \mathbb{R}^d \setminus \eta, \tag{3.2}$$

$$d(x, \eta), b(x, \eta) < \infty, \quad \eta \in \Gamma_0, x \in \mathbb{R}^d \setminus \eta, \tag{3.3}$$

$$\int_M (d(x, \eta) + b(x, \eta)) d\lambda(\eta) < \infty, \quad M \in \mathcal{B}(\Gamma_0) \text{ bounded, a.a. } x \in \mathbb{R}^d, \quad (3.4)$$

$$\int_\Lambda (d(x, \eta) + b(x, \eta)) dx < \infty, \quad \eta \in \Gamma_0, \Lambda \in \mathcal{B}_b(\mathbb{R}^d). \quad (3.5)$$

Proposition 3.1 *Let conditions (3.2)–(3.5) hold. The for any $G \in B_{bs}(\Gamma_0)$ and $F = KG$ one has $LF \in \mathcal{F}_0(\Gamma)$.*

Proof By (2.23), there exist $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$, $N \in \mathbb{N}$, $C > 0$ (dependent on G) such that

$$\begin{aligned} |F(\gamma \setminus x) - F(\gamma)| &\leq C \mathbb{1}_\Lambda(x) (1 + |\gamma_\Lambda|)^N, & x \in \gamma, \gamma \in \Gamma, \\ |F(\gamma \cup x) - F(\gamma)| &\leq C \mathbb{1}_\Lambda(x) (2 + |\gamma_\Lambda|)^N, & \gamma \in \Gamma, x \in \mathbb{R}^d \setminus \gamma. \end{aligned}$$

Then, by (3.3), (3.5), for any $\eta \in \Gamma_0$,

$$|(LF)(\eta)| \leq C (2 + |\eta_\Lambda|)^N \left(\sum_{x \in \eta_\Lambda} d(x, \eta \setminus x) + \int_\Lambda b(x, \eta) dx \right) < \infty.$$

The statement is proved. □

We start from the deriving of the expression for $\widehat{L} = K^{-1}LK$.

Proposition 3.2 *For any $G \in B_{bs}(\Gamma_0)$ the following formula holds:*

$$\begin{aligned} (\widehat{L}G)(\eta) &= - \sum_{\xi \subset \eta} G(\xi) \sum_{x \in \xi} (K^{-1}d(x, \cdot \cup \xi \setminus x))(\eta \setminus \xi) \\ &\quad + \sum_{\xi \subset \eta} \int_{\mathbb{R}^d} G(\xi \cup x) (K^{-1}b(x, \cdot \cup \xi))(\eta \setminus \xi) dx, \quad \eta \in \Gamma_0. \end{aligned} \quad (3.6)$$

Proof First of all, note that, by (3.3) and (2.22), the expressions $(K^{-1}b(x, \cdot \cup \xi))(\eta)$ and $(K^{-1}d(x, \cdot \cup \xi))(\eta)$ have sense. Recall that $G \in B_{bs}(\Gamma_0)$ implies $F \in \mathcal{F}_{cyl}(\Gamma) \subset \mathcal{F}_0(\Gamma)$; then, by (2.21),

$$\begin{aligned} F(\gamma \setminus x) - F(\gamma) &= \sum_{\eta \in \gamma \setminus x} G(\eta) - \sum_{\eta \in \gamma} G(\eta) \\ &= - \sum_{\eta \in \gamma \setminus x} G(\eta \cup x) = -(K(G(\cdot \cup x)))(\gamma \setminus x). \end{aligned} \quad (3.7)$$

In the same way, for $x \notin \gamma$, we derive

$$F(\gamma \cup x) - F(\gamma) = (K(G(\cdot \cup x)))(\gamma). \quad (3.8)$$

By Proposition 3.1, the values of $(\widehat{L}G)(\eta)$ are finite, and, by (2.22), one can interchange order of summations and integration in the following computations, that takes into account (3.7), (3.8):

$$\begin{aligned} (\widehat{L}G)(\eta) &= - \sum_{\zeta \subset \eta} (-1)^{|\eta \setminus \zeta|} \sum_{x \in \zeta} d(x, \zeta \setminus x) \sum_{\xi \subset \zeta \setminus x} G(\xi \cup x) \\ &\quad + \int_{\mathbb{R}^d} \sum_{\zeta \subset \eta} (-1)^{|\eta \setminus \zeta|} b(x, \zeta) \sum_{\xi \subset \zeta} G(\xi \cup x) dx, \end{aligned}$$

and making substitution $\xi' = \xi \cup x \subset \zeta$, one may continue

$$\begin{aligned} &= - \sum_{\zeta \subset \eta} (-1)^{|\eta \setminus \zeta|} \sum_{\xi' \subset \zeta} \sum_{x \in \xi'} d(x, \zeta \setminus x) G(\xi') \\ &\quad + \int_{\mathbb{R}^d} \sum_{\zeta \subset \eta} (-1)^{|\eta \setminus \zeta|} b(x, \zeta) \sum_{\xi \subset \zeta} G(\xi \cup x) dx. \end{aligned}$$

Next, for any measurable $H : \Gamma_0 \times \Gamma_0 \rightarrow \mathbb{R}$, one has

$$\sum_{\zeta \subset \eta} \sum_{\xi \subset \zeta} H(\xi, \zeta) = \sum_{\xi \subset \eta} \sum_{\substack{\zeta \subset \eta \\ \xi \subset \zeta}} H(\xi, \zeta) = \sum_{\xi \subset \eta} \sum_{\zeta' \subset \eta \setminus \xi} H(\xi, \zeta' \cup \xi).$$

Using this changing of variables rule, we continue:

$$\begin{aligned} (\widehat{L}G)(\eta) &= - \sum_{\xi \subset \eta} \sum_{\zeta' \subset \eta \setminus \xi} (-1)^{|\eta \setminus (\xi \cup \zeta')|} \sum_{x \in \xi} d(x, \zeta' \cup \xi \setminus x) G(\xi) \\ &\quad + \int_{\mathbb{R}^d} \sum_{\xi \subset \eta} \sum_{\zeta' \subset \eta \setminus \xi} (-1)^{|\eta \setminus (\zeta' \cup \xi)|} b(x, \zeta' \cup \xi) G(\xi \cup x) dx, \end{aligned}$$

that yields (3.6), using the equality $|\eta \setminus (\xi \cup \zeta')| = |(\eta \setminus \xi) \setminus \zeta'|$ and (2.22). □

Remark 3.3 The initial value problem (2.47) can be considered in the following matrix form, cf. (2.43),

$$\begin{aligned} \frac{\partial}{\partial t} G_t^{(n)}(x_1, \dots, x_n) &= (\widehat{L}_{n,m} G_t^{(n)})(x_1, \dots, x_n), \quad t > 0, \quad n \in \mathbb{N}_0, \\ G_t^{(n)}(x_1, \dots, x_n)|_{t=0} &= G_0^{(n)}(x_1, \dots, x_n). \end{aligned}$$

The expression (3.6) shows that the matrix above has on the main diagonal the collection of operators $\widehat{L}_{n,n}$, $n \in \mathbb{N}_0$ which forms the following operator on functions on Γ_0 :

$$(\widehat{L}_{\text{diag}}G)(\eta) = -D(\eta)G(\eta) + \sum_{y \in \eta} \int_{\mathbb{R}^d} G((\eta \setminus y) \cup x) [b(x, \eta) - b(x, \eta \setminus y)] dx, \tag{3.9}$$

where the term in the square brackets is equal, by (2.22), to $(K^{-1}b(x, \cdot \cup (\eta \setminus y)))(\{y\})$. Next, by (3.6), there exists only one non-zero upper diagonal in the matrix. The corresponding operator is

$$(\widehat{L}_{\text{upper}}G)(\eta) = \int_{\mathbb{R}^d} G(\eta \cup x) b(x, \eta) dx, \tag{3.10}$$

since $(K^{-1}b(x, \cdot \cup \eta))(\emptyset) = b(x, \eta)$. The rest part of the expression (3.6) corresponds to the low diagonals.

As we mentioned above, to derive the expression for \widehat{L}^* we need some discrete analog of the integration by parts formula. As such, we will use the partial case of the well-known lemma (see, e.g. [56]):

Lemma 3.4 For any measurable function $H : \Gamma_0 \times \Gamma_0 \times \Gamma_0 \rightarrow \mathbb{R}$

$$\int_{\Gamma_0} \sum_{\xi \subset \eta} H(\xi, \eta \setminus \xi, \eta) d\lambda(\eta) = \int_{\Gamma_0} \int_{\Gamma_0} H(\xi, \eta, \eta \cup \xi) d\lambda(\xi) d\lambda(\eta) \tag{3.11}$$

if at least one side of the equality is finite for $|H|$.

In particular, if $H(\xi, \cdot, \cdot) \equiv 0$ if only $|\xi| \neq 1$ we obtain an analog of (2.8), namely

$$\int_{\Gamma_0} \sum_{x \in \eta} h(x, \eta \setminus x, \eta) d\lambda(\eta) = \int_{\Gamma_0} \int_{\mathbb{R}^d} h(x, \eta, \eta \cup x) dx d\lambda(\eta), \tag{3.12}$$

for any measurable function $h : \mathbb{R}^d \times \Gamma_0 \times \Gamma_0 \rightarrow \mathbb{R}$ such that both sides make sense.

Using this, one can derive the explicit form of \widehat{L}^* .

Proposition 3.5 For any $k \in B_{bs}(\Gamma_0)$ the following formula holds:

$$\begin{aligned} (\widehat{L}^*k)(\eta) &= - \sum_{x \in \eta} \int_{\Gamma_0} k(\zeta \cup \eta)(K^{-1}d(x, \cdot \cup \eta \setminus x))(\zeta) d\lambda(\zeta) \\ &\quad + \sum_{x \in \eta} \int_{\Gamma_0} k(\zeta \cup (\eta \setminus x))(K^{-1}b(x, \cdot \cup \eta \setminus x))(\zeta) d\lambda(\zeta), \end{aligned} \tag{3.13}$$

where \widehat{L}^*k is defined by (2.42).

Proof Using Lemma 3.4, (2.42), (3.6), we obtain for any $G \in B_{bs}(\Gamma_0)$

$$\begin{aligned} &\int_{\Gamma_0} G(\eta)(\widehat{L}^*k)(\eta) d\lambda(\eta) \\ &= - \int_{\Gamma_0} \sum_{\xi \subset \eta} G(\xi) \sum_{x \in \xi} (K^{-1}d(x, \cdot \cup \xi \setminus x))(\eta \setminus \xi) k(\eta) d\lambda(\eta) \\ &\quad + \int_{\Gamma_0} \sum_{\xi \subset \eta} \int_{\mathbb{R}^d} G(\xi \cup x) (K^{-1}b(x, \cdot \cup \xi))(\eta \setminus \xi) dx k(\eta) d\lambda(\eta) \\ &= - \int_{\Gamma_0} \int_{\Gamma_0} G(\xi) \sum_{x \in \xi} (K^{-1}d(x, \cdot \cup \xi \setminus x))(\eta) k(\eta \cup \xi) d\lambda(\eta) d\lambda(\xi) \\ &\quad + \int_{\Gamma_0} \int_{\Gamma_0} \int_{\mathbb{R}^d} G(\xi \cup x) (K^{-1}b(x, \cdot \cup \xi))(\eta) dx k(\eta \cup \xi) d\lambda(\eta) d\lambda(\xi). \end{aligned}$$

Applying (3.12) for the second term, we easily obtain the statement. The correctness of using (2.8) and (3.12) follows from the assumptions that $G, k \in B_{bs}(\Gamma_0)$; therefore, all integrals over Γ_0 will be taken, in fact, over some bounded $M \in \mathcal{B}(\Gamma_0)$. Then, using (3.4), (3.5), we obtain that the all integrals are finite. \square

Remark 3.6 Accordingly to Remark 3.3 [or just directly from (3.13)], we have that the matrix corresponding to (2.43) has the main diagonal given by

$$\begin{aligned} (\widehat{L}_{diag}^*k)(\eta) &= -D(\eta)k(\eta) \\ &\quad + \sum_{x \in \eta} \int_{\mathbb{R}^d} k((\eta \setminus x) \cup y) [b(x, (\eta \setminus x) \cup y) - b(x, \eta \setminus x)] dy, \end{aligned} \tag{3.14}$$

where we have used (3.12). Next, this matrix has only one non-zero low diagonal, given by the expression

$$(\widehat{L}_{low}^*k)(\eta) = \sum_{x \in \eta} k(\eta \setminus x) b(x, \eta \setminus x). \tag{3.15}$$

The rest part of expression (3.13) corresponds to the upper diagonals.

Let us consider now several examples of rates b and d which will appear in the following considerations (concrete examples of birth-and-death dynamics, with such rates, important for applications will be presented later). As we see from (3.6), (3.13), we always need to calculate expressions like $(K^{-1}a(x, \cdot \cup \xi))(\eta), \eta \cap \xi = \emptyset$, where a equal to b or d . We consider the following kinds of function $a : \mathbb{R}^d \times \Gamma \rightarrow \mathbb{R}$:

– *Constant rate:*

$$a(x, \gamma) \equiv m > 0. \tag{3.16}$$

If we substitute $f \equiv 0$ into (2.26), we obtain that

$$(K^{-1}m)(\eta) = m0^{|\eta|}, \quad \eta \in \Gamma_0, \tag{3.17}$$

where as usual $0^0 := 1$, and, of course, in this case $K^{-1}a(x, \cdot \cup \xi)(\eta)$ also equal to $m0^{|\eta|}$ for any $\xi \in \Gamma_0$;

– *Linear rate:*

$$a(x, \gamma) = \langle c(x - \cdot), \gamma \rangle = \sum_{y \in \gamma} c(x - y), \tag{3.18}$$

where c is a potential like in Example 2.2. Any such c for a given $x \in \mathbb{R}^d$ defines a function $C_x : \Gamma_0 \rightarrow \mathbb{R}$ such that $C_x(\eta) = 0$ for all $\eta \notin \Gamma^{(1)}$ and, for any $\eta \in \Gamma^{(1)}$, $y \in \mathbb{R}^d$ with $\eta = \{y\}$, we have $C_x(\eta) = c(x - y)$. Then, in this case, taking into account (3.17) and the obvious equality

$$\langle c(x - \cdot), \eta \cup \xi \rangle = \langle c(x - \cdot), \eta \rangle + \langle c(x - \cdot), \xi \rangle, \tag{3.19}$$

we obtain

$$(K^{-1}a(x, \cdot \cup \xi))(\eta) = a(x, \xi)0^{|\eta|} + C_x(\eta), \quad \eta \in \Gamma_0. \tag{3.20}$$

– *Exponential rate:*

$$a(x, \gamma) = e^{\langle c(x-\cdot), \gamma \rangle} = \exp \left\{ \sum_{y \in \gamma} c(x - y) \right\}, \tag{3.21}$$

where c as above. Taking into account (3.19) and (2.26), we obtain that in this case

$$(K^{-1}a(x, \cdot \cup \xi))(\eta) = a(x, \xi)e_\lambda(e^{c(x-\cdot)} - 1, \eta), \quad \eta \in \Gamma_0. \tag{3.22}$$

– *Product of linear and exponential rates:*

$$a(x, \gamma) = \langle c_1(x - \cdot), \gamma \rangle e^{c_2(x-\cdot), \gamma}, \tag{3.23}$$

where c_1 and c_2 are potentials as before. Then we have

$$a(x, \eta \cup \xi) = a(x, \eta)e^{c_2(x-\cdot), \xi} + a(x, \xi)e^{c_2(x-\cdot), \eta}. \tag{3.24}$$

Next, by (2.22),

$$\begin{aligned} (K^{-1}a(x, \cdot))(\eta) &= \sum_{\zeta \subset \eta} (-1)^{|\eta \setminus \zeta|} \sum_{y \in \zeta} c_1(x - y) e^{c_2(x-\cdot), \zeta} \\ &= \sum_{y \in \eta} c_1(x - y) \sum_{\zeta \subset \eta \setminus y} (-1)^{|\eta \setminus y \setminus \zeta|} e^{c_2(x-\cdot), \zeta \cup y}, \end{aligned}$$

and taking into account (2.26),

$$= \sum_{y \in \eta} c_1(x - y) e^{c_2(x-y)} e_\lambda(e^{c_2(x-\cdot)} - 1, \eta \setminus y). \tag{3.25}$$

By (3.24) and (3.25), we finally obtain that in this case

$$\begin{aligned} (K^{-1}a(x, \cdot \cup \xi))(\eta) &= e^{c_2(x-\cdot), \xi} \sum_{y \in \eta} c_1(x - y) e^{c_2(x-y)} e_\lambda(e^{c_2(x-\cdot)} - 1, \eta \setminus y) \\ &\quad + a(x, \xi) e_\lambda(e^{c_2(x-\cdot)} - 1, \eta), \quad \eta \in \Gamma_0. \end{aligned} \tag{3.26}$$

– *Mixing of linear and exponential rates:*

$$a(x, \gamma) = \sum_{y \in \gamma} c_1(x - y) e^{c_2(y-\cdot), \gamma \setminus y}. \tag{3.27}$$

We have

$$\begin{aligned} a(x, \eta \cup \xi) &= \sum_{y \in \eta} c_1(x - y) e^{c_2(y-\cdot), \eta \setminus y} e^{c_2(y-\cdot), \xi} \\ &\quad + \sum_{y \in \xi} c_1(x - y) e^{c_2(y-\cdot), \eta} e^{c_2(y-\cdot), \xi \setminus y}. \end{aligned}$$

Then, similarly to (3.26), we easily derive

$$\begin{aligned} (K^{-1}a(x, \cdot \cup \xi))(\eta) &= \sum_{y \in \eta} c_1(x - y)e_\lambda(e^{c_2(y-\cdot)} - 1, \eta \setminus y)e^{(c_2(y-\cdot), \xi)} \\ &+ \sum_{y \in \xi} c_1(x - y)e_\lambda(e^{c_2(y-\cdot)} - 1, \eta)e^{(c_2(y-\cdot), \xi \setminus y)}. \end{aligned}$$

Using the similar arguments one can consider polynomial rates and their compositions with exponents as well.

3.3 Semigroup evolutions in the space of quasi-observables

We proceed now to the construction of a semigroup in the space \mathcal{L}_C , $C > 0$, see (2.46), which has a generator, given by \widehat{L} , with a proper domain. To define such domain, let us set

$$D(\eta) := \sum_{x \in \eta} d(x, \eta \setminus x) \geq 0, \quad \eta \in \Gamma_0; \tag{3.28}$$

$$\mathcal{D} := \{G \in \mathcal{L}_C \mid D(\cdot)G \in \mathcal{L}_C\}. \tag{3.29}$$

Note that $B_{bs}(\Gamma_0) \subset \mathcal{D}$ and $B_{bs}(\Gamma_0)$ is a dense set in \mathcal{L}_C . Therefore, \mathcal{D} is also a dense set in \mathcal{L}_C . We will show now that $(\widehat{L}, \mathcal{D})$ given by (3.6), (3.29) generates C_0 -semigroup on \mathcal{L}_C if only ‘the full energy of death’, given by (3.28), is big enough.

Theorem 3.7 *Suppose that there exists $a_1 \geq 1, a_2 > 0$ such that for all $\xi \in \Gamma_0$ and a.a. $x \in \mathbb{R}^d$*

$$\sum_{x \in \xi} \int_{\Gamma_0} |K^{-1}d(x, \cdot \cup \xi \setminus x)|(\eta) C^{|\eta|} d\lambda(\eta) \leq a_1 D(\xi), \tag{3.30}$$

$$\sum_{x \in \xi} \int_{\Gamma_0} |K^{-1}b(x, \cdot \cup \xi \setminus x)|(\eta) C^{|\eta|} d\lambda(\eta) \leq a_2 D(\xi). \tag{3.31}$$

and, moreover,

$$a_1 + \frac{a_2}{C} < \frac{3}{2}. \tag{3.32}$$

Then $(\widehat{L}, \mathcal{D})$ is the generator of a holomorphic semigroup $\widehat{T}(t)$ on \mathcal{L}_C .

Remark 3.8 Having in mind Remark 3.3 one can say that the idea of the proof is to show that the multiplication part of the diagonal operator (3.9) will dominate on the rest part of the operator matrix $(\widehat{L}_{n,m})$ provided the conditions (3.30), (3.31) hold. Note also that, by (2.20), (2.18), (2.19), (3.28), the l.h.s of (3.30) is equal to

$$D(\xi) + \sum_{x \in \xi} \int_{\Gamma_0 \setminus \{\emptyset\}} |K^{-1}d(x, \cdot \cup \xi \setminus x)|(\eta) C^{|\eta|} d\lambda(\eta).$$

This is the reason to demand that a_1 should be not less than 1.

Proof of Theorem 3.7 Let us consider the multiplication operator (L_0, \mathcal{D}) on \mathcal{L}_C given by

$$(L_0G)(\eta) = -D(\eta)G(\eta), \quad G \in \mathcal{D}, \quad \eta \in \Gamma_0. \tag{3.33}$$

We recall that a densely defined closed operators A on \mathcal{L}_C is called sectorial of angle $\omega \in (0, \frac{\pi}{2})$ if its resolvent set $\rho(A)$ contains the sector

$$\text{Sect}\left(\frac{\pi}{2} + \omega\right) := \left\{z \in \mathbb{C} \mid |\arg z| < \frac{\pi}{2} + \omega\right\} \setminus \{0\},$$

and for each $\varepsilon \in (0; \omega)$ there exists $M_\varepsilon \geq 1$ such that

$$\|R(z, A)\| \leq \frac{M_\varepsilon}{|z|} \tag{3.34}$$

for all $z \neq 0$ with $|\arg z| \leq \frac{\pi}{2} + \omega - \varepsilon$. Here and below we will use notation

$$R(z, A) := (z\mathbb{I} - A)^{-1}, \quad z \in \rho(A).$$

The set of all sectorial operators of angle $\omega \in (0, \frac{\pi}{2})$ in \mathcal{L}_C we denote by $\mathcal{H}_C(\omega)$. Any $A \in \mathcal{H}_C(\omega)$ is a generator of a bounded semigroup $T(t)$ which is holomorphic in the sector $|\arg t| < \omega$ (see, e.g. [19, Theorem II.4.6]). One can prove the following lemma:

Lemma 3.9 *The operator (L_0, \mathcal{D}) given by (3.33) is a generator of a contraction semigroup on \mathcal{L}_C . Moreover, $L_0 \in \mathcal{H}_C(\omega)$ for all $\omega \in (0, \frac{\pi}{2})$ and (3.34) holds with $M_\varepsilon = \frac{1}{\cos \omega}$ for all $\varepsilon \in (0; \omega)$.*

Proof of Lemma 3.9 It is not difficult to show that the densely defined operator L_0 is closed in \mathcal{L}_C . Let $0 < \omega < \frac{\pi}{2}$ be arbitrary and fixed. Clear, that for all $z \in \text{Sect}(\frac{\pi}{2} + \omega)$

$$|D(\eta) + z| > 0, \quad \eta \in \Gamma_0.$$

Therefore, for any $z \in \text{Sect}(\frac{\pi}{2} + \omega)$ the inverse operator $R(z, L_0) = (z\mathbb{I} - L_0)^{-1}$, the action of which is given by

$$(R(z, L_0)G)(\eta) = \frac{1}{D(\eta) + z} G(\eta), \tag{3.35}$$

is well defined on the whole space \mathcal{L}_C . Moreover,

$$|D(\eta) + z| = \sqrt{(D(\eta) + \text{Re } z)^2 + (\text{Im } z)^2} \geq \begin{cases} |z|, & \text{if } \text{Re } z \geq 0 \\ |\text{Im } z|, & \text{if } \text{Re } z < 0 \end{cases},$$

and for any $z \in \text{Sect}(\frac{\pi}{2} + \omega)$

$$|\text{Im } z| = |z| \sin \arg z \geq |z| \left| \sin \left(\frac{\pi}{2} + \omega \right) \right| = |z| \cos \omega.$$

As a result, for any $z \in \text{Sect}(\frac{\pi}{2} + \omega)$

$$\|R(z, L_0)\| \leq \frac{1}{|z| \cos \omega}, \tag{3.36}$$

that implies the second assertion. Note also that $|D(\eta) + z| \geq \text{Re } z$ for $\text{Re } z > 0$; hence,

$$\|R(z, L_0)\| \leq \frac{1}{\text{Re } z} \tag{3.37}$$

that proves the first statement by the classical Hille–Yosida theorem. □

For any $G \in B_{\text{bs}}(\Gamma_0)$ we define

$$\begin{aligned} (L_1 G)(\eta) &:= (\widehat{L}G)(\eta) - (L_0 G)(\eta) \\ &= - \sum_{\xi \subsetneq \eta} G(\xi) \sum_{x \in \xi} (K^{-1}d(x, \cdot \cup \xi \setminus x))(\eta \setminus \xi) \\ &\quad + \sum_{\xi \subset \eta} \int_{\mathbb{R}^d} G(\xi \cup x) (K^{-1}b(x, \cdot \cup \xi))(\eta \setminus \xi) dx. \end{aligned} \tag{3.38}$$

Next Lemma shows that, under conditions (3.30), (3.31) above, the operator L_1 is relatively bounded by the operator L_0 .

Lemma 3.10 *Let (3.30), (3.31) hold. Then (L_1, \mathcal{D}) is a well-defined operator in \mathcal{L}_C such that*

$$\|L_1 R(z, L_0)\| \leq a_1 - 1 + \frac{a_2}{C}, \quad \text{Re } z > 0 \tag{3.39}$$

and

$$\|L_1 G\| \leq \left(a_1 - 1 + \frac{a_2}{C} \right) \|L_0 G\|, \quad G \in \mathcal{D}. \tag{3.40}$$

Proof of Lemma 3.10 By Lemma 3.4, we have for any $G \in \mathcal{L}_C$, $\operatorname{Re} z > 0$

$$\begin{aligned} & \int_{\Gamma_0} \left| - \sum_{\xi \not\subset \eta} \frac{1}{z + D(\xi)} G(\xi) \sum_{x \in \xi} (K^{-1}d(x, \cdot \cup \xi \setminus x))(\eta \setminus \xi) \right| C^{|\eta|} d\lambda(\eta) \\ & \leq \int_{\Gamma_0} \sum_{\xi \not\subset \eta} \frac{1}{|z + D(\xi)|} |G(\xi)| \sum_{x \in \xi} |K^{-1}d(x, \cdot \cup \xi \setminus x)|(\eta \setminus \xi) C^{|\eta|} d\lambda(\eta) \\ & = \int_{\Gamma_0} \frac{1}{|z + D(\xi)|} |G(\xi)| \sum_{x \in \xi} \int_{\Gamma_0} |K^{-1}d(x, \cdot \cup \xi \setminus x)|(\eta) C^{|\eta|} d\lambda(\eta) C^{|\xi|} d\lambda(\xi) \\ & \quad - \int_{\Gamma_0} \frac{1}{|z + D(\eta)|} D(\eta) |G(\eta)| C^{|\eta|} d\lambda(\eta) \\ & \leq (a_1 - 1) \int_{\Gamma_0} \frac{1}{\operatorname{Re} z + D(\eta)} D(\eta) |G(\eta)| C^{|\eta|} d\lambda(\eta) \leq (a_1 - 1) \|G\|_C, \end{aligned}$$

and

$$\begin{aligned} & \int_{\Gamma_0} \left| \sum_{\xi \subset \eta} \int_{\mathbb{R}^d} \frac{1}{z + D(\xi \cup x)} G(\xi \cup x) (K^{-1}b(x, \cdot \cup \xi))(\eta \setminus \xi) dx \right| C^{|\eta|} d\lambda(\eta) \\ & \leq \int_{\Gamma_0} \int_{\Gamma_0} \int_{\mathbb{R}^d} \frac{1}{|z + D(\xi \cup x)|} |G(\xi \cup x)| |K^{-1}b(x, \cdot \cup \xi)|(\eta) dx C^{|\eta|} C^{|\xi|} d\lambda(\xi) d\lambda(\eta) \\ & = \frac{1}{C} \int_{\Gamma_0} \frac{1}{|z + D(\xi)|} |G(\xi)| \sum_{x \in \xi} \int_{\Gamma_0} |K^{-1}b(x, \cdot \cup \xi \setminus x)|(\eta) C^{|\eta|} d\lambda(\eta) C^{|\xi|} d\lambda(\xi) \\ & \leq \frac{a_2}{C} \int_{\Gamma_0} \frac{1}{\operatorname{Re} z + D(\xi)} |G(\xi)| D(\xi) C^{|\xi|} d\lambda(\xi) \leq \frac{a_2}{C} \|G\|_C. \end{aligned}$$

Combining these inequalities we obtain (3.39). The same considerations yield

$$\begin{aligned} & \int_{\Gamma_0} \left| - \sum_{\xi \not\subset \eta} G(\xi) \sum_{x \in \xi} (K^{-1}d(x, \cdot \cup \xi \setminus x))(\eta \setminus \xi) \right| C^{|\eta|} d\lambda(\eta) \\ & \quad + \int_{\Gamma_0} \left| \sum_{\xi \subset \eta} \int_{\mathbb{R}^d} G(\xi \cup x) (K^{-1}b(x, \cdot \cup \xi))(\eta \setminus \xi) dx \right| C^{|\eta|} d\lambda(\eta) \\ & \leq \left((a_1 - 1) + \frac{a_2}{C} \right) \int_{\Gamma_0} |G(\eta)| D(\eta) C^{|\eta|} d\lambda(\eta), \end{aligned}$$

that proves (3.40) as well.

And now we proceed to finish the proof of the Theorem 3.7. Let us set

$$\theta := a_1 + \frac{a_2}{C} - 1 \in \left(0; \frac{1}{2} \right).$$

Then $\frac{\theta}{1-\theta} \in (0; 1)$. Let $\omega \in (0; \frac{\pi}{2})$ be such that $\cos \omega < \frac{\theta}{1-\theta}$. Then, by the proof of Lemma 3.9, $L_0 \in \mathcal{H}_C(\omega)$ and $\|R(z, L_0)\| \leq \frac{M}{|z|}$ for all $z \neq 0$ with $|\arg z| \leq \frac{\pi}{2} + \omega$, where $M := \frac{1}{\cos \omega}$. Then

$$\theta = \frac{1}{1 + \frac{1-\theta}{\theta}} < \frac{1}{1 + \frac{1}{\cos \omega}} = \frac{1}{1 + M}.$$

Hence, by (3.40) and the proof of [19, Theorem III.2.10], we have that $(\widehat{L} = L_0 + L_1, \mathcal{D})$ is a generator of holomorphic semigroup on \mathcal{L}_C . □



Remark 3.11 By (3.28), the estimates (3.30), (3.31) are satisfied if

$$\int_{\Gamma_0} |K^{-1}d(x, \cdot \cup \xi)|(\eta) C^{|\eta|} d\lambda(\eta) \leq a_1 d(x, \xi), \tag{3.41}$$

$$\int_{\Gamma_0} |K^{-1}b(x, \cdot \cup \xi)|(\eta) C^{|\eta|} d\lambda(\eta) \leq a_2 d(x, \xi). \tag{3.42}$$

3.4 Evolutions in the space of correlation functions

In this Subsection we will use the semigroup $\widehat{T}(t)$ acting on the space of quasi-observables for a construction of solution to the evolution Eq. (2.41) on the space of correlation functions.

We denote $d\lambda_C := C^{|\cdot|}d\lambda$; and the dual space $(\mathcal{L}_C)' = (L^1(\Gamma_0, d\lambda_C))' = L^\infty(\Gamma_0, d\lambda_C)$. As was mentioned before the space $(\mathcal{L}_C)'$ is isometrically isomorphic to the Banach space \mathcal{K}_C considered in (2.44)–(2.45). The isomorphism is given by the isometry R_C

$$(\mathcal{L}_C)' \ni k \mapsto R_C k := k \cdot C^{|\cdot|} \in \mathcal{K}_C. \tag{3.43}$$

Recall, one may consider the duality between the Banach spaces \mathcal{L}_C and \mathcal{K}_C given by (2.39) with

$$|\langle\langle G, k \rangle\rangle| \leq \|G\|_C \cdot \|k\|_{\mathcal{K}_C}.$$

Let $(\widehat{L}', \text{Dom}(\widehat{L}'))$ be an operator in $(\mathcal{L}_C)'$ which is dual to the closed operator $(\widehat{L}, \mathcal{D})$. We consider also its image on \mathcal{K}_C under the isometry R_C . Namely, let $\widehat{L}^* = R_C \widehat{L}' R_{C^{-1}}$ with the domain $\text{Dom}(\widehat{L}^*) = R_C \text{Dom}(\widehat{L}')$. Similarly, one can consider the adjoint semigroup $\widehat{T}'(t)$ in $(\mathcal{L}_C)'$ and its image $\widehat{T}^*(t)$ in \mathcal{K}_C .

The space \mathcal{L}_C is not reflexive: hence, $\widehat{T}^*(t)$ is not C_0 -semigroup in whole \mathcal{K}_C . By, e.g. [19, Subsection II.2.5], the last semigroup will be weak*-continuous, weak*-differentiable at 0 and \widehat{L}^* will be weak*-generator of $\widehat{T}^*(t)$. Therefore, one has an evolution in the space of correlation functions. In fact, we have a solution to the evolution Eq. (2.41), in a weak*-sense. This subsection is devoted to the study of a classical solution to this equation. By, e.g. [19, Subsection II.2.6], the restriction $\widehat{T}^\circ(t)$ of the semigroup $\widehat{T}^*(t)$ onto its invariant Banach subspace $\text{Dom}(\widehat{L}^*)$ (here and below all closures are in the norm of the space \mathcal{K}_C) is a strongly continuous semigroup. Moreover, its generator \widehat{L}° will be a part of \widehat{L}^* , namely

$$\text{Dom}(\widehat{L}^\circ) = \left\{ k \in \text{Dom}(\widehat{L}^*) \mid \widehat{L}^* k \in \overline{\text{Dom}(\widehat{L}^*)} \right\} \tag{3.44}$$

and $\widehat{L}^\circ k = \widehat{L}^* k$ for any $k \in \text{Dom}(\widehat{L}^\circ)$.

Proposition 3.12 *Let (3.30), (3.31) be satisfied. Suppose that there exists $A > 0$, $N \in \mathbb{N}_0$, $\nu \geq 1$ such that for $\xi \in \Gamma_0$ and $x \notin \xi$*

$$d(x, \xi) \leq A(1 + |\xi|)^N \nu^{|\xi|}, \tag{3.45}$$

Then for any $\alpha \in (0; \frac{1}{\nu})$

$$\mathcal{K}_{\alpha C} \subset \text{Dom}(\widehat{L}^*). \tag{3.46}$$

Proof In order to show (3.46) it is enough to verify that for any $k \in \mathcal{K}_{\alpha C}$ there exists $k^* \in \mathcal{K}_C$ such that for any $G \in \text{Dom}(\widehat{L})$

$$\langle\langle \widehat{L}G, k \rangle\rangle = \langle\langle G, k^* \rangle\rangle. \tag{3.47}$$

By the same calculations as in the proof of Proposition 3.5, it is easy to see that (3.47) is valid for any $k \in \mathcal{K}_{\alpha C}$ with $k^* = \widehat{L}^* k$, where \widehat{L}^* is given by (3.13), provided $k^* \in \mathcal{K}_C$.

To prove the last inclusion, one can estimate, by (3.30), (3.31), (3.45) that

$$\begin{aligned} & C^{-|\eta|} |(\widehat{L}^* k)(\eta)| \\ & \leq C^{-|\eta|} \sum_{x \in \eta} \int_{\Gamma_0} |k(\zeta \cup \eta)| |K^{-1}d(x, \cdot \cup \eta \setminus x)|(\zeta) d\lambda(\zeta) \end{aligned}$$

$$\begin{aligned}
 &+ C^{-|\eta|} \sum_{x \in \eta} \int_{\Gamma_0} |k(\zeta \cup (\eta \setminus x))| |K^{-1}b(x, \cdot \cup \eta \setminus x)|(\zeta) d\lambda(\zeta) \\
 &\leq \|k\|_{\mathcal{K}_{\alpha C}} \alpha^{|\eta|} \sum_{x \in \eta} \int_{\Gamma_0} (\alpha C)^{|\zeta|} |K^{-1}d(x, \cdot \cup \eta \setminus x)|(\zeta) d\lambda(\zeta) \\
 &\quad + \frac{1}{\alpha C} \|k\|_{\mathcal{K}_{\alpha C}} \alpha^{|\eta|} \sum_{x \in \eta} \int_{\Gamma_0} (\alpha C)^{|\zeta|} |K^{-1}b(x, \cdot \cup \eta \setminus x)|(\zeta) d\lambda(\zeta) \\
 &\leq \|k\|_{\mathcal{K}_{\alpha C}} \left(a_1 + \frac{a_2}{\alpha C}\right) \alpha^{|\eta|} \sum_{x \in \eta} d(x, \eta \setminus x) \\
 &\leq A \|k\|_{\mathcal{K}_{\alpha C}} \left(a_1 + \frac{a_2}{\alpha C}\right) \alpha^{|\eta|} (1 + |\eta|)^{N+1} \nu^{|\eta|-1}.
 \end{aligned}$$

Using elementary inequality

$$(1 + t)^b a^t \leq \frac{1}{a} \left(\frac{b}{-e \ln a}\right)^b, \quad b \geq 1, a \in (0; 1), t \geq 0, \tag{3.48}$$

we have for $\alpha \nu < 1$

$$\operatorname{ess\,sup}_{\eta \in \Gamma_0} C^{-|\eta|} |(\widehat{L}^*k)(\eta)| \leq \|k\|_{\mathcal{K}_{\alpha C}} \left(a_1 + \frac{a_2}{\alpha C}\right) \frac{A}{\alpha \nu^2} \left(\frac{N + 1}{-e \ln(\alpha \nu)}\right)^{N+1} < \infty.$$

The statement is proved. □

Lemma 3.13 *Let (3.45) hold. We define for any $\alpha \in (0; 1)$*

$$\mathcal{D}_\alpha := \{G \in \mathcal{L}_{\alpha C} \mid D(\cdot)G \in \mathcal{L}_{\alpha C}\}.$$

Then for any $\alpha \in (0; \frac{1}{\nu})$

$$\mathcal{D} \subset \mathcal{L}_C \subset \mathcal{D}_\alpha \subset \mathcal{L}_{\alpha C} \tag{3.49}$$

Proof The first and last inclusions are obvious. To prove the second one, we use (3.45), (3.48) and obtain for any $G \in \mathcal{L}_C$

$$\begin{aligned}
 \int_{\Gamma_0} D(\eta) |G(\eta)| (\alpha C)^{|\eta|} d\lambda(\eta) &\leq \int_{\Gamma_0} \alpha^{|\eta|} \sum_{x \in \eta} A(1 + |\eta|)^{N+1} \nu^{|\eta|-1} |G(\eta)| C^{|\eta|} d\lambda(\eta) \\
 &\leq \operatorname{const} \int_{\Gamma_0} |G(\eta)| C^{|\eta|} d\lambda(\eta) < \infty.
 \end{aligned}$$

The statement is proved. □

Proposition 3.14 *Let (3.30), (3.31), and (3.45) hold with*

$$a_1 + \frac{a_2}{\alpha C} < \frac{3}{2} \tag{3.50}$$

for some $\alpha \in (0; 1)$. Then $(\widehat{L}, \mathcal{D}_\alpha)$ is a generator of a holomorphic semigroup $\widehat{T}_\alpha(t)$ on $\mathcal{L}_{\alpha C}$.

Proof The proof is similar to the proof of Theorem 3.7, taking into account that bounds (3.31), (3.30) imply the same bounds for αC instead of C . Note also that (3.50) is stronger than (3.32). □

Proposition 3.15 *Let (3.30), (3.31), and (3.45) hold with*

$$1 \leq \nu < \frac{C}{a_2} \left(\frac{3}{2} - a_1\right). \tag{3.51}$$

Then, for any α with

$$\frac{a_2}{C(\frac{3}{2} - a_1)} < \alpha < \frac{1}{\nu}, \tag{3.52}$$

the set $\overline{\mathcal{K}_{\alpha C}}$ is a $\widehat{T}^\circ(t)$ -invariant Banach subspace of \mathcal{K}_C . Moreover, the set $\mathcal{K}_{\alpha C}$ is $\widehat{T}^\circ(t)$ -invariant too.



Proof First of all note that the condition on α implies (3.50). Next, we prove that $\widehat{T}_\alpha(t)G = \widehat{T}(t)G$ for any $G \in \mathcal{L}_C \subset \mathcal{L}_{\alpha C}$. Let $\widehat{L}_\alpha = (\widehat{L}, \mathcal{D}_\alpha)$ be the operator in $\mathcal{L}_{\alpha C}$. There exists $\omega > 0$ such that $(\omega; +\infty) \subset \rho(\widehat{L}) \cap \rho(\widehat{L}_\alpha)$, see, e.g. [19, Section III.2]. For some fixed $z \in (\omega; +\infty)$ we denote by $R(z, \widehat{L}) = (z\mathbb{I} - \widehat{L})^{-1}$ the resolvent of $(\widehat{L}, \mathcal{D})$ in \mathcal{L}_C and by $R(z, \widehat{L}_\alpha) = (z\mathbb{I} - \widehat{L}_\alpha)^{-1}$ the resolvent of \widehat{L}_α in $\mathcal{L}_{\alpha C}$. Then for any $G \in \mathcal{L}_C$ we have $R(z, \widehat{L})G \in \mathcal{D} \subset \mathcal{D}_\alpha$ and

$$R(z, \widehat{L})G - R(z, \widehat{L}_\alpha)G = R(z, \widehat{L}_\alpha)((z\mathbb{I} - \widehat{L}_\alpha) - (z\mathbb{I} - \widehat{L}))R(z, \widehat{L})G = 0,$$

since $\widehat{L}_\alpha = \widehat{L}$ on \mathcal{D} . As a result, $\widehat{T}_\alpha(t)G = \widehat{T}(t)G$ on \mathcal{L}_C .

Note that for any $G \in \mathcal{L}_C \subset \mathcal{L}_{\alpha C}$ and for any $k \in \mathcal{K}_{\alpha C} \subset \mathcal{K}_C$ we have $\widehat{T}_\alpha(t)G \in \mathcal{L}_{\alpha C}$ and

$$\langle \widehat{T}_\alpha(t)G, k \rangle = \langle G, \widehat{T}_\alpha^*(t)k \rangle,$$

where, by the same construction as before, $\widehat{T}_\alpha^*(t)k \in \mathcal{K}_{\alpha C}$. But $G \in \mathcal{L}_C, k \in \mathcal{K}_C$ implies

$$\langle \widehat{T}_\alpha(t)G, k \rangle = \langle \widehat{T}(t)G, k \rangle = \langle G, \widehat{T}^*(t)k \rangle.$$

Hence, $\widehat{T}^*(t)k = \widehat{T}_\alpha^*(t)k \in \mathcal{K}_{\alpha C}$ that proves the statement due to continuity of the family $\widehat{T}^*(t)$. □

By, e.g. [19, Subsection II.2.3], one can consider the restriction $\widehat{T}^{\circ\alpha}(t)$ of the semigroup $\widehat{T}^\circ(t)$ onto $\overline{\mathcal{K}_{\alpha C}}$. It will be strongly continuous semigroup with the generator $\widehat{L}^{\circ\alpha}$ which is a restriction of \widehat{L}° onto $\overline{\mathcal{K}_{\alpha C}}$. Namely cf. 3.44,

$$\text{Dom}(\widehat{L}^{\circ\alpha}) = \left\{ k \in \overline{\mathcal{K}_{\alpha C}} \mid \widehat{L}^*k \in \overline{\mathcal{K}_{\alpha C}} \right\}, \tag{3.53}$$

and $\widehat{L}^{\circ\alpha}k = \widehat{L}^\circ k = \widehat{L}^*k$ for any $k \in \overline{\mathcal{K}_{\alpha C}}$. In the other words, $\widehat{L}^{\circ\alpha}$ is a part of \widehat{L}^* .

And now we may proceed to the main statement of this Subsection.

Theorem 3.16 *Let (3.30), (3.31), (3.45), and (3.51) hold, and let α be chosen as in (3.52). Then for any $k_0 \in \overline{\mathcal{K}_{\alpha C}}$ there exists a unique classical solution to (2.41) in the space $\overline{\mathcal{K}_{\alpha C}}$, and this solution is given by $k_t = \widehat{T}^{\circ\alpha}(t)k_0$. Moreover, $k_0 \in \mathcal{K}_{\alpha C}$ implies $k_t \in \mathcal{K}_{\alpha C}$.*

Proof We recall that $(\widehat{L}, \mathcal{D})$ is a densely defined closed operator on \mathcal{L}_C (as a generator of a C_0 -semigroup $\widehat{T}(t)$). Then, by, e.g. [79, Lemma 1.4.1], for the dual operator $(\widehat{L}^*, \text{Dom}(\widehat{L}^*))$ we have that $\rho(\widehat{L}^*) = \rho(\widehat{L})$ and, for any $z \in \rho(\widehat{L}), R(z, \widehat{L}^*) = R(z, \widehat{L})^*$. In particular,

$$\|R(z, \widehat{L}^*)\| = \|R(z, \widehat{L})^*\| = \|R(z, \widehat{L})\|. \tag{3.54}$$

Next, if we denote by $R(z, \widehat{L})^\circ$ the restriction of $R(z, \widehat{L})^*$ onto $R(z, \widehat{L})^*$ -invariant space $\overline{\text{Dom}(\widehat{L}^*)}$ then, by, e.g. [79, Theorem 1.4.2], $\rho(\widehat{L}^\circ) = \rho(\widehat{L}^*)$ and, for any $z \in \rho(\widehat{L}^*) = \rho(\widehat{L}), R(z, \widehat{L}^\circ) = R(z, \widehat{L})^\circ$. Therefore, by (3.54),

$$\|R(z, \widehat{L}^\circ)\| \leq \|R(z, \widehat{L})\|.$$

Then, taking into account that by Theorem 3.7 the operator $(\widehat{L}, \mathcal{D})$ is a generator of the holomorphic semigroup $\widehat{T}(t)$, we immediately conclude that the same property has the semigroup $\widehat{T}^\circ(t)$ with the generator $(\widehat{L}^\circ, \text{Dom}(\widehat{L}^\circ))$ in the space $\overline{\text{Dom}(\widehat{L}^*)}$.

As a result, by, e.g. [70, Corollary 4.1.5], the initial value problem (2.41) in the Banach space $\overline{\text{Dom}(\widehat{L}^*)}$ has a unique classical solution for any $k_0 \in \overline{\text{Dom}(\widehat{L}^*)}$. In particular, it means that the solution $k_t = \widehat{T}^\circ(t)k_0$ is continuously differentiable in t w.r.t. the norm of $\overline{\text{Dom}(\widehat{L}^*)}$ that is the norm $\|\cdot\|_{\mathcal{K}_C}$ and also $k_t \in \text{Dom}(\widehat{L}^\circ)$. But by Proposition 3.15, the space $\overline{\mathcal{K}_{\alpha C}}$ is $\widehat{T}^\circ(t)$ -invariant. Hence, if we consider now the initial value $k_0 \in \overline{\mathcal{K}_{\alpha C}} \subset \overline{\text{Dom}(\widehat{L}^*)}$ we obtain with a necessity that $k_t = \widehat{T}^\circ(t)k_0 = \widehat{T}^{\circ\alpha}(t)k_0 \in \overline{\mathcal{K}_{\alpha C}}$. Therefore, $k_t \in \overline{\mathcal{K}_{\alpha C}} \cap \text{Dom}(\widehat{L}^\circ) = \text{Dom}(\widehat{L}^{\circ\alpha})$ (see again [19, Subsection II.2.3]) and, recall, k_t is continuously differentiable in t w.r.t. the norm $\|\cdot\|_{\mathcal{K}_C}$ that is the norm in $\overline{\mathcal{K}_{\alpha C}}$. This completes the proof of the first statement. The second one follows directly now from Proposition 3.15. □

3.5 Examples of dynamics

We proceed now to describing the concrete birth-and-death dynamics which are important for different application. We will consider the explicit conditions on parameters of systems which imply the general conditions on rates b and d above. For simplicity of notations we denote the l.h.s. of (3.30) and (3.31) by $I_d(\xi)$ and $I_b(\xi)$, $\xi \in \Gamma_0$, respectively.

Example 3.17 (Surgailis dynamics) Let the rates d and b are independent on configuration variable, namely

$$d(x, \gamma) = m(x), \quad b(x, \gamma) = z(x), \quad x \in \mathbb{R}^d, \gamma \in \Gamma, \tag{3.55}$$

where $0 < m, z \in L^\infty(\mathbb{R}^d)$. Then, by (3.17) we obtain that

$$I_d(\xi) = \langle m, \xi \rangle = D(\xi), \quad I_b(\xi) = \langle z, \xi \rangle, \quad \xi \in \Gamma_0.$$

Therefore, (3.30), (3.31), (3.32) hold if only

$$z(x) \leq am(x), \quad x \in \mathbb{R}^d \tag{3.56}$$

with any

$$0 < a < \frac{C}{2}. \tag{3.57}$$

Clearly, in this case (3.45) holds with $N = 0, \nu = 1$; therefore, the condition (3.52) is just

$$\frac{2a}{C} < \alpha < 1. \tag{3.58}$$

The case of constant (in space) m and σ was considered in [23]. Similarly to that results, one can derive the explicit expression for the solution to the initial value problem (2.41) considered point-wise in Γ_0 , namely

$$k_t(\eta) = e_\lambda(e^{-tm}, \eta) \sum_{\xi \subset \eta} e_\lambda\left(\frac{z}{m}(e^{tm} - 1), \xi\right) k_0(\eta \setminus \xi), \quad \eta \in \Gamma_0. \tag{3.59}$$

Note that, using (3.59), it can be possible to show directly that the statement of Theorem 3.16 still holds if we drop 2 in (3.57) and (3.58).

Example 3.18 (Glauber-type dynamics). Let L be given by (3.1) with

$$d(x, \gamma \setminus x) = m(x) \exp \left\{ s \sum_{y \in \gamma \setminus x} \phi(x - y) \right\}, \quad x \in \gamma, \gamma \in \Gamma, \tag{3.60}$$

$$b(x, \gamma) = z(x) \exp \left\{ (s - 1) \sum_{y \in \gamma} \phi(x - y) \right\}, \quad x \in \mathbb{R}^d \setminus \gamma, \gamma \in \Gamma, \tag{3.61}$$

where $\phi : \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{R}_+$ is a pair potential, $\phi(-x) = \phi(x), 0 < z, m \in L^\infty(\mathbb{R}^d)$, and $s \in [0; 1]$. Note that in the case $m(x) \equiv 1, z(x) \equiv z > 0$ and for any $s \in [0; 1]$ the operator L is well defined and, moreover, symmetric in the space $L^2(\Gamma, \mu)$, where μ is a Gibbs measure, given by the pair potential ϕ and activity parameter z (see, e.g. [55] and references therein). This gives possibility to study the corresponding semigroup in $L^2(\Gamma, \mu)$. If, additionally, $s = 0$, the corresponding dynamics was also studied in another Banach spaces, see, e.g. [28, 34, 53]. Below we show that one of the main results of the paper stated in Theorem 3.16 can be applied to the case of arbitrary $s \in [0; 1]$ and non-constant m and z . Set

$$\beta_\tau := \int_{\mathbb{R}^d} |e^{\tau\phi(x)} - 1| dx \in [0; \infty], \quad \tau \in [-1; 1]. \tag{3.62}$$

Let s be arbitrary and fixed. Suppose that $\beta_s < \infty, \beta_{s-1} < \infty$. Then, by (3.60), (3.61), (3.22), and (2.28),

$$I_d(\xi) = \sum_{x \in \xi} d(x, \xi \setminus x) e^{C\beta_s} = D(\xi) e^{C\beta_s},$$

and, analogously, taking into account that $\phi \geq 0$,

$$I_b(\xi) = \sum_{x \in \xi} b(x, \xi \setminus x) e^{C\beta_{s-1}} \leq \sum_{x \in \xi} \frac{z(x)}{m(x)} d(x, \xi \setminus x) e^{C\beta_{s-1}}$$

Therefore, to apply Theorem 3.7 we should assume that there exists $\sigma > 0$ such that

$$z(x) \leq \sigma m(x), \quad x \in \mathbb{R}^d, \tag{3.63}$$

and

$$e^{C\beta_s} + \frac{\sigma}{C} e^{C\beta_{s-1}} < \frac{3}{2}. \tag{3.64}$$

In particular, for $s = 0$ we need

$$\frac{\sigma}{C} e^{C\beta_{-1}} < \frac{1}{2}. \tag{3.65}$$

Next, to have (3.45) and (3.51), we will distinguish two cases. For $s = 0$ we obtain (3.45) since $m \in L^\infty(\mathbb{R}^d)$. In this case, $\nu = 1$ that fulfills (3.51) as well. For $s \in (0, 1]$, we should assume that

$$0 \leq \phi \in L^\infty(\mathbb{R}^d). \tag{3.66}$$

Then, by (3.60), $\nu = e^{s\bar{\phi}} \geq 1$, where $\bar{\phi} := \|\phi\|_{L^\infty(\mathbb{R}^d)}$. Therefore, to have (3.51), we need the following improvement of (3.64):

$$e^{C\beta_s} + \frac{\sigma}{C} e^{s\bar{\phi} + C\beta_{s-1}} < \frac{3}{2}. \tag{3.67}$$

Example 3.19 (Bolker–Dieckman–Law–Pacala (BDLP) model) This example describes a generalization of the model of plant ecology (see [26] and references therein). Let L be given by (3.1) with

$$d(x, \gamma \setminus x) = m(x) + \varkappa^-(x) \sum_{y \in \gamma \setminus x} a^-(x - y), \quad x \in \gamma, \gamma \in \Gamma, \tag{3.68}$$

$$b(x, \gamma) = \varkappa^+(x) \sum_{y \in \gamma} a^+(x - y), \quad x \in \mathbb{R}^d \setminus \gamma, \gamma \in \Gamma, \tag{3.69}$$

where $0 < m \in L^\infty(\mathbb{R}^d)$, $0 \leq \varkappa^\pm \in L^\infty(\mathbb{R}^d)$, $0 \leq a^\pm \in L^1(\mathbb{R}^d, dx) \cap L^\infty(\mathbb{R}^d, dx)$, $\int_{\mathbb{R}^d} a^\pm(x) dx = 1$. Then, by (3.17), (3.20), and (2.18)–(2.19),

$$I_d(\xi) = \sum_{x \in \xi} d(x, \xi \setminus x) + \sum_{x \in \xi} C\varkappa^-(x), \quad I_b(\xi) = \sum_{x \in \xi} b(x, \xi \setminus x) + \sum_{x \in \xi} C\varkappa^+(x).$$

Let us suppose, cf. [26], that there exists $\delta > 0$ such that

$$(4 + \delta)C\varkappa^-(x) \leq m(x), \quad x \in \mathbb{R}^d, \tag{3.70}$$

$$(4 + \delta)\varkappa^+(x) \leq m(x), \quad x \in \mathbb{R}^d, \tag{3.71}$$

$$4\varkappa^+(x)a^+(x) \leq C\varkappa^-(x)a^-(x), \quad x \in \mathbb{R}^d, \tag{3.72}$$

Then

$$d(x, \xi) + C\varkappa^-(x) \leq d(x, \xi) + \frac{m(x)}{4 + \delta} \leq \left(1 + \frac{1}{4 + \delta}\right) d(x, \xi),$$

$$b(x, \xi) + C\varkappa^+(x) \leq \frac{C}{4} \varkappa^-(x) \sum_{y \in \xi} a^-(x - y) + \frac{Cm(x)}{4 + \delta} < \frac{C}{4} d(x, \xi),$$

Hence, (3.30), (3.31) hold with

$$a_1 = 1 + \frac{1}{4 + \delta}, \quad a_2 = \frac{C}{4},$$

that fulfills (3.32). Next, under conditions (3.70), (3.72), we have

$$d(x, \xi) \leq \|m\|_{L^\infty(\mathbb{R}^d)} + \|\varkappa^-\|_{L^\infty(\mathbb{R}^d)} \|a^-\|_{L^\infty(\mathbb{R}^d)} |\xi|, \quad \xi \in \Gamma_0,$$

and hence (3.45) holds with $\nu = 1$, which makes (3.51) obvious.

Remark 3.20 It was shown in [26] that, for the case of constant m, κ^\pm , the condition like (3.70) is essential. Namely, if $m > 0$ is arbitrary small the operator \widehat{L} will not be even accretive in \mathcal{L}_C .

Example 3.21 (Contact model with establishment) Let L be given by (3.1) with $d(x, \gamma) = m(x)$ for all $\gamma \in \Gamma$ and

$$b(x, \gamma) = \kappa(x) \exp \left\{ \sum_{y \in \gamma} \phi(x - y) \right\} \sum_{y \in \gamma} a(x - y), \quad \gamma \in \Gamma, x \in \mathbb{R}^d \setminus \gamma. \tag{3.73}$$

Here $0 < m \in L^\infty(\mathbb{R}^d), 0 \leq \kappa \in L^\infty(\mathbb{R}^d), 0 \leq a \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d), \int_{\mathbb{R}^d} a(x) dx = 1$.

3.6 Stationary equation

In this subsection we study the question about stationary solutions to (2.41). For any $s \geq 0$, we consider the following subset of \mathcal{K}_C

$$\mathcal{K}_{\alpha C}^{(s)} := \{k \in \mathcal{K}_{\alpha C} \mid k(\emptyset) = s\}.$$

We define $\widetilde{\mathcal{K}}$ to be the closure of $\mathcal{K}_{\alpha C}^{(0)}$ in the norm of \mathcal{K}_C . It is clear that $\widetilde{\mathcal{K}}$ with the norm of \mathcal{K}_C is a Banach space.

Proposition 3.22 *Let (3.30), (3.31), and (3.45) be satisfied with*

$$a_1 + \frac{a_2}{C} < 2. \tag{3.74}$$

Assume, additionally, that

$$d(x, \emptyset) > 0, \quad x \in \mathbb{R}^d. \tag{3.75}$$

Then for any $\alpha \in (0; \frac{1}{v})$ the stationary equation

$$\widehat{L}^*k = 0 \tag{3.76}$$

has a unique solution k_{inv} from $\mathcal{K}_{\alpha C}^{(1)}$ which is given by the expression

$$k_{\text{inv}} = 1^* + (\mathbb{I} - S)^{-1}E. \tag{3.77}$$

Here 1^ denotes the function defined by $1^*(\eta) = 0^{|\eta|}, \eta \in \Gamma_0$; the function $E \in \mathcal{K}_{\alpha C}^{(0)}$ is such that*

$$E(\eta) = \mathbb{1}_{\Gamma^{(1)}}(\eta) \sum_{x \in \eta} \frac{b(x, \emptyset)}{d(x, \emptyset)}, \quad \eta \in \Gamma_0,$$

and S is a generalized Kirkwood–Salzburg operator on $\widetilde{\mathcal{K}}$, given by

$$\begin{aligned} (Sk)(\eta) = & -\frac{1}{D(\eta)} \sum_{x \in \eta} \int_{\Gamma_0 \setminus \{\emptyset\}} k(\zeta \cup \eta)(K^{-1}d(x, \cdot \cup \eta \setminus x))(\zeta) d\lambda(\zeta) \\ & + \frac{1}{D(\eta)} \sum_{x \in \eta} \int_{\Gamma_0} k(\zeta \cup (\eta \setminus x))(K^{-1}b(x, \cdot \cup \eta \setminus x))(\zeta) d\lambda(\zeta), \end{aligned} \tag{3.78}$$

for $\eta \neq \emptyset$ and $(Sk)(\emptyset) = 0$. In particular, if $b(x, \emptyset) = 0$ for a.a. $x \in \mathbb{R}^d$ then this solution is such that

$$k_{\text{inv}}^{(n)} = 0, \quad n \geq 1. \tag{3.79}$$

Remark 3.23 It is worth noting that (3.41), (3.42) imply (3.75).

Proof Suppose that (3.76) holds for some $k \in \mathcal{K}_{\alpha C}^{(1)}$. Then

$$\begin{aligned}
 D(\eta)k(\eta) &= -\sum_{x \in \eta} \int_{\Gamma_0 \setminus \{\emptyset\}} k(\zeta \cup \eta)(K^{-1}d(x, \cdot \cup \eta \setminus x))(\zeta) d\lambda(\zeta) \\
 &\quad + \sum_{x \in \eta} \int_{\Gamma_0} k(\zeta \cup (\eta \setminus x))(K^{-1}b(x, \cdot \cup \eta \setminus x))(\zeta) d\lambda(\zeta).
 \end{aligned}
 \tag{3.80}$$

The equality (3.80) is satisfied for any $k \in \mathcal{K}_{\alpha C}^{(1)}$ at the point $\eta = \emptyset$. Using the fact that $D(\emptyset) = 0$ one may rewrite (3.80) in terms of the function $\tilde{k} = k - 1^* \in \mathcal{K}_{\alpha C}^{(0)}$, namely

$$\begin{aligned}
 D(\eta)\tilde{k}(\eta) &= -\sum_{x \in \eta} \int_{\Gamma_0 \setminus \{\emptyset\}} \tilde{k}(\zeta \cup \eta)(K^{-1}d(x, \cdot \cup \eta \setminus x))(\zeta) d\lambda(\zeta) \\
 &\quad + \sum_{x \in \eta} \int_{\Gamma_0} \tilde{k}(\zeta \cup (\eta \setminus x))(K^{-1}b(x, \cdot \cup \eta \setminus x))(\zeta) d\lambda(\zeta) \\
 &\quad + \sum_{x \in \eta} 0^{|\eta \setminus x|} b(x, \eta \setminus x).
 \end{aligned}
 \tag{3.81}$$

As a result,

$$\tilde{k}(\eta) = (S\tilde{k})(\eta) + E(\eta), \quad \eta \in \Gamma_0.$$

Next, for $\eta \neq \emptyset$

$$\begin{aligned}
 &C^{-|\eta|} |(Sk)(\eta)| \\
 &\leq \frac{C^{-|\eta|}}{D(\eta)} \sum_{x \in \eta} \int_{\Gamma_0 \setminus \{\emptyset\}} |k(\zeta \cup \eta)| |(K^{-1}d(x, \cdot \cup \eta \setminus x))(\zeta)| d\lambda(\zeta) \\
 &\quad + \frac{C^{-|\eta|}}{D(\eta)} \sum_{x \in \eta} \int_{\Gamma_0} |k(\zeta \cup (\eta \setminus x))| |(K^{-1}b(x, \cdot \cup \eta \setminus x))(\zeta)| d\lambda(\zeta) \\
 &\leq \frac{\|k\|_{\mathcal{K}_C}}{D(\eta)} \sum_{x \in \eta} \int_{\Gamma_0 \setminus \{\emptyset\}} C^{|\zeta|} |(K^{-1}d(x, \cdot \cup \eta \setminus x))(\zeta)| d\lambda(\zeta) \\
 &\quad + \frac{\|k\|_{\mathcal{K}_C}}{D(\eta)} \frac{1}{C} \sum_{x \in \eta} \int_{\Gamma_0} C^{|\zeta|} |(K^{-1}b(x, \cdot \cup \eta \setminus x))(\zeta)| d\lambda(\zeta) \\
 &\leq \frac{\|k\|_{\mathcal{K}_C}}{D(\eta)} D(\eta) \left(a_1 - 1 + \frac{a_2}{C} \right) = \left(a_1 - 1 + \frac{a_2}{C} \right) \|k\|_{\mathcal{K}_C}.
 \end{aligned}$$

Hence,

$$\|S\| = a_1 + \frac{a_2}{C} - 1 < 1$$

in \tilde{K} . This finishes the proof. □

Remark 3.24 The name of the operator (3.78) is motivated by Example 3.18. Namely, if $s = 0$ then the operator (3.78) has form

$$(Sk)(\eta) = \frac{1}{m|\eta|} \sum_{x \in \eta} e_\lambda(e^{-\phi(x-\cdot)}, \eta \setminus x) \int_{\Gamma_0} k(\zeta \cup (\eta \setminus x)) e_\lambda(e^{-\phi(x-\cdot)} - 1, \zeta) d\lambda(\zeta),$$

that is quite similar of the so-called Kirkwood–Salsburg operator known in mathematical physics (see, e.g. [49, 75]). For $s = 0$ condition (3.74) has form $\frac{z}{C} e^{C\beta-1} < 1$. Under this condition, the stationary solution to (3.76) is unique and coincides with the correlation function of the Gibbs measure, corresponding to potential ϕ and activity z .

Remark 3.25 It is worth pointing out that $b(x, \emptyset) = 0$ in the case of Example 3.19. Therefore, if we suppose [cf. (3.70), (3.72)] that $2\kappa^-C < m$ and $2\kappa^+a^+(x) \leq C\kappa^-a^-(x)$, for $x \in \mathbb{R}^d$, condition (3.74) will be satisfied. However, the unique solution to (3.76) will be given by (3.79). In the next example we improve this statement.

Example 3.26 Let us consider the following natural modification of BDLP-model coming from Example 3.19: let d be given by (3.68) and

$$b(x, \gamma) = \kappa + \kappa^+ \sum_{y \in \gamma} a^+(x - y), \quad x \in \mathbb{R}^d \setminus \gamma, \quad \gamma \in \Gamma, \tag{3.82}$$

where κ^+, a^+ are as before and $\kappa > 0$. Then, under assumptions

$$2 \max \left\{ \kappa^- C; \frac{2\kappa}{C} \right\} < m \tag{3.83}$$

and

$$2\kappa^+a^+(x) \leq C\kappa^-a^-(x), \quad x \in \mathbb{R}^d, \tag{3.84}$$

we obtain for some $\delta > 0$

$$\begin{aligned} \int_{\Gamma_0} |K^{-1}d(x, \cdot \cup \xi)|(\eta) C^{|\eta|} d\lambda(\eta) &= d(x, \xi) + C\kappa^- \leq \left(1 + \frac{1}{2 + \delta}\right) d(x, \xi) \\ \int_{\Gamma_0} |K^{-1}b(x, \cdot \cup \xi)|(\eta) C^{|\eta|} d\lambda(\eta) &= b(x, \xi) + C\kappa^+ \\ &\leq \kappa + \frac{1}{2} C\kappa^- \sum_{y \in \xi} a^-(x - y) + \frac{m}{4} C < \frac{C}{2} d(x, \xi). \end{aligned}$$

The latter inequalities imply (3.74). In this case, $E(\eta) = \mathbb{1}_{\Gamma(\cup)}(\eta) \frac{\kappa}{m}$.

Remark 3.27 If $a^+(x) = a^-(x)$, $x \in \mathbb{R}^d$ and $\kappa^+ = z\kappa^-$, $\kappa = zm$ for some $z > 0$ then $b(x, \gamma) = zd(x, \gamma)$ and the Poisson measure π_z with the intensity z will be symmetrizing measure for the operator L . In particular, it will be invariant measure. This fact means that its correlation function $k_z(\eta) = z^{|\eta|}$ is a solution to (3.76). Conditions (3.83) and (3.84) in this case are equivalent to $4z < C$ and $2\kappa^-C < m$. As a result, due to uniqueness of such solution,

$$1^*(\eta) + z(\mathbb{1} - S)^{-1} \mathbb{1}_{\Gamma(\cup)}(\eta) = z^{|\eta|}, \quad \eta \in \Gamma_0.$$

4 Approximative approach for the Glauber dynamics

In this section we consider an approximative approach for the construction of the Glauber-type dynamics described in Example 3.18 for

$$s = 0, \quad m(x) \equiv 1, \quad z(x) \equiv z > 0.$$

Therefore, in such a case, (3.1) has the form

$$\begin{aligned} (LF)(\gamma) &:= \sum_{x \in \gamma} [F(\gamma \setminus x) - F(\gamma)] \\ &+ z \int_{\mathbb{R}^d} [F(\gamma \cup x) - F(\gamma)] \exp\{-E^\phi(x, \gamma)\} dx, \quad \gamma \in \Gamma, \end{aligned} \tag{4.1}$$

with E^ϕ given by (2.11).

Let $G \in B_{bs}(\Gamma_0)$ then $F = KG \in \mathcal{F}_{cyl}(\Gamma)$. By (3.6), (3.17), (3.22), one has the following explicit form for the mapping $\widehat{L} := K^{-1}LK$ on $B_{bs}(\Gamma_0)$

$$\begin{aligned} (\widehat{L}G)(\eta) &= -|\eta|G(\eta) \\ &+ z \sum_{\xi \subset \eta} \int_{\mathbb{R}^d} e^{-E^\phi(x, \xi)} G(\xi \cup x) e_\lambda(e^{-\phi(x^-)} - 1, \eta \setminus \xi) dx, \end{aligned} \tag{4.2}$$

where e_λ is given by (2.25).

Let us denote, for any $\eta \in \Gamma_0$,

$$(L_0G)(\eta) := -|\eta|G(\eta); \tag{4.3}$$

$$(L_1G)(\eta) := z \sum_{\xi \subset \eta} \int_{\mathbb{R}^d} e^{-E^\phi(x,\xi)} G(\xi \cup x) e_\lambda(e^{-\phi(x-\cdot)} - 1, \eta \setminus \xi) dx. \tag{4.4}$$

To simplify notation we continue to write C_ϕ for β_{-1} . In contrast to (3.29), we will not work the maximal domain of the operator L_0 , namely the following statement will be used

Proposition 4.1 *The expression (4.2) defines a linear operator \widehat{L} in \mathcal{L}_C with the dense domain $\mathcal{L}_{2C} \subset \mathcal{L}_C$.*

Proof For any $G \in \mathcal{L}_{2C}$

$$\|L_0G\|_C = \int_{\Gamma_0} |G(\eta)| |\eta| C^{|\eta|} d\lambda(\eta) < \int_{\Gamma_0} |G(\eta)| 2^{|\eta|} C^{|\eta|} d\lambda(\eta) < \infty$$

and, by Lemma 3.4,

$$\begin{aligned} \|L_1G\|_C &\leq z \int_{\Gamma_0} \sum_{\xi \subset \eta} \int_{\mathbb{R}^d} e^{-E^\phi(x,\xi)} |G(\xi \cup x)| e_\lambda \left(\left| e^{-\phi(x-\cdot)} - 1 \right|, \eta \setminus \xi \right) dx C^{|\eta|} d\lambda(\eta) \\ &= z \int_{\Gamma_0} \int_{\Gamma_0} \int_{\mathbb{R}^d} e^{-E^\phi(x,\xi)} |G(\xi \cup x)| e_\lambda \left(\left| e^{-\phi(x-\cdot)} - 1 \right|, \eta \right) dx C^{|\eta|} C^{|\xi|} d\lambda(\xi) d\lambda(\eta) \\ &\leq \frac{z}{C} \exp\{CC_\phi\} \int_{\Gamma_0} |G(\xi)| |\xi| C^{|\xi|} d\lambda(\xi) < \frac{z}{C} \exp\{CC_\phi\} \int_{\Gamma_0} |G(\xi)| 2^{|\xi|} C^{|\xi|} d\lambda(\xi) \\ &< \infty. \end{aligned}$$

Embedding $\mathcal{L}_{2C} \subset \mathcal{L}_C$ is dense since $B_{bs}(\Gamma_0) \subset \mathcal{L}_{2C}$. □

4.1 Description of approximation

In this section we will use the symbol K_0 to denote the restriction of K onto functions on Γ_0 .

Let $\delta \in (0; 1)$ be arbitrary and fixed. Consider for any $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ the following linear mapping on functions $F \in K_0(B_{bs}(\Gamma_0)) \subset \mathcal{F}_{cyl}(\Gamma)$

$$\begin{aligned} (P_\delta^\Lambda F)(\gamma) &= \sum_{\eta \subset \gamma} \delta^{|\eta|} (1 - \delta)^{|\gamma \setminus \eta|} (\Xi_\delta^\Lambda(\gamma))^{-1} \\ &\times \int_{\Gamma_\Lambda} (z\delta)^{|\omega|} \prod_{y \in \omega} e^{-E^\phi(y,\gamma)} F((\gamma \setminus \eta) \cup \omega) d\lambda(\omega), \quad \gamma \in \Gamma_0, \end{aligned} \tag{4.5}$$

where

$$\Xi_\delta^\Lambda(\gamma) = \int_{\Gamma_\Lambda} (z\delta)^{|\omega|} \prod_{y \in \omega} e^{-E^\phi(y,\gamma)} d\lambda(\omega). \tag{4.6}$$

Clearly, P_δ^Λ is a positive preserving mapping and

$$(P_\delta^\Lambda 1)(\gamma) = \sum_{\eta \subset \gamma} \delta^{|\eta|} (1 - \delta)^{|\gamma \setminus \eta|} = 1, \quad \gamma \in \Gamma_0.$$

Operator (4.5) is constructed as a transition operator of a Markov chain, which is a time discretization of a continuous time process with the generator (4.1) and discretization parameter $\delta \in (0; 1)$. Roughly speaking, according to the representation (4.5), the probability of transition $\gamma \rightarrow (\gamma \setminus \eta) \cup \omega$ (which describes removing of subconfiguration $\eta \subset \gamma$ and birth of a new subconfiguration $\omega \in \Gamma(\Lambda)$) after small time δ is equal to

$$(\Xi_\delta^\Lambda(\gamma))^{-1} \delta^{|\eta|} (1 - \delta)^{|\gamma \setminus \eta|} (z\delta)^{|\omega|} \prod_{y \in \omega} e^{-E^\phi(y,\gamma)}.$$

We may rewrite (4.5) in another manner.

Proposition 4.2 For any $F \in \mathcal{F}_{\text{cyl}}(\Gamma_0)$ the following equality holds:

$$(P_\delta^\Lambda F)(\gamma) = \sum_{\xi \subset \gamma} (1 - \delta)^{|\xi|} \int_{\Gamma_\Lambda} (z\delta)^{|\omega|} \prod_{y \in \omega} e^{-E^\phi(y, \gamma)} \times (K_0^{-1} F)(\xi \cup \omega) d\lambda(\omega). \tag{4.7}$$

Proof Let $G := K_0^{-1} F \in B_{\text{bs}}(\Gamma_0)$. Since Ξ_δ^Λ doesn't depend on η , for $\gamma \in \Gamma_0$ we have

$$(P_\delta^\Lambda F)(\gamma) = (\Xi_\delta^\Lambda(\gamma))^{-1} \int_{\Gamma_\Lambda} (z\delta)^{|\omega|} \prod_{y \in \omega} e^{-E^\phi(y, \gamma)} \times \sum_{\eta \subset \gamma} \delta^{|\gamma \setminus \eta|} (1 - \delta)^{|\eta|} F(\eta \cup \omega) d\lambda(\omega). \tag{4.8}$$

To rewrite (4.5), we have used also that any $\eta \subset \gamma$ corresponds to a unique $\gamma \setminus \eta \subset \gamma$. Applying the definition of K_0 to $F = K_0 G$ we obtain

$$\sum_{\eta \subset \gamma} \delta^{|\gamma \setminus \eta|} (1 - \delta)^{|\eta|} F(\eta \cup \omega) = \sum_{\eta \subset \gamma} \delta^{|\gamma \setminus \eta|} (1 - \delta)^{|\eta|} \sum_{\zeta \subset \eta} \sum_{\beta \subset \omega} G(\zeta \cup \beta) = \sum_{\zeta \subset \gamma} \sum_{\beta \subset \omega} G(\zeta \cup \beta) \sum_{\eta' \subset \gamma \setminus \zeta} \delta^{|\gamma \setminus (\eta' \cup \zeta)|} (1 - \delta)^{|\eta' \cup \zeta|}, \tag{4.9}$$

where after changing summation over $\eta \subset \gamma$ and $\zeta \subset \eta$ we have used the fact that for any configuration $\eta \subset \gamma$ which contains fixed $\zeta \subset \gamma$ there exists a unique $\eta' \subset \gamma \setminus \zeta$ such that $\eta = \eta' \cup \zeta$. But by the binomial formula

$$\sum_{\eta' \subset \gamma \setminus \zeta} \delta^{|\gamma \setminus (\eta' \cup \zeta)|} (1 - \delta)^{|\eta' \cup \zeta|} = (1 - \delta)^{|\zeta|} \sum_{\eta' \subset \gamma \setminus \zeta} \delta^{|\gamma \setminus \zeta \setminus \eta'|} (1 - \delta)^{|\eta'|} = (1 - \delta)^{|\zeta|} (\delta + 1 - \delta)^{|\gamma \setminus \zeta|} = (1 - \delta)^{|\zeta|}. \tag{4.10}$$

Combining (4.8), (4.9), (4.10), we get

$$(P_\delta^\Lambda F)(\gamma) = (\Xi_\delta^\Lambda(\gamma))^{-1} \int_{\Gamma_\Lambda} (z\delta)^{|\omega|} \prod_{y \in \omega} e^{-E^\phi(y, \gamma)} \times \sum_{\zeta \subset \gamma} \sum_{\beta \subset \omega} G(\zeta \cup \beta) (1 - \delta)^{|\zeta|} d\lambda(\omega).$$

Next, Lemma 3.4 yields

$$(P_\delta^\Lambda F)(\gamma) = (\Xi_\delta^\Lambda(\gamma))^{-1} \int_{\Gamma_\Lambda} \int_{\Gamma_\Lambda} (z\delta)^{|\omega \cup \beta|} \prod_{y \in \omega \cup \beta} e^{-E^\phi(y, \gamma)} \times \sum_{\zeta \subset \gamma} G(\zeta \cup \beta) (1 - \delta)^{|\zeta|} d\lambda(\omega) d\lambda(\beta) = \int_{\Gamma_\Lambda} (z\delta)^{|\beta|} \prod_{y \in \beta} e^{-E^\phi(y, \gamma)} \sum_{\zeta \subset \gamma} G(\zeta \cup \beta) (1 - \delta)^{|\zeta|} d\lambda(\beta),$$

which proves the statement. □

In the next proposition we describe the image of P_δ^Λ under the K_0 -transform.

Proposition 4.3 Let $\widehat{P}_\delta^\Lambda = K_0^{-1} P_\delta^\Lambda K_0$. Then for any $G \in B_{\text{bs}}(\Gamma_0)$ the following equality holds:

$$(\widehat{P}_\delta^\Lambda G)(\eta) = \sum_{\xi \subset \eta} (1 - \delta)^{|\xi|} \int_{\Gamma_\Lambda} (z\delta)^{|\omega|} G(\xi \cup \omega) \times \prod_{y \in \xi} e^{-E^\phi(y, \omega)} \prod_{y' \in \eta \setminus \xi} (e^{-E^\phi(y', \omega)} - 1) d\lambda(\omega), \quad \eta \in \Gamma_0. \tag{4.11}$$

Proof By (4.7) and the definition of K_0^{-1} , we have

$$\begin{aligned} & (\widehat{P}_\delta^\Lambda G)(\eta) \\ &= \sum_{\zeta \subset \eta} (-1)^{|\eta \setminus \zeta|} \sum_{\xi \subset \zeta} (1 - \delta)^{|\xi|} \int_{\Gamma_\Lambda} (z\delta)^{|\omega|} \prod_{y \in \omega} e^{-E^\phi(y, \zeta)} G(\xi \cup \omega) d\lambda(\omega) \\ &= \sum_{\xi \subset \eta} (1 - \delta)^{|\xi|} \sum_{\zeta \subset \eta \setminus \xi} (-1)^{|\eta \setminus \xi \setminus \zeta|} \int_{\Gamma_\Lambda} (z\delta)^{|\omega|} \prod_{y \in \omega} e^{-E^\phi(y, \zeta \cup \xi)} G(\xi \cup \omega) d\lambda(\omega). \end{aligned}$$

By the definition of the relative energy

$$\prod_{y \in \omega} e^{-E^\phi(y, \zeta \cup \xi)} = \prod_{y \in \xi} e^{-E^\phi(y, \omega)} \prod_{y' \in \zeta} e^{-E^\phi(y', \omega)}.$$

The well-known equality (see, e.g. [36])

$$\begin{aligned} \sum_{\zeta \subset \eta \setminus \xi} (-1)^{|\eta \setminus \xi \setminus \zeta|} \prod_{y' \in \zeta} e^{-E^\phi(y', \omega)} &= \left(K_0^{-1} \prod_{y' \in \cdot} e^{-E^\phi(y', \omega)} \right) (\eta \setminus \xi) \\ &= \prod_{y' \in \eta \setminus \xi} \left(e^{-E^\phi(y', \omega)} - 1 \right) \end{aligned}$$

completes the proof. □

4.2 Construction of the semigroup on \mathcal{L}_C

By analogy with (4.11), we consider the following linear mapping on measurable functions on Γ_0 :

$$\begin{aligned} (\widehat{P}_\delta G)(\eta) &:= \sum_{\xi \subset \eta} (1 - \delta)^{|\xi|} \int_{\Gamma_0} (z\delta)^{|\omega|} G(\xi \cup \omega) \\ &\quad \times \prod_{y \in \xi} e^{-E^\phi(y, \omega)} \prod_{y' \in \eta \setminus \xi} \left(e^{-E^\phi(y', \omega)} - 1 \right) d\lambda(\omega), \quad \eta \in \Gamma_0. \end{aligned} \tag{4.12}$$

Proposition 4.4 *Let*

$$ze^{CC\phi} \leq C. \tag{4.13}$$

Then \widehat{P}_δ , given by (4.12), is a well-defined linear operator in \mathcal{L}_C , such that

$$\|\widehat{P}_\delta\| \leq 1. \tag{4.14}$$

Proof Since $\phi \geq 0$ we have

$$\begin{aligned} \|\widehat{P}_\delta G\|_C &\leq \int_{\Gamma_0} \sum_{\xi \subset \eta} (1 - \delta)^{|\xi|} \int_{\Gamma_0} (z\delta)^{|\omega|} |G(\xi \cup \omega)| \\ &\quad \times \prod_{y \in \xi} e^{-E^\phi(y, \omega)} \prod_{y' \in \eta \setminus \xi} \left| e^{-E^\phi(y', \omega)} - 1 \right| d\lambda(\omega) C^{|\eta|} d\lambda(\eta) \\ &= \int_{\Gamma_0} \int_{\Gamma_0} (1 - \delta)^{|\xi|} \int_{\Gamma_0} (z\delta)^{|\omega|} |G(\xi \cup \omega)| \\ &\quad \times \prod_{y \in \xi} e^{-E^\phi(y, \omega)} \prod_{y' \in \eta} \left| e^{-E^\phi(y', \omega)} - 1 \right| d\lambda(\omega) C^{|\eta|} C^{|\xi|} d\lambda(\xi) d\lambda(\eta) \\ &= \int_{\Gamma_0} \int_{\Gamma_0} (1 - \delta)^{|\xi|} (z\delta)^{|\omega|} |G(\xi \cup \omega)| \end{aligned}$$

$$\times \prod_{y \in \xi} e^{-E^\phi(y, \omega)} \exp \left\{ C \int_{\mathbb{R}^d} \left(1 - e^{-E^\phi(y', \omega)} \right) dy' \right\} d\lambda(\omega) C^{|\xi|} d\lambda(\xi).$$

It is easy to see by the induction principle that for $\phi \geq 0, \omega \in \Gamma_0, y \notin \omega$

$$1 - e^{-E^\phi(y, \omega)} = 1 - \prod_{x \in \omega} e^{-\phi(x-y)} \leq \sum_{x \in \omega} \left(1 - e^{-\phi(x-y)} \right). \tag{4.15}$$

Then

$$\begin{aligned} \|\widehat{P}_\delta G\|_C &\leq \int_{\Gamma_0} \int_{\Gamma_0} (1 - \delta)^{|\xi|} (z\delta)^{|\omega|} |G(\xi \cup \omega)| \\ &\quad \times \exp \left\{ C \sum_{x \in \omega} \int_{\mathbb{R}^d} \left(1 - e^{-\phi(x-y)} \right) dy \right\} d\lambda(\omega) C^{|\xi|} d\lambda(\xi) \\ &= \int_{\Gamma_0} \int_{\Gamma_0} (1 - \delta)^{|\xi|} (z\delta)^{|\omega|} |G(\xi \cup \omega)| e^{CC_\phi|\omega|} C^{|\xi|} d\lambda(\omega) d\lambda(\xi) \\ &= \int_{\Gamma_0} \left[(1 - \delta) C + z\delta e^{CC_\phi} \right]^{|\omega|} |G(\omega)| d\lambda(\omega) \leq \|G\|_C. \end{aligned}$$

For the last inequality we have used that (4.13) implies $(1 - \delta) C + z\delta e^{CC_\phi} \leq C$. Note that, for λ -a.a. $\eta \in \Gamma_0$

$$(\widehat{P}_\delta G)(\eta) < \infty, \tag{4.16}$$

and the statement is proved. □

Proposition 4.5 *Let the inequality (4.13) be fulfilled and define*

$$\widehat{L}_\delta := \frac{1}{\delta} (\widehat{P}_\delta - \mathbb{1}), \quad \delta \in (0; 1),$$

where $\mathbb{1}$ is the identity operator in \mathcal{L} . Then for any $G \in \mathcal{L}_{2C}$

$$\|(\widehat{L}_\delta - \widehat{L})G\|_C \leq 3\delta \|G\|_{2C}. \tag{4.17}$$

Proof Let us denote

$$(\widehat{P}_\delta^{(0)} G)(\eta) = \sum_{\xi \subset \eta} (1 - \delta)^{|\xi|} G(\xi) 0^{|\eta \setminus \xi|} = (1 - \delta)^{|\eta|} G(\eta); \tag{4.18}$$

$$(\widehat{P}_\delta^{(1)} G)(\eta) = z\delta \sum_{\xi \subset \eta} (1 - \delta)^{|\xi|} \int_{\mathbb{R}^d} G(\xi \cup x) \tag{4.19}$$

$$\times \prod_{y \in \xi} e^{-\phi(y-x)} \prod_{y \in \eta \setminus \xi} \left(e^{-\phi(y-x)} - 1 \right) dx; \tag{4.20}$$

and

$$\widehat{P}_\delta^{(\geq 2)} = \widehat{P}_\delta - \left(\widehat{P}_\delta^{(0)} + \widehat{P}_\delta^{(1)} \right). \tag{4.21}$$

Clearly

$$\begin{aligned} \|(\widehat{L}_\delta - \widehat{L})G\|_C &= \left\| \frac{1}{\delta} (\widehat{P}_\delta G - G) - \widehat{L}G \right\|_C \\ &\leq \left\| \frac{1}{\delta} (\widehat{P}_\delta^{(0)} G - G) - L_0 G \right\|_C + \left\| \frac{1}{\delta} \widehat{P}_\delta^{(1)} G - L_1 G \right\|_C + \frac{1}{\delta} \left\| \widehat{P}_\delta^{(\geq 2)} G \right\|_C. \end{aligned} \tag{4.22}$$

Now we estimate each of the terms in (4.22) separately. By (4.3) and (4.18), we have

$$\left\| \frac{1}{\delta} (\widehat{P}_\delta^{(0)} G - G) - L_0 G \right\|_C = \int_{\Gamma_0} \left| \frac{(1 - \delta)^{|\eta|} - 1}{\delta} + |\eta| \right| |G(\eta)| C^{|\eta|} d\lambda(\eta).$$

But, for any $|\eta| \geq 2$

$$\begin{aligned} \left| \frac{(1-\delta)^{|\eta|} - 1}{\delta} + |\eta| \right| &= \left| \sum_{k=2}^{|\eta|} \binom{|\eta|}{k} (-1)^k \delta^{k-1} \right| \\ &= \delta \left| \sum_{k=2}^{|\eta|} \binom{|\eta|}{k} (-1)^k \delta^{k-2} \right| \leq \delta \sum_{k=2}^{|\eta|} \binom{|\eta|}{k} < \delta \cdot 2^{|\eta|}. \end{aligned}$$

Therefore,

$$\left\| \frac{1}{\delta} \left(\widehat{P}_\delta^{(0)} G - G \right) - L_0 G \right\|_C \leq \delta \|G\|_{2C}. \tag{4.23}$$

Next, by (4.4) and (4.20), one can write

$$\begin{aligned} \left\| \frac{1}{\delta} \widehat{P}_\delta^{(1)} G - L_1 G \right\|_C &= z \int_{\Gamma_0} \left| \sum_{\xi \subset \eta} \left((1-\delta)^{|\xi|} - 1 \right) \int_{\mathbb{R}^d} G(\xi \cup x) \prod_{y \in \xi} e^{-\phi(y-x)} \right. \\ &\quad \times \left. \prod_{y \in \eta \setminus \xi} \left(e^{-\phi(y-x)} - 1 \right) dx \right| C^{|\eta|} d\lambda(\eta) \\ &\leq z \int_{\Gamma_0} \int_{\Gamma_0} \left(1 - (1-\delta)^{|\xi|} \right) \int_{\mathbb{R}^d} |G(\xi \cup x)| \prod_{y \in \xi} e^{-\phi(y-x)} \\ &\quad \times \prod_{y \in \eta} \left(1 - e^{-\phi(y-x)} \right) dx C^{|\xi|} C^{|\eta|} d\lambda(\xi) d\lambda(\eta), \end{aligned}$$

where we have used Lemma 3.4. Note that for any $|\xi| \geq 1$

$$1 - (1-\delta)^{|\xi|} = \delta \sum_{k=0}^{|\xi|-1} (1-\delta)^k \leq \delta |\xi|$$

Then, by (4.13) and (2.12), one may estimate

$$\begin{aligned} \left\| \frac{1}{\delta} \widehat{P}_\delta^{(1)} G - L_1 G \right\|_C &\leq z \delta \int_{\Gamma_0} |\xi| \int_{\mathbb{R}^d} |G(\xi \cup x)| dx C^{|\xi|} e^{CC_\phi} d\lambda(\xi) \\ &\leq z \delta \int_{\Gamma_0} |\xi| (|\xi| - 1) |G(\xi)| C^{|\xi|-1} e^{CC_\phi} d\lambda(\xi). \end{aligned} \tag{4.24}$$

Since $n(n-1) \leq 2^n, n \geq 1$ and by (4.13), the latter expression can be bounded by

$$\delta \int_{\Gamma_0} |G(\xi)| (2C)^{|\xi|} \lambda(d\xi).$$

Finally, Lemma 3.4, (4.15) and bound $e^{-E^\phi(y,\omega)} \leq 1$, imply (we set here $\Gamma_0^{(\geq 2)} := \bigsqcup_{n \geq 2} \Gamma^{(n)}$)

$$\begin{aligned} \left\| \frac{1}{\delta} \widehat{P}_\delta^{(\geq 2)} G \right\|_C &\leq \frac{1}{\delta} \int_{\Gamma_0} \sum_{\xi \subset \eta} (1-\delta)^{|\xi|} \int_{\Gamma_0^{(\geq 2)}} (z\delta)^{|\omega|} |G(\xi \cup \omega)| \\ &\quad \times \prod_{y \in \xi} e^{-E^\phi(y,\omega)} \prod_{y \in \eta \setminus \xi} \left(1 - e^{-E^\phi(y,\omega)} \right) d\lambda(\omega) C^{|\eta|} d\lambda(\eta) \\ &\leq \delta \int_{\Gamma_0} \sum_{\xi \subset \eta} (1-\delta)^{|\xi|} \int_{\Gamma_0^{(\geq 2)}} z^{|\omega|} |G(\xi \cup \omega)| \\ &\quad \times \prod_{y \in \xi} e^{-E^\phi(y,\omega)} \prod_{y \in \eta \setminus \xi} \left(1 - e^{-E^\phi(y,\omega)} \right) d\lambda(\omega) C^{|\eta|} d\lambda(\eta) \end{aligned}$$

$$\begin{aligned}
&\leq \delta \int_{\Gamma_0} \sum_{\xi \subset \eta} (1 - \delta)^{|\xi|} \int_{\Gamma_0} z^{|\omega|} |G(\xi \cup \omega)| \\
&\quad \times \prod_{y \in \xi} e^{-E^\phi(y, \omega)} \prod_{y \in \eta \setminus \xi} (1 - e^{-E^\phi(y, \omega)}) d\lambda(\omega) C^{|\eta|} d\lambda(\eta) \\
&\leq \delta \int_{\Gamma_0} \int_{\Gamma_0} (1 - \delta)^{|\xi|} z^{|\omega|} |G(\xi \cup \omega)| \\
&\quad \times \int_{\Gamma_0} \prod_{y \in \eta} (1 - e^{-E^\phi(y, \omega)}) C^{|\eta|} d\lambda(\eta) d\lambda(\omega) C^{|\xi|} d\lambda(\xi) \\
&\leq \delta \int_{\Gamma_0} \int_{\Gamma_0} (1 - \delta)^{|\xi|} z^{|\omega|} |G(\xi \cup \omega)| e^{CC_\phi|\omega|} d\lambda(\omega) C^{|\xi|} d\lambda(\xi) \\
&\leq \delta \int_{\Gamma_0} \left[(1 - \delta)C + ze^{CC_\phi} \right]^{|\omega|} |G(\omega)| d\lambda(\omega) \\
&\leq \delta \int_{\Gamma_0} [(2 - \delta)C]^{|\omega|} |G(\omega)| d\lambda(\omega) \leq \delta \int_{\Gamma_0} |G(\omega)| (2C)^{|\omega|} d\lambda(\omega). \quad (4.25)
\end{aligned}$$

Combining inequalities (4.23)–(4.25) we obtain the assertion of the proposition. \square

We will need the following results in the sequel:

Lemma 4.6 (cf. [20, Corollary 3.8]) *Let A be a linear operator on a Banach space L with $D(A)$ dense in L , and let $\|\cdot\|$ be a norm on $D(A)$ with respect to which $D(A)$ is a Banach space. For $n \in \mathbb{N}$ let T_n be a linear $\|\cdot\|$ -contraction on L such that $T_n : D(A) \rightarrow D(A)$, and define $A_n = n(T_n - 1)$. Suppose there exist $\omega \geq 0$ and a sequence $\{\varepsilon_n\} \subset (0; +\infty)$ tending to zero such that for $n \in \mathbb{N}$*

$$\|(A_n - A)f\| \leq \varepsilon_n \|f\|, \quad f \in D(A) \quad (4.26)$$

and

$$\|T_n \upharpoonright_{D(A)}\| \leq 1 + \frac{\omega}{n}. \quad (4.27)$$

Then A is closable and the closure of A generates a strongly continuous contraction semigroup on L .

Lemma 4.7 (cf. [20, Theorem 6.5]) *Let $L, L_n, n \in \mathbb{N}$ be Banach spaces, and $p_n : L \rightarrow L_n$ be bounded linear transformation, such that $\sup_n \|p_n\| < \infty$. For any $n \in \mathbb{N}$, let T_n be a linear contraction on L_n , let $\varepsilon_n > 0$ be such that $\lim_{n \rightarrow \infty} \varepsilon_n = 0$, and put $A_n = \varepsilon_n^{-1}(T_n - \mathbb{1})$. Let $T(t)$ be a strongly continuous contraction semigroup on L with generator A and let D be a core for A . Then the following are equivalent:*

1. For each $f \in L$, $T_n^{[t/\varepsilon_n]} p_n f \rightarrow p_n T(t)f$ in L_n for all $t \geq 0$ uniformly on bounded intervals. Here and below $[\cdot]$ mean the entire part of a real number.
2. For each $f \in D$, there exists $f_n \in L_n$ for each $n \in \mathbb{N}$ such that $f_n \rightarrow p_n f$ and $A_n f_n \rightarrow p_n A f$ in L_n .

And now we are able to show the existence of the semigroup on \mathcal{L}_C .

Theorem 4.8 *Let*

$$z \leq \min\{Ce^{-CC_\phi}; 2Ce^{-2CC_\phi}\}. \quad (4.28)$$

Then $(\widehat{L}, \mathcal{L}_{2C})$ from Proposition 4.1 is a closable linear operator in \mathcal{L}_C and its closure $(\widehat{L}, D(\widehat{L}))$ generates a strongly continuous contraction semigroup \widehat{T}_t on \mathcal{L}_C .

Proof We apply Lemma 4.6 for $L = \mathcal{L}_C$, $(A, D(A)) = (\widehat{L}, \mathcal{L}_{2C})$, $\|\cdot\| := \|\cdot\|_{2C}$; $T_n = \widehat{P}_\delta$ and $A_n = n(T_n - 1) = \frac{1}{\delta}(\widehat{P}_\delta - \mathbb{1}) = \widehat{L}_\delta$, where $\delta = \frac{1}{n}$, $n \geq 2$.

Condition $ze^{CC_\phi} \leq C$, Proposition 4.4, and Proposition 4.5 provide that $T_n, n \geq 2$ are linear $\|\cdot\|_C$ -contractions and (4.26) holds with $\varepsilon_n = \frac{3}{n} = 3\delta$. On the other hand, in addition, Proposition 4.4 applied to the constant $2C$ instead of C gives (4.27) for $\omega = 0$ under condition $ze^{2CC_\phi} \leq 2C$. \square

Moreover, since we proved the existence of the semigroup \widehat{T}_t on \mathcal{L}_C one can apply contractions \widehat{P}_δ defined above by (4.12) to approximate the semigroup \widehat{T}_t .



Corollary 4.9 *Let (4.13) hold. Then for any $G \in \mathcal{L}_C$*

$$(\widehat{P}_{\frac{1}{n}})^{[nt]}G \rightarrow \widehat{T}_t G, \quad n \rightarrow \infty$$

for all $t \geq 0$ uniformly on bounded intervals.

Proof The statement is a direct consequence of Theorem 4.8, convergence (4.17), and Lemma 4.7 (if we set $L_n = L = \mathcal{L}_C, p_n = \mathbb{1}, n \in \mathbb{N}$). □

4.3 Finite-volume approximation of \widehat{T}_t

Note that \widehat{P}_δ defined by (4.12) is a formal point-wise limit of $\widehat{P}_\delta^\Lambda$ as $\Lambda \uparrow \mathbb{R}^d$. We have shown in (4.16) that this definition is correct. Corollary 4.9 claims additionally that the linear contractions \widehat{P}_δ approximate the semigroup \widehat{T}_t , when $\delta \downarrow 0$. One may also show that mappings $\widehat{P}_\delta^\Lambda$ have a similar property when $\Lambda \uparrow \mathbb{R}^d, \delta \downarrow 0$.

Let us fix a system $\{\Lambda_n\}_{n \geq 2}$, where $\Lambda_n \in \mathcal{B}_b(\mathbb{R}^d), \Lambda_n \subset \Lambda_{n+1}, \bigcup_n \Lambda_n = \mathbb{R}^d$. We set

$$T_n := \widehat{P}_{\frac{1}{n}}^{\Lambda_n}.$$

Note that any T_n is a linear mapping on $B_{bs}(\Gamma_0)$. We consider also the system of Banach spaces of measurable functions on Γ_0

$$\mathcal{L}_{C,n} := \left\{ G : \Gamma(\Lambda_n) \rightarrow \mathbb{R} \mid \|G\|_{C,n} := \int_{\Gamma(\Lambda_n)} |G(\eta)| C^{|\eta|} d\lambda(\eta) < \infty \right\}.$$

Let $p_n : \mathcal{L}_C \rightarrow \mathcal{L}_{C,n}$ be a cut-off mapping, namely for any $G \in \mathcal{L}_C$

$$(p_n G)(\eta) = \mathbb{1}_{\Gamma(\Lambda_n)}(\eta) G(\eta).$$

Then, obviously, $\|p_n G\|_{C,n} \leq \|G\|_C$. Hence, $p_n : \mathcal{L}_C \rightarrow \mathcal{L}_{C,n}$ is a linear bounded transformation with $\|p_n\| = 1$.

Proposition 4.10 *Let (4.13) hold. Then for any $G \in \mathcal{L}_C$*

$$\|(T_n)^{[nt]} p_n G - p_n \widehat{T}_t G\|_{C,n} \rightarrow 0, \quad n \rightarrow \infty$$

for all $t \geq 0$ uniformly on bounded intervals.

Proof The proof of the proposition is completed by showing that all conditions of Lemma 4.7 hold. Using completely the same arguments as in the proof of Proposition 4.4 one gets that each $T_n = \widehat{P}_{\frac{1}{n}}^{\Lambda_n}$ is a linear contraction on $\mathcal{L}_{C,n}, n \geq 2$ (note that for any $n \geq 2, (2.12)$ implies $\int_{\Lambda_n} (1 - e^{-\phi(x)}) dx \leq C_\phi < \infty$). Next, we set $A_n = n(T_n - \mathbb{1}_n)$ where $\mathbb{1}_n$ is a unit operator on $\mathcal{L}_{C,n}$ and let us expand T_n in three parts analogously to the proof of Proposition 4.5: $T_n = T_n^{(0)} + T_n^{(1)} + T_n^{(\geq 2)}$. As a result, $A_n = n(T_n^{(0)} - \mathbb{1}_n) + nT_n^{(1)} + nT_n^{(\geq 2)}$. For any $G \in \mathcal{L}_{2C}$ we set $G_n = p_n G \in \mathcal{L}_{2C,n} \subset \mathcal{L}_{C,n}$. To finish the proof we have to verify that for any $G \in \mathcal{L}_{2C}$

$$\|A_n G_n - p_n \widehat{L} G\|_{C,n} \rightarrow 0, \quad n \rightarrow \infty. \tag{4.29}$$

For any $G \in \mathcal{L}_{2C}$

$$\begin{aligned} \|A_n G_n - p_n L G\|_{C,n} &\leq \|n(T_n^{(0)} - \mathbb{1}_n)G_n - p_n L_0 G\|_{C,n} \\ &\quad + \|nT_n^{(1)}G_n - p_n L_1 G\|_{C,n} + \|nT_n^{(\geq 2)}G_n\|_{C,n}. \end{aligned} \tag{4.30}$$

Note that $p_n L_0 G = L_0 G_n$. Using the same arguments as in the proof of Proposition 4.5 we obtain

$$\|n(T_n^{(0)} - \mathbb{1}_n)G_n - p_n L_0 G\|_{C,n} + \|nT_n^{(\geq 2)}G_n\|_{C,n} \leq \frac{2}{n} \|G\|_{2C,n} \leq \frac{2}{n} \|G\|_{2C}.$$

Next,

$$\begin{aligned}
 & \|nT_n^{(1)}G_n - p_nL_1G\|_{C,n} \\
 & \leq z \int_{\Gamma_{\Lambda_n}} \sum_{\xi \subset \eta} \int_{\mathbb{R}^d} \left| \left(1 - \frac{1}{n}\right)^{|\xi|} \mathbb{1}_{\Lambda_n}(x) - 1 \right| |G(\xi \cup x)| \\
 & \quad \times \prod_{y \in \xi} e^{-\phi(y-x)} \prod_{y \in \eta \setminus \xi} \left(1 - e^{-\phi(y-x)}\right) dx C^{|\eta|} d\lambda(\eta) \\
 & \leq z \int_{\Gamma(\Lambda_n)} \int_{\Gamma(\Lambda_n)} \int_{\mathbb{R}^d} \left[1 - \left(1 - \frac{1}{n}\right)^{|\xi|} \mathbb{1}_{\Lambda_n}(x) \right] |G(\xi \cup x)| \\
 & \quad \times \prod_{y \in \eta} \left(1 - e^{-\phi(y-x)}\right) dx C^{|\eta \cup \xi|} d\lambda(\eta) d\lambda(\xi) \\
 & \leq C \int_{\Gamma(\Lambda_n)} \int_{\mathbb{R}^d} \left[1 - \left(1 - \frac{1}{n}\right)^{|\xi|} \mathbb{1}_{\Lambda_n}(x) \right] |G(\xi \cup x)| dx C^{|\xi|} d\lambda(\xi),
 \end{aligned}$$

where we have used (2.12) and (4.13). Using the same estimates as for (4.24) we may continue

$$\begin{aligned}
 & \leq C \int_{\Gamma(\Lambda_n)} \int_{\Lambda_n} \left[1 - \left(1 - \frac{1}{n}\right)^{|\xi|} \right] |G(\xi \cup x)| dx C^{|\xi|} d\lambda(\xi) \\
 & \quad + C \int_{\Gamma(\Lambda_n)} \int_{\Lambda_n^c} |G(\xi \cup x)| dx C^{|\xi|} d\lambda(\xi) \\
 & \leq \frac{1}{n} \|G\|_{2C,n} + C \int_{\Gamma_0} \int_{\Lambda_n^c} |G(\xi \cup x)| dx C^{|\xi|} d\lambda(\xi).
 \end{aligned}$$

But by the Lebesgue dominated convergence theorem,

$$\int_{\Gamma_0} \int_{\Lambda_n^c} |G(\xi \cup x)| dx C^{|\xi|} d\lambda(\xi) \rightarrow 0, \quad n \rightarrow \infty.$$

Indeed, $\mathbb{1}_{\Lambda_n^c}(x)|G(\xi \cup x)| \rightarrow 0$ point-wisely and may be estimated on $\Gamma_0 \times \mathbb{R}^d$ by $|G(\xi \cup x)|$ which is integrable:

$$C \int_{\Gamma_0} \int_{\mathbb{R}^d} |G(\xi \cup x)| dx C^{|\xi|} d\lambda(\xi) = \int_{\Gamma_0} |\xi| |G(\xi)| C^{|\xi|} d\lambda(\xi) \leq \|G\|_{2C} < \infty.$$

Therefore, by (4.30), the convergence (4.29) holds for any $G \in \mathcal{L}_{2C}$, which completes the proof. □

4.4 Evolution of correlation functions

Under condition (4.28), we proceed now to the same arguments as in Subsect. 3.4. Namely, one can construct the restriction $\widehat{T}^\circ(t)$ of the semigroup of $\widehat{T}^*(t)$ onto the Banach space $\overline{D(\widehat{L}^*)}$ (recall that the closure is in the norm of \mathcal{K}_C). Note that the domain of the dual operator to $(\widehat{L}, \mathcal{L}_{2C})$ might be bigger than the domain considered in Subsect. 3.4. Nevertheless, $\widehat{T}^\circ(t)$ will be a C_0 -semigroup on $\overline{D(\widehat{L}^*)}$ and its generator \widehat{L}° will be a part of \widehat{L}^* , namely (3.44) holds and $\widehat{L}^*k = \widehat{L}^\circ k$ for any $k \in D(\widehat{L}^\circ)$.

The next statement is a straightforward consequence of Proposition 3.12.

Proposition 4.11 *For any $\alpha \in (0; 1)$ the following inclusions hold: $\mathcal{K}_{\alpha C} \subset D(\widehat{L}^*) \subset \overline{D(\widehat{L}^*)} \subset \mathcal{K}_C$.*



Then, by Proposition 3.5, we immediately obtain that, for $k \in \mathcal{K}_{\alpha C}$,

$$\begin{aligned}
 (\widehat{L}^*k)(\eta) &= -|\eta|k(\eta) \\
 &+ z \sum_{x \in \eta} e^{-E^\phi(x, \eta \setminus x)} \int_{\Gamma_0} e_\lambda(e^{-\phi(x^-)} - 1, \xi) k((\eta \setminus x) \cup \xi) d\lambda(\xi).
 \end{aligned}
 \tag{4.31}$$

The next statement is an analog of Proposition 3.15.

Proposition 4.12 *Suppose that (4.28) is satisfied. Furthermore, we additionally assume that*

$$z < Ce^{-CC_\phi}, \text{ if } CC_\phi \leq \ln 2. \tag{4.32}$$

Then there exists $\alpha_0 = \alpha_0(z, \phi, C) \in (0; 1)$ such that for any $\alpha \in (\alpha_0; 1)$ the set $\mathcal{K}_{\alpha C}$ is the $\widehat{T}^(t)$ -invariant linear subspace of \mathcal{K}_C .*

Proof Let us consider function $f(x) := xe^{-x}$, $x \geq 0$. It has the following properties: f is increasing on $[0; 1]$ from 0 to e^{-1} and it is asymptotically decreasing on $[1; +\infty)$ from e^{-1} to 0; $f(x) < f(2x)$ for $x \in (0, \ln 2)$; $x = \ln 2$ is the only non-zero solution to $f(x) = f(2x)$.

By assumption (4.28), $zC_\phi \leq \min\{CC_\phi e^{-CC_\phi}, 2CC_\phi e^{-2CC_\phi}\}$. Therefore, if $CC_\phi e^{-CC_\phi} \neq 2CC_\phi e^{-2CC_\phi}$ then (4.28) with necessity implies

$$zC_\phi < e^{-1}. \tag{4.33}$$

This inequality remains also true if $CC_\phi = \ln 2$ because of (4.32). Under condition (4.33), the equation $f(x) = zC_\phi$ has exactly two roots, say, $0 < x_1 < 1 < x_2 < +\infty$. Then, (4.32) implies $x_1 < CC_\phi < 2CC_\phi \leq x_2$.

If $CC_\phi > 1$ then we set $\alpha_0 := \max\left\{\frac{1}{2}; \frac{1}{CC_\phi}; \frac{1}{C}\right\} < 1$. This yields $2\alpha CC_\phi > CC_\phi$ and $\alpha CC_\phi > 1 > x_1$.

If $x_1 < CC_\phi \leq 1$ then we set $\alpha_0 := \max\left\{\frac{1}{2}; \frac{x_1}{CC_\phi}; \frac{1}{C}\right\} < 1$ that gives $2\alpha CC_\phi > CC_\phi$ and $\alpha CC_\phi > x_1$.

As a result,

$$x_1 < \alpha CC_\phi < CC_\phi < 2\alpha CC_\phi < 2CC_\phi \leq x_2 \tag{4.34}$$

and $1 < \alpha C < C < 2\alpha C < 2C$. The last inequality shows that $\mathcal{L}_{2C} \subset \mathcal{L}_{2\alpha C} \subset \mathcal{L}_C \subset \mathcal{L}_{\alpha C}$. Moreover, by (4.34), we may prove that the operator $(\widehat{L}, \mathcal{L}_{2\alpha C})$ is closable in $\mathcal{L}_{\alpha C}$ and its closure is a generator of a contraction semigroup $\widehat{T}_\alpha(t)$ on $\mathcal{L}_{\alpha C}$. The proof is identical to the proofs above.

It is easy to see that $\widehat{T}_\alpha(t)G = \widehat{T}(t)G$ for any $G \in \mathcal{L}_C$. Indeed, from the construction of the semigroup $\widehat{T}(t)$ and analogous construction for the semigroup $\widehat{T}_\alpha(t)$, we have that there exists family of mappings \widehat{P}_δ , $\delta > 0$ independent of α and C , given by (4.12), such that $\widehat{P}_\delta^{\lfloor \frac{t}{\delta} \rfloor}$ for any $t \geq 0$ strongly converges to $\widehat{T}(t)$ and $\widehat{T}_\alpha(t)$ in \mathcal{L}_C and $\mathcal{L}_{\alpha C}$, correspondingly, as $\delta \rightarrow 0$. Here and below $\lfloor \cdot \rfloor$ means the entire part of a number. Then for any $G \in \mathcal{L}_C \subset \mathcal{L}_{\alpha C}$ we have that $\widehat{T}(t)G \in \mathcal{L}_C \subset \mathcal{L}_{\alpha C}$ and $\widehat{T}_\alpha(t)G \in \mathcal{L}_{\alpha C}$ and

$$\begin{aligned}
 \|\widehat{T}(t)G - \widehat{T}_\alpha(t)G\|_{\alpha C} &\leq \|\widehat{T}(t)G - \widehat{P}_\delta^{\lfloor \frac{t}{\delta} \rfloor}G\|_{\alpha C} + \|\widehat{T}_\alpha(t)G - \widehat{P}_\delta^{\lfloor \frac{t}{\delta} \rfloor}G\|_{\alpha C} \\
 &\leq \|\widehat{T}(t)G - \widehat{P}_\delta^{\lfloor \frac{t}{\delta} \rfloor}G\|_C + \|\widehat{T}_\alpha(t)G - \widehat{P}_\delta^{\lfloor \frac{t}{\delta} \rfloor}G\|_{\alpha C} \rightarrow 0,
 \end{aligned}$$

as $\delta \rightarrow 0$. Therefore, $\widehat{T}(t)G = \widehat{T}_\alpha(t)G$ in $\mathcal{L}_{\alpha C}$ (recall that $G \in \mathcal{L}_C$) that yields $\widehat{T}(t)G(\eta) = \widehat{T}_\alpha(t)G(\eta)$ for λ -a.a. $\eta \in \Gamma_0$ and, therefore, $\widehat{T}(t)G = \widehat{T}_\alpha(t)G$ in \mathcal{L}_C .

Note that for any $G \in \mathcal{L}_C \subset \mathcal{L}_{\alpha C}$ and for any $k \in \mathcal{K}_{\alpha C} \subset \mathcal{K}_C$ we have $\widehat{T}_\alpha(t)G \in \mathcal{L}_{\alpha C}$ and

$$\langle \widehat{T}_\alpha(t)G, k \rangle = \langle G, \widehat{T}_\alpha^*(t)k \rangle,$$

where, by construction, $\widehat{T}_\alpha^*(t)k \in \mathcal{K}_{\alpha C}$. But $G \in \mathcal{L}_C, k \in \mathcal{K}_C$ implies

$$\langle \widehat{T}_\alpha(t)G, k \rangle = \langle \widehat{T}(t)G, k \rangle = \langle G, \widehat{T}^*(t)k \rangle.$$

Hence, $\widehat{T}^*(t)k = \widehat{T}_\alpha^*(t)k \in \mathcal{K}_{\alpha C}, k \in \mathcal{K}_{\alpha C}$ that proves the statement. □

Remark 4.13 As a result, (4.28) implies that for any $k_0 \in \overline{D(\widehat{L}^*)}$ the Cauchy problem in \mathcal{K}_C

$$\begin{cases} \frac{\partial}{\partial t} k_t = \widehat{L}^* k_t \\ k_t|_{t=0} = k_0 \end{cases} \tag{4.35}$$

has a unique mild solution: $k_t = \widehat{T}^*(t)k_0 = \widehat{T}^\circ(t)k_0 \in \overline{D(\widehat{L}^*)}$. Moreover, $k_0 \in \mathcal{K}_{\alpha C}$ implies $k_t \in \mathcal{K}_{\alpha C}$ provided (4.32) is satisfied.

Remark 4.14 The Cauchy problem (4.35) is well-posed in $\widehat{\mathcal{K}}_C = \overline{D(\widehat{L}^*)}$, i.e. for every $k_0 \in D(\widehat{L}^\circ)$ there exists a unique solution $k_t \in \widehat{\mathcal{K}}_C$ of (4.35).

Let (4.28) and (4.32) be satisfied and let α_0 be chosen as in the proof of Proposition 4.12 and fixed. Suppose that $\alpha \in (\alpha_0; 1)$. Then, Propositions 4.11 and 4.12 imply $\overline{\mathcal{K}_{\alpha C}} \subset \overline{D(\widehat{L}^*)}$ and the Banach subspace $\overline{\mathcal{K}_{\alpha C}}$ is $\widehat{T}^*(t)$ - and, therefore, $\widehat{T}^\circ(t)$ -invariant due to the continuity of these operators.

Let now $\widehat{T}^{\circ\alpha}(t)$ be the restriction of the strongly continuous semigroup $\widehat{T}^\circ(t)$ onto the closed linear subspace $\overline{\mathcal{K}_{\alpha C}}$. By general result (see, e.g., [19]), $\widehat{T}^{\circ\alpha}(t)$ is a strongly continuous semigroup on $\overline{\mathcal{K}_{\alpha C}}$ with generator $\widehat{L}^{\circ\alpha}$ which is the restriction of the operator \widehat{L}° . Namely

$$D(\widehat{L}^{\circ\alpha}) = \left\{ k \in \overline{\mathcal{K}_{\alpha C}} \mid \widehat{L}^* k \in \overline{\mathcal{K}_{\alpha C}} \right\}, \tag{4.36}$$

and

$$\widehat{L}^{\circ\alpha} k = \widehat{L}^\circ k = \widehat{L}^* k, \quad k \in D(\widehat{L}^{\circ\alpha}) \tag{4.37}$$

Since $\widehat{T}(t)$ is a contraction semigroup on \mathcal{L}_C , then, $\widehat{T}^\circ(t)$ is also a contraction semigroup on $(\mathcal{L}_C)'$; but isomorphism (3.43) is isometrical; therefore, $\widehat{T}^*(t)$ is a contraction semigroup on \mathcal{K}_C . As a result, its restriction $\widehat{T}^{\circ\alpha}(t)$ is a contraction semigroup on $\overline{\mathcal{K}_{\alpha C}}$. Note also, that by (4.36),

$$D_{\alpha C} := \left\{ k \in \mathcal{K}_{\alpha C} \mid \widehat{L}^* k \in \overline{\mathcal{K}_{\alpha C}} \right\}$$

is a core for $\widehat{L}^{\circ\alpha}$ in $\overline{\mathcal{K}_{\alpha C}}$.

By (4.12), for any $k \in \mathcal{K}_{\alpha C}$, $G \in B_{bs}(\Gamma_0)$ we have

$$\begin{aligned} & \int_{\Gamma_0} (\widehat{P}_\delta G)(\eta) k(\eta) d\lambda(\eta) \\ &= \int_{\Gamma_0} \sum_{\xi \subset \eta} (1 - \delta)^{|\xi|} \int_{\Gamma_0} (z\delta)^{|\omega|} G(\xi \cup \omega) \prod_{y \in \xi} e^{-E^\phi(y, \omega)} \\ & \quad \times \prod_{y \in \eta \setminus \xi} \left(e^{-E^\phi(y, \omega)} - 1 \right) d\lambda(\omega) k(\eta) d\lambda(\eta) \\ &= \int_{\Gamma_0} \int_{\Gamma_0} (1 - \delta)^{|\xi|} \int_{\Gamma_0} (z\delta)^{|\omega|} G(\xi \cup \omega) \prod_{y \in \xi} e^{-E^\phi(y, \omega)} \\ & \quad \times \prod_{y \in \eta} \left(e^{-E^\phi(y, \omega)} - 1 \right) d\lambda(\omega) k(\eta \cup \xi) d\lambda(\xi) d\lambda(\eta) \\ &= \int_{\Gamma_0} \int_{\Gamma_0} \sum_{\omega \subset \xi} (1 - \delta)^{|\xi \setminus \omega|} (z\delta)^{|\omega|} G(\xi) \prod_{y \in \xi \setminus \omega} e^{-E^\phi(y, \omega)} \\ & \quad \times \prod_{y \in \eta} \left(e^{-E^\phi(y, \omega)} - 1 \right) k(\eta \cup \xi \setminus \omega) d\lambda(\xi) d\lambda(\eta); \end{aligned}$$

therefore,

$$(\widehat{P}_\delta^* k)(\eta) = \sum_{\omega \subset \eta} (1 - \delta)^{|\eta \setminus \omega|} (z\delta)^{|\omega|} \prod_{y \in \eta \setminus \omega} e^{-E^\phi(y, \omega)}$$



$$\times \int_{\Gamma_0} \prod_{y \in \xi} \left(e^{-E^\phi(y, \omega)} - 1 \right) k(\xi \cup \eta \setminus \omega) d\lambda(\xi). \tag{4.38}$$

Proposition 4.15 *Suppose that (4.28) and (4.32) are fulfilled. Then, for any $k \in D_{\alpha C}$ and $\alpha \in (\alpha_0, 1)$, where α_0 is chosen as in the proof of Proposition 4.12,*

$$\lim_{\delta \rightarrow 0} \left\| \frac{1}{\delta} (\widehat{P}_\delta^* - \mathbb{1})k - \widehat{L}^{\odot \alpha} k \right\|_{\mathcal{K}_C} = 0. \tag{4.39}$$

Proof Let us recall (4.31) and define

$$\begin{aligned} (\widehat{P}_\delta^{*,(0)} k)(\eta) &= (1 - \delta)^{|\eta|} k(\eta); \\ (\widehat{P}_\delta^{*,(1)} k)(\eta) &= z\delta \sum_{x \in \eta} (1 - \delta)^{|\eta|-1} e_\lambda \left(e^{-\phi(x-\cdot)}, \eta \setminus x \right) \\ &\quad \times \int_{\Gamma_0} e_\lambda \left(e^{-\phi(x-\cdot)} - 1, \xi \right) k(\xi \cup \eta \setminus x) d\lambda(\xi) \end{aligned}$$

and $\widehat{P}_\delta^{*,(\geq 2)} = \widehat{P}_\delta^* - \widehat{P}_\delta^{*,(0)} - \widehat{P}_\delta^{*,(1)}$.

We will use the following elementary inequality, for any $n \in \mathbb{N} \cup \{0\}$, $\delta \in (0, 1)$:

$$0 \leq n - \frac{1 - (1 - \delta)^n}{\delta} \leq \delta \frac{n(n - 1)}{2}.$$

Then, for any $k \in \mathcal{K}_{\alpha C}$ and λ -a.a. $\eta \in \Gamma_0$, $\eta \neq \emptyset$

$$\begin{aligned} & C^{-|\eta|} \left| \frac{1}{\delta} (\widehat{P}_{\delta, \varepsilon}^{*,(0)} - \mathbb{1})k(\eta) + |\eta|k(\eta) \right| \\ & \leq \|k\|_{\mathcal{K}_{\alpha C}} \alpha^{|\eta|} \left| |\eta| - \frac{1 - (1 - \delta)^{|\eta|}}{\delta} \right| \leq \frac{\delta}{2} \|k\|_{\mathcal{K}_{\alpha C}} \alpha^{|\eta|} |\eta| (|\eta| - 1) \end{aligned} \tag{4.40}$$

and the function $\alpha^x x(x - 1)$ is bounded for $x \geq 1$, $\alpha \in (0, 1)$. Next, for any $k \in \mathcal{K}_{\alpha C}$ and λ -a.a. $\eta \in \Gamma_0$, $\eta \neq \emptyset$

$$\begin{aligned} & C^{-|\eta|} \left| \frac{1}{\delta} \widehat{P}_\delta^{*,(1)} k(\eta) - z \sum_{x \in \eta} \int_{\Gamma_0} e_\lambda \left(e^{-\phi(x-\cdot)}, \eta \setminus x \right) \right. \\ & \quad \left. \times e_\lambda \left(e^{-\phi(x-\cdot)} - 1, \xi \right) k(\xi \cup \eta \setminus x) d\lambda(\xi) \right| \\ & \leq \|k\|_{\mathcal{K}_{\alpha C}} \frac{z}{\alpha C} \alpha^{|\eta|} \sum_{x \in \eta} (1 - (1 - \delta)^{|\eta|-1}) \int_{\Gamma_0} e_\lambda \left(\alpha C (e^{-\phi(x-\cdot)} - 1), \xi \right) d\lambda(\xi) \\ & \leq \|k\|_{\mathcal{K}_{\alpha C}} \frac{z}{\alpha C} \alpha^{|\eta|} \sum_{x \in \eta} (1 - (1 - \delta)^{|\eta|-1}) \exp \{ \alpha C C_\phi \} \\ & \leq \|k\|_{\mathcal{K}_{\alpha C}} \frac{z}{\alpha C} \alpha^{|\eta|} \delta |\eta| (|\eta| - 1) \exp \{ \alpha C C_\phi \}. \end{aligned} \tag{4.41}$$

which is small in δ uniformly by $|\eta|$. Now, using inequality

$$1 - e^{-E^\phi(y, \omega)} = 1 - \prod_{x \in \omega} e^{-\phi(x-y)} \leq \sum_{x \in \omega} \left(1 - e^{-\phi(x-y)} \right),$$

we obtain

$$\begin{aligned} & \frac{1}{\delta} C^{-|\eta|} \sum_{\substack{\omega \subset \eta \\ |\omega| \geq 2}} (1 - \delta)^{|\eta \setminus \omega|} (z\delta)^{|\omega|} e_\lambda \left(e^{-E^\phi(\cdot, \omega)}, \eta \setminus \omega \right) \\ & \quad \times \int_{\Gamma_0} e_\lambda \left(\left| e^{-E^\phi(\cdot, \omega)} - 1 \right|, \xi \right) |k(\xi \cup \eta \setminus \omega)| d\lambda(\xi) \end{aligned}$$

$$= \|k\|_{\mathcal{K}_{\alpha C}} \alpha^{|\eta|} \frac{1}{\delta} \sum_{\substack{\omega \subset \eta \\ |\omega| \geq 2}} (1 - \delta)^{|\eta \setminus \omega|} \left(\frac{z\delta}{\alpha C} \exp\{\alpha C C_\phi\} \right)^{|\omega|};$$

recall that $\alpha > \alpha_0$, therefore, $z \exp\{\alpha C C_\phi\} \leq \alpha C$, and one may continue

$$\begin{aligned} &\leq \|k\|_{\mathcal{K}_{\alpha C}} \alpha^{|\eta|} \frac{1}{\delta} \sum_{\substack{\omega \subset \eta \\ |\omega| \geq 2}} (1 - \delta)^{|\eta \setminus \omega|} \delta^{|\omega|} \\ &= \|k\|_{\mathcal{K}_{\alpha C}} \delta \alpha^{|\eta|} \sum_{k=2}^{|\eta|} \frac{|\eta|!}{k! (|\eta| - k)!} (1 - \delta)^{|\eta| - k} \delta^{k-2} \\ &= \|k\|_{\mathcal{K}_{\alpha C}} \delta \alpha^{|\eta|} \sum_{k=0}^{|\eta|-2} \frac{|\eta|!}{(k+2)! (|\eta| - k - 2)!} (1 - \delta)^{|\eta| - k - 2} \delta^k \\ &= \|k\|_{\mathcal{K}_{\alpha C}} \delta \alpha^{|\eta|} |\eta| (|\eta| - 1) \sum_{k=0}^{|\eta|-2} \frac{(|\eta| - 2)!}{(k+2)! (|\eta| - k - 2)!} (1 - \delta)^{|\eta| - 2 - k} \delta^k \\ &\leq \|k\|_{\mathcal{K}_{\alpha C}} \delta \alpha^{|\eta|} |\eta| (|\eta| - 1) \sum_{k=0}^{|\eta|-2} \frac{(|\eta| - 2)!}{k! (|\eta| - k - 2)!} (1 - \delta)^{|\eta| - 2 - k} \delta^k \\ &= \|k\|_{\mathcal{K}_{\alpha C}} \delta \alpha^{|\eta|} |\eta| (|\eta| - 1). \end{aligned} \tag{4.42}$$

Combining inequalities (4.40)–(4.42) we obtain (4.39). □

As a result, we obtain an approximation for the semigroup.

Theorem 4.16 *Let α_0 be chosen as in the proof of the Proposition 4.12 and be fixed. Let $\alpha \in (\alpha_0; 1)$ and $k \in \overline{\mathcal{K}_{\alpha C}}$ be given. Then*

$$(\widehat{P}_\delta^*)^{[t/\delta]} k \rightarrow \widehat{T}^{\odot \alpha}(t)k, \quad \delta \rightarrow 0$$

in the space $\overline{\mathcal{K}_{\alpha C}}$ with norm $\|\cdot\|_{\mathcal{K}_C}$ for all $t \geq 0$ uniformly on bounded intervals.

Proof We may apply Proposition 4.15 to use Lemma 4.7 in the case $L_n = L = \overline{\mathcal{L}_{\alpha C}}$, $p_n = \mathbb{1}$, $f_n = f = k$, $\varepsilon_n = \delta \rightarrow 0$, $n \in \mathbb{N}$. □

4.5 Positive definiteness

We consider a small modification of the notion of positive definite functions considered in Proposition 2.12. Namely, we denote by $L_{\text{ls}}^0(\Gamma_0)$ the set of all measurable functions on Γ_0 which have a local support, i.e. $G \in L_{\text{ls}}^0(\Gamma_0)$ if there exists $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ such that $G \upharpoonright_{\Gamma_0 \setminus \Gamma(\Lambda)} = 0$. We will say that a measurable function $k : \Gamma_0 \rightarrow \mathbb{R}$ is a positive defined function if, for any $G \in L_{\text{ls}}^0(\Gamma_0)$ such that $KG \geq 0$ and $G \in \mathcal{L}_C$ for some $C > 1$ the inequality (2.30) holds.

For a given $C > 1$, we set $\mathcal{L}_C^{\text{ls}} = L_{\text{ls}}^0(\Gamma_0) \cap \mathcal{L}_C$. Since $B_{\text{bs}}(\Gamma_0) \subset \mathcal{L}_C^{\text{ls}}$, for any $C > 1$, Proposition 2.12 (see also the second part of Remark 2.13) implies that if k is a positive definite function as above then there exists a unique measure $\mu \in \mathcal{M}_{\text{fm}}^1(\Gamma)$ such that $k = k_\mu$ be its correlation function in the sense of (2.24). Our aim is to show that the evolution $k \mapsto \widehat{T}^{\odot}(t)k$ preserves this property of the positive definiteness.

Theorem 4.17 *Let (4.28) hold and $k \in \overline{D(\widehat{L}^*)} \subset \mathcal{K}_C$ be a positive definite function. Then $k_t := \widehat{T}^{\odot}(t)k \in \overline{D(\widehat{L}^*)} \subset \mathcal{K}_C$ will be a positive definite function for any $t \geq 0$.*

Proof Let $C > 0$ be arbitrary and fixed. For any $G \in \mathcal{L}_C^{\text{ls}}$ we have

$$\int_{\Gamma_0} G(\eta) k_t(\eta) d\lambda(\eta) = \int_{\Gamma_0} (\widehat{T}(t)G)(\eta) k(\eta) d\lambda(\eta). \tag{4.43}$$

By Proposition 4.10, under condition (4.28), we obtain that

$$\lim_{n \rightarrow 0} \int_{\Gamma(\Lambda_n)} |T_n^{[nt]} \mathbb{1}_{\Gamma(\Lambda_n)} G(\eta) - \mathbb{1}_{\Gamma(\Lambda_n)}(\eta) (\widehat{T}(t)G)(\eta)| C^{|\eta|} d\lambda(\eta) = 0,$$

where for $n \geq 2$

$$T_n = \widehat{P}_{\frac{1}{n}}^{\Lambda_n}$$

and $\Lambda_n \nearrow \mathbb{R}^d$. Note that, by the dominated convergence theorem,

$$\begin{aligned} \int_{\Gamma_0} (\widehat{T}(t)G)(\eta) k(\eta) d\lambda(\eta) &= \lim_{n \rightarrow \infty} \int_{\Gamma_0} \mathbb{1}_{\Gamma(\Lambda_n)}(\eta) (\widehat{T}(t)G)(\eta) k(\eta) d\lambda(\eta) \\ &= \lim_{n \rightarrow \infty} \int_{\Gamma(\Lambda_n)} (\widehat{T}(t)G)(\eta) k(\eta) d\lambda(\eta). \end{aligned}$$

Next,

$$\begin{aligned} &\left| \int_{\Gamma(\Lambda_n)} (\widehat{T}(t)G)(\eta) k(\eta) d\lambda(\eta) - \int_{\Gamma(\Lambda_n)} T_n^{[nt]} \mathbb{1}_{\Gamma(\Lambda_n)} G(\eta) k(\eta) d\lambda(\eta) \right| \\ &\leq \int_{\Gamma(\Lambda_n)} |T_n^{[nt]} \mathbb{1}_{\Gamma(\Lambda_n)} G(\eta) - \mathbb{1}_{\Gamma(\Lambda_n)}(\eta) (\widehat{T}(t)G)(\eta)| k(\eta) d\lambda(\eta) \\ &\leq \|k\|_{\mathcal{K}_C} \int_{\Gamma(\Lambda_n)} |T_n^{[nt]} \mathbb{1}_{\Gamma(\Lambda_n)} G(\eta) - \mathbb{1}_{\Gamma(\Lambda_n)}(\eta) (\widehat{T}(t)G)(\eta)| C^{|\eta|} d\lambda(\eta) \rightarrow 0, \quad n \rightarrow \infty. \end{aligned}$$

Therefore,

$$\int_{\Gamma_0} (\widehat{T}(t)G)(\eta) k(\eta) d\lambda(\eta) = \lim_{n \rightarrow \infty} \int_{\Gamma(\Lambda_n)} T_n^{[nt]} \mathbb{1}_{\Gamma(\Lambda_n)} G(\eta) k(\eta) d\lambda(\eta). \tag{4.44}$$

Our aim is to show that for any $G \in \mathcal{L}_C^{\text{ls}}$ the inequality $KG \geq 0$ implies

$$\int_{\Gamma_0} G(\eta) k_t(\eta) d\lambda(\eta) \geq 0.$$

By (4.43) and (4.44), it is enough to show that for any $m \in \mathbb{N}$ and for any $G \in \mathcal{L}_C^{\text{ls}}$ such that $KG \geq 0$ the following inequality holds:

$$\int_{\Gamma_0} \mathbb{1}_{\Gamma(\Lambda_n)} T_n^m \mathbb{1}_{\Gamma(\Lambda_n)} G(\eta) k(\eta) d\lambda(\eta) \geq 0, \quad m \in \mathbb{N}_0. \tag{4.45}$$

The inequality (4.45) is fulfilled if only

$$K \mathbb{1}_{\Gamma(\Lambda_n)} T_n^m G_n \geq 0, \tag{4.46}$$

where $G_n := \mathbb{1}_{\Gamma(\Lambda_n)} G$. Note that

$$\begin{aligned} (K \mathbb{1}_{\Gamma(\Lambda_n)} T_n^m G_n)(\gamma) &= \sum_{\eta \in \gamma} \mathbb{1}_{\Gamma(\Lambda_n)}(\eta) (T_n^m G_n)(\eta) \\ &= \sum_{\eta \subset \gamma \wedge \Lambda_n} (T_n^m G_n)(\eta) = (K T_n^m G_n)(\gamma_{\Lambda_n}) \end{aligned} \tag{4.47}$$

for any $m \in \mathbb{N}_0$. In particular,

$$(K G_n)(\gamma) = (K \mathbb{1}_{\Gamma(\Lambda_n)} G)(\gamma) = (KG)(\gamma_{\Lambda_n}) \geq 0. \tag{4.48}$$

Let us now consider any $\tilde{G} \in \mathcal{L}_C^{\text{ls}}$ [stress that \tilde{G} is not necessary equal to 0 outside of $\Gamma(\Lambda_n)$] and suppose that $(K\tilde{G})(\gamma) \geq 0$ for any $\gamma \in \Gamma(\Lambda_n)$. Then

$$\begin{aligned} (KT_n\tilde{G})(\gamma_{\Lambda_n}) &= (K\widehat{P}_{\frac{1}{n}}^{\Lambda_n}\tilde{G})(\gamma_{\Lambda_n}) = (P_{\frac{1}{n}}^{\Lambda_n}K\tilde{G})(\gamma_{\Lambda_n}) \\ &= \left(\Xi_{\frac{1}{n}}^{\Lambda_n}(\gamma_{\Lambda_n})\right)^{-1} \sum_{\eta \subset \gamma_{\Lambda_n}} \left(\frac{1}{n}\right)^{|\eta|} \left(1 - \frac{1}{n}\right)^{|\gamma \setminus \eta|} \\ &\quad \times \int_{\Gamma(\Lambda_n)} \left(\frac{z}{n}\right)^{|\omega|} \prod_{y \in \omega} e^{-E^\phi(y,\gamma)} (K\tilde{G})((\gamma_{\Lambda_n} \setminus \eta) \cup \omega) d\lambda(\omega) \geq 0. \end{aligned} \tag{4.49}$$

By (4.48), setting $\tilde{G} = G_n \in \mathcal{L}_C^{\text{ls}}$ we obtain, because of (4.49), $KT_nG_n \geq 0$. Next, setting $\tilde{G} = T_nG_n \in \mathcal{L}_C^{\text{ls}}$ we obtain, by (4.49), $KT_n^2G_n \geq 0$. Then, using an induction mechanism, we obtain that

$$(KT_n^m G_n)(\gamma_{\Lambda_n}) \geq 0, \quad m \in \mathbb{N}_0,$$

that, by (4.46) and (4.47), yields (4.45). This completes the proof. □

4.6 Ergodicity

Let $k \in \overline{\mathcal{K}_{\alpha C}}$ be such that $k(\emptyset) = 0$; then, by (4.38), $(\widehat{P}_\delta^* k)(\emptyset) = 0$. Class of all such functions we denote by \mathcal{K}_α^0 .

Proposition 4.18 *Assume that there exists $v \in (0; 1)$ such that*

$$z \leq \min\left\{vCe^{-CC\phi}; 2Ce^{-2CC\phi}\right\}. \tag{4.50}$$

Let, additionally, $\alpha \in (\alpha_0; 1)$, where α_0 is chosen as in the proof of the Proposition 4.12. Then for any $\delta \in (0; 1)$ the following estimate holds:

$$\left\|\widehat{P}_\delta^* \upharpoonright_{\mathcal{K}_\alpha^0}\right\| \leq 1 - (1 - v)\delta. \tag{4.51}$$

Proof It is easily seen that for any $k \in \mathcal{K}_\alpha^0$ the following inequality holds:

$$|k(\eta)| \leq 1_{|\eta|>0} \|k\|_{\mathcal{K}_C} C^{|\eta|}, \quad \lambda\text{-a.a. } \eta \in \Gamma_0.$$

Then, using (4.38), we have

$$\begin{aligned} &C^{-|\eta|} |(\widehat{P}_\delta^* k)(\eta)| \\ &\leq C^{-|\eta|} \sum_{\omega \subset \eta} (1 - \delta)^{|\eta \setminus \omega|} (z\delta)^{|\omega|} \int_{\Gamma_0} \prod_{y \in \xi} (1 - e^{-E^\phi(y,\omega)}) |k(\xi \cup \eta \setminus \omega)| d\lambda(\xi) \\ &\leq \|k\|_{\mathcal{K}_C} \sum_{\omega \subset \eta} (1 - \delta)^{|\eta \setminus \omega|} \left(\frac{z\delta}{C}\right)^{|\omega|} \int_{\Gamma_0} \prod_{y \in \xi} (1 - e^{-E^\phi(y,\omega)}) C^{|\xi|} \mathbb{1}_{|\xi|+|\eta \setminus \omega|>0} d\lambda(\xi) \\ &= \|k\|_{\mathcal{K}_C} \sum_{\omega \subsetneq \eta} (1 - \delta)^{|\eta \setminus \omega|} \left(\frac{z\delta}{C}\right)^{|\omega|} \int_{\Gamma_0} \prod_{y \in \xi} (1 - e^{-E^\phi(y,\omega)}) C^{|\xi|} d\lambda(\xi) \\ &\quad + \|k\|_{\mathcal{K}_C} \left(\frac{z\delta}{C}\right)^{|\eta|} \int_{\Gamma_0} \prod_{y \in \xi} (1 - e^{-E^\phi(y,\omega)}) C^{|\xi|} \mathbb{1}_{|\xi|>0} d\lambda(\xi) \\ &= \|k\|_{\mathcal{K}_C} \sum_{\omega \subsetneq \eta} (1 - \delta)^{|\eta \setminus \omega|} \left(\frac{z\delta}{C}\right)^{|\omega|} \int_{\Gamma_0} \prod_{y \in \xi} (1 - e^{-E^\phi(y,\omega)}) C^{|\xi|} d\lambda(\xi) \end{aligned}$$

$$\begin{aligned}
 & + \|k\|_{\mathcal{K}_C} \left(\frac{z\delta}{C}\right)^{|\eta|} \int_{\Gamma_0} \prod_{y \in \xi} (1 - e^{-E^\phi(y,\omega)}) C^{|\xi|} d\lambda(\xi) - \|k\|_{\mathcal{K}_C} \left(\frac{z\delta}{C}\right)^{|\eta|} \\
 & = \|k\|_{\mathcal{K}_C} \sum_{\omega \subset \eta} (1 - \delta)^{|\eta \setminus \omega|} \left(\frac{z\delta}{C}\right)^{|\omega|} \int_{\Gamma_0} \prod_{y \in \xi} (1 - e^{-E^\phi(y,\omega)}) C^{|\xi|} d\lambda(\xi) \\
 & \quad - \|k\|_{\mathcal{K}_C} \left(\frac{z\delta}{C}\right)^{|\eta|} \\
 & = \|k\|_{\mathcal{K}_C} \sum_{\omega \subset \eta} (1 - \delta)^{|\eta \setminus \omega|} \left(\frac{z\delta}{C}\right)^{|\omega|} \exp \left\{ C \int_{\mathbb{R}^d} (1 - e^{-E^\phi(y,\omega)}) dy \right\} \\
 & \quad - \|k\|_{\mathcal{K}_C} \left(\frac{z\delta}{C}\right)^{|\eta|} \\
 & \leq \|k\|_{\mathcal{K}_C} \sum_{\omega \subset \eta} (1 - \delta)^{|\eta \setminus \omega|} \left(\frac{z\delta}{C}\right)^{|\omega|} \exp \{CC_\beta |\omega|\} - \|k\|_{\mathcal{K}_C} \left(\frac{z\delta}{C}\right)^{|\eta|} \\
 & \leq \|k\|_{\mathcal{K}_C} \sum_{\omega \subset \eta} (1 - \delta)^{|\eta \setminus \omega|} (\nu\delta)^{|\omega|} - \|k\|_{\mathcal{K}_C} \left(\frac{z\delta}{C}\right)^{|\eta|} \\
 & = \|k\|_{\mathcal{K}_C} \left((1 - (1 - \nu)\delta)^{|\eta|} - \left(\frac{z\delta}{C}\right)^{|\eta|} \right) \\
 & = \|k\|_{\mathcal{K}_C} \left(1 - (1 - \nu)\delta - \frac{z\delta}{C} \right)^{|\eta|-1} \sum_{j=0}^{|\eta|-1} (1 - (1 - \nu)\delta)^{|\eta|-1-j} \left(\frac{z\delta}{C}\right)^j \\
 & \leq \|k\|_{\mathcal{K}_C} \left(1 - (1 - \nu)\delta - \frac{z\delta}{C} \right)^{|\eta|-1} \sum_{j=0}^{|\eta|-1} \left(\frac{z\delta}{C}\right)^j \\
 & = \|k\|_{\mathcal{K}_C} \left(1 - (1 - \nu)\delta - \frac{z\delta}{C} \right) \frac{1 - \left(\frac{z\delta}{C}\right)^{|\eta|}}{1 - \frac{z\delta}{C}} \\
 & \leq \|k\|_{\mathcal{K}_C} \left(1 - (1 - \nu)\delta - \frac{z\delta}{C} \right) \frac{1}{1 - \frac{z\delta}{C}} \\
 & = \|k\|_{\mathcal{K}_C} \left(1 - \frac{(1 - \nu)\delta}{1 - \frac{z\delta}{C}} \right) \leq \|k\|_{\mathcal{K}_C} (1 - (1 - \nu)\delta),
 \end{aligned}$$

where we have used that, clearly, $z < \nu C < C$. The statement is proved. □

Remark 4.19 Condition (4.50) is equivalent to (4.28) and (4.32).

As it was mentioned in Example 3.18, under condition (cf. (4.33))

$$zC_\phi < (2e)^{-1}, \tag{4.52}$$

there exists (see, e.g., [35] for details) a Gibbs measure μ on $(\Gamma, \mathcal{B}(\Gamma))$ corresponding to the potential $\phi \geq 0$ and activity parameter z . We denote the corresponding correlation function by k_μ . The measure μ is reversible (symmetrizing) for the operator defined by (4.1) (see, e.g., [35, 54]). Therefore, for any $F \in KB_{bs}(\Gamma_0)$

$$\int_{\Gamma} LF(\gamma) d\mu(\gamma) = 0. \tag{4.53}$$

Theorem 4.20 *Let (4.52) and (4.50) hold and let $\alpha \in (\alpha_0; 1)$, where α_0 is chosen as in the proof of Proposition 4.12. Let $k_0 \in \mathcal{K}_{\alpha C}$, $k_t = \widehat{T}^{\odot \alpha}(t)k_0$. Then for any $t \geq 0$*

$$\|k_t - k_\mu\|_{\mathcal{K}_C} \leq e^{-(1-\nu)t} \|k_0 - k_\mu\|_{\mathcal{K}_C}. \tag{4.54}$$

Proof First of all, let us note that for any $\alpha \in (\alpha_0; 1)$ the inequality (4.34) implies $z \leq \alpha C \exp\{-\alpha C C_\phi\}$. Hence $k_\mu(\eta) \leq (\alpha C)^{|\eta|}$, $\eta \in \Gamma_0$. Therefore, $k_\mu \in \mathcal{K}_{\alpha C} \subset \overline{\mathcal{K}_{\alpha C}} \cap D(\widehat{L}^*)$. By (4.53), for any $G \in B_{bs}(\Gamma_0)$ we have $\langle \widehat{L}G, k_\mu \rangle = 0$. It means that $\widehat{L}^*k_\mu = 0$. Therefore, $\widehat{L}^{\circ\alpha}k_\mu = 0$. As a result, $\widehat{T}^{\circ\alpha}(t)k_\mu = k_\mu$. Let $r_0 = k_0 - k_\mu \in \overline{\mathcal{K}_{\alpha C}}$. Then $r_0 \in \mathcal{K}_a^0$ and

$$\begin{aligned} \|k_t - k_\mu\|_{\mathcal{K}_C} &= \|\widehat{T}^{\circ\alpha}(t)r_0\|_{\mathcal{K}_C} \\ &\leq \|(\widehat{P}_\delta^*)^{[\frac{t}{\delta}]}r_0\|_{\mathcal{K}_C} + \|\widehat{T}^{\circ\alpha}(t)r_0 - (\widehat{P}_\delta^*)^{[\frac{t}{\delta}]}r_0\|_{\mathcal{K}_C} \\ &\leq \|\widehat{P}_\delta^* \upharpoonright_{\mathcal{K}_\alpha^0}\|^{[\frac{t}{\delta}]} \cdot \|r_0\|_{\mathcal{K}_C} + \|\widehat{T}^{\circ\alpha}(t)r_0 - (\widehat{P}_\delta^*)^{[\frac{t}{\delta}]}r_0\|_{\mathcal{K}_C} \\ &\leq (1 - (1 - \nu)\delta)^{\frac{t}{\delta}-1} \|r_0\|_{\mathcal{K}_C} + \|\widehat{T}^{\circ\alpha}(t)r_0 - (\widehat{P}_\delta^*)^{[\frac{t}{\delta}]}r_0\|_{\mathcal{K}_C}, \end{aligned}$$

since $0 < 1 - (1 - \nu)\delta < 1$ and $\frac{t}{\delta} < [\frac{t}{\delta}] + 1$. Taking the limit as $\delta \downarrow 0$ in the right hand side of this inequality we obtain (4.54). \square

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Albeverio, S.; Kondratiev, Y.; Röckner, M.: Analysis and geometry on configuration spaces. *J. Funct. Anal.* **154**(2), 444–500 (1998)
- Albeverio, S.; Kondratiev, Y.; Röckner, M.: Analysis and geometry on configuration spaces: the Gibbsian case. *J. Funct. Anal.* **157**(1), 242–291 (1998)
- Arendt, W.; Grabosch, A.; Greiner, G.; Groh, U.; Lotz, H.P.; Moustakas, U.; Nagel, R.; Neubrander, F.; Schlotterbeck, U.: One-parameter semigroups of positive operators. *Lecture Notes in Mathematics*, vol. 1184. Springer, Berlin (1986)
- Berezansky, Y.; Kondratiev, Y.; Kuna, T.; Lytvynov, E.: On a spectral representation for correlation measures in configuration space analysis. *Methods Funct. Anal. Topol.* **5**(4), 87–100 (1999)
- Berns, C.; Kondratiev, Y.; Kozitsky, Y.; Kutoviy, O.: Kawasaki dynamics in continuum: micro- and mesoscopic descriptions. *J. Dyn. Diff. Eqn.* **25**(4), 1027–1056 (2013)
- Bertini, L.; Cancrini, N.; Cesi, F.: The spectral gap for a Glauber-type dynamics in a continuous gas. *Ann. Inst. H. Poincaré Probab. Stat.* **38**(1), 91–108 (2002)
- Bogoliubov, N.N.: Problems of a dynamical theory in statistical physics. In: *Studies in Statistical Mechanics*, vol. I, pp. 1–118. North-Holland, Amsterdam (1962)
- Bolker, B.; Pacala, S.W.: Using moment equations to understand stochastically driven spatial pattern formation in ecological systems. *Theor. Popul. Biol.* **52**(3), 179–197 (1997)
- Bolker, B.; Pacala, S.W.: Spatial moment equations for plant competitions: understanding spatial strategies and the advantages of short dispersal. *Am. Nat.* **153**, 575–602 (1999)
- Borodin, A.; Olshanski, G.: Point processes and the infinite symmetric group. *Math. Res. Lett.* **5**(6), 799–816 (1998)
- Borodin, A.; Olshanski, G.: Distributions on partitions, point processes, and the hypergeometric kernel. *Commun. Math. Phys.* **211**(2), 335–358 (2000)
- Borodin, A.; Olshanski, G.: Harmonic analysis on the infinite-dimensional unitary group and determinantal point processes. *Ann. Math. (2)*, **161**(3), 1319–1422 (2005)
- Borodin, A.; Olshanski, G.: Representation theory and random point processes. In: *European Congress of Mathematics*, pp. 73–94. European Mathematical Society, Zürich (2005)
- Borodin, A.; Olshanski, G.; Strahov, E.: Giambelli compatible point processes. *Adv. Appl. Math.* **37**(2), 209–248 (2006)
- Campbell, N.R.: The study of discontinuous problem. *Proc. Camb. Philos. Soc.* **15**, 117–136 (1909)
- Campbell, N.R.: Discontinuities in light emission. *Proc. Camb. Philos. Soc.* **15**, 310–328 (1910)
- Dieckmann, U.; Law, R.: Relaxation projections and the method of moments. In: *The Geometry of Ecological Interactions*, pp. 412–455. Cambridge University Press, Cambridge (2000)
- Dobrushin, R.L.; Sinai, Y.G.; Sukhov, Y.M.: Dynamical systems of statistical mechanics. In: Sinai, Y.G. (ed.) *Ergodic Theory with Applications to Dynamical Systems and Statistical Mechanics*, *Encyclopaedia of Mathematical Sciences*, vol. II. Springer, Heidelberg (1989)
- Engel, K.-J.; Nagel, R.: *One-parameter Semigroups for Linear Evolution Equations*, *Graduate Texts in Mathematics*, vol. 194. Springer, New York (2000)
- Ethier, S.N.; Kurtz, T.G.: *Markov Processes: Characterization and Convergence*. Wiley Series in Probability and Mathematical Statistics. Wiley, New York (1986)
- Filonenko, D.; Finkelshtein, D.; Kondratiev, Y.: On two-component contact model in continuum with one independent component. *Methods Funct. Anal. Topol.* **14**(3), 209–228 (2008)



22. Finkelshtein, D.: Measures on two-component configuration spaces. *Condens. Matter Phys.* **12**(1), 5–18 (2009)
23. Finkelshtein, D.: Functional evolutions for homogeneous stationary death-immigration spatial dynamics. *Methods Funct. Anal. Topol.* **17**(4), 300–318 (2011)
24. Finkelshtein, D.; Kondratiev, Y.: Measures on configuration spaces defined by relative energies. *Methods Funct. Anal. Topol.* **11**(2), 126–155 (2005)
25. Finkelshtein, D.; Kondratiev, Y.; Kozitsky, Y.: Glauber dynamics in continuum: a constructive approach to evolution of states. *Discret. Contin. Dyn. Syst. Ser. A.* **33**(4), 1431–1450 (2013)
26. Finkelshtein, D.; Kondratiev, Y.; Kutoviy, O.: Individual based model with competition in spatial ecology. *SIAM J. Math. Anal.* **41**(1), 297–317 (2009)
27. Finkelshtein, D.; Kondratiev, Y.; Kutoviy, O.: Vlasov scaling for the Glauber dynamics in continuum. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **14**(4), 537–569 (2011)
28. Finkelshtein, D.; Kondratiev, Y.; Kutoviy, O.: Correlation functions evolution for the Glauber dynamics in continuum. *Semigroup Forum.* **85**, 289–306 (2012)
29. Finkelshtein, D.; Kondratiev, Y.; Kutoviy, O.: Semigroup approach to birth-and-death stochastic dynamics in continuum. *J. Funct. Anal.* **262**(3), 1274–1308 (2012)
30. Finkelshtein, D.; Kondratiev, Y.; Kutoviy, O.: Establishment and fecundity in spatial ecological models: statistical approach and kinetic equations. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **16**(2), 1350014 (2013)
31. Finkelshtein, D.; Kondratiev, Y.; Kutoviy, O.: An operator approach to Vlasov scaling for some models of spatial ecology. *Methods Funct. Anal. Topol.* **19**(2), 108–126 (2013)
32. Finkelshtein, D.; Kondratiev, Y.; Kutoviy, O.; Lytvynov, E.: Binary jumps in continuum. I. Equilibrium processes and their scaling limits. *J. Math. Phys.* **52**(063304), 1–25 (2011)
33. Finkelshtein, D.; Kondratiev, Y.; Kutoviy, O.; Lytvynov, E.: Binary jumps in continuum. II. Non-equilibrium process and a Vlasov-type scaling limit. *J. Math. Phys.* **52**(113301), 1–27 (2011)
34. Finkelshtein, D.; Kondratiev, Y.; Kutoviy, O.; Zhizhina, E.: An approximative approach for construction of the Glauber dynamics in continuum. *Math. Nachr.* **285**(2–3), 223–235 (2012)
35. Finkelshtein, D.; Kondratiev, Y.; Lytvynov, E.: Equilibrium Glauber dynamics of continuous particle systems as a scaling limit of Kawasaki dynamics. *Random Oper. Stoch. Equ.* **15**(2), 105–126 (2007)
36. Finkelshtein, D.; Kondratiev, Y.; Oliveira, M. J.: Markov evolutions and hierarchical equations in the continuum. I. One-component systems. *J. Evol. Equ.* **9**(2), 197–233 (2009)
37. Finkelshtein, D.; Kondratiev, Y.; Oliveira, M.J.: Glauber dynamics in the continuum via generating functionals evolution. *Complex Anal. Oper. Theory* **6**(4), 923–945 (2012)
38. Finkelshtein, D.; Kondratiev, Y.; Oliveira, M.J.: Kawasaki dynamics in the continuum via generating functionals evolution. *Methods Funct. Anal. Topol.* **18**(1), 55–67 (2012)
39. Finkelshtein, D.; Kondratiev, Y.; Oliveira, M.J.: Markov evolutions and hierarchical equations in the continuum. II: Multi-component systems. *Rep. Math. Phys.* **71**(1), 123–148 (2013)
40. Fisher, M.E.; Ruelle, D.: The stability of many-particle systems. *J. Math. Phys.* **7**, 260–270 (1966)
41. Fournier, N.; Meleard, S.: A microscopic probabilistic description of a locally regulated population and macroscopic approximations. *Ann. Appl. Probab.* **14**(4), 1880–1919 (2004)
42. Garcia, N.L.: Birth and death processes as projections of higher dimensional poisson processes. *Adv. Appl. Probab.* **27**, 911–930 (1995)
43. Garcia, N.L.; Kurtz, T.G.: Spatial birth and death processes as solutions of stochastic equations. *ALEA Lat. Am. J. Probab. Math. Stat.* **1**, 281–303 (electronic) (2006)
44. Garcia, N.L.; Kurtz, T.G.: Spatial point processes and the projection method. *Prog. Probab.* **60**, 271–298 (2008)
45. Georgii, H.-O.: Canonical and grand canonical Gibbs states for continuum systems. *Commun. Math. Phys.* **48**(1), 31–51 (1976)
46. Holley, R.A.; Stroock, D.W.: Nearest neighbor birth and death processes on the real line. *Acta Math.* **140**(1–2), 103–154 (1978)
47. Kendall, D.G.: *Stochastic Geometry*, chapter Foundations of a theory of random sets, pp. 322–376. Wiley, New York (1974)
48. Kondratiev, Y.; Kuna, T.: Harmonic analysis on configuration space. I. General theory. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **5**(2), 201–233 (2002)
49. Kondratiev, Y.; Kuna, T.: Correlation functionals for Gibbs measures and Ruelle bounds. *Methods Funct. Anal. Topol.* **9**(1), 9–58 (2003)
50. Kondratiev, Y.; Kutoviy, O.: On the metrical properties of the configuration space. *Math. Nachr.* **279**(7), 774–783 (2006)
51. Kondratiev, Y.; Kutoviy, O.; Minlos, R.: On non-equilibrium stochastic dynamics for interacting particle systems in continuum. *J. Funct. Anal.* **255**(1), 200–227 (2008)
52. Kondratiev, Y.; Kutoviy, O.; Pirogov, S.: Correlation functions and invariant measures in continuous contact model. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **11**(2), 231–258 (2008)
53. Kondratiev, Y.; Kutoviy, O.; Zhizhina, E.: Nonequilibrium Glauber-type dynamics in continuum. *J. Math. Phys.* **47**(11), 113501 (2006)
54. Kondratiev, Y.; Lytvynov, E.: Glauber dynamics of continuous particle systems. *Ann. Inst. H. Poincaré Probab. Stat.* **41**(4), 685–702 (2005)
55. Kondratiev, Y.; Lytvynov, E.; Röckner, M.: Equilibrium Kawasaki dynamics of continuous particle systems. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **10**(2), 185–209 (2007)
56. Kondratiev, Y.; Minlos, R.; Zhizhina, E.: One-particle subspace of the Glauber dynamics generator for continuous particle systems. *Rev. Math. Phys.* **16**(9), 1073–1114 (2004)
57. Kuna, T.: *Studies in configuration space analysis and applications*. Bonner Mathematische Schriften [Bonn Mathematical Publications], 324. Universität Bonn Mathematisches Institut, Bonn, p. ii+187. Dissertation, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn (1999)



58. Lenard, A.: Correlation functions and the uniqueness of the state in classical statistical mechanics. *Commun. Math. Phys.* **30**, 35–44 (1973)
59. Lenard, A.: States of classical statistical mechanical systems of infinitely many particles. I. *Arch. Ration. Mech. Anal.* **59**(3), 219–239 (1975)
60. Lenard, A.: States of classical statistical mechanical systems of infinitely many particles. II. Characterization of correlation measures. *Arch. Ration. Mech. Anal.* **59**(3), 241–256 (1975)
61. Levin, S.A.: Complex adaptive systems: exploring the known, the unknown and the unknowable. *Bull. AMS*, **40**(1), 3–19 (2002)
62. Liggett, T.M.: Interacting particle systems, *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 276. Springer, New York (1985)
63. Liggett, T.M.: Stochastic interacting systems: contact, voter and exclusion processes, *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 324. Springer, Berlin (1999)
64. Lotz, H.P.: Uniform convergence of operators on L^∞ and similar spaces. *Math. Z.* **190**(2), 207–220 (1985)
65. Mecke, J.: Eine charakteristische Eigenschaft der doppelt stochastischen Poissonschen Prozesse. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, **11**, 74–81 (1968)
66. Murrell, U.; David, J.; Dieckmann; Law, R.: On moment closures for population dynamics in continuous space. *J. Theor. Biol.* **229**(3), 421–432 (2004)
67. Nguyen, X.-X.; Zessin, H.: Integral and differential characterizations of the Gibbs process. *Math. Nachr.* **88**, 105–115 (1979)
68. Ovaskainen, O.; Finkelshtein, D.; Kutoviy, O.; Cornell, S.; Bolker, B.; Kondratiev, Y.: A mathematical framework for the analysis of spatial-temporal point processes. *Theor. Ecol.* **7**(1), 101–113 (2014)
69. Ovsjannikov, L.V.: Singular operator in the scale of Banach spaces. *Dokl. Akad. Nauk SSSR*, **163**, 819–822 (1965)
70. Pazy, A.: *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Applied Mathematical Sciences, vol. 44. Springer, New York (1983)
71. Penrose, M.D.: Existence and spatial limit theorems for lattice and continuum particle systems. *Probab. Surv.* **5**, 1–36 (2008)
72. Phillips, R.S.: The adjoint semi-group. *Pac. J. Math.* **5**, 269–283 (1955)
73. Preston, C.: Spatial birth-and-death processes. *Bull. Inst. Int. Stat.* **46**(2), 371–391, 405–408 (1975)
74. Rényi, A.: Remarks on the Poisson process. *Studia Sci. Math. Hung.* **2**, 119–123 (1970)
75. Ruelle, D.: *Statistical mechanics: Rigorous results*, p. xi+219. W.A. Benjamin, Inc., New York (1969)
76. Ruelle, D.: Superstable interactions in classical statistical mechanics. *Commun. Math. Phys.* **18**, 127–159 (1970)
77. Safonov, M.V.: The abstract Cauchy-Kovalevskaya theorem in a weighted Banach space. *Commun. Pure Appl. Math.* **48**(6), 629–637 (1995)
78. Trèves, F.: Ovcyannikov theorem and hyperdifferential operators. *Notas de Matemática*, No. 46. Instituto de Matemática Pura e Aplicada, Conselho Nacional de Pesquisas, Rio de Janeiro, p. iii+238 (1968)
79. van Neerven, J.: *The Adjoint of a Semigroup of Linear Operators*, Lecture Notes in Mathematics, vol. 1529. Springer, Berlin (1992)

