



MEETING ABSTRACT

Open Access

PW01-026 – Validation of pediatric diagnostic criteria in FMF

E Demirkaya¹, S Ozen², C Saglam^{1*}, T Turker³, A Duzova⁴, P Woo⁵, I Konè-Paut⁶, M Doglio⁷, G Amaryan⁸, J Frenkel⁹, Y Uziel¹⁰, A Insalaco¹¹, L Cantarini¹², M Hofer¹³, S Boiu¹⁴, C Modesto¹⁵, A Bryant⁵, D Rigante¹⁶, E Papadopoulou-Alataki¹⁷, S Guillaume-Czitrom¹⁸, N Ruperto⁷, M Gattorno⁷, PRINTO and Eurofever Project

From 7th Congress of International Society of Systemic Auto-Inflammatory Diseases (ISSAID) Lausanne, Switzerland. 22-26 May 2013

Introduction

The diagnosis of FMF is made clinically and may be confirmed by identifying mutations in the *MEFV* gene. The most commonly used diagnostic criteria for FMF are those of Tel Hashomer, which have been established in the Jewish adult population. Recently, a Turkish group (Yalcinkaya-Ozen's diagnostic criteria) proposed new criteria for diagnosis of FMF in children.

Objectives

We analyzed the validity and reliability of Yalcinkaya-Ozen's diagnostic criteria for childhood FMF in a large international registry.

Methods

The study group consisted of 339 FMF patients diagnosed according to Tel Hashomer criteria. A control group of 377 patients were diagnosed other periodic fever syndromes including MKD, TRAPS, CAPS and PFAPA syndromes. Both groups were evaluated according to the Tel Hashomer criteria and the new set of diagnostic criteria proposed to use in childhood FMF. The diagnostic performance of both criteria was assessed by multiple logistic regression analysis.

Results

The sensitivity and specificity of Tel Hashomer criteria in our study were 35.1% and 97.7 %, respectively. The presence of two or more of these new five criteria diagnosed FMF with a high and sufficient sensitivity of 87.4 % and the NPV of 74.8%. When we used at least three Yalcinkaya-Ozen's criteria, the discrimination of the

diseases other than FMF reached the highest specificity of 88.2% and the PPV of 82.9% however the sensitivity was compromised. In case of all the new set of criteria were met, the sensitivity and specificity were 99.6% and 5.6%, respectively with a PPV of 94.1% and an NPV of 49.2%. Our study showed that ethnicity had no impact on the validation.

Conclusion

Tel Hashomer diagnostic criteria was found to have high specificity, whereas Yalcinkaya-Ozen's criteria has a higher sensitivity for the diagnosis of FMF with the combination of at least any two out of these five criteria are met. The small number of patients with amyloidosis or erysipelas like erythema and the response to colchicine therapy constituted the drawbacks in assessing the patients with Tel Hashomer criteria.

Disclosure of interest

None declared.

Authors' details

¹FMF Arthritis Vasculitis and Orphan Disease Research Center (FAVOR), Gulhane Military Medical Faculty, Ankara, Turkey. ²Pediatric Rheumatology, Hacettepe University, School of Medicine, Ankara, Turkey. ³Epidemiology, Gulhane Military Medical Faculty, Ankara, Turkey. ⁴Rheumatology, Istanbul University, Istanbul, Turkey. ⁵Pediatric Rheumatology, UCL, London, UK. ⁶Pediatric Rheumatology, University of Paris SUD, Paris, France. ⁷Pediatric Rheumatology, Ospedale Gaslini, Genoa, Italy. ⁸National Pediatric Familial Mediterranean Fever Centre, Institute of Child and Adolescent Health, Yerevan, Armenia. ⁹Pediatrics, University Medical Center Utrecht, Utrecht, the Netherlands. ¹⁰Pediatrics, Meir Medical Centre, Kfar Saba, Israel. ¹¹Reumatologia, Ospedale Pediatrico Bambin Gesù, Rome, Italy. ¹²Rheumatology, Policlinico le Scotte, University of Siena, Siena, Italy. ¹³Pediatric Rheumatology, Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland. ¹⁴Pediatric Rheumatology, Université Paris-Descartes, Paris, France. ¹⁵Reumatologia, Hospital Valle de Hebron, Barcelona, Spain. ¹⁶Pediatrics, Università Cattolica Sacro Cuore, Rome, Italy. ¹⁷Fourth Department of Pediatrics, Aristotle University of Thessaloniki Papageorgiou

¹FMF Arthritis Vasculitis and Orphan Disease Research Center (FAVOR), Gulhane Military Medical Faculty, Ankara, Turkey
Full list of author information is available at the end of the article

Hospital, Thessaloniki, Greece. ¹⁸Pediatric Rheumatology, Bicêtre-Hôpitaux
Universitaires Paris-Sud, Bicetre, France.

Published: 8 November 2013

doi:10.1186/1546-0096-11-S1-A79

Cite this article as: Demirkaya *et al.*: PW01-026 – Validation of pediatric
diagnostic criteria in FMF. *Pediatric Rheumatology* 2013 11(Suppl 1):A79.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

