

Research article

Open Access

Transcriptomic signature of Bexarotene (Rexinoid LGD1069) on mammary gland from three transgenic mouse mammary cancer models

Martin C Abba^{1,4}, Yuhui Hu¹, Carla C Levy¹, Sally Gaddis¹, Frances S Kittrell², Yun Zhang², Jamal Hill², Reid P Bissonnette³, Daniel Medina², Powel H Brown² and C Marcelo Aldaz*¹

Address: ¹Department of Carcinogenesis, The University of Texas M.D. Anderson Cancer Center, Science Park-Research Division, Smithville, 78957, TX, USA, ²Breast Center, Departments of Medicine and Molecular and Cellular Biology, Baylor College of Medicine, Houston, 77030, TX, USA, ³Department of Molecular Oncology, Ligand Pharmaceuticals Inc., San Diego, California, 92121, USA and ⁴CINIBA, Facultad de Ciencias Médicas, Universidad Nacional de La Plata, La Plata, Argentina

Email: Martin C Abba - mabba777@hotmail.com; Yuhui Hu - maaldaz@mdanderson.org; Carla C Levy - maaldaz@mdanderson.org; Sally Gaddis - salgaddis@mdanderson.org; Frances S Kittrell - dmedina@bcm.tmc.edu; Yun Zhang - yunz@bcm.edu; Jamal Hill - jhill@bcm.edu; Reid P Bissonnette - rbissonnette@ligand.com; Daniel Medina - dmedina@bcm.tmc.edu; Powel H Brown - pbrown@bcm.tmc.edu; C Marcelo Aldaz* - maaldaz@mdanderson.org

* Corresponding author

Published: 11 September 2008

Received: 22 May 2008

BMC Medical Genomics 2008, 1:40 doi:10.1186/1755-8794-1-40

Accepted: 11 September 2008

This article is available from: <http://www.biomedcentral.com/1755-8794/1/40>

© 2008 Abba et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The rexinoid bexarotene (LGD1069, Targretin) is a highly selective retinoid × receptor (RXR) agonist that inhibits the growth of pre-malignant and malignant breast cells. Bexarotene was shown to suppress the development of breast cancer in transgenic mice models without side effects. The chemopreventive effects of bexarotene are due to transcriptional modulation of cell proliferation, differentiation and apoptosis. Our goal in the present study was to obtain a profile of the genes modulated by bexarotene on mammary gland from three transgenic mouse mammary cancer models in an effort to elucidate its molecular mechanism of action and for the identification of biomarkers of effectiveness.

Methods: Serial analysis of gene expression (SAGE) was employed to profile the transcriptome of p53-null, MMTV-ErbB2, and C3(1)-SV40 mammary cells obtained from mice treated with bexarotene and their corresponding controls.

Results: This resulted in a dataset of approximately 360,000 transcript tags representing over 20,000 mRNAs from a total of 6 different SAGE libraries. Analysis of gene expression changes induced by bexarotene in mammary gland revealed that 89 genes were dysregulated among the three transgenic mouse mammary models. From these, 9 genes were common to the three models studied.

Conclusion: Analysis of the indicated core of transcripts and protein-protein interactions of this commonly modulated genes indicate two functional modules significantly affected by rexinoid bexarotene related to protein biosynthesis and bioenergetics signatures, in addition to the targeting of cancer-causing genes related with cell proliferation, differentiation and apoptosis.

Background

The American Cancer Society estimates that 212,920 new cases of invasive breast cancer and 40,970 deaths were expected to occur in the United States in 2006 [1]. Approximately two-thirds of all breast cancers are ER α (+) at the time of diagnosis and expression of this receptor is determinant of a tumor phenotype that is associated with hormone-responsiveness. Patients with tumors that express ER α have a longer disease-free interval and overall survival than patients with tumors that lack ER α expression [2]. Despite the effectiveness of anti-estrogen selective ER modulators (tamoxifen and raloxifene) for ER α (+) breast cancer treatment, there is a clear need to develop agents for the prevention and treatment of ER α (-) breast cancer.

Genetically engineered mouse mammary cancer models are defined by a known genetic background and develop tumors after a predictable time course [3]. Importantly, mammary tumors arising in transgenic mice are generally ER α (-) providing a useful system for testing chemopreventive agents against hormonally non-responsive tumors.

Retinoids are biologically active derivatives of vitamin A that play essential roles in embryonic or adult cell behavior modulating cell proliferation, differentiation and apoptosis. Signal transduction is mediated by two classes of nuclear receptors retinoid-dependent transcriptional activators: the retinoic acid receptor (RAR α , β , γ) and the retinoid \times receptor (RXR α , β , γ). These ligand-dependent transcription factors bind to response elements (RAREs or RXREs) in the promoter region of modulated genes [4]. The RXR protein can also dimerize with other nuclear hormone receptors such as vitamin D receptor, thyroid hormone receptors, PPAR (α , γ) and orphan receptors conferring retinoids responsiveness to additional subset of target genes [5].

We previously analyzed the chemopreventive effectiveness of a highly selective RXR agonist, the retinoid bexarotene (LGD1069) in three different transgenic mouse mammary models [6,7]. These studies showed a significant decrease in mammary tumorigenicity when MMTV-ErbB2, p53-null and C3(1)-SV40 tag mammary gland recipient virgin mice were treated with bexarotene (100 mg/kg dose). Although, bexarotene is more effective against c-erbB2 induced mammary tumors than against p53-null or SV40Tag mammary tumors; this data demonstrated that bexarotene is effective against the early stages of premalignant development independently of the genetic model assessed. More importantly, if specific gene expression signatures modulated by bexarotene across mammary cancer models could be identified, they might point to core transcriptional program/s on which attention should be focused.

In an effort to elucidate the molecular mechanism of action of chemopreventive retinoid bexarotene and to identify potential biomarkers of significance, here we report a comparative transcriptome profiling of three mouse mammary cancer models by *Serial Analysis of Gene Expression* (SAGE). We focused our analysis on untreated mammary gland and on retinoid bexarotene treated mammary gland at time periods prior to the histopathologic identification of premalignant progression. These studies identified a series of retinoid-regulated genes and molecular pathways that may be critical for the cancer preventive activity of bexarotene.

Methods

Retinoid LGD1069 and transgenic mouse mammary models

The RXR-selective retinoid used in this study bexarotene (LGD1069, Targretin) was obtained from Ligand Pharmaceutical, Inc (San Diego, CA).

Female MMTV-erbB2 mice [8], (obtained from The Jackson Lab., Bar Harbor ME) and C3(1)/SV40 T-antigen strain mice [9] (obtained from The National Cancer Institute, Frederick, MD) were housed in the institutional animal facilities. Balb/c p53-null mammary epithelium transplanted into the cleared mammary fat pads of three-week old mice p53 wt Balb/c mice [10] were initiated and maintained at BMC. Each group included age-matched untreated controls and bexarotene-treated mice. All mice were treated 6 days/week during 2 months starting at 8 weeks of age with bexarotene suspended in purified sesame oil (Croda, Inc., Mill Hall, PA). The retinoid was administered by gastric gavage using a 20-gauge gavage needle in a volume of 0.1-ml containing vehicle 100 mg/kg of bexarotene. Virgin animals were used to avoid confounding effects of hormonal surges during pregnancy. All animal research was conducted in AAALAC accredited facilities, following international guidelines and all research was approved by the corresponding Institutional bioethics committees.

SAGE methodology

The six mouse SAGE libraries were generated following standard procedures as described previously [11]. Briefly total RNA was extracted from frozen samples using TRIzol (Invitrogen, Carlsbad, CA, USA). SAGE library construction was performed with the I-SAGE kit (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol and introducing only minor modifications. The anchoring enzyme was *Nla*III and the tagging enzyme used was *Bsm*FI. Concatemered ditags were cloned into pZERO-1 and sequenced with an ABI 3700 DNA Analyzer (Applied Biosystems, Foster City, CA, USA). To decrease the chances potential artifacts due to sample heterogeneity, each control or bexarotene treatment SAGE library represents a pool of three mammary epithelial samples

from three age-matched separate mice. For the studies on the p53-null mammary cancer model we used mammary epithelial enriched preparations as previously described [12], for the MMTV-erbB2 and C3(1)/SV40 T-antigen models we used total mammary gland preparations. SAGE libraries were generated at an approximate resolution of 60,000 SAGE tags per library.

SAGE data processing and statistical analysis

SAGE tag extraction from sequencing files was performed by using the SAGE2000 software version 4.0 (a kind gift of Dr. Kenneth Kinzler, John Hopkins University, Baltimore, MD). SAGE data management, tag to gene matching, as well as additional gene annotations and links to publicly available resources such as Gene Ontology (GO), UniGene, and Entrez gene ID, were performed using a suite of web-based SAGE library tools developed by us. In our analyses we only considered tags with single tag-to gene reliable matches. To compare the control (vehicle) vs. bexarotene treatment SAGE libraries in each transgenic mice model, we utilized the Audic and Claverie's significance test [13]. Statistical analysis and scatter plot visualization of SAGE libraries were done with the Discovery Space 4 software (Genome Science Centre, BC Cancer Agency, Canada, Vancouver) <http://www.bcgsc.ca/platform/bioinfo/software/ds>.

Bexarotene molecular signature determination

The main strategy of this analysis was to identify commonly deregulated genes by bexarotene treatment among the different mammary cancer models tested (Figure 1). Differentially expressed genes were compiled into one Excel spreadsheet pivot Table for comparison of overlapping data between p53-null, MMTV-erbB2 and C3(1)/SV40 T-antigen transgenic mouse mammary models. Any combination of two lists was compared for matching gene-identity. The number and identity of genes commonly affected in two models (*e.g.* MMTV-erbB2 vs. p53-null) was determined. We used the normal approximation to the binomial distribution as previously described [14] to calculate whether the number of matching genes derived from each pairwise comparison was of statistical significance ($p < 0.05$). To enable illustration of the commonly deregulated genes between mammary cancer models, we used the TIGR MultiExperiment Viewer (MeV 3.0) software. This tool was used for average clustering of SAGE based on the fold change of tag counts for each transcript comparing bexarotene treatment to control (vehicle) in each transgenic mice mammary model. For automated functional annotation and classification of genes of interest based on Gene Ontology (GO) terms, we used the EASE [15] available at the *Database for Annotation, Visualization and Integrated Discovery (DAVID)* [16]. All of the raw SAGE data reported as additional files in this article are publicly available and also can be viewed at




Figure 1
Candidate genes and pathways modulated in normal mammary epithelium by rexinoid bexarotene in three different transgenic mice mammary cell models were identified through a three-stage process:
A. Identification of differentially expressed genes in mammary gland as a result of treatment with bexarotene comparing with vehicle control, in each of the mammary cancer models
B. Inter-model comparison for the identification of overlapping gene expression profiles.
C. Identification of associated functional modules and pathways affected by bexarotene treatment.

http://sciencepark.mdanderson.org/labs/ggeg/SAGE_Proj_14.htm.

In order to identify the molecular pathways that are mainly affected by the rexinoid bexarotene, we look for protein/gene interaction networks in the common core of modulated genes. The protein-protein interaction network associating genes of the three transgenic mouse mammary models was generated using the database STRING ('Search Tool for the Retrieval of Interacting Genes/Proteins') <http://string.embl.de/> [17]. The database STRING aims to collect, predict and unify most type of protein-protein associations, including direct and indirect associations. STRING runs a set of prediction algorithms, and transfers known interactions from model organisms to other species based on predicted orthology of the respective proteins [18]. In order to identify each gene in the database, we used both mouse gene name and Entrez gene ID in the 'protein-mode' application. The analysis input options were 'co-occurrence', 'co-expression', 'experiments', 'databases', and 'text mining' data at high

confidence level of predicted human orthology groups. Pathways are discriminated by different colors based on up-modulated (red node) or down-modulated (green node) transcripts in order to indicate protein-protein networks modulated by bexarotene across mammary cancer models.

Results and discussion

RXR-selective rexinoids inhibit the proliferation of normal, pre-malignant and malignant breast cells suppressing mammary tumor development in MMTV-erbB2, p53-Null, and C3(1)/SV40 T-antigen transgenic mice models [[6,7] and Medina et al., unpublished]. The chemopreventive effects of bexarotene are likely due to transcriptional modulation of genes related to repression of cell proliferation and stimulation of apoptosis and cell differentiation [19].

In order to identify rexinoid-regulated biomarkers, we generated six mouse SAGE libraries corresponding to mammary gland samples from control and bexarotene treatment from three transgenic mouse mammary cancer models: p53-Null [10], MMTV-erbB2 [8] and C3(1)/SV40 T-antigen [9]. This resulted in the sequencing of 360,000 tags (60,000 tags per library), thus monitoring the behavior of more than 20,000 transcript tags. Our statistical analyses revealed 236 transcripts differentially regulated by bexarotene treatment in mammary epithelium from p53-null background, 283 transcripts in mammary gland from the MMTV-erbB2 model, and 290 transcripts in the C3(1)/SV40 T-antigen transgenic mice mammary model (Figure 2A; see Additional file 1). Table 1 shown the most highly bexarotene deregulated transcripts from each transgenic mice mammary cancer model (Fold change \geq 7; $p < 0.01$).

In order to identify co-occurring differentially expressed genes among the three transgenic mice analyses, we performed an inter-model comparison between the above-described SAGE datasets (Figure 1). Among the three mice mammary models, a total of 711 transcripts were identified as deregulated by the rexinoid bexarotene treatment. Eighty-nine genes were identified in more than one mammary cancer model (Figure 3A; see Additional file 2). Interestingly, nine of these 89 genes were deregulated by bexarotene in mammary gland tissue from all three transgenic models: Muc15 (Mucin 15), Cdo1 (Cystein dioxygenase 1), Rps8 (Ribosomal protein S28), Rps27 (Ribosomal protein S27), Rps24 (Ribosomal protein S24), Hspa5 (Heat shock 70 kD protein 5), Csrp1 (Cysteine and glycine-rich protein 1), Npm1 (Nucleophosmin 1), and Cycs (Cytochrome c somatic). Gene Ontology annotation of the 89 deregulated genes that were common in any two models showed that approximately 18% of the transcripts are involved in tricarboxylic

acid cycle/oxidative phosphorylation, 14% are related to signal transduction/transcriptional regulation, 14% are related to protein metabolism and 12% are related to cell proliferation/differentiation and apoptosis.

A probabilistic analysis showed that 56 genes were co-deregulated in MMTV-erbB2 and C3(1)/SV40 T-antigen mice models, representing a non-random significant number of overlapping genes based on normal approximation to the binomial distribution ($p < 0.001$) (Figure 3B). Thirty-five genes were identified as co-deregulated in MMTV-erbB2 and p53-null mice models ($p < 0.001$). The set of 16 genes overlapping between p53-null and C3(1)/SV40 T-antigen were not statistical significant, i.e. the overlapping could be simply by chance ($p > 0.05$) (Figure 3B). In other words, it appears that a better correlation was observed between MMTV-ErbB2 with the other two models, than between p53-null and C3(1)-SV40 tag transgenic mouse mammary gland models. These data suggest that mammary tumors derived from different primary oncogenic pathways could respond differently to the same chemoprevention agent. In addition, these results indicates that transcripts modulated by bexarotene in the MMTV-ErbB2 mammary gland share almost all the common features among the transgenic mouse models analyzed. As mentioned above, we have previously shown that bexarotene suppresses mammary tumor development in the MMTV-ErbB2, p53-null and C3(1)-SV40 tag transgenic mouse mammary gland models [6,7]. Interestingly, the specific response of these three transgenic mouse mammary models to bexarotene treatment varies with the genetic background assessed. For instance, the bexarotene treatment is much more effective against MMTV-ErbB2 induced mammary tumors than against C3(1)-SV40 or p53-null mammary tumors [Medina et al., unpublished]. In the MMTV-ErbB2 mammary gland, bexarotene reduced tumor incidence by 75% and lengthened median tumor latency from 234 days to over 420 days [7]. However, in the p53-null and C3(1)-SV40 mammary gland where p53 or p53/Rb activities are affected respectively, bexarotene treatment showed modest chemoprevention activity. Both these molecules exert primary functions downstream of the CDKs, loci of targets activity. In this sense, human breast cancer is a complex disease caused by dysregulation of many different oncogenes, tumor suppressor genes and growth factor pathways. The MMTV-ErbB2, p53-null and C3(1)-SV40 tag mouse mammary gland cancer models are valuable tools for the elucidation of the mechanisms of mammary tumorigenesis [3]. However, it is important to recognize that no one model represents the heterogeneity of human breast cancer.

We present in Figure 4 a protein-protein interaction network associating the common core of non-random bexar-

Table 1: Most highly deregulated transcripts in mammary gland induced by bexarotene treatment on each transgenic mice mammary cancer model (Fold change ≥ 7 ; $p < 0.01$).

Tag	Gene	Description	Entrez Gene	Fold Change*
p53-null				
GTTTGCTGTA	<i>Serpinb6a</i>	Serine (or cysteine) peptidase inhibitor	20719	17.0
AGTCTCGAGG	<i>Slc1a5</i>	Solute carrier family 1	20514	12.0
GGTTTGGGGG	<i>Jup</i>	Junction plakoglobin	16480	11.0
TGCGTGCTGG	<i>Timp2</i>	Tissue inhibitor of metalloproteinase 2	21858	11.0
TTGAAATTAC	<i>BC061494</i>	CDNA sequence	381832	11.0
GATTTCTTTG	<i>Gpc3</i>	Glypican 3	14734	10.0
TAACCAAAAA	<i>Itgb4</i>	Integrin beta 4	192897	10.0
CCCAGTCCCT	<i>Ltbp4</i>	Latent transforming growth factor bindin. prot. 4	108075	8.0
GACTCTATAT	<i>Csn2</i>	Casein beta	12991	-15.0
CAATAAAACA	<i>Sar1b</i>	SAR1 gene homolog B (S. Cerevisiae)	66397	-11.0
GCAGCGATTC	<i>Nme2</i>	Expressed in non-metastatic cells 2	18103	-10.0
TGTTCTATGG	<i>Laptm5</i>	Lysosomal-associated protein transmembrane 5	16792	-9.0
GTGTTTTGCT	<i>AI451557</i>	Expressed sequence	102084	-9.0
CTAGGTGGTG	<i>Glycam1</i>	Glycosylation dependent cell adhesion molecule 1	14663	-8.8
TAAAGTCAAT	<i>Muc15</i>	Mucin 15	269328	-8.0
TCAGAGTGAG	<i>Igh-6</i>	Immunoglobulin heavy chain 6	16019	-7.5
MMTV-erbB2				
AGACCCTGTC	<i>Pnpla3</i>	Patatin-like phospholipase domain containing 3	116939	44.0
TATGAGATAG	<i>Timm9</i>	Translocase of inner mitochondrial membrane 9	30056	15.0
AGCCCTCGGA	<i>Acads</i>	Acyl-Coenzyme A dehydrogenase	11409	12.0
ACCGGGCTGG	<i>Elovl6</i>	ELOVL family member 6	170439	12.0
TGACAGAAGA	<i>Tnnc2</i>	Troponin C2	21925	10.0
TCTCTCAGTC	<i>Anxa5</i>	Annexin A5	11747	9.0
CACAGAACCA	<i>0610031J06</i>	RIKEN cDNA 0610031J06 gene	56700	7.0
CCTGCAGCAG	<i>2900073H19</i>	RIKEN cDNA 2900073H19 gene	68205	7.0
GCCACTTAAG	<i>Cd79a</i>	CD79A antigen (immunoglobulin-associated alpha)	12518	-15.0
AGCCATCATA	<i>2610042014</i>	RIKEN cDNA 2610042014 gene	66460	-13.0
AGCGAAATAA	<i>Gmfg</i>	Glia maturation factor gamma	63986	-11.0
CTGCAGCCTA	<i>Stx5a</i>	Syntaxin 5A	56389	-11.0
TTACAAGCCT	<i>Cks1b</i>	CDC28 protein kinase 1b	54124	-10.0
GTGGACTCAA	<i>Iftm1</i>	Interferon induced transmembrane protein 1	68713	-10.0
CATAGTTTAA	<i>Nol7</i>	Nucleolar protein 7	70078	-10.0
AAGTCTTCA	<i>Csn1s2a</i>	Casein alpha s2-like A	12993	-9.0
C3(1) SV40 T-antigen				
AGCAGTGCTT	<i>Ccdc3</i>	Coiled-coil domain containing 3	74186	13.0
CAGTTTGTA	<i>Pdha1</i>	Pyruvate dehydrogenase E1 alpha 1	18597	10.0
AATGTGTATG	<i>Abca8a</i>	ATP-binding cassette sub-family A (ABC1)	217258	9.0
ATTCCTGTT	<i>Krtap8-1</i>	Keratin associated protein 8-1	16703	8.0
CCGAAAAAAA	<i>Pink1</i>	PTEN induced putative kinase 1	68943	7.0
ACTCTAAAAA	<i>Tmem55b</i>	Transmembrane protein 55b	219024	7.0
CTGTAGTGTC	<i>Ltf</i>	Lactotransferrin	17002	7.0
CTGTCCAAGG	<i>Bhlhb2</i>	Basic helix-loop-helix domain containing class B2	20893	7.0
GAAAAATAAA	<i>Fndc3a</i>	Fibronectin type III domain containing 3a	319448	-20.0
TAAATTAAGA	<i>Hexb</i>	Hexosaminidase B	15212	-16.0
TTAGAAGTGA	<i>Savl</i>	Salvador homolog 1 (Drosophila)	64010	-15.0
GGGGTGAGG	<i>Hisppd1</i>	Histidine acid phosphatase domain containing 1	227399	-15.0
TAACAAAGGA	<i>Ahcy1l</i>	S-adenosylhomocysteine hydrolase-like 1	229709	-14.0
GATTAATAACA	<i>4931406120</i>	RIKEN cDNA 4931406120 gene	66743	-11.0
TTAACTACTGT	<i>Rab35</i>	RAB35, member RAS oncogene family	77407	-10.0
CAGATTAATA	<i>Gbp6</i>	Guanylate binding protein 6	229900	-9.0

*Up-regulated transcripts in bexarotene treatment are represented by positive fold changes and down-regulated transcripts are represented by negative fold changes.

otene modulated genes across transgenic mouse mammary models. The graph was generated employing the STRING on-line resource based on high confidence data related with 'co-expression/co-occurrence', 'experimental/biochemical data' and 'association in curated database/text mining' [17]. STRING is a comprehensive tool integrating protein association information with the capability to transfer known interactions from model organisms to other species (*e.g.*: from mouse to human orthology genes/proteins) based on predicted orthology of the respective proteins. The generated graph (Figure 4) indicates strong interactions among a set of 33 proteins transcriptionally modulated by bexarotene. Furthermore, the network architecture suggests the existence of two functional modules in this figure, involving the down-modulation of genes related with *protein biosynthesis* pathway, and up-modulation of genes related with *tricarboxylic acid cycle/oxidative phosphorylation* pathways.

Protein biosynthesis signature

A common observation in cancer gene expression profiling is the systematic up-regulation of ribosomal genes among the most abundant transcripts in human and mouse mammary carcinomas compared with normal tissues [20-24]. The up-regulation of ribosomal genes was significantly correlated with variation in the cell doubling time *in vitro*, supporting the notion that these genes are up-regulated in relation to the increase of cell proliferation rate or growth rate during malignant transformation. Interestingly and in an opposite manner, bexarotene treatment cause in 'normal' mammary gland the down-regulated expression of more than 10 genes related to protein biosynthesis including numerous ribosomal proteins (*Rpl19*, *Rpl37*, *Rps4x*, *Rps8*, *Rps24*, *Rps27*, *Rps29*), *Eef1b2* (*Eukaryotic translation elongation factor 1 beta 2*), *Eif2s3x* (*Eukaryotic translation initiation factor 2*), *Fau* (*Finkel-Biskis-Reilly murine sarcoma virus*) and *Tpt1* (*tumor protein, translationally-controlled 1*). The inhibition of mRNA synthesis for genes encoding ribosomal proteins has been suggested as a mechanism that could reprogram the cancer cell to recover some of its normal functions in a tumor reversion process [25].

Tpt1 (also known as *Tctp*) encodes a GDP dissociation inhibitor protein of the translation elongation factor eEF1A [26]. The human *TPT1* gene is overexpressed in cancerous cell lines compared with cell lines derived from normal tissues. Tuynder et al. (2002) demonstrated that the expression levels of *TPT1* were strongly down-regulated at the mRNA and protein levels during tumor reversion/suppression. MCF7 and T47D cell lines transfected with *Tpt1* siRNA showed a more organized ductal-like structures similar to those generated by down-regulation of $\beta 1$ integrin [25]. Here we observed that bexarotene sig-

nificantly downregulated expression of *Tpt1* in mammary epithelium.

Bioenergetics signatures

More than 50 years ago, Warburg proposed that malignant phenotype might be caused by a decrease in mitochondrial energy metabolism paralleled by increased glycolytic flux [27]. Increasing evidence is in line with this hypothesis suggesting a close link between metabolic and genetic changes observed during malignant growth [28,29]. Recently it has been demonstrated that impaired bioenergetic function of mitochondria is a hallmark of carcinogenesis in breast, gastric, lung and oesophageal cancer [30,31]. Moreover, Schulz et al. (2006) showed that induction of mitochondrial oxidative metabolism efficiently suppresses malignant growth *in vitro* and *in vivo*. Interestingly, we identified a systematic up-regulation of transcripts related to oxidative phosphorylation induced by bexarotene treatment in mammary gland (Figure 4). The transcripts commonly up-regulated by bexarotene treatment in at least two of the models were *Atp5b* (*ATP synthase F1 complex beta subunit*), *Atp5e* (*ATP synthase F1 complex epsilon subunit*), *Cyc1* (*Cytochrome c-1*), *Cycc* (*Cytochrome c somatic*), *Cox5b* (*Cytochrome c oxidase, subunit Vb*), *Cox7b* (*Cytochrome c oxidase subunit VIIb*), *Cox8a* (*Cytochrome c oxidase, subunit VIIa*), *Ndufa1* (*NADH dehydrogenase 1 alpha subcomplex*), *Ndufc1* (*NADH dehydrogenase 1*), *Ndufb8* (*NADH dehydrogenase 1 beta subcomplex 8*), *Ndufa11* (*NADH dehydrogenase 1 alpha subcomplex 11*) and *Uqcrrh* (*Ubiquinol-cytochrome c reductase hinge protein*) (see Additional files 1 and 2). Consistent with a significant increase of oxidative phosphorylation enzymes, we observed that *Atpif1* gene (*ATPase inhibitory factor 1*) was significantly down-regulated by bexarotene treatment in the MMTV-erbB2 and C3(1)/SV40 T-antigen transgenic mice models. In this sense, Isidoro et al. (2005) showed that down-regulation of ATPase β -F1 per se allowed the identification of a subgroup of breast cancer patients with significant worse prognosis. Finally, is important to note that mitochondrial oxidative phosphorylation is required for efficient execution of apoptosis. Cells which are unable to carry on oxidative phosphorylation have a resistant apoptotic phenotype [32]. Overall, these findings suggest the oxidative phosphorylation induction (prevention impaired bioenergetic function) as a novel mechanism of bexarotene's chemopreventive effects.

Fatty acid metabolic signature

Lipid metabolism and the intracellular transport of bioactive species is a critical component in the process by which these molecules continuously stimulate proliferation through interactions with nuclear receptors. Bexarotene treatment of MMTV-ErbB2 and C3(1)/SV40 transgenic mammary gland up-regulated various genes related with lipid/fatty acid metabolism (Figure 4) such as: *Fabp3*




Figure 2
Deregulated transcripts in mammary gland by systemic treatment with bexarotene in the three transgenic mice mammary cancer models. **A.** Scatter-plot representation of differentially expressed genes between bexarotene treated mice and vehicle control SAGE libraries ($p < 0.05$). **B.** Gene ontology (GO) classification of bexarotene induced differentially expressed transcripts on mammary gland from the different transgenic models. Relative representation of the deregulated transcripts with specific GO term annotations related to biological processes or molecular function.




Figure 3

Co-occurring differentially expressed genes among transgenic mouse mammary models. Eighty-nine genes were identified as modulated in more than one transgenic mice model. **A.** Heat map of the 89 deregulated transcripts. Color scale at the bottom depicts the approximate fold change in expression for each transcript and library relative to control mammary gland. Negative fold change (transcripts with decreased expression in bexarotene treated animals) is represented in green, and positive fold change (transcripts with overexpression in bexarotene treated mice) is represented in red. **B.** Statistical comparison between MMTV-erbB2 vs. p53-null and MMTV-erbB2 vs. C3(1)/SV40 T-antigen transgenic mice models showing a highly significant number of overlapping genes ($p < 0.001$). The number of overlapping genes between p53-null and C3(1) SV40 models it is not statistical significant ($p > 0.05$).

(Fatty acid binding protein 3), *Fabp4* (Fatty acid binding protein 4), *Dgat2* (Diacylglycerol O-acyltransferase 2), *Dbi* (Diazepam binding inhibitor), *Hadhb* (Hydroxyacyl-Coenzyme A dehydrogenase), *Acca2* (Acetyl-Coenzyme A acyltransferase 2), and *Cpt2* (Carnitine palmitoyltransferase 2). Interestingly, a family of cytoplasmic proteins known as FABPs mediates transport and utilization of lipids, and different FABP types have been implicated in control of cell proliferation and cancer progression. Recently, FABP1 and FABP2 were shown to be up-regulated in breast cancer cell lines while FABP3 and FABP4 were down-regulated in breast cancer cells [33]. Moreover, FABP4 is a marker protein for differentiated mammary gland that is expressed only in normal lactating cells and not in tumor mammary cells. Transfection of cDNA clone of FABP4 into MCF7 cells results in growth inhibition and lower tumorigenicity in nude mice [34].

Cell proliferation and apoptosis signatures

Rexinoid bexarotene down-regulated several genes related with cell cycle/proliferation in mammary gland from the different transgenic mice models (Figure 2A; see Additional files 1 and 2).

Among this functional group we find *Npm1* (also known as Nucleophosmin/B23) protein that belongs to a nuclear chaperone family of phosphoproteins that take part in various cellular processes such as cell proliferation and transformation [35]. Human *NPM1* is overexpressed in various tumors types, and it has been proposed as a marker for gastric, colon, ovarian and prostate carcinomas [35]. *NPM1* overexpression promotes cell survival in several cell types through the inhibition of distinct pro-apoptotic pathways [36]. We detected a systematic down-regulation of *Npm1* gene in mammary gland from Bexarotene treated mice on the three models studied (average fold change = -2.6; $p < 0.01$). Interestingly, proteomic analyses identified NPM1 as a protein associated with acquired estrogen-independence in human breast cancer cells [37]. Moreover, down-regulation of *NPM1* mRNA delay cell-cycle progression and the entry into mitosis [38], whereas *NPM1* overexpression decreases the sensitivity of human leukaemia cells to retinoic-acid-induced differentiation and apoptosis [39,40].

Another gene in this category includes *Stmn1* which encodes an 18 kDa cytosolic phosphoprotein (also known as Stathmin 1 or Oncoprotein 18) that is regulated during cell cycle by transcriptional and posttranscriptional mechanisms. *STMN1* overexpression has been demonstrated at mRNA and protein levels in a significant proportion of human breast carcinomas (about 30%) [41]. Moreover, *STMN1* overexpression was correlated with the loss of ER α and with histological grade III breast carcinomas. *STMN1* has been suggested as a key regulator

of the cell division through its influence on microtubule dynamics. We identified a statistical significant decrease of *Stmn1* expression (average fold change = -5.4; $p < 0.05$) caused by bexarotene treatment in mammary gland from MMTV-erbB2 and C3(1)/SV40 T-antigen mice. Interestingly, we previously demonstrated that mouse *Stmn1* and human homologue *STMN1* genes are overexpressed in invasive breast carcinomas by northern and real time RT-PCR analyses [24].

Numerous studies have shown that down-regulation of p27^{Kip1}, an inhibitor of cyclin-dependent kinase, is associated with poor prognosis in many cancers such as: breast, colorectal, prostate, and lung carcinomas. We previously detected the overexpression of *CDC28 protein kinase regulatory subunit 1B (Cks1b)* in human and mouse mammary tumors [24]. Interestingly, rexinoid bexarotene strongly down-regulated *Cks1b* expression in the MMTV-erbB2 model (Fold change = -10.0; $p = 0.006$) (see Additional file 1). Human *CKS1B* functions as an important adaptor of SCF Skp2 ubiquitin ligase and facilitates SCF Skp2 targeting of the cell proliferation inhibitor p27 (Kip1) for ubiquitination and subsequent degradation [42]. It was also suggested that *CKS1B* may be involved in p21 degradation in a similar fashion [43]. Overexpression of *CKS1B* has been observed associated to poorly differentiated tumors (histological grade III) and with the loss of ER/PR status [Slotky et al., 2005]. In addition, *CKS1B* overexpression was strongly and independently associated with poor overall survival in human breast cancer [44].

On the other hand, bexarotene treatment up-modulated two apoptosis related genes (*Cidec* and *Cycs*) in 'normal' mouse mammary gland from two of the models (*Cidec*) and in all three models (*Cycs*) (Figure 3). *Cidec* (also known as *Fsp27*) encode a novel family member of the cell-death-inducing DFF45-like-effectors (CIDEs) [45]. Although, its well known that DFF45 is a subunit of the DNA fragmentation factor that is cleaved by caspase-3 during apoptosis [46]. The molecular mechanism by which *Cidec* induces apoptosis remains to be elucidated.

Cell adhesion and invasion signatures

During their metastatic conversion, epithelial cells acquire the ability to invade the surrounding tissues and later disseminate to secondary organs mostly via lymphatic vessels. Epithelial cell adhesions, including intercellular (junctional) and cell-extracellular matrix adhesions, are critical to the maintenance of structural integrity, polarity, and cell-cell communication. We detected a significant decrease in *Cldn3 (Claudin 3)* (Average fold change = -6), *Glycam1 (Glycosylation dependent cell adhesion molecule 1)* (Average fold change = -7), *Pscd6p (Pleckstrin homology Sec7 binding protein)* (Average fold change = -6) gene expression modulated by bexarotene treatment among transgenic

mice models. The *Claudin* genes (*Cldn*) encode a family of proteins important in epithelial cell tight junction, which are critical to the maintenance of cell polarity and permeability [47,48]. Most Claudin genes appear with decreased expression in cancer however *CLDN3* and *CLDN4* genes have been found frequently up-regulated in ovarian, breast, prostate and pancreatic cancers [49-52]. Recently, has been suggested that Claudins may be involved in survival and invasion of cancer cells [48]. We detected down-regulation of *Cldn3* gene in mammary gland from bexarotene treated mice in the MMTV-erbB2 and C3(1)/SV40 models. The role of *Gycam1* and *Pscdbp* genes in breast cancer progression remains unknowns.

Conclusion

The present study showed that the rexinoid bexarotene (LGD1069) exerts its chemopreventive activity by affecting multiple cellular pathways, not only targeting cancer-causing genes related with cell proliferation, differentiation and apoptosis, but also by modulating protein biosynthesis and mitochondrial bioenergetics. Further analysis and validation of the identified genes will be required to determine the prognostic value as biomarkers of bexarotene treatment response, and to determine whether some of them and their protein products may constitute novel candidates for additional targeted therapeutic interventions.

We have recently completed a Phase II biomarker modulation trial in which women at high risk of breast cancer were treated with placebo or bexarotene. Using breast tissue from these high risk women, we are now studying whether these newly identified rexinoid-regulated biomarkers are also being modulated in human breast tissue. Results from these human studies will reveal whether these new biomarkers will be useful for predicting a cancer preventive response from rexinoids or as targets for future therapies.

Abbreviations

SAGE: Serial Analysis of Gene Expression; ER: Estrogen Receptor; RAR: Retinoic Acid Receptor; RXR: Retinoid × Receptor; GO: Gene Ontology database; DAVID: Database for Annotation, Visualization and Integratid Discovery; STRING: Search Tool for the Retrieval of Interacting Genes/Proteins.

Competing interests

The authors declare that they have no competing financial interests. As indicated author Reid Bissonnette works for Ligand Pharmaceuticals.

Authors' contributions

MCA conducted the analysis of the data; wrote the article; YH and CCL conducted SAGE studies; SG in charge of bioinformatics resources; FSK, YZ, JH in charge of animal

experiments; RPB provided the compound LGD1069 and contributed to the writing of the study, DM contributed to the writing of the manuscript and directed the p53 null studies, PHB and CMA directed the studies and contributed to the writing of the article.

Additional material

Additional file 1

Differentially expressed genes in mammary gland as the result of systemic bexarotene treatment versus control. The data provided represent the statistical analysis of SAGE libraries from p53-Null, MMTV-erbB2 and C3(1)/SV40 transgenic mice mammary cancer models (p < 0.05).

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1755-8794-1-40-S1.xls>]

Additional file 2

Co-occurring mammary gland deregulated transcripts as the result of systemic bexarotene treatment among transgenics mice mammary cancer models. The data provided represent the inter-model comparison for the identification of overlapping gene expression profiles among transgenics mice mammary cancer models (p < 0.05).

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1755-8794-1-40-S2.xls>]

Acknowledgements

This work was supported by NIH-NCI Grants R01 CA101211 (PB), U19 CA84978 (CMA), U01CA84243 and NIEHS center grant ES-07784.

References

- Smigal C, Jemal A, Ward E, Cokkinides V, Smith R, Howe HL, Thun M: **Trends in breast cancer by race and ethnicity: update 2006.** *CA Cancer J Clin* 2006, **56**:168-183.
- Shek LL, Dodolphin W: **Survival with breast cancer: the importance of estrogen receptor quantity.** *Eur J Cancer Clin Oncol* 1989, **25**:243-250.
- Shen Q, Brown PH: **Transgenic mouse models for the prevention of breast cancer.** *Mutat Res* 2005, **576**:93-110.
- Mehta K: **Retinoids as regulators of gene transcription.** *J Biol Regul Homeost Agents* 2003, **17**:1-12.
- Pemrick SM, Lucas DA, Grippo JF: **The retinoid receptors.** *Leukemia* 1994, **8**:1797-1806.
- Wu K, Kim H, Rodriguez JL, Hilsenbeck SG, Mohsin SK, Xu X, Lamph WW, Kuhn JG, Green JE, Brown PH: **Suppression of mammary tumorigenesis in transgenic mice by the RXR-selective retinoid, LGD1069.** *Cancer Epidemiol Biomarkers Prev* 2002, **11**:467-474.
- Wu K, Zhang Y, Xu X, Hill J, Celestino J, Kim H, Mohsin SK, Hilsenbeck SG, Lamph WW, Bissonette R, Brown PH: **The retinoid × receptor-selective retinoid, LGD prevent the development of estrogen receptor-negative mammary tumors in transgenic mice.** *Cancer Res* 2002, **62**:6376-80.
- Guy CT, Webster MA, Schaller M, Parsons TJ, Cardiff RD, Muller WJ: **Expression of the neu proto-oncogene in the mammary epithelium of transgenic mice induces metastatic disease.** *Proc Natl Acad Sci USA* 1992, **89**:10578-82.
- Green J, Shibata M, Yoshidome K, Liu M, Jorcyk C, Anver M, Wigington J, Wiltrout R, Shibata E, Kaczmarczyk S, Wang W, Liu Z, Calvo A, Couldrey C: **The C3(1)/SV40 T-antigen transgenic mouse model of mammary cancer: ductal epithelial cell targeting with multistage progression to carcinoma.** *Oncogene* 2000, **19**:1020-27.
- Jerry DJ, Kittrell FS, Kuperwasser C, Laucirica R, Dickinson ES, Bonilla PJ, Butel JS, Medina D: **A mammary-specific model demon-**

- strates the role of the p53 tumor suppressor gene in tumor development. *Oncogene* 2000, **19**:1052-8.
11. Charpentier AH, Bednarek AK, Daniel RL, Hawkins KA, Laflin KJ, Gaddis S, MacLeod MC, Aldaz CM: **Effects of estrogen on global gene expression: identification of novel targets of estrogen action.** *Cancer Res* 2000, **60**:5977-83.
 12. Aldaz CM, Hu Y, Daniel R, Gaddis S, Kittrell F, Medina D: **Serial analysis of gene expression in normal p53 null mammary epithelium.** *Oncogene* 2002, **21**:6366-6376.
 13. Audic S, Claverie J: **The significance of digital gene expression profiles.** *Genome Res* 1997, **7**:986-995.
 14. Smid M, Dorssers LCJ, Jenster G: **Venn Mapping: clustering of heterologous microarray data based on the number of co-occurring differentially expressed genes.** *Bioinformatic* 2003, **19**:2065-2071.
 15. Hosack DA, Dennis G, Sherman BT, Lane HC, Lempicki RA: **Identifying biological themes within lists of genes with EASE.** *Genome Biol* 2003, **4**:R70.
 16. Dennis G, Sherman BT, Hosack DA, Yang J, Gao W, Lane HC, Lempicki RA: **DAVID: Database for Annotation, Visualization, and Integrated Discovery.** *Genome Biol* 2003, **4**:R60.
 17. Von Mering C, Jensen LJ, Snel B, Hooper SD, Krupp M, Foglierini M, Jouffre N, Huynen MA, Bork P: **STRING: known and predicted protein-protein associations, integrated and transferred across organisms.** *Nucleic Acids Res* 2005, **33**:D433-D437.
 18. Von Mering C, Jensen LJ, Kuhn M, Chaffron S, Doerks T, Krüger B, Snel B, Bork P: **STRING 7-recent developments in the integration and prediction of protein interactions.** *Nucleic Acids Res* 2007, **35**:D358-D362.
 19. Kim H, Kong G, DeNardo D, Li Y, Uray I, Pal S, Mohsin S, Hilsenbeck SG, Bissonnette R, Lamph WW, Johnson K, Brown PH: **Identification of biomarkers modulated by the rexinoid LGD1069 (Bexarotene) in human breast cells using oligonucleotide arrays.** *Cancer Res* 2006, **66**:12009-18.
 20. Zhang L, Zhou W, Velculescu VE, Kern SE, Hruban RH, Hamilton SR, Vogelstein B, Kinzler KW: **Gene expression profiles in normal and cancer cells.** *Science* 1997, **276**:1268-1272.
 21. Nacht M, Ferguson AT, Zhang W, Petroziello JM, Cook BP, Gao YH, Maguire S, Riley D, Coppola G, Landes GM, Madden SL, Sukumar S: **Combining serial analysis of gene expression and array technologies to identify genes differentially expressed in breast cancer.** *Cancer Res* 1999, **59**:5464-5470.
 22. Porter DA, Krop IE, Nasser S, Sgroi D, Kaelin CM, Marks JR, Riggins G, Polyak K: **A SAGE (Serial Analysis of Gene Expression) view of breast tumor progression.** *Cancer Res* 2001, **61**:5697-5702.
 23. Abba MC, Drake JA, Hawkins KA, Hu Y, Sun H, Notcovich C, Gaddis S, Sahin A, Baggerly K, Aldaz CM: **Transcriptomic changes in human breast cancer progression as determined by serial analysis of gene expression.** *Breast Cancer Res* 2004, **6**:R499-R513.
 24. Hu Y, Sun H, Drake J, Kittrell F, Abba MC, Deng L, Gaddis S, Sahin A, Baggerly K, Medina D, Aldaz CM: **From mice to humans: identification of commonly deregulated genes in mammary cancer via comparative SAGE studies.** *Cancer Res* 2004, **64**:7748-55.
 25. Tuynder M, Susini L, Prieur S, Besse S, Ficuci G, Amson R, Telerman A: **Biological models and genes of tumor reversion: cellular reprogramming through tpt1/TCTP and SIAH-1.** *Proc Natl Acad Sci USA* 2002, **99**:14976-14981.
 26. Cans C, Passer BJ, Shalak V, Nancy-Portebois V, Crible V, Amzallag N, Allanic D, Tufino R, Argentinini M, Moras D, Ficuci G, Goud B, Mirande M, Amson R, Telerman A: **Translationally controlled tumor protein acts as a guanine nucleotide dissociation inhibitor on the translation elongation factor eEF1A.** *Proc Natl Acad Sci USA* 2003, **100**:13892-13897.
 27. Warburg O: **On the origin of cancer cells.** *Science* 1956, **123**:309-314.
 28. Ramanathan A, Wang C, Schreiber SL: **Perturbation profiling of a cell-line model of tumorigenesis by using metabolic measurements.** *Proc Natl Acad Sci USA* 2005, **102**:5992-5997.
 29. Schulz TJ, Thierbach R, Voigt A, Drewes G, Mietzner B, Steinberg P, Pfeiffer AFH, Ristow M: **Induction of oxidative metabolism by mitochondrial frataxin inhibits cancer growth.** *J Biol Chem* 2006, **281**:977-981.
 30. Isidoro A, Martínez M, Fernández PL, Ortega AD, Santamaría G, Chamorro M, Reed JC, Cuezva JM: **Alteration of the bioenergetic phenotype of mitochondria is a hallmark of breast, gastric, lung and oesophageal cancer.** *Biochem J* 2004, **378**:17-20.
 31. Isidoro A, Casado E, Redondo A, Acebo P, Espinosa E, Alonso AM, Cejas P, Hardisson D, Vara JAF, Belda-Iniesta C, González-Barón M, Cuezva JM: **Breast carcinoma fulfill the Warburg hypothesis and provide metabolic markers of cancer prognosis.** *Carcinogenesis* 2005, **26**:2095-2104.
 32. Dey R, Moraes CT: **Lack of oxidative phosphorylation and low-mitochondrial membrane potential decrease susceptibility to apoptosis and do not modulate the protective effect of Bcl-x(L) in osteosarcoma cells.** *J Biol Chem* 2000, **275**:7087-7094.
 33. Hammamieh R, Chakraborty N, Barmada M, Das R, Jett M: **Expresión patterns of fatty acid binding proteins in breast cancer cells.** *J Exp Ther Oncol* 2005, **5**:133-43.
 34. Buhlmann C, Borchers T, Pollak M, Spener F: **Fatty acid metabolism in human breast cancer cells (MCF7) transfected with heart-type fatty acid binding protein.** *Mol Cell Biochem* 1999, **199**:41-8.
 35. Grisendi S, Mecucci C, Falini B, Pandolfi PP: **Nucleophosmin and cancer.** *Nature Rev Cancer* 2006, **6**:493-505.
 36. Ye K: **Nucleophosmin/B23, a multifunctional protein that can regulate apoptosis.** *Cancer Biol Ther* 2005, **4**:918-923.
 37. Skaar TC, Prasad SC, Sharareh S, Lippman ME, Brunner N, Clarke R: **Two-dimensional gel electrophoresis analyses identify nucleophosmin as an estrogen regulated protein associated with acquired estrogen-independence in human breast cancer cells.** *J Steroid Biochem Mol Biol* 1998, **67**:391-402.
 38. Jianq PS, Yung BY: **Down-regulation of nucleophosmin/B23 mRNA delays the entry of cells into mitosis.** *Biochem Biophys Res Commun* 1999, **257**:865-70.
 39. Wu HL, Hsu CY, Liu WH, Yung BY: **Berberine-induced apoptosis of human leukemia HL-60 cells is associated with down-regulation of nucleophosmin/B23 and telomerase activity.** *Int J Cancer* 1999, **81**:923-9.
 40. Hsu CY, Yung BY: **Over-expression of nucleophosmin/B23 decreases the susceptibility of human leukemia HL-60 cells to retinoic acid-induced differentiation and apoptosis.** *Int J Cancer* 2000, **88**:392-400.
 41. Bieche I, Lachkar S, Becette V, Cifuentes-Diaz C, Sobel A, Lidereau R, Curmi P: **Overexpression of the stathmin gene in a subset of human breast cancer.** *Br J Cancer* 1998, **78**:701-709.
 42. Barket J, Lukas J: **p27 destruction: Cks I pulls the trigger.** *Nat Cell Biol* 2001, **3**:E95-8.
 43. Bornstein G, Bloom J, Sitry-Shevah D, Nakayama K, Pagano M, Hershko A: **Role of the SCFSkp2 ubiquitin ligase in the degradation of p21Cip1 in S phase.** *J Biol Chem* 2003, **278**:25752-7.
 44. Slotky M, Shapira M, Ben-Izhak O, Linn S, Futerman B, Tsalic M, Hershko DD: **The expression of the ubiquitin ligase subunit Cks I in human breast cancer.** *Breast Cancer Res* 2005, **7**:R737-R744.
 45. Liang L, Zhao M, Xu Z, Yokoyama KK, Li T: **Molecular cloning and characterization of CIDE-3, a novel member of the cell-death-inducing DNA-fragmentation-factor (DFF45)-like effector family.** *Biochem J* 2003, **370**:195-203.
 46. Inohara N, Koseki T, Chen S, Wu X, Núñez G: **CIDE, a novel family of cell death activators with homology to the 45 kDa subunit of the DNA fragmentation factor.** *EMBO J* 1998, **17**:2526-2533.
 47. Morita K, Furuse M, Fujimoto K, Tsukita S: **Claudin multigene family encoding four-transmembrane domain protein components of tight junction strands.** *Proc Natl Acad Sci USA* 1999, **96**:511-516.
 48. Agarwal R, D'Souza T, Morin PJ: **Claudin-3 and claudin-4 expression in ovarian epithelial cells enhances invasion and is associated with increased matrix metalloproteinase-2 activity.** *Cancer Res* 2005, **65**:7378-7385.
 49. Long H, Crean CD, Lee WH, Cummings OW, Gabig TG: **Expression of Clostridium perfringens enterotoxin receptors claudin-3 and claudin-4 in prostate cancer epithelium.** *Cancer Res* 2001, **61**:7878-7881.
 50. Rangel LBA, Agarwal R, D'Souza T, Pizer ES, Alò PL, Lancaster WD, Gregoire L, Schwartz DR, Cho KR, Morin PJ: **Tight junction proteins Claudin-3 and Claudin-4 are frequently overexpressed in ovarian cancer but not in ovarian cystadenomas.** *Clin Cancer Res* 2003, **9**:2567-2575.

51. Hewitt KJ, Agarwal R, Morin PJ: **The claudin gene family: expression in normal and neoplastic tissues.** *BMC Cancer* 2006, **6**:186.
52. Heinzelmann-Schwarz VA, Gardiner-Garden M, Henshall SM, Scurry J, Scolyer RA, Davies MJ, Heinzelmann M, Kalish LH, Bali A, Kench JG, Edwards LS, Bergh PM Vanden, Hacker NF, Sutherland RL, O'Brien PM: **Overexpression of the cell adhesion molecules DDRI, Claudin 3, and Ep-CAM in metaplastic ovarian epithelium and ovarian cancer.** *Clin Cancer Res* 2004, **10**:4427-4436.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1755-8794/1/40/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

