

RESEARCH

Open Access

Zeros of some difference polynomials

Shuangting Lan and Zongxuan Chen*

*Correspondence:
chzx@vip.sina.com
School of Mathematical Sciences,
South China Normal University,
Guangzhou, 510631, People's
Republic of China

Abstract

In this paper, we study zeros of some difference polynomials in $f(z)$ and their shifts, where $f(z)$ is a finite order meromorphic function having deficient value ∞ . These results improve previous findings.

1 Introduction and results

In this paper, we use the basic notions of Nevanlinna's theory (see [1]). In addition, we use the notations $\sigma(f)$ to denote the order of growth of the meromorphic function $f(z)$, and $\lambda(f)$ to denote the exponent of convergence of zeros of $f(z)$.

Hayman [2] studied the Picard values of a meromorphic function and their derivatives, and he obtained the following result.

Theorem A *If $f(z)$ is a transcendental entire function, $n \geq 3$ is an integer, and $a (\neq 0)$ is a constant, then $f'(z) - af(z)^n$ assumes all finite values infinitely often.*

Yang [3, 4] extended Theorem A to some differential polynomial in $f(z)$, where $f(z)$ is a transcendental meromorphic function satisfying $N(r, f) + N(r, \frac{1}{f}) = S(r, f)$.

Recently, many papers have focused on complex differences. They have obtained many new results on difference utilizing the value distribution theory of meromorphic functions. Some findings may be viewed as the corresponding difference analogues. First, recall some definitions. We call

$$f(z)^{i_{\lambda,0}} f(z+c_1)^{i_{\lambda,1}} \cdots f(z+c_k)^{i_{\lambda,k}}$$

is a monomial in $f(z)$ and its shifts $f(z+c_1), \dots, f(z+c_k)$, where c_1, \dots, c_k are distinct nonzero complex constants, and $d(\lambda) = i_{\lambda,0} + i_{\lambda,1} + \cdots + i_{\lambda,k}$ is its degree. Further,

$$P(z, f) = \sum_{\lambda \in I} a_{\lambda}(z) f(z)^{i_{\lambda,0}} f(z+c_1)^{i_{\lambda,1}} \cdots f(z+c_k)^{i_{\lambda,k}} \quad (1.1)$$

is called a difference polynomial in $f(z)$ and its shifts, and $d(P) = \max_{\lambda \in I} \{d(\lambda)\}$ is its degree, where I is a finite set of the index $\lambda = (i_{\lambda,0}, \dots, i_{\lambda,k})$, and $a_{\lambda}(z)$ are meromorphic coefficients being small with respect to f in the sense that $T(r, a_{\lambda}) = S(r, f)$, $\lambda \in I$.

Zheng and Chen [5] obtained the following theorem, which may be considered as a difference counterpart of Theorem A.

Theorem B *Let $f(z)$ be a transcendental entire function of finite order, and let a, c be nonzero constants. Then, for any integer $n \geq 3$, $f(z + c) - af(z)^n$ assumes all finite values $b \in \mathbb{C}$ infinitely often.*

Chen [6] obtained the following Theorems C-E, which may be considered as another difference counterpart of Theorem A.

Theorem C *Let $f(z)$ be a transcendental entire function of finite order, and let $a, c \in \mathbb{C} \setminus \{0\}$ be constants with c such that $f(z + c) \not\equiv f(z)$. Set $\Psi_n(z) = \Delta f(z) - af(z)^n$, where $\Delta f(z) = f(z + c) - f(z)$ and $n \geq 3$ is an integer. Then Ψ_n assumes all finite values infinitely often, and for every $b \in \mathbb{C}$ one has $\lambda(\Psi_n - b) = \sigma(f)$.*

Theorem D *Let $f(z)$ be a transcendental entire function of finite order with a Borel exceptional value 0, and let $a, c \in \mathbb{C} \setminus \{0\}$ be constants, with c such that $f(z + c) \not\equiv f(z)$. Then $\Psi_2(z)$ assumes all finite values infinitely often, and for every $b \in \mathbb{C}$ one has $\lambda(\Psi_2 - b) = \sigma(f)$.*

Zheng and Chen [7] extended the above Theorems B-D to a difference polynomial and proved the following theorem.

Theorem E *Let $f(z)$ be a transcendental meromorphic function of finite order satisfying*

$$N(r, f) + N\left(r, \frac{1}{f}\right) = S(r, f). \tag{1.2}$$

Suppose that $P(z, f)$ is a difference polynomial of the form (1.1) and contains exactly one term of maximal total degree in $f(z)$ and its shifts. Then, for any given $a \in \mathbb{C} \setminus \{0\}$, the difference polynomial

$$Q(z, f) = f(z)^n + P(z, f), \quad n \geq d(P) + 2$$

satisfies $\delta(a, Q(z, f)) < 1$. Therefore, $Q(z, f)$ takes every nonzero finite value infinitely often.

We observe that condition (1.2) can be weakened. It is well known that deficient value plays an important role in the theory of value distribution of meromorphic functions. In this paper, we consider the meromorphic function $f(z)$ having deficient value ∞ and obtain the following improvements of Theorem E, which may be stated as follows.

Theorem 1.1 *Let $f(z)$ be a transcendental meromorphic function of finite order, and let $P(z, f) (\not\equiv 0)$ be a difference polynomial of the form (1.1) with k different shifts. Assume that $\delta(\infty, f) > 1 - \frac{1}{2k+7}$. Then the difference polynomial*

$$Q(z, f) = f(z)^n + P(z, f), \quad n \geq d(P) + 2, \tag{1.3}$$

has infinitely many zeros and satisfies $\delta(0, Q(z, f)) < 1$.

If we add the condition $\overline{N}(r, \frac{1}{f}) = S(r, f)$, then the condition $\delta(\infty, f) > 1 - \frac{1}{2k+7}$ in Theorem 1.1 can be weakened as $\delta(\infty, f) > 1 - \frac{1}{2k+5}$, and we obtain the following theorem.

Theorem 1.2 Let $f(z)$ be a transcendental meromorphic function of finite order, and let $P(z, f) (\neq 0)$ be a difference polynomial of the form (1.1) with k different shifts. Assume that $\delta(\infty, f) > 1 - \frac{1}{2k+5}$ and $\overline{N}(r, \frac{1}{f}) = S(r, f)$, then the difference polynomial

$$Q(z, f) = f(z)^n + P(z, f), \quad n \geq d(P) + 1 \tag{1.4}$$

has infinitely many zeros and satisfies $\delta(0, Q(z, f)) < 1$.

Example 1.1 For $f(z) = e^z + 1$, $c = \log 3$, we have $\Psi_2(z) = -f(z)^2 + \Delta f(z) = -e^{2z} - 1$. Here $\Psi_2(z) \neq -1$, which shows that the condition $\overline{N}(r, \frac{1}{f}) = S(r, f)$ in Theorem 1.2 cannot be replaced with $\overline{N}(r, \frac{1}{f-c}) = S(r, f)$, where c is a nonzero constant.

If we add the additional condition $\overline{N}(r, f) = S(r, f)$, then we obtain the following.

Theorem 1.3 Let $f(z)$ be a transcendental meromorphic function of finite order satisfying

$$\delta(\infty, f) > \frac{2}{3}, \quad \overline{N}(r, f) = S(r, f).$$

Then the difference polynomial $Q(z, f)$ defined by (1.3) has infinitely many zeros and satisfies $\delta(0, Q(z, f)) < 1$.

Theorem 1.4 Let $f(z)$ be a transcendental meromorphic function of finite order with $\delta(\infty, f) > 0$ satisfying

$$\overline{N}(r, f) + \overline{N}\left(r, \frac{1}{f}\right) = S(r, f). \tag{1.5}$$

Then the difference polynomial $Q(z, f)$ defined by (1.4) has infinitely many zeros and satisfies $\delta(0, Q(z, f)) < 1$.

Remark 1.1 For any given $b \in \mathbb{C}$, applying Theorems 1.1-1.4 to the difference polynomial $Q^*(z, f) = Q(z, f) - b$, then we have $\delta(b, Q(z, f)) < 1$.

Remark 1.2 If $Q(z, f) = a_n(z)f(z)^n + P(z, f)$, where $a_n(z)$ is small with respect to f , applying Theorems 1.1-1.4 to the difference polynomial $Q^*(z, f) = \frac{Q(z, f)}{a_n(z)}$, then we also have $\delta(0, Q(z, f)) < 1$.

Applying Theorems 1.1-1.4 to the difference polynomial $\Psi_n = \Delta f(z) - af(z)^n$, we get the following Corollaries 1.5-1.8, which extend Theorems C, D.

Corollary 1.5 Let $f(z)$ be a transcendental meromorphic function of finite order with $\delta(\infty, f) > \frac{8}{9}$, and let $a, c \in \mathbb{C} \setminus \{0\}$ be constants with c such that $\Delta f(z)$ does not reduce to any constant and $n \geq 3$ is an integer. Then, for every $b \in \mathbb{C}$, one has $\delta(b, \Psi_n) < 1$.

Corollary 1.6 Let $f(z)$ be a transcendental meromorphic function of finite order satisfying

$$\delta(\infty, f) > \frac{2}{3}, \quad \overline{N}(r, f) = S(r, f),$$

and let $a, c \in \mathbb{C} \setminus \{0\}$ be constants, with c such that $\Delta f(z)$ does not reduce to any constant and $n \geq 3$ is an integer. Then, for every $b \in \mathbb{C}$, one has $\delta(b, \Psi_n) < 1$.

Corollary 1.7 Let $f(z)$ be a transcendental meromorphic function of finite order satisfying

$$\delta(\infty, f) > \frac{6}{7}, \quad \overline{N}\left(r, \frac{1}{f}\right) = S(r, f),$$

and let $a, c \in \mathbb{C} \setminus \{0\}$ be constants, with c such that $\Delta f(z)$ does not reduce to any constant. Then, for every $b \in \mathbb{C}$, one has $\delta(b, \Psi_2) < 1$.

Corollary 1.8 Let $f(z)$ be a transcendental meromorphic function of finite order satisfying $\delta(\infty, f) > 0$ and

$$\overline{N}(r, f) + \overline{N}\left(r, \frac{1}{f}\right) = S(r, f),$$

and let $a, c \in \mathbb{C} \setminus \{0\}$ be constants with c such that $\Delta f(z)$ does not reduce to any constant. Then, for every $b \in \mathbb{C}$, one has $\delta(b, \Psi_2) < 1$.

2 Lemmas for the proof of the theorem

Lemma 2.1 [8] Given two distinct complex constants η_1, η_2 , let f be a meromorphic function of finite order. Then we have

$$m\left(r, \frac{f(z + \eta_1)}{f(z + \eta_2)}\right) = S(r, f).$$

Lemma 2.2 [9] Let $f(z)$ be a nonconstant meromorphic function of finite order, then for any given constant $c (\neq 0)$, we have

$$N(r, f(z + c)) = N(r, f) + S(r, f), \quad \overline{N}(r, f(z + c)) = \overline{N}(r, f) + S(r, f)$$

possibly outside of an exceptional set of finite logarithmic measure.

Lemma 2.3 [4] Let $f(z)$ be a transcendental meromorphic function, and let $P(f)$ be an algebraic polynomial in f of the form

$$P(f) = a_n(z)f(z)^n + a_{n-1}(z)f(z)^{n-1} + \dots + a_1(z)f(z) + a_0(z), \tag{2.1}$$

where $a_n(z) \not\equiv 0$, $a_j (j = 0, \dots, n)$ are small with respect to f . Then

$$T(r, P(f)) = nT(r, f) + S(r, f) \quad \text{as } r \rightarrow \infty.$$

Using the same method as in the proof of Lemma 2.3, we show the following.

Lemma 2.4 Let $f(z)$ be a transcendental meromorphic function with $\delta(\infty, f) > 0$, and let $P(f)$ be an algebraic polynomial in f of the form (2.1), where $a_n(z) \not\equiv 0$, $a_j(z)$ satisfy $m(r, a_j) = S(r, f)$, $j = 0, \dots, n$, then

$$m(r, P(f)) \leq nm(r, f) + S(r, f).$$

Proof Since $a_j(z)$ satisfy $m(r, a_j) = S(r, f)$, $j = 0, \dots, n$, then

$$\begin{aligned} m(r, P(f)) &\leq m(r, f(a_n f^{n-1} + a_{n-1} f^{n-2} + \dots + a_2 f + a_1)) + m(r, a_0) + O(1) \\ &\leq m(r, f) + m(r, (a_n f^{n-1} + a_{n-1} f^{n-2} + \dots + a_2 f + a_1)) + m(r, a_0) + O(1) \\ &\leq m(r, f) + m(r, f(a_n f^{n-2} + a_{n-1} f^{n-3} + \dots + a_2)) + m(r, a_1) + m(r, a_0) + O(1) \\ &\leq 2m(r, f) + m(r, f(a_n f^{n-3} + a_{n-1} f^{n-4} + \dots + a_3)) + \sum_{j=0}^2 m(r, a_j) + O(1) \\ &\dots \\ &\leq nm(r, f) + \sum_{j=0}^n m(r, a_j) + O(1) \\ &= nm(r, f) + S(r, f). \end{aligned}$$

Thus,

$$m(r, P(f)) \leq nm(r, f) + S(r, f). \quad \square$$

Lemma 2.5 [10] *Let f be a transcendental meromorphic solution of finite order of a difference equation of the form*

$$U(z, f)P(z, f) = Q(z, f), \tag{2.2}$$

where $U(z, f)$, $P(z, f)$ and $Q(z, f)$ are difference polynomials such that the total degree $\deg_f U(z, f) = n$ in $f(z)$ and its shifts, and $\deg_f Q(z, f) \leq n$. Moreover, we assume that all coefficients $a_\lambda(z)$ in (2.2) are small in the sense that $T(r, a_\lambda) = S(r, f)$ and that $U(z, f)$ contains just one term of maximal total degree in $f(z)$ and its shifts. Then, for each $\varepsilon > 0$, we have

$$m(r, P(z, f)) = S(r, f)$$

possibly outside of an exceptional set of finite logarithmic measure.

Yang and Laine in [11] further pointed out the following.

Remark 2.1 If $P(z, f)$ and $Q(z, f)$ are differential-difference polynomials in $f(z)$ and its shifts with coefficients $a_\lambda(z)$ satisfy $m(r, a_\lambda) = S(r, f)$, then Lemma 2.5 still holds.

3 Proof of Theorems 1.1-1.4

Proof of Theorem 1.1 We assert that $Q(z, f)$ does not reduce to any constant. In fact if $Q(z, f) \equiv C_1$, where C_1 is some constant, then

$$f(z)^n = -P(z, f) + C_1. \tag{3.1}$$

Applying Lemma 2.5 to (3.1), since $n \geq d(P) + 2$, we have

$$m(r, f) = S(r, f). \tag{3.2}$$

Since $\delta(\infty, f) = \delta > 0$, then for any given ε_0 ($0 < \varepsilon_0 < \delta$) and sufficiently large r , we have

$$m(r, f) \geq (\delta - \varepsilon_0)T(r, f). \tag{3.3}$$

By (3.2) and (3.3), we get

$$T(r, f) = S(r, f),$$

which is impossible. So, $Q(z, f)$ does not reduce to any constant.

Next we rearrange the expression difference polynomial $Q(z, f)$ by collecting together all terms having the same total degree and then writing $Q(z, f)$ in the form

$$Q(z, f) = f(z)^n + \sum_{j=0}^{d(P)} \tilde{a}_j(z) f(z)^j, \tag{3.4}$$

where the coefficients $\tilde{a}_j(z)$, $j = 0, 1, \dots, d(P)$ are the sum of finitely many terms of the type

$$a_\lambda(z) \prod_{j=1}^{\tau_\lambda} \left(\frac{f(z + c_j)}{f(z)} \right)^{i_{\lambda,j}}.$$

By Lemma 2.1 and our assumption concerning the coefficients a_λ , we obtain

$$m(r, \tilde{a}_j) = S(r, f), \quad j = 0, 1, \dots, d(P). \tag{3.5}$$

By (3.4), (3.5) and Lemma 2.4, we have

$$m(r, Q(z, f)) \leq nm(r, f) + S(r, f).$$

By Lemma 2.2, we have

$$\begin{aligned} N(r, Q(z, f)) &\leq nN(r, f(z)) + d(P) \sum_{j=1}^k N(r, f(z + c_j)) + S(r, f) \\ &= (n + kd(P))N(r, f) + S(r, f) \end{aligned}$$

and

$$\begin{aligned} \bar{N}(r, Q(z, f)) &\leq \bar{N}(r, f(z)) + \sum_{j=1}^k \bar{N}(r, f(z + c_j)) + O(1) \\ &\leq (k + 1)\bar{N}(r, f) + S(r, f). \end{aligned} \tag{3.6}$$

Thus,

$$\begin{aligned} T(r, Q(z, f)) &= m(r, Q(z, f)) + N(r, Q(z, f)) \\ &\leq (n + kd(P))T(r, f) + S(r, f). \end{aligned} \tag{3.7}$$

So, $S(r, Q(z, f)) \leq S(r, f)$.

Differentiating both sides of (1.3), we obtain that

$$Q'(z, f) = \eta f^{n-1} f' + P'(z, f). \tag{3.8}$$

Multiplying both sides of (1.3) by $\frac{Q'}{Q}$ and subtracting from (3.8), we get

$$f^{n-1} \left(\eta f' - \frac{Q'}{Q} f \right) = \frac{Q'}{Q} P(z, f) - P'(z, f) \tag{3.9}$$

or

$$f^{n-2} f \left(\eta f' - \frac{Q'}{Q} f \right) = \frac{Q'}{Q} P(z, f) - P'(z, f). \tag{3.10}$$

Now assert $\eta f' - \frac{Q'}{Q} f \neq 0$. Otherwise $\frac{Q'}{Q} = \eta \frac{f'}{f}$, i.e., $Q = C_2 f^n$ for a suitable constant C_2 leading to

$$(C_2 - 1)f(z)^n = P(z, f). \tag{3.11}$$

We can see that $C_2 \neq 1$, for $P(z, f) \neq 0$. Since $n > d(P)$ by assumption and Lemma 2.5, (3.11) implies $m(r, f) = S(r, f)$, which shows that $T(r, f) = S(r, f)$, again by (3.3). It is a contradiction.

Applying Lemma 2.5 and Remark 2.1 to (3.9) and (3.10) respectively, again by assumption $n \geq d(P) + 2$, we obtain

$$m \left(r, \eta f' - \frac{Q'}{Q} f \right) = S(r, f) \tag{3.12}$$

and

$$m \left(r, f \left(\eta f' - \frac{Q'}{Q} f \right) \right) = S(r, f). \tag{3.13}$$

By (3.6), we get that

$$\begin{aligned} N \left(r, \eta f' - \frac{Q'}{Q} f \right) &\leq N(r, f) + \bar{N}(r, f) + \bar{N}(r, Q) + \bar{N} \left(r, \frac{1}{Q} \right) + O(1) \\ &\leq N(r, f) + (k + 2)\bar{N}(r, f) + \bar{N} \left(r, \frac{1}{Q} \right) + S(r, f) \end{aligned} \tag{3.14}$$

and

$$N \left(r, f \left(\eta f' - \frac{Q'}{Q} f \right) \right) \leq 2N(r, f) + \bar{N}(r, f) + \bar{N}(r, Q) + \bar{N} \left(r, \frac{1}{Q} \right) + O(1)$$

$$\leq 2N(r, f) + (k + 2)\bar{N}(r, f) + \bar{N}\left(r, \frac{1}{Q}\right) + S(r, f). \tag{3.15}$$

By (3.12)-(3.15), we obtain

$$\begin{aligned} T\left(r, nf' - \frac{Q'}{Q}f\right) &= m\left(r, nf' - \frac{Q'}{Q}f\right) + N\left(r, nf' - \frac{Q'}{Q}f\right) \\ &\leq N(r, f) + (k + 2)\bar{N}(r, f) + \bar{N}\left(r, \frac{1}{Q}\right) + S(r, f) \end{aligned} \tag{3.16}$$

and

$$\begin{aligned} T\left(r, f\left(nf' - \frac{Q'}{Q}f\right)\right) &= m\left(r, f\left(nf' - \frac{Q'}{Q}f\right)\right) + N\left(r, f\left(nf' - \frac{Q'}{Q}f\right)\right) \\ &\leq 2N(r, f) + (k + 2)\bar{N}(r, f) + \bar{N}\left(r, \frac{1}{Q}\right) + S(r, f). \end{aligned} \tag{3.17}$$

Therefore,

$$\begin{aligned} T(r, f) &\leq T\left(r, nf' - \frac{Q'}{Q}f\right) + T\left(r, f\left(nf' - \frac{Q'}{Q}f\right)\right) + O(1) \\ &\leq 3N(r, f) + 2(k + 2)\bar{N}(r, f) + 2\bar{N}\left(r, \frac{1}{Q}\right) + S(r, f), \end{aligned} \tag{3.18}$$

or

$$T(r, f) \leq (2k + 7)N(r, f) + 2\bar{N}\left(r, \frac{1}{Q}\right) + S(r, f).$$

For any given ε ($0 < \varepsilon < \delta - 1 + \frac{1}{2k+7}$), we have

$$T(r, f) \leq (2k + 7)(1 - \delta + \varepsilon)T(r, f) + 2\bar{N}\left(r, \frac{1}{Q}\right) + S(r, f),$$

or

$$(1 - (2k + 7)(1 - \delta + \varepsilon) + o(1))T(r, f) \leq 2\bar{N}\left(r, \frac{1}{Q}\right). \tag{3.19}$$

By (3.7) and (3.19), we have

$$(1 - (2k + 7)(1 - \delta + \varepsilon) + o(1))T(r, Q) \leq 2(n + kd(P))\bar{N}\left(r, \frac{1}{Q}\right),$$

which leads to

$$\overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}\left(r, \frac{1}{Q}\right)}{T(r, Q)} \geq \frac{1 - (2k + 7)(1 - \delta + \varepsilon)}{2(n + kd(P))} > 0.$$

Therefore,

$$\delta(0, Q) = 1 - \overline{\lim}_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{Q}\right)}{T(r, Q)} < 1.$$

□

Proof of Theorem 1.3 Using the same method as in the proof of Theorem 1.1, we see that (3.1)-(3.18) hold. By assumption $\bar{N}(r, f) = S(r, f)$ and (3.16)-(3.18), we have

$$\begin{aligned} T(r, f) &\leq 3N(r, f) + 2(k + 2)\bar{N}(r, f) + 2\bar{N}\left(r, \frac{1}{Q}\right) + S(r, f), \\ &= 3N(r, f) + 2\bar{N}\left(r, \frac{1}{Q}\right) + S(r, f). \end{aligned}$$

For any given ε , $0 < \varepsilon < \delta - \frac{2}{3}$, where $\delta = \delta(\infty, f) > \frac{2}{3}$, we have

$$(1 - 3(1 - \delta + \varepsilon) + o(1))T(r, f) \leq 2\bar{N}\left(r, \frac{1}{Q}\right).$$

By (3.7), we have

$$(1 - 3(1 - \delta + \varepsilon) + o(1))T(r, Q) \leq 2(n + kd(P))\bar{N}\left(r, \frac{1}{Q}\right),$$

which leads to

$$\overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}(r, \frac{1}{Q})}{T(r, Q)} \geq \frac{1 - 3(1 - \delta + \varepsilon)}{2(n + kd(P))} > 0.$$

Therefore,

$$\delta(0, Q) = 1 - \overline{\lim}_{r \rightarrow \infty} \frac{N(r, \frac{1}{Q})}{T(r, Q)} < 1. \quad \square$$

Proof of Theorem 1.2 The proof is similar to the proof of Theorem 1.1.

We also obtain (3.1)-(3.9), (3.12), (3.14) and (3.16), by assumption that $n \geq d(P) + 1$.

By logarithmic derivative lemma, we get

$$m\left(r, n\frac{f'}{f} - \frac{Q'}{Q}\right) \leq S(r, f) + S(r, Q) = S(r, f). \quad (3.20)$$

Since $\bar{N}(r, \frac{1}{f}) = S(r, f)$, by (3.6), we get

$$\begin{aligned} N\left(r, n\frac{f'}{f} - \frac{Q'}{Q}\right) &\leq \bar{N}(r, f) + \bar{N}\left(r, \frac{1}{f}\right) + \bar{N}(r, Q) + \bar{N}\left(r, \frac{1}{Q}\right) + O(1) \\ &\leq (k + 2)\bar{N}(r, f) + \bar{N}\left(r, \frac{1}{Q}\right) + S(r, f). \end{aligned} \quad (3.21)$$

By (3.20) and (3.21), we have

$$\begin{aligned} T\left(r, n\frac{f'}{f} - \frac{Q'}{Q}\right) &= m\left(r, n\frac{f'}{f} - \frac{Q'}{Q}\right) + N\left(r, n\frac{f'}{f} - \frac{Q'}{Q}\right) \\ &\leq (k + 2)\bar{N}(r, f) + \bar{N}\left(r, \frac{1}{Q}\right) + S(r, f). \end{aligned} \quad (3.22)$$

Then,

$$\begin{aligned} T(r, f) &\leq T\left(r, nf' - \frac{Q'}{Q}f\right) + T\left(r, n\frac{f'}{f} - \frac{Q'}{Q}\right) + O(1) \\ &\leq N(r, f) + 2(k+2)\overline{N}(r, f) + 2\overline{N}\left(r, \frac{1}{Q}\right) + S(r, f), \end{aligned} \tag{3.23}$$

or

$$T(r, f) \leq (2k+5)N(r, f) + 2\overline{N}\left(r, \frac{1}{Q}\right) + S(r, f).$$

For any given ε , $0 < \varepsilon < \delta - 1 + \frac{1}{2k+5}$, where $\delta = \delta(\infty, f) > 1 - \frac{1}{2k+5}$, by (3.7), we obtain

$$(1 - (2k+5)(1 - \delta + \varepsilon))T(r, Q) \leq 2(n + kd(P))\overline{N}\left(r, \frac{1}{Q}\right),$$

which leads to

$$\overline{\lim}_{r \rightarrow \infty} \frac{\overline{N}(r, \frac{1}{Q})}{T(r, Q)} \geq \frac{1 - (2k+5)(1 - \delta + \varepsilon)}{2(n + kd(P))} > 0.$$

Therefore,

$$\delta(0, Q) = 1 - \overline{\lim}_{r \rightarrow \infty} \frac{N(r, \frac{1}{Q})}{T(r, Q)} < 1. \quad \square$$

Proof of Theorem 1.4 Using the same method as in the proof of Theorem 1.2, we obtain (3.23). By (1.5) and (3.23), we have

$$T(r, f) \leq N(r, f) + 2\overline{N}\left(r, \frac{1}{Q}\right) + S(r, f).$$

For any given ε , $0 < \varepsilon < \delta$, where $\delta = \delta(\infty, f)$, we have

$$(\delta - \varepsilon + o(1))T(r, f) \leq 2\overline{N}\left(r, \frac{1}{Q}\right).$$

Then by (3.7), we have

$$(\delta - \varepsilon + o(1))T(r, Q) \leq 2(n + kd(P))\overline{N}\left(r, \frac{1}{Q}\right),$$

which leads to

$$\overline{\lim}_{r \rightarrow \infty} \frac{\overline{N}(r, \frac{1}{Q})}{T(r, Q)} \geq \frac{\delta - \varepsilon}{2(n + kd(P))} > 0.$$

Therefore,

$$\delta(0, Q) = 1 - \overline{\lim}_{r \rightarrow \infty} \frac{N(r, \frac{1}{Q})}{T(r, Q)} < 1. \quad \square$$

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors drafted the manuscript, read and approved the final manuscript.

Acknowledgements

The authors would like to thank the editor and the referees for their constructive comments to improve the readability of our paper. The project was supported by the National Natural Science Foundation of China (11171119, 11126145).

Received: 25 March 2013 Accepted: 14 June 2013 Published: 1 July 2013

References

1. Yang, L: Value Distribution Theory and Its New Research. Science Press, Beijing (1982) (in Chinese)
2. Hayman, WK: Picard values of meromorphic functions and their derivatives. *Ann. Math.* **70**(1), 9-42 (1959)
3. Yang, CC: On deficiencies of differential polynomials. *Math. Z.* **116**, 197-204 (1970)
4. Yang, CC: On deficiencies of differential polynomials II. *Math. Z.* **125**, 107-112 (1972)
5. Zheng, XM, Chen, ZX: On the value distribution of some difference polynomials. *J. Math. Anal. Appl.* **397**, 814-821 (2013)
6. Chen, ZX: On value distribution of difference polynomials of meromorphic functions. *Abstr. Appl. Anal.* **2011**, Article ID 239853 (2011)
7. Zheng, XM, Chen, ZX: On deficiencies of some difference polynomials. *Acta Math. Sin.* **54**, 983-992 (2011) (in Chinese)
8. Halburd, RG, Korhonen, RJ: Difference analogue of the lemma on the logarithmic derivative with applications to difference equations. *J. Math. Anal. Appl.* **314**, 477-487 (2006)
9. Chiang, YM, Feng, SJ: On the Nevanlinna characteristic of $f(z + \eta)$ and difference equations in the complex plane. *Ramanujan J.* **16**, 105-129 (2008)
10. Laine, I, Yang, CC: Clunie theorems for difference and q -difference polynomials. *J. Lond. Math. Soc.* **2**(76), 556-566 (2007)
11. Yang, CC, Laine, I: On analogies between nonlinear difference and differential equations. *Proc. Jpn. Acad., Ser. A, Math. Sci.* **86**(1), 10-14 (2010)

doi:10.1186/1687-1847-2013-194

Cite this article as: Lan and Chen: Zeros of some difference polynomials. *Advances in Difference Equations* 2013 2013:194.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
