

POSTER PRESENTATION

Open Access

Entorhinal phase precession revisited – single-run analysis of *in-vivo* grid cell data

Eric T Reifenstein^{1*}, Martin B Stemmler², Andreas VM Herz², Susanne Schreiber¹*From* Twentieth Annual Computational Neuroscience Meeting: CNS*2011 Stockholm, Sweden. 23-28 July 2011

When a rat explores its environment, grid cells in the medial entorhinal cortex show increased activity at specific locations that constitute a regular hexagonal grid. As the rat enters and progresses through one of these “grid fields” on a linear track, spikes occur at successively earlier phases in the LFP’s theta rhythm. This phenomenon is called phase precession. For rats foraging in two-dimensional environments, however, phase precession has not yet been quantified. Unlike on the linear track, a rat does not repeat the same path over and over again in an open arena; pooling different runs to assess phase precession becomes fraught with difficulty. Instead, we analyze grid cell spike trains recorded by [1] on a run-by-run basis, and do the same also for linear track data (by [2]). Surprisingly, even on the linear track, phase precession during single runs is stronger than the average phase precession in the pooled data. We show that a grid cell spike’s theta phase allows one to estimate the animal’s position on a linear track to within about 10 % of the size of a typical grid field. The spectrum of grid cell spike trains recorded in a two-dimensional environment reveals a peak that is shifted by approximately 1 Hz relative to the peak in the LFP theta rhythm, indicative of robust phase precession. We investigate the dependence of spike phase on the radial distance from the grid field center and how phase precession depends on whether the path is transverse or tangential to the grid field, in order to create a model for decoding position in two dimensions from spike phases.

Acknowledgements

We thank the Moser lab for providing us with the opportunity to analyze their data. This work was supported by the DFG (GRK 1123) and the BMBF (BCCN Berlin, BPCN).

* Correspondence: eric.reifenstein@bccn-berlin.de¹Institute for Theoretical Biology, Humboldt-Universität zu Berlin, Berlin, Germany

Full list of author information is available at the end of the article

Author details

¹Institute for Theoretical Biology, Humboldt-Universität zu Berlin, Berlin, Germany. ²Department Biology II, Ludwig-Maximilians-Universität, Planegg-Martinsried, Germany.

Published: 18 July 2011

References

1. Sargolini F, Fyhn M, Hafting T, McNaughton BL, Witter MP, Moser MB, Moser EI: **Conjunctive representation of position, direction, and velocity in entorhinal cortex.** *Science* 2006, **312**(5774):758-762.
2. Hafting T, Fyhn M, Bonnevie T, Moser MB, Moser EI: **Hippocampus-independent phase precession in entorhinal grid cells.** *Nature* 2008, **453**:1248-1252.

doi:10.1186/1471-2202-12-S1-P288

Cite this article as: Reifenstein *et al.*: Entorhinal phase precession revisited – single-run analysis of *in-vivo* grid cell data. *BMC Neuroscience* 2011 **12**(Suppl 1):P288.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

