

ORAL PRESENTATION

Open Access

The relationship of the transmural extent of T2-edema compared with the transmural extent of infarction: implications for the assessment of the area-at-risk

Han W Kim^{*}, Ben Wince, Lowie van Assche, Wolfgang G Rehwald, Lubna B Bhatti, Deneen M Spatz, Igor Klem, Raymond J Kim

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Objective

To compare the transmural extent of hyperintensity (“edema”) on T2-weighted (T2W) CMR to the transmural extent of infarction by delayed-enhancement(DE)-CMR.

Background

It has been reported that hyperintensity on T2W-CMR represents the area-at-risk. This conclusion is based upon the observation that the region of T2-hyperintensity is always larger than region of infarction. However, differences in measured size may be due to many factors, including the choice of different image intensity thresholds (Nat Rev Cardiol. 2010 Oct;7(10):547-9.). Additionally, basic physiology studies have established that the area-at-risk is not simply larger than the region of infarction, but is nearly always transmural regardless of the transmural extent of infarct size. The relationship between the transmural extent of T2-hyperintensity and the transmural extent of infarction is unknown.

Methods

34 canines underwent coronary occlusion of the LAD or the LCx followed by reperfusion. Occlusion time was varied between 45-120 minutes to produce a variable TEI. T2W-TSE and DE-CMR were performed at median of 3 days post MI. A standard T2W-TSE pulse sequence was used with vendor supplied coil normalization (TE 80 ms, slice thickness 7mm, no gap) to obtain short-axis slices of the LV. T2W-TSE and DE-CMR images were separately scored on a per-infarct basis (blinded to the

other technique) for the presence/absence of transmural T2-hyperintensity or transmural hyperenhancement, respectively, at any location within the heart. Additionally, the T2W and DE images were compared using a 6-point scale (0=0%, 1=1-25%, 2=26-50%, 3=51-75%, 4=76-99%, 5=100%) to determine the relative transmural extents.

Results

On a per-infarct basis, 59% (20/34) subjects were scored to have transmural T2-hyperintensity, which was similar to the proportion with transmural hyperenhancement (56%, 19/34). On a per-slice basis (n=244), transmural T2-hyperintensity at any location was found in 30.3% of slices, which was similar to the proportion of transmural hyperenhancement (32.0%, p=0.53. Table 1). Similarly, there was no difference in mean transmural extent score between DE (2.21.7) and T2W-CMR (2.21.7, p=0.38). Representative images from a subendocardial infarct are shown in Figure 1.


Conclusion

T2-hyperintensity is frequently not transmural and is highly associated with the transmural extent of

Table 1

Edema Transmural	HE Transmural	
	No	Yes
No	153	17
Yes	13	61

Duke University, Durham, NC, USA



infarction. This is problematic for the use of T2W-TSE as a representation of the area-at-risk.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-O67

Cite this article as: Kim *et al.*: The relationship of the transmural extent of T2-edema compared with the transmural extent of infarction: implications for the assessment of the area-at-risk. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1):O67.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 **BioMed Central**