

## RESEARCH

## Open Access



# Molecular profiling of advanced solid tumors and patient outcomes with genotype-matched clinical trials: the Princess Margaret IMPACT/COMPACT trial

Tracy L. Stockley<sup>1,2,3</sup>, Amit M. Oza<sup>4,5</sup>, Hal K. Berman<sup>1,2</sup>, Natasha B. Leighl<sup>4,5</sup>, Jennifer J. Knox<sup>4,5</sup>, Frances A. Shepherd<sup>4,5</sup>, Eric X. Chen<sup>4,5</sup>, Monika K. Krzyzanowska<sup>4,5</sup>, Neesha Dhani<sup>4,5</sup>, Anthony M. Joshua<sup>4,5</sup>, Ming-Sound Tsao<sup>1,2,6</sup>, Stefano Serra<sup>1,2</sup>, Blaise Clarke<sup>1,2</sup>, Michael H. Roehrl<sup>1,2,6</sup>, Tong Zhang<sup>1</sup>, Mahadeo A. Sukhai<sup>1</sup>, Nadia Califaretti<sup>7,8</sup>, Mateya Trinkaus<sup>9</sup>, Patricia Shaw<sup>1,2</sup>, Theodorus van der Kwast<sup>1,2</sup>, Lisa Wang<sup>10</sup>, Carl Virtanen<sup>3,11</sup>, Raymond H. Kim<sup>3,4,5</sup>, Albiruni R. A. Razak<sup>4,5</sup>, Aaron R. Hansen<sup>4,5</sup>, Celeste Yu<sup>3</sup>, Trevor J. Pugh<sup>3,6,11</sup>, Suzanne Kamel-Reid<sup>1,2,3,6</sup>, Lillian L. Siu<sup>3,4,5</sup> and Philippe L. Bedard<sup>3,4,5\*</sup>

## Abstract

**Background:** The clinical utility of molecular profiling of tumor tissue to guide treatment of patients with advanced solid tumors is unknown. Our objectives were to evaluate the frequency of genomic alterations, clinical “actionability” of somatic variants, enrollment in mutation-targeted or other clinical trials, and outcome of molecular profiling for advanced solid tumor patients at the Princess Margaret Cancer Centre (PM).

**Methods:** Patients with advanced solid tumors aged  $\geq 18$  years, good performance status, and archival tumor tissue available were prospectively consented. DNA from archival formalin-fixed paraffin-embedded tumor tissue was tested using a MALDI-TOF MS hotspot panel or a targeted next generation sequencing (NGS) panel. Somatic variants were classified according to clinical actionability and an annotated report included in the electronic medical record. Oncologists were provided with summary tables of their patients’ molecular profiling results and available mutation-specific clinical trials. Enrolment in genotype-matched versus genotype-unmatched clinical trials following release of profiling results and response by RECIST v1.1 criteria were evaluated.

**Results:** From March 2012 to July 2014, 1893 patients were enrolled and 1640 tested. After a median follow-up of 18 months, 245 patients (15 %) who were tested were subsequently treated on 277 therapeutic clinical trials, including 84 patients (5 %) on 89 genotype-matched trials. The overall response rate was higher in patients treated on genotype-matched trials (19 %) compared with genotype-unmatched trials (9 %;  $p < 0.026$ ). In a multi-variable model, trial matching by genotype ( $p = 0.021$ ) and female gender ( $p = 0.034$ ) were the only factors associated with increased likelihood of treatment response.

(Continued on next page)

\* Correspondence: [philippe.bedard@uhn.ca](mailto:philippe.bedard@uhn.ca)

<sup>3</sup>Cancer Genomics Program, Princess Margaret Cancer Centre, Toronto, Canada

<sup>4</sup>Division of Medical Oncology and Hematology, Princess Margaret Cancer Centre, 610 University Avenue, Toronto M5G 2M9, Canada

Full list of author information is available at the end of the article



(Continued from previous page)

**Conclusions:** Few advanced solid tumor patients enrolled in a prospective institutional molecular profiling trial were treated subsequently on genotype-matched therapeutic trials. In this non-randomized comparison, genotype-enrichment of early phase clinical trials was associated with an increased objective tumor response rate.

**Trial registration:** NCT01505400 (date of registration 4 January 2012).

**Keywords:** Molecular profiling, DNA sequencing, Clinical trials, Solid tumors, Precision medicine

## Background

Molecular profiling can provide diagnostic, prognostic, or treatment-related information to guide cancer patient management. Advances in next-generation sequencing (NGS) have enabled multiplex testing to overcome the constraints associated with sequential single-analyte testing [1–3]. Large-scale research projects have elucidated genomic landscapes of many cancers but have provided limited insight into the clinical utility of genomic testing. Our aim was to evaluate if targeted DNA profiling improves outcomes for patients assigned to clinical trials based on knowledge of actionable somatic mutations.

At the Princess Margaret Cancer Centre (PM), the Integrated Molecular Profiling in Advanced Cancers Trial (IMPACT) and Community Molecular Profiling in Advanced Cancers Trial (COMPACT) are prospective studies that provide molecular characterization data to oncologists to match patients with advanced solid tumors to clinical trials with targeted therapies. Here, we report the frequency of alterations, clinical “actionability” of the somatic variants, clinical trial enrollment, and outcome based upon molecular profiling results.

## Methods

### Patient cohort

For IMPACT, patients with advanced solid tumors treated at PM were prospectively consented for molecular profiling during a routine clinical visit. For COMPACT, patients with advanced solid tumors treated at other hospitals in Ontario were referred to a dedicated weekly clinic at PM for eligibility review, consent, and blood sample collection. Eligible patients had advanced solid tumors, were aged  $\geq 18$  years, had Eastern Cooperative Oncology Group (ECOG) performance status  $\leq 1$ , and had available formalin-fixed paraffin-embedded (FFPE) archival tumor tissue. The University Health Network Research Ethics Board approved this study (#11-0962-CE). Enrollment for IMPACT began on 1 March 2012 and for COMPACT on 16 November 2012 and ended on 31 July 2014 for this analysis.

### Specimens

DNA was extracted from sections of FFPE tumor specimens from biopsies or surgical resections. If multiple archival tumor specimens were available, the most recent

archival FFPE specimen was reviewed, with a minimum acceptable tumor cellularity of 10 %. Tumor regions were isolated by 1–2  $\times$  1 mm punch from FFPE blocks or manual macrodissection of unstained material from 15–20 slides. FFPE samples were deparaffinized, cells lysed with proteinase K, and DNA extracted using the QIAmp DNA FFPE Tissue Kit (Qiagen, Germantown, MD, USA). DNA was quantified using the Qubit dsDNA Assay kit on the Qubit 2.0 Fluorometer (ThermoFisher Scientific, Waltham, MA, USA).

Participants provided a peripheral blood sample (5 mL in EDTA-coated tubes) as a source of matched germline DNA. DNA was extracted using either standard manual phenol/chloroform extraction methods or automated extraction (MagAttract DNA Mini M48 kit; Qiagen). Patients were offered return of pathogenic germline results at the time of consent and asked to identify a family member delegate who could receive results on their behalf if required.

### Molecular profiling assays

All testing was performed in a laboratory accredited by the College of American Pathologists (CAP) and certified to meet Clinical Laboratory Improvement Amendments (CLIA). Three molecular profiling assays were used over the study period: a custom multiplex genotyping panel on a matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass-spectrometry platform (MassARRAY, Agena Bioscience, San Diego, CA, USA) to genotype 279 mutations within 23 genes (Additional file 1: Table S1); the TruSeq Amplicon Cancer Panel (TSACP, Illumina) on the MiSeq sequencer (Illumina) covering regions of 48 genes (Additional file 1: Table S2); and the Ion AmpliSeq Cancer Panel (ASCP, ThermoFisher Scientific) on the Ion Proton sequencer (ThermoFisher Scientific) covering regions of 50 genes (Additional file 1: Table S3). For more in-depth methodology on molecular profiling assays, including sequence alignment and base calling, see Additional file 1: Supplementary Methods.

### Variant assessment and classification

Variants were assessed and classified according to the classification scheme of Sukhai et al. [4]. Briefly, a five-class scheme was used to sort variants according to actionability (defined as providing information on prognosis,

prediction, diagnosis, or treatment), recurrence of variants in specific tumor sites, and known or predicted deleterious effects on protein function. Interpretation and data integration were performed using Alamut v.2.4.5 (Interactive Biosoftware, Rouen, France). Primary review, assessment, and classification of all variants were independently performed by a minimum of two assessors followed by a third review prior to reporting, with cases where assessors disagreed resolved by group discussion.

### Immunohistochemistry (IHC)

Phosphatase and tensin homolog (PTEN) IHC was performed using rabbit monoclonal Ab 138G6 (Cell Signaling Technology, Danvers, MA, USA) on a Dako platform using a dilution of 1:50 and Flex + 30 protocol. Complete absence of tumor cell staining with positive staining of surrounding tumor stroma fibroblasts/endothelial cells was used to denote PTEN deficiency [5].

### Return of testing results

The molecular profiling report was included in the electronic medical record and returned to the treating oncologist. The clinical significance of profiling results was discussed with PM patients during a routine clinic visit by their treating oncologist. A PM oncologist reviewed results with patients treated at other hospitals by telephone. All oncologists were provided with regular summary tables of testing results and mutation-specific clinical trial listings available at PM. A monthly genomic tumor board was convened at PM to establish consensus treatment recommendations for patients with complex profiling results. A committee consisting of a molecular geneticist, medical geneticist, genetic counselor, and medical oncologist reviewed pathogenic germline variants before return of germline testing results. Germline results were disclosed to the patient or designate by a genetic counselor or medical geneticist.

### Clinical data collection

For each patient, baseline patient and tumor characteristics, treatment regimen(s), time on treatment(s) and survival were retrieved from medical records and updated every three months. Therapeutic clinical trial enrollment was evaluated from the date of reporting molecular profiling results until 9 January 2015. Genotype-matched trials were defined as studies with eligibility criteria restricted to patients with specific somatic mutations, those with a targeted drug with enriched clinical or pre-clinical activity in a patient's genotype, or those with a drug that inhibited a pathway directly linked to the somatic mutation. Decisions about trial enrollment were based upon trial availability, patient or physician preference, and did not follow a pre-specified algorithm.

Targeted lesion measurements and RECIST 1.1 [6] assessments were performed by radiologists.

### Statistics

Descriptive statistics were used to summarize patient characteristics, profiling results, and anti-tumor activity. Comparisons between patients with profiling results treated on genotype-matched and genotype-unmatched trials were performed using a generalized estimating equation (GEE) model [7]. A multi-variable GEE model for response included trial matching by genotype, gender, trial phase, number of lines of prior systemic therapy, investigational agent class, age, tumor type, and sequencing platform. A mixed model was used to compare time on treatment, defined as the date of trial enrollment until the date of discontinuation of investigational treatment. A robust score test was used to compare overall survival following trial enrolment between genotype-matched and genotype-unmatched groups [8]. These comparisons accounted for individual patients who were included on multiple therapeutic trials [8]. Differences with  $p$  values of  $< 0.05$  were considered statistically significant.

## Results

### Patient cohort


A total of 1893 patients were enrolled, including gynecological (23 %), breast (18 %), lung (18 %), colorectal (17 %), pancreatobiliary (8 %), upper aerodigestive (6 %), genitourinary (5 %), and other (5 %) cancers (Table 1). The median age was 59 years (age range, 18–89 years); patients were predominantly female (69 %); had received a median of 2 prior systemic treatments (range, 1–18), and had excellent performance status (43 % PS0 and 56 % PS1). Of 253 (13 %) screen failures, 10 % were for insufficient tissue or DNA and 3 % for clinical deterioration or other reasons. The median follow-up from reporting results was 18 months (range, 1–33 months). A total of 651 (40 %) patients were deceased at the time of the database lock.

### Molecular profiling

Successful molecular profiling was achieved in 1640 patients (87 %), 827 (50 %) had samples tested by MALDI-TOF MS, 792 (48 %) by TSACP, and 21 (1 %) by ASCP (Fig. 1). One or more somatic mutations were detected in 341 (41 %) patients tested by MALDI-TOF MS, 583 (74 %) by TSACP, and 14 (67 %) by ASCP. Median laboratory turnaround time (sample receipt to report) was 32 days (range, 6–228 days). Of patient samples tested by MALDI-TOF MS, *KRAS* (21 %) was the most frequently mutated gene, followed by *PIK3CA* (12 %), with additional genes in the range of 1–5 % frequency. Of samples tested by the TSACP, *TP53* had the highest mutation frequency (47 % of all identified

**Table 1** Characteristics of patients enrolled into IMPACT/COMPACT (n = 1893)

Patients enrolled	Patients enrolled	Patients profiled	Patients profiled enrolled on any therapeutic trials	Patients profiled enrolled on genotype-matched trials
Median age (range)	59 (18–89)	58 (18–89)	58 (18–81)	58 (24–81)
Female/Male	1303/590 (59 %)/(31 %)	1166/479 (71 %)/(29 %)	205/72 (74 %)/(26 %)	64/25 (72 %)/(28 %)
Median lines of prior treatment (range)	2 (1–18)	2 (1–18)	2 (1–16)	2 (1–11)
ECOG performance status (0/1/2)	43 %/56 %/<1 %	44 %/55 %/<1 %	78 %/22 %/0 %	78 %/22 %/0 %
Median time from collection of archival tumor sample to profiling report in years (range)		1.6 (0.1-24.9)	1.7 (0.1-18.9)	1.9 (0.1-18.9)
Primary lesion/Metastatic lesion profiled		1080/560 (66 %)/(34 %)	193/84 (70 %)/(30 %)	66/23 (74 %)/(26 %)
<b>Tumor types</b>				
Breast	341 (18 %)	310 (19 %)	41/310 (13 %)	19/310 (6 %)
Colorectal	326 (17 %)	299 (18 %)	38/299 (13 %)	18/299 (6 %)
Gynecological	430 (23 %)	405 (25 %)	80/405 (20 %)	20/405 (5 %)
Lung	339 (18 %)	256 (16 %)	43/256 (17 %)	18/256 (7 %)
Genitourinary	92 (5 %)	74 (5 %)	9/74 (12 %)	4/74 (5 %)
Pancreatobiliary	151 (8 %)	104 (6 %)	9/104 (9 %)	1/104 (1 %)
Upper aerodigestive	115 (6 %)	102 (6 %)	8/102 (8 %)	2/102 (2 %)
Other	99 (5 %)	81 (5 %)	17/81 (21 %)	2/81 (2 %)
TOTAL	1893	1640	245/1640 (15 %)	84/1640 (5 %)




variants), followed by *KRAS*, *PIK3CA*, and *APC* with mutation frequencies in the range of 5–15 % and the remainder <5 % (Fig. 2). We attribute the difference in mutation landscape between these two platforms to inclusion of *TP53* in the TSACP assay but not in MALDI-TOF (see Additional file 1: Supplemental Methods).

Class 1 and 2 variants are the most clinically significant with known actionability for the specific variant in the tumor site tested (Class 1) or a different tumor site (Class 2) [4]. More than 20 % of patients with breast, colorectal, gynecological, lung, or pancreatobiliary cancers had Class 1 or 2 variants detected by TSACP or MALDI-TOF (Fig. 3). Of patients with genitourinary cancers, only 9 % had actionable variants identified on TSACP and 3 % on MALDI-TOF. For patients with other solid tumors, 25 % had actionable variants identified on TSACP and 18 % on MALDI-TOF. PTEN protein expression was lost by IHC


for 122/788 (15 %) tumors tested. *PTEN* gene mutations were detected by NGS in 14/122 (11.5 %) tumors that were PTEN-negative by IHC.

**Clinical trials and outcomes**

Of the 1640 patients with molecular profiling results, 245 (15 %) were subsequently enrolled in 277 therapeutic clinical trials, including 84 (5 %) treated on 89 genotype-matched trials (Table 2). Patients with pancreatobiliary, upper aerodigestive tract, and other solid tumors were least likely to be treated on genotype-matched trials. Somatic mutations in four genes (*PIK3CA*, *KRAS*, *BRAF*, and *EGFR*) accounted for 76/89 (85 %) of genotype-matched trial enrollments: including *PIK3CA* for breast cancer (20/22); *BRAF* (5/18) and *KRAS* (9/18) for colorectal cancer; *KRAS* (9/18) and *EGFR* (7/18) for non-small cell lung cancer; and *KRAS* (14/22) and *PIK3CA* (7/22) for gynecological cancers. A



**Fig. 2** Mutation frequency by gene from results of (a) MALDI-TOF, n = 827, and (b) TruSeq Amplicon Cancer Panel, n = 792. Mutation frequency was calculated as number of variant occurrences within each gene divided by the total number of patients



complete list of genotype-matched clinical trials by drug class, somatic genotype (variant level), and tumor type are summarized in Table 3.

The age and sex distribution, as well as the number of lines of prior systemic therapy, were similar between the genotype-matched and genotype-unmatched trial patient cohorts (Table 2). There was no difference in the proportion of trials that were genotype-matched between patients profiled on MALDI-TOF MS (61/176 [35 %]) compared with TSACP (28/101 [28 %];  $p = 0.24$ ). A higher proportion of genotype-matched trial patients were treated in phase I studies (81 %) compared with genotype-unmatched trials (46 %;  $p < 0.001$ ). Genotype-matched trial patients were more likely to be treated with targeted drug combinations without chemotherapy or immunotherapy. The overall response rate was higher in patients treated on genotype-matched trials (19 %) compared with genotype-unmatched trials (9 %;  $p = 0.026$ ) (Fig. 4). In

multi-variable analysis, trial matching according to genotype ( $p = 0.021$ ) and female gender ( $p = 0.034$ ) were the only statistically significant factors associated with response (Additional file 1: Table S4). Genotype-matched trial patients were more likely to achieve a best response of any shrinkage in the sum of their target lesions (62 %) compared with genotype-unmatched trial patients (32 %;  $p < 0.001$ ). There was no difference in the time on treatment (15 months versus 15 months;  $p = 0.12$ ) or overall survival (16 months versus 13 months;  $p = 0.10$ ) for patients treated on genotype-matched versus genotype-unmatched trials.

### Germline testing

Of the patients who were asked during consent about return of incidental pathogenic germline mutations, 658/698 (94.3 %) indicated that they wished to receive these results. Two patients were identified with *TP53* variants



**Table 2** Characteristics of patients enrolled in therapeutic trials following molecular profiling

	All trials	Genotype-matched	Genotype-unmatched	<i>p</i> value
Median age (range)	58 (18–81)	58 (24–81)	58.5 (18–80)	NS
Female/Male	205/72	64/25	141/47	NS
Median prior systemic therapies (range)	2 (1–16)	2 (1–11)	2 (1–16)	NS
Tumor type (number of patients)				
Breast	47	22	25	NS
Colorectal	43	18	25	
Lung	48	18	30	
Gynecological	91	22	69	
Other	48	9	39	
Genotyping platform (number of patients)				
MALDI-TOF MS panel	176	61	115	NS
TruSeq Amplicon cancer panel	101	28	73	
Ampliseq cancer panel	0	0	0	
Trial phase (number of patients)				
Phase I	158	72	86	<0.001
Phase II	67	9	58	
Phase III	52	8	44	
Investigational agent(s) (number of patients)				
Targeted monotherapy	112	23	89	<0.001
Targeted drug combination	87	59	28	
Targeted drug and chemotherapy	43	7	36	
Immunotherapy	34	0	34	
Radiotherapy	1	0	1	

NS not significant

in DNA extracted from blood. The first patient was a 36-year-old woman diagnosed with metastatic breast cancer, with a prior papillary thyroid cancer at the age of 28 years, who had a heterozygous germline *TP53* c.817C > T (p.Arg273Cys) pathogenic mutation. Her family history was notable for her mother who died from cancer of unknown primary at the age of 63 years and a maternal aunt with breast cancer at the age of 62 years. The second patient, a 77-year-old woman diagnosed with metastatic cholangiocarcinoma, had no family history of malignancy. We detected a heterozygous *TP53* c.524G > A (p.Arg175His) pathogenic mutation at 15 % allele frequency in the blood that was not present in tumor. This finding is not consistent with inherited Li-Fraumeni syndrome (LFS), but may represent either clonal mosaicism or an age-related or treatment-related mutation limited to blood.

## Discussion

We demonstrated that molecular profiling with mass-spectrometry-based genotyping or targeted NGS can be implemented in a large academic cancer center to identify patients with advanced solid tumors who are candidates for genotype-matched clinical trials. The rapid enrolment to our study reflects the high level of motivation of patients and their oncologists to pursue genomic testing that has been previously reported by our group [9, 10] and others [1, 11–13]. Disappointingly, only 5 % of patients who underwent successful molecular profiling in our study were subsequently treated on genotype-matched clinical trials, consistent with other centers. For comparison, the MD Anderson institutional genomic testing protocol matched 83/2000 (4 %) of patients [1], the SAFIR-01 breast cancer trial matched 28/423 (7 %) [14], and the British Columbia Cancer Agency Personalized Oncogenomics Trial matched 1/100 (1 %) [15]. To facilitate trial accrual, we incorporated multidisciplinary tumor board discussions, physician-directed email alerts with genotype-matched trial listings available at our institution, and individual physician summaries of profiling results. In spite of these efforts, the rate of genotype-matched clinical trial enrolment was low, due to patient deterioration, lack of available clinical trials, and unwillingness of patients to travel for clinical trial participation. There was no difference in proportion of patients treated on genotype-matched trials who underwent profiling using MALDI-TOF or a larger targeted NGS panel. This highlights how few somatic mutations are truly “druggable” through clinical trial matching, even in a large academic cancer center with a broad portfolio of phase I/II trials.

A key finding of our study is that patients in genotype-matched trials were more likely to achieve response than patients in genotype-unmatched trials. Albeit a non-randomized comparison, this finding comprises an important metric and distinguishes our molecular profiling program from other prospective studies that have not tracked longitudinal clinical outcome [1, 16, 17]. Von Hoff and colleagues were the first to report clinical outcome from a prospective molecular profiling (MP) study, with 18/66 (27 %) of patients who received treatment guided by MP data, including RNA-expression profiling and immunohistochemistry (IHC) or fluorescence in situ hybridization (FISH) testing for 11 markers, achieving a progression-free survival (PFS) ratio on MP-selected therapy/PFS on prior therapy) of  $\geq 1.3$  [18]. This study was performed prior to the era of multiplex mutation testing and many patients received MP-guided therapy with cytotoxic therapy using biomarker data that has not been shown to influence treatment response. An analysis of 1114 patients treated on investigational clinical trials at the Clinical Center for Targeted Therapy at MD

**Table 3** Genotype-matched clinical trials by drug class, somatic genotypes (variant level), and tumor type (n = 89)

Tumor type	Somatic genotype (variant)	Genotype-matching trial drug class	Target lesion percent change
Colorectal	No mutations (KRAS wildtype)	EGFR	-100 %
Breast	PIK3CA E545K	PI3K, Endocrine	-90 %
Colorectal	BRAF V600E	BRAF, PI3K, EGFR	-85 %
Colorectal	BRAF V600E TP53 S215G	BRAF, PI3K, EGFR	-81 %
Gynecological	KRAS G13D	PI3K, MEK	-70 %
Gynecological	PIK3CA H1047R KRAS G12D	VEGF	-64 %
Gynecological	KRAS G12V	PI3K, MEK	-63 %
Breast	PIK3CA H1047R	PI3K, IGF1R	-61 %
Lung	EGFR E746_A750del EGFR T790M	EGFR	-58 %
Lung	KRAS G13D	MEK	-54 %
Breast	PIK3CA H1047R	PI3K	-50 %
Lung	EGFR L858R EGFR T790M CTNNB1 S37C	EGFR	-47 %
Gynecological	KRAS G12D	MEK	-47 %
Gynecological	KRAS G12V NRAS Q61R	PI3K, IGF1R	-45 %
Gynecological	KRAS G12A	PI3K, MEK	-38 %
Breast	PIK3CA N345K	ANG2, MTOR	-37 %
Lung	KRAS G12D	PI3K, MEK	-37 %
Gynecological	PIK3CA H1047R KRAS G12D	PI3K, MEK	-37 %
Lung	KRAS G12C	MEK	-28 %
Gynecological	TP53 K132N	WEE1	-26 %
Gynecological	KRAS G12D	PI3K, MEK	-25 %
Colorectal	BRAF V600E TP53 R273C PIK3CA Q546K APC E1544X	BRAF, PI3K, EGFR	-24 %
Breast	ERBB2 D769H PIK3CA N345K	HER2	-23 %
Upper aerodigestive	BRAF V600E	MEK	-20 %
Lung	KRAS G12S	MEK	-20 %
Gynecological	KRAS G12D	MEK, PI3K	-20 %
Colorectal	BRAF V600E TP53 R175H TP53 Q165X	BRAF, PI3K, EGFR	-20 %
Breast	FGFR2 Y376C	FGFR	-19 %
Breast	PIK3CA H1047L	PI3K	-18 %

**Table 3** Genotype-matched clinical trials by drug class, somatic genotypes (variant level), and tumor type (n = 89) (Continued)

Other	GNAQ Q209P	MEK	-18 %
Lung	EGFR L858R	HER3, EGFR	-18 %
Lung	KRAS G12A	PI3K, MEK	-17 %
Lung	EGFR L858R	HER3, EGFR	-17 %
Colorectal	KRAS G12D PIK3CA E545K	PI3K, MEK	-16 %
Gynecological	KRAS G12V	PI3K, MEK	-16 %
Gynecological	BRAF V600E	MEK	-15 %
Gynecological	KRAS G12D	PI3K, MEK	-15 %
Lung	KRAS G12V	PI3K, MEK	-13 %
Colorectal	KRAS G12S	MEK, EGFR	-13 %
Gynecological	NRAS Q61K	PI3K, MEK	-13 %
Pancreatobiliary	KRAS G12V	PI3K, IGF1R	-13 %
Gynecological	PIK3CA H1047R	ANG2, mTOR	-9 %
Breast	PIK3CA E545K	AKT	-7 %
Colorectal	KRAS G12V	MEK, EGFR	-7 %
Breast	ERBB2 D769H PIK3CA N345K	PI3K, IGF1R	-7 %
Lung	KRAS G12V	MEK	-6 %
Colorectal	ERBB2 L755S BRAF N581S ERBB2 L755S APC Q1429fs	VEGF, ANG2	-6 %
Genitourinary	No mutations PTEN negative on IHC	PI3K	-5 %
Lung	EGFR L858R	EGFR	-4 %
Lung	EGFR L858R	EGFR	-4 %
Lung	EGFR E746_A750del	EGFR	-3 %
Breast	PIK3CA H1047R	AKT	-3 %
Upper aerodigestive	KRAS G12V	PI3K, MEK	-2 %
Gynecological	KRAS G12D	PI3K, MEK	-2 %
Colorectal	KRAS G13D	MEK, EGFR	-0.6 %
Genitourinary	No mutations PTEN negative on IHC	PI3K	0 %
Breast	PIK3CA H1047R	PI3K	0 %
Colorectal	PIK3CA E542K KRAS G12V	PI3K	+0.5 %
Colorectal	No Mutations (KRAS wildtype)	MEK, EGFR	+2 %
Colorectal	KRAS G13D PIK3CA E545K	PI3K, MEK	+2 %
Breast	PIK3CA H1047L	PI3K, IGF1R	+4 %
Genitourinary	No mutations	PI3K	+4 %




**Table 3** Genotype-matched clinical trials by drug class, somatic genotypes (variant level), and tumor type (*n* = 89) (Continued)

	<i>PTEN</i> negative on IHC		
Gynecological	<i>PIK3CA</i> E545K	MTOR	+5 %
Breast	<i>PIK3CA</i> N345K	PI3K, IGF1R	+6 %
Breast	<i>PIK3CA</i> N345K	PI3K, MEK	+6 %
	<i>NRAS</i> G12D		
Gynecological	<i>PIK3CA</i> C420R	FGFR, PI3K	+8 %
Colorectal	<i>KRAS</i> G12D	MEK, EGFR	+9 %
Lung	<i>PIK3CA</i> E545K	PI3K, IGF1R	+11 %
Colorectal	No mutations ( <i>KRAS</i> wildtype)	MEK, EGFR	+11 %
Colorectal	<i>KRAS</i> G12D	PI3K, MEK	+12 %
	<i>PIK3CA</i> Q546K		
Gynecological	<i>PIK3CA</i> H1047R	PI3K, IGF1R	+12 %
Breast	<i>PIK3CA</i> p.Glu545Gly	AKT	+28 %
	<i>PTEN</i> p.Leu320X		
	<i>PIK3CA</i> p.Arg93Gln		
Gynecological	<i>TP53</i> R175H	PI3K	+29 %
	<i>PIK3CA</i> R93W		
	<i>FBXW7</i> R479Q		
Colorectal	<i>KRAS</i> G12D	PI3K, MEK	+30 %
	<i>PIK3CA</i> E545K		
Genitourinary	<i>PIK3CA</i> p.Asn345Lys	PI3K	+31 %
Breast	<i>PIK3CA</i> E545K	FGFR, PI3K	+32 %
Breast	<i>PIK3CA</i> H1047R	PI3K, IGF1R	+39 %
Colorectal	No mutations ( <i>KRAS</i> wildtype)	MEK, EGFR	+55 %
Breast	<i>PIK3CA</i> E542K	PI3K, Endocrine	+66 %
Breast	<i>PIK3CA</i> N345K	PI3K, IGF1R	NE
Lung	<i>KRAS</i> G12C	PI3K, MEK	NE
Breast	<i>PIK3CA</i> H1047R	PI3K, IGF1R	NE
Lung	<i>KRAS</i> G12A	MEK	NE
	<i>PIK3CA</i> H1047R		
Gynecological	<i>KRAS</i> G12A	PI3K, IGF1R	NE
Breast	<i>PIK3CA</i> E545K	PI3K	NE
	<i>TP53</i> L252del		
	<i>BRAF</i> c.1315-4C > G		
Gynecological	<i>KRAS</i> G12V	MEK	NE
Lung	<i>BRAF</i> V600E	MEK	NE
Breast	<i>PIK3CA</i> H1047L	PI3K	NE
	<i>TP53</i> C238Y		
Other	<i>KIT</i> V559A	PI3K, EGFR	NE

Anderson Cancer Center reported that the response rate for patients with  $\geq 1$  molecular alteration treated on trials with matched therapy was higher (27 % versus 5 %,  $p < 0.0001$ ) and the time to treatment failure was longer (5.2 versus 3.1 months;  $p < 0.0001$ ) than those who received non-matched therapy [19]. Limitations of this study were that some patients underwent molecular testing after trial assignment and different sequential molecular tests such as polymerase chain reaction-based sequencing, IHC, and FISH, were performed based upon the patient's tumor type.

The same investigators from MD Anderson recently reported the results of their prospective genomic profiling study that enrolled 500 patients with advanced refractory solid tumors assessed in their phase I program [20]. They utilized the FoundationOne™ 236-gene targeted sequencing panel and standard of care biomarker test results (such as ER, PR, and HER2 IHC for breast cancer) to inform treatment selection for commercially available therapies and clinical trial enrollment. A numerically higher rate of prolonged disease control (complete response, partial response, or stable disease  $\geq 6$  months) was observed in patients who received matched therapy (122/500) compared with those who received unmatched therapy (66/500) (19 % versus 8 %,  $p = 0.061$ ). Higher matching scores, calculated based on the number of drug matches and genomic aberrations per patient, were independently associated with a greater frequency of prolonged disease control (22 % [high scores] versus 9 % [low scores],  $p = 0.024$ ), longer time-to-treatment failure (hazard ratio [HR] = 0.52, 95 % confidence interval [CI] = 0.36–0.74,  $p = 0.0003$ ), and survival (HR = 0.65, CI = 0.43–1.0,  $p = 0.05$ ). Likewise, a retrospective review of 347 consecutive patients with advanced solid malignancies treated at the UC San Diego Moores Cancer Center who had targeted sequencing of archival tumor tissue using an earlier version of Foundation One™ (182-gene panel) reported a higher rate of disease control  $\geq 6$  months (34.5 %) for patients (87/342) treated with matched therapy compared with patients (93/342) treated with unmatched therapy [21]. In both of these studies, the rate of treatment matching (25 %) was significantly greater than our study (5 %). This may be due to the use of larger gene panels that include copy number alterations and recurrent translocations that may identify more “drug-gable” alterations for matched therapy; analysis of patient outcomes beyond therapeutic clinical trials that included off-label treatment matching; and varying definitions of genomic alteration and treatment-matching pairs. For instance, the UC San Diego Moores matched therapy cohort included 11 patients (13 %) with breast cancer who received endocrine therapy based on ER expression and 11 patients (13 %) with breast cancer who received HER2-directed



**Fig. 4 a** Waterfall plot of best tumor shrinkage of target lesions by RECIST for patients treated on (a) genotype-matched clinical trials (n = 79) and (b) genotype-unmatched clinical trials (n = 150)

therapy based on *ERBB2* (HER2) amplification. Since ER and HER2 testing are routinely performed in breast cancer patients to guide standard therapies, these patients would not have been included in our matched therapy cohort if the ER and HER2 status were known prior to enrollment in our molecular profiling study.

The only randomized trial that has prospectively assessed the utility of molecular profiling (SHIVA) reported no difference in objective response or PFS for patients treated with genotype-matched versus standard treatments [13]. More than 40 % of patients randomized in the SHIVA trial did not have genomic alterations identified and were included based upon expression of hormone receptors. Patients were matched to a limited range of approved targeted agents following a predefined algorithm that did not include best-in-class investigational agents that are being tested in early phase clinical trials. Despite the negative results of SHIVA, enthusiasm

to conduct genomic-based clinical trials such as NCI-MATCH [12] [NCT02465060], and LUNG-MAP [22] [NCT02154490] remains strong to further define the value of precision medicine. The findings of our study, in which the majority of patients treated on genotype-matched trials were enrolled in phase I targeted therapy trials, are consistent with a recent meta-analysis of phase I trials that demonstrated a higher overall response rate (30.6 % versus 4.9 %,  $p < 0.001$ ) and median PFS (5.7 months versus 2.95 months,  $p < 0.001$ ) for targeted therapy trials that used biomarker-selection compared with those that did not [23].

Measuring the clinical utility of molecular profiling is difficult [3]. We did not comprehensively capture how testing results influenced clinical decisions outside of therapeutic clinical trial enrolment, such as reclassification of tumor subtype and site of primary based on mutation results. For example, we enrolled a patient

with an unknown primary cancer with intra-abdominal metastases that was found to harbor a somatic *IDH1* p.Arg132Cys variant, leading to the reclassification as a likely intrahepatic cholangiocarcinoma. We also did not fully evaluate the use of testing results to avoid ineffective standard treatments (i.e. *KRAS* exon 4 somatic variants in colorectal cancer to inform decision not to use EGFR monoclonal antibody treatment) and treatment with approved targeted agents outside of their approved indications. Few patients in our study received targeted treatments based upon profiling results outside of clinical trials, due to limited access to targeted drugs outside of publicly funded standard-of-care indications in Ontario.

New technological advances are being studied in molecular profiling programs—including larger gene panels [2, 17]; whole exome [16], whole genome (WGS) or RNA sequencing (RNA-Seq) [24, 25]; and integrative systems biology analyses of deregulated cellular pathways [26]. Greater access to clinical trials for genomically characterized patients, such as umbrella and basket trial designs [27], may also improve the success of genotype-treatment matching. To assess whether decision support tools integrated at the point of care can improve enrollment of patients on genotype-matched trials, we are piloting a smart phone application to help physicians identify genotype-matched trials for their patients with profiling data.

There are several limitations of our study. Only a single archival sample was profiled for each patient, often obtained many years prior to molecular testing. Fresh biopsy of a current metastatic lesion for molecular profiling at the time of study enrolment may have yielded different results due to clonal evolution or tumor heterogeneity [28]. Our genomic testing was limited to hotspot point mutation testing or limited targeted sequencing and did not include gene copy number alterations or recurrent translocations that may be important for the selection of genotype-matched therapy. There were patients identified with potentially “druggable” mutations who were candidates for genotype-matched trials; however, they could not be enrolled because of the constraints of slot allocation in early phase clinical trials across multiple institutions or were deemed ineligible due to trial-specific exclusion criteria. Our study population also included many patients with heavily pre-treated metastatic disease who were not well enough for further therapy when results of molecular testing were reported. In addition, tumor response is an imperfect surrogate endpoint to assess therapeutic benefit in early phase clinical trials that should be interpreted with caution [28]. We did not observe a difference in time on treatment or overall survival for patients treated on genotype-matched versus genotype-unmatched clinical trials. PFS data were not available in our cohort precluding a comparison of the outcome of genotype-matched therapy with the

immediate prior line of treatment, as has been reported by other investigators [13, 14, 21].

## Conclusions

We provide preliminary evidence that genotype-matched trial treatment selected on the basis of molecular profiling was associated with increased tumor shrinkage, although only a small proportion of profiled patients benefitted from this approach. Through this initiative, we have created a valuable repository of data and tumor samples that are amenable to additional research and data sharing initiatives. Greater efforts should be made to expand opportunities for genotype-trial matching and further studies are needed to evaluate the clinical utility of targeted NGS profiling.

## Additional file

**Additional file 1:** Supplementary methods and Tables S1–S4. (DOCX 148 kb)

## Acknowledgements

The authors acknowledge Swati Garg, PhD, and Mariam Thomas, PhD, Princess Margaret Cancer Centre, for their contributions to variant data analysis. They are also thankful to all of the medical oncologists, pathologists, laboratory technicians, clinical data coordinators, and correlative studies coordinators who participated in this research study.

## Funding

This work was supported by the Princess Margaret Cancer Foundation; the Cancer Care Ontario Applied Clinical Research Unit [to LLS]; the University of Toronto Division of Medical Oncology Strategic Innovation [to PLB]; and the Ontario Ministry of Health and Long-Term Care Academic Health Sciences Centre Alternate Funding Plan Innovation Award [to PLB].

## Availability of data and materials

The dataset supporting the results of this article is available in the American Association for Cancer Research Project Genomics, Evidence, Neoplasia, Information, Exchange [GENIE] (<http://www.aacr.org/Research/Research/Pages/aacr-project-genie.aspx#.WAWFw2XqeDo>).

## Authors' contributions

TLS and PLB had full access to all of the data in the study and take responsibility for the integrity of the data and accuracy of the data analysis. LLS, PLB, SK-R, and CY conceived of the study concept and wrote the protocol. All authors participated in the acquisition, analysis, or interpretation of data. TS, SK-R, LLS, CY, and PLB drafted the manuscript for initial review by all authors. LW performed statistical analysis. All authors read and approved the final manuscript.

## Authors' information

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## Consent for publication

Not applicable.

## Ethics approval and consent to participate

The University Health Network Research Ethics Board approved this study (#11-0962-CE).

**Author details**

<sup>1</sup>Laboratory Medicine Program, University Health Network, Toronto, Canada. <sup>2</sup>Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Canada. <sup>3</sup>Cancer Genomics Program, Princess Margaret Cancer Centre, Toronto, Canada. <sup>4</sup>Division of Medical Oncology and Hematology, Princess Margaret Cancer Centre, 610 University Avenue, Toronto M5G 2M9, Canada. <sup>5</sup>Department of Medicine, University of Toronto, Toronto, Canada. <sup>6</sup>Department of Medical Biophysics, University of Toronto, Toronto, Canada. <sup>7</sup>Department of Oncology, Grand River Regional Cancer Centre, Kitchener-Waterloo, Canada. <sup>8</sup>Department of Oncology, McMaster University, Faculty of Health Sciences, Hamilton, Canada. <sup>9</sup>Department of Medicine, Markham Stouffville Hospital, Markham, Canada. <sup>10</sup>Department of Biostatistics, Princess Margaret Cancer Centre, Toronto, Canada. <sup>11</sup>Princess Margaret Research Institute, Princess Margaret Cancer Centre, Toronto, Canada.

Received: 19 June 2016 Accepted: 11 October 2016

Published online: 25 October 2016

**References**

- Meric-Bernstam F, Brusco L, Shaw K, Horombe C, Kopetz S, Davies MA, et al. Feasibility of large-scale genomic testing to facilitate enrollment onto genomically matched clinical trials. *J Clin Oncol*. 2015;33:2753–62.
- Lacombe D, Tejpar S, Salgado R, Cardoso F, Golfopoulos V, Aust D, et al. European perspective for effective cancer drug development. *Nat Rev Clin Oncol*. 2014;11(8):492–8.
- Hyman DM, Solit DB. Tumor genetic screening programs: a call to action. *J Clin Oncol*. 2015;33(25):2725–6.
- Sukhai MA, Craddock KJ, Thomas M, Hansen AR, Zhang T, Siu L, et al. A classification system for clinical relevance of somatic variants identified in molecular profiling of cancer. *Genet Med*. 2016;18:128–36.
- Yanagawa N, Leduc C, Kohler D, Saieg MA, John T, Sykes J, et al. Loss of phosphatase and tensin homolog protein expression is an independent poor prognostic marker in lung adenocarcinoma. *J Thorac Oncol*. 2012;7(10):1513–21.
- Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer*. 2009;45(2):228–47.
- Touloumis A, Agresti A, Kateri M. GEE for multinomial responses using a local odds ratios parameterization. *Biometrics*. 2013;69(3):633–40.
- Borgan Ø. (2001), *Modeling Survival Data: Extending the Cox Model*. Terry M. Therneau and Patricia M. Grambsch, Springer-Verlag, New York, 2000. doi:10.1002/sim.956.
- Miller FA, Hayeems RZ, Bytautas JP, Bedard PL, Ernst S, Hirte H, et al. Testing personalized medicine: patient and physician expectations of next-generation genomic sequencing in late-stage cancer care. *Eur J Hum Genet*. 2014;22(3):391–5.
- Blanchette PS, Spreafico A, Miller FA, Chan K, Bytautas J, Kang S, et al. Genomic testing in cancer: Patient knowledge, attitudes, and expectations. *Cancer*. 2014;120(19):3066–73.
- André F, Bachelot T, Commo F, Campone M, Arnedos M, Dieras V, et al. Comparative genomic hybridisation array and DNA sequencing to direct treatment of metastatic breast cancer: a multicentre, prospective trial (SAFIR01/UNICANCER). *Lancet Oncol*. 2014;15(3):267–74.
- Conley BA, Doroshow JH. Molecular analysis for therapy choice: NCI MATCH. In: *Seminars in oncology*; 2014. Elsevier; 2014: p. 297–299.
- Le Tourneau C, Delord J-P, Gonçalves A, Gavoille C, Dubot C, Isambert N, et al. Molecularly targeted therapy based on tumour molecular profiling versus conventional therapy for advanced cancer (SHIVA): a multicentre, open-label, proof-of-concept, randomised, controlled phase 2 trial. *Lancet Oncol*. 2015;16:1324–34.
- Von Hoff DD, Stephenson JJ, Rosen P, Loesch DM, Borad MJ, Anthony S, et al. Pilot study using molecular profiling of patients' tumors to find potential targets and select treatments for their refractory cancers. *J Clin Oncol*. 2010;28(33):4877–83.
- Laskin J, Jones S, Aparicio S, Chia S, Ch'ng C, Deyell R, et al. Lessons learned from the application of whole-genome analysis to the treatment of patients with advanced cancers. *Mol Case Stud*. 2015;1(1):a000570.
- Van Allen EM, Wagle N, Stojanov P, Perrin DL, Cibulskis K, Marlow S, et al. Whole-exome sequencing and clinical interpretation of formalin-fixed, paraffin-embedded tumor samples to guide precision cancer medicine. *Nat Med*. 2014;20(6):682–8.
- Cheng DT, Mitchell TN, Zehir A, Shah RH, Benayed R, Syed A, et al. Memorial Sloan Kettering-Integrated Mutation Profiling of Actionable Cancer Targets (MSK-IMPACT): a hybridization capture-based next-generation sequencing clinical assay for solid tumor molecular oncology. *J Mol Diagn*. 2015;17(3):251–64.
- Von Hoff D, Stephenson Jr J, Rosen P, Loesch D, Borad M, Anthony S, et al. Pilot study using molecular profiling of patients' tumors to find potential targets and select treatments for their refractory cancers. *J Clin Oncol Off J Am Soc Clin Oncol*. 2010;28(33):4877.
- Tsimberidou A-M, Iskander NG, Hong DS, Wheler JJ, Falchook GS, Fu S, et al. Personalized medicine in a phase I clinical trials program: the MD Anderson Cancer Center initiative. *Clin Cancer Res*. 2012;18(22):6373–83.
- Wheler JJ, Janku F, Naing A, Li Y, Stephen B, Zinner RG, et al. Cancer therapy directed by comprehensive genomic profiling: a single center study. *Cancer Res*. 2016;76:3690–701.
- Schwaederle M, Parker BA, Schwab RB, Daniels GA, Piccioni DE, Kesari S, et al. Precision Oncology: The UC San Diego Moores Cancer Center PREDICT Experience. *Mol Cancer Ther*. 2016;15(4):743–52.
- Herbst RS, Gandara DR, Hirsch FR, Redman MW, LeBlanc M, Mack PC, et al. Lung Master Protocol (Lung-MAP)—a biomarker-driven protocol for accelerating development of therapies for squamous cell lung cancer: SWOG S1400. *Clin Cancer Res*. 2015;21(7):1514–24.
- Schwaederle M, Zhao M, Lee JJ, Lazar V, Leyland-Jones B, Schilsky RL, et al. Association of biomarker-based treatment strategies with response rates and progression-free survival in refractory malignant neoplasms: a meta-analysis. *JAMA Oncol*. 2016. DOI: 10.1001/jamaoncol.2016.2129.
- Mody RJ, Wu Y-M, Lonigro RJ, Cao X, Roychowdhury S, Vats P, et al. Integrative clinical sequencing in the management of refractory or relapsed cancer in youth. *JAMA*. 2015;314(9):913–25.
- Roychowdhury S, Iyer MK, Robinson DR, Lonigro RJ, Wu Y-M, Cao X, et al. Personalized oncology through integrative high-throughput sequencing: a pilot study. *Sci Transl Med*. 2011;3(111):111ra121.
- Rodon J, Soria J, Berger R, Batist G, Tsimberidou A, Bresson C, et al. Challenges in initiating and conducting personalized cancer therapy trials: perspectives from WINTHER, a Worldwide Innovative Network (WIN) Consortium trial. *Ann Oncol*. 2015;26:1791–8.
- Slejfer S, Bogaerts J, Siu LL. Designing transformative clinical trials in the cancer genome era. *J Clin Oncol*. 2013;31(15):1834–41.
- Gerlinger M, Rowan AJ, Horswell S, Larkin J, Endesfelder D, Gronroos E, et al. Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *N Engl J Med*. 2012;366(10):883–92.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
www.biomedcentral.com/submit

