

RESEARCH

Open Access

The p38alpha mitogen-activated protein kinase limits the CNS proinflammatory cytokine response to systemic lipopolysaccharide, potentially through an IL-10 dependent mechanism

Adam D Bachstetter¹, Bin Xing^{1,3} and Linda J Van Eldik^{1,2*}

Abstract

Background: The p38α mitogen-activated protein kinase (MAPK) is a well-characterized intracellular kinase involved in the overproduction of proinflammatory cytokines from glia. As such, p38α appears to be a promising therapeutic target for neurodegenerative diseases associated with neuroinflammation. However, the *in vivo* role of p38α in cytokine production in the CNS is poorly defined, and prior work suggests that p38α may be affecting a yet to be identified negative feedback mechanism that limits the acute, injury-induced proinflammatory cytokine surge in the CNS.

Methods: To attempt to define this negative feedback mechanism, we used two *in vitro* and two *in vivo* models of neuroinflammation in a mouse where p38α is deficient in cells of the myeloid lineage.

Results: We found that p38α in myeloid cells has an important role in limiting amplitude of the acute proinflammatory cytokine response to a systemic inflammatory challenge. Moreover, we identified IL-10 as a potential negative feedback mechanism regulated by p38α.

Conclusions: Our data suggest that p38α regulates a proper balance between the pro- and anti-inflammatory cytokine responses to systemic inflammation, and that if circulating IL-10 levels are not elevated to counter-balance the increased systemic proinflammatory responses, the spread of the inflammatory response from the periphery to the CNS is exaggerated.

Keywords: cytokines, glia, interleukin, neuroinflammation, signal transduction, tumor necrosis factor alpha

Introduction

Inflammation is a complex response that requires tight regulatory control of its initiation and resolution. In the central nervous system (CNS), neuroinflammation has gained increasing support as a contributing mechanism in a number of neurodegenerative diseases [1,2], including Alzheimer's disease (AD) [3] and Parkinson's disease [4], as well as acute neurodegenerative conditions such as traumatic brain injury (TBI) [5], spinal cord injury [6] and stroke [7]. As one aspect of neuroinflammation, proinflammatory cytokines have been linked to tissue injury and

disease processes [8,9]. For example, overproduction of IL-1β by activated microglia can induce pathological tau phosphorylation and a decline in synaptophysin levels in neurons [10,11]. Cytokine production is regulated at multiple levels; however, one intracellular mechanism involved in upregulated cytokine production is the p38 mitogen-activated protein kinase (MAPK). The p38 MAPK family consists of at least four isoforms (p38α,β,δ,γ), which are encoded by separate genes, expressed in different tissues and cell types, and are often functionally distinct [12]. Mechanistically, the p38 MAPKs phosphorylate protein substrates that are downstream targets involved in the regulation of proinflammatory cytokine gene expression through effects on transcription and translation (for review see: [12]).

Since the discovery of the pyridinyl imidazole class of small molecule anti-inflammatory compounds that inhibit

* Correspondence: linda.vaneldik@uky.edu

¹Sanders-Brown Center on Aging, University of Kentucky, 800 S. Limestone Street, Lexington, KY 40536, USA

²Department of Anatomy and Neurobiology, University of Kentucky, 800 S. Limestone Street, Lexington, KY 40536, USA

Full list of author information is available at the end of the article

the activity of p38 (commonly known as 'SB' compounds) [13], extensive resources have been allocated to the development and refinement of p38 inhibitors for the treatment of peripheral inflammatory diseases [14,15]. A few studies have used small molecule p38 inhibitors *in vivo* to demonstrate that pharmacological inhibition of p38 α is neuroprotective in animal models of CNS disorders, such as AD [16,17], global and focal ischemia [18-21], neuropathic pain [22], and seizures [23]. However, human clinical trial experience with p38 α inhibitors has yielded mixed results. Published data on the effects of p38 inhibition in cardiovascular disease, chronic obstructive pulmonary disease, and neuropathic pain show encouraging results, whereas trials in rheumatoid arthritis and Crohn's disease show limited, if any, efficacy (for review see: [24]). These results emphasize the importance of a better understanding of the *in vivo* role of p38 α in various inflammatory diseases, especially neuroinflammatory disorders of the CNS.

While p38 α appears promising as a therapeutic target for neurodegenerative diseases [17,25,26], more preclinical work is necessary to determine the appropriate disease indications, therapeutic window, and potential side effects of inhibiting the p38 α pathway. For example, *in vitro* studies have demonstrated that the p38 α isoform contributes to approximately 50% of the IL-1 β and TNF α produced by microglia in response to inflammatory stimuli [27]. In contrast, following a diffuse TBI in mice with a genetic knockdown of p38 α in cells of the myeloid lineage (p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice), an enhanced CNS acute cytokine surge (more IL-1 β , IL-6, TNF α) was found at 6 hr post-injury; however by 7 days post-injury, cytokine levels in the p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice were below that of the injured wild-type (WT) mice [28]. These results suggest that p38 α may be affecting a yet to be identified negative feedback mechanism that limits the acute cytokine surge occurring during the first hours after an *in vivo* insult, which is not evident *in vitro*. The goal of this project was to explore the potential mechanism by which p38 α could suppress the acute proinflammatory cytokine response in the CNS. We report here that the CNS proinflammatory response to systemic lipopolysaccharide (LPS) administration is enhanced in the p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice compared to WT mice, and that p38 α is required by myeloid cells to produce IL-10 in response to LPS. Our results suggest that p38 α has a critical role in limiting the spread of the peripheral inflammatory response to the CNS through regulating the production of the anti-inflammatory cytokine IL-10.

Materials and methods

Reagents

Lipopolysaccharides (LPS) from *Salmonella enterica serotype typhimurium* (Sigma-Aldrich, St. Louis, MO, USA; Cat. no. L6143-1MG; EU/MG of LPS is 600,000)

was prepared in sterile 0.9% sodium chloride that was free of preservatives (Hospira, Inc., Lake Forest, IL; cat. no. NDC 0409-4888-10). Sterile 0.9% sodium chloride was used as the vehicle (veh) control in all experiments.

Animals

Experiments were conducted in accordance with the principles of animal care and experimentation in the Guide For the Care and Use of Laboratory Animals. The Institutional Animal Care and Use Committee of the University of Kentucky approved the use of animals in this study (protocol #2010-0615). The p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice were generated as previously described [29]. The first exon of the p38 α gene (*MAPK14*) was flanked by two loxP sites. The p38 α floxed mice were maintained as homozygotes: p38 $\alpha^{\text{fl/fl}}$. To induce cell-specific deletion of p38 α MAPK in myeloid cells including macrophages and microglia, the p38 $\alpha^{\text{fl/fl}}$ mice were crossed with mice that contain Cre driven by the lysozyme promoter. The restricted cell-type expression of the lysozyme promoter [30,31] results in myeloid cell-specific deletion, as we have previously validated [27,28,32]. The mice were backcrossed to generate mice that were null for mouse WT p38 α , homozygous for floxed p38 α , and hemizygous for LysM-Cre: p38 $\alpha^{\text{fl/fl};\text{LysM-Cre}}$. The p38 $\alpha^{\text{fl/fl}}$ were crossed with the p38 $\alpha^{\text{fl/fl};\text{LysM-Cre}}$ to generate litters where approximately 50% of the mice are microglia/macrophage conditional p38 α KO (p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO) and approximately 50% are p38 $\alpha^{\text{fl/fl}}$ littermates (used as WT controls). All experiments used both genders of mice at approximately a 50:50 ratio. Genotyping was performed by Transnetyx, Inc (Cordova, TN, USA). On the basis of preliminary data and prior published work [27], we find cytokine levels to be consistently below or at the limit of detection in unstimulated mice. Therefore, to reduce the total number of mice needed for the experiments, the number of mice treated with veh alone was kept to the minimum (n = 2 per group/time point) necessary to confirm there was no elevated basal cytokine response due to experimental manipulation or other idiopathic cause.

Lipopolysaccharide administration

For intracerebroventricular (ICV) administration of LPS, three to four month old male and female mice were anesthetized with 5% isoflurane prior to stabilizing the head using ear bars in a digital mouse stereotaxic frame (Stoelting Co, Wood Dale, IL, USA). Anesthesia was maintained with continuous inhalation of isoflurane (3.5%, 1 liter/min). A midline incision was made in the scalp to expose the skull. A hole was drilled into the skull over the right lateral ventricle at the following coordinates: AP = -0.5 mm; ML = -1.0 mm. With a 10- μ l Hamilton syringe with a blunt 28-gauge needle, veh or LPS was injected at the following coordinates: AP = -0.5 mm; ML = -1.0 mm;

DV = -1.8 mm. The Quintessential Stereotaxic Injector (Stoelting Co, Wood Dale, IL, USA) was used to inject 2 μ l at a rate of 0.5 μ l per min. After injection, the needle was left in place for 2 min before being slowly withdrawn. The incision was closed using staples, and the animal was kept on a heat pad until return of normal activity, at which time the mouse was returned to its home cage. Serum and brain tissues were harvested at 6 hr after treatment. Intra-peritoneal (IP) administrations of LPS (5 mg/kg) or veh were done as previously described [27]. Serum and brain tissues were harvested at 1, 6, and 24 hr after treatment.

Tissue collection and processing

Mice were injected with an overdose of sodium pentobarbital (Pentazol powder; Vibrac Animal Health, Ft Worth, TX, USA: cat. no. NDC-051311-103-25). Blood was collected by cardiac puncture, prior to transcardiac perfusion with ice-cold phosphate buffered saline (PBS) for 5 min. The mice were then decapitated. The brain was dissected on ice, snap-frozen in liquid nitrogen, and stored at 80°C until time of use. Brain tissue (cortex or hippocampus) was homogenized using high shear homogenizer, in a 1:10 (w/v) of ice-cold freshly prepared lysis buffer consisting of PBS containing 1 μ g/ml Leupeptin, 1 mM PMSE, and 1 mM EDTA. The tissue homogenate was centrifuged at 14,000 \times g for 20 min at 4°C in a microcentrifuge, and supernatants were collected and stored at -80°C until use.

Microglia and mixed glia cultures

Microglia cultures were prepared as previously described [32]. Briefly, mixed glial cultures (approximately 95% astrocytes, approximately 5% microglia) were prepared from the cerebral cortices of 1 to 3 day old mice. The tissue was trypsinized, and the cells were resuspended in glia complete medium (α -minimum essential medium (α -MEM; Mediatech, Manassas, VA, USA) supplemented with 10% fetal bovine serum (FBS) (US Characterized FBS; Hyclone; Cat no. SH30071.03), 100 IU/ml penicillin, 100 μ g/ml streptomycin, and 2 mM L-Glutamine). After 10 to 14 days in culture, microglia were isolated from the mixed glial cultures by the shake-off procedure [33]. Loosely adherent microglia were shaken off at 250 rpm for 2 hr at 37°C. The cell-containing medium was centrifuged at 180 \times g for 3 min, and the cells were seeded onto 48-well plate at a density of 2×10^4 . Alternatively, the mixed glia cultures were trypsinized and seeded onto a 48-well plate at a density of 2×10^4 . Cells were maintained for 24 hr in glia complete medium. LPS (3 ng/ml) or saline veh was then added directly to the glia complete medium for 30 min, after which the LPS- or veh-containing media was removed, the cells were washed once in glia complete medium, and then maintained for 24 hr in glia complete medium. Aliquots (5 μ l) of the conditioned medium from the primary microglia and the

mixed glia cultures were harvested at different times for cytokine measurements.

Peritoneal macrophages and cortical microglia isolation for gene expression

Peritoneal macrophages were isolated following standard procedures as previously described [34,35]. Briefly, the mice were anesthetized, and 10 ml of PBS was injected into and recollected from the peritoneal cavity. The mice were then transcardiac-perfused with ice-cold PBS. The brain was harvested and the cortex was collected for microglia isolation. Microglia were isolated from the brain using a Percoll gradient following standard procedures as previously described [36]. Briefly, the brain tissue was homogenized in 6.5 ml of ice-cold PBS using a dounce homogenizer. The brain homogenate was added to 3 ml of 100% isotonic Percoll (ISP) in a 15-ml conical tube and mixed by inversion. A 2-ml layer of 50% ISP and 2-ml layer of 70% ISP were layered under the 100% ISP/brain homogenate (30% ISP in solution). The 15-ml conical tube containing the Percoll gradient was centrifuged at 500 \times g for 20 min at room temperature. Microglia were collected at the 70% to 50% ISP interface. Peritoneal macrophages and cortical microglia were washed once in PBS before RNA extraction. RNA was isolated using an RNeasy kit (Qiagen, Valencia, CA, USA) according to the manufacturer's protocols. RNA quantity and quality were determined using A260/A280 readings by NanoDrop (Thermo Scientific). Reverse transcription (RT) was done following the manufacturer's protocol using a High Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Life Technologies, Grand Island, NY, USA). Controls included no template and no RT conditions. Real-time PCR was performed using the TaqMan Gene Expression Assay kit (Applied Biosystems, Life Technologies, Grand Island, NY, USA) according to the manufacturer's instructions on a ViiA™ 7 Real-Time PCR System (Applied Biosystems, Life Technologies, Grand Island, NY, USA). The following TaqMan probes (Applied Biosystems) were used: CCR2 (Mm00438270-m1); CX3CR1 (Mm02620111_s1); MAPK14 (Mm00442497_m1); 18S (Hs99999901_s1). Relative gene expression was calculated by the $2^{-\Delta\Delta CT}$ method.

Cytokine measurements

IL-1ra levels were measured with an R&D Systems DuoSet ELISA kit (R&D Systems, Minneapolis, MN, USA: cat. no. DY480) according to the manufacturer's specifications. All other cytokine levels were measured as previously described [27,37] using Meso Scale Discovery (MSD: Rockville, MD, USA) custom multiplex high-sensitivity ELISA kits according to the manufacturer's instructions. Tissue cytokine levels were normalized to the total amount of protein in the sample, as determined by BCA Protein Assay (Pierce, Rockford, IL, USA).

Statistics

Statistical analysis was conducted using GraphPad prism software V.6 (GraphPad Software, La Jolla, CA, USA: <http://www.graphpad.com>). Comparisons between LPS-stimulated WT and p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice were made by unpaired t-test. Comparisons were not made between naïve, or vehicle-injected mice as the cytokine levels were found to be consistently below or at the limit of detection. Statistical significance was defined as $P < 0.05$. Values are expressed as mean \pm SEM.

Results

The *in vitro* levels of TNF α in response to lipopolysaccharide are reduced in the p38 $\alpha^{\Delta\text{LysM-Cre}}$ knockout mice

We have previously demonstrated using primary microglia cultures that the induction of TNF α in response to TLR activation is governed by p38 α , such that loss or inhibition of p38 α results in less TNF α in response to stimuli [27,32]. However, *in vivo*, we recently found that the role of p38 α in regulating proinflammatory cytokine responses, including TNF α , is more complicated than that predicted from our *in vitro* results. Specifically, using a LysM-Cre mouse line to selectively knock out p38 α in cells of the myeloid lineage (p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice), we found that the acute (6 hr post-injury) proinflammatory cytokine response to a diffuse TBI was enhanced in the KO mice compared to WT controls, even though the chronic cytokine levels were reduced in the p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice [28]. These *in vivo* results suggest that p38 α can both stimulate and unexpectedly limit the CNS proinflammatory cytokine response to injury. However, the mechanism by which p38 α may suppress the acute CNS proinflammatory cytokine response has yet to be identified.

To begin to elucidate the mechanism by which p38 α could limit the CNS proinflammatory cytokine response, we returned to our *in vitro* primary microglia and mixed glia assays. Our prior work measured the levels of TNF α that accumulated in the conditioned media with LPS present in cultures for a minimum of 18 hr. However, our *in vivo* TBI results showed that the acute cytokine response was higher in the p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice at 6 hr post-injury, but lower by 7 d post-injury [28]. We hypothesized that p38 α may have different effects temporally in the TNF α response (that is, an acute spike during the first 6 hr in TNF α levels), which we did not detect in our prior *in vitro* assays when LPS was continuously present, and TNF α was allowed to accumulate in the well for 18 hr. To test this idea, we treated primary microglia cultures with LPS for only 30 min, then removed the media, washed the cells once, and then added fresh medium (without LPS). The purpose of removing the LPS was to eliminate the chronic TLR4 pathway activation, which we know in microglia is partially dependent

on p38 α [27,32]; in this way we could look more selectively at feedback mechanism. We sampled 5 μl of microglia-conditioned media every 2 hr for the first 8 hr and at 24 hr after LPS stimulation, and measured TNF α levels. As shown in Figure 1A, at all time points from 2 to




Figure 1 Microglia and mixed glia cultures from p38 $\alpha^{\Delta\text{LysM-Cre}}$ knockout (KO) are less responsive to lipopolysaccharide (LPS). Wild-type (WT) (white) or p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO (black) primary microglia (A) and mixed glia (B) were stimulated with LPS for 30 min, cells washed, and then conditioned media was harvested at select time points for TNF α Meso Scale Discovery (MSD) assays. The data are representative of three independent experiments. (* $P < 0.05$; *** $P < 0.0005$).

24 hr, significantly lower TNF α levels were found in the p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO primary microglia compared to the WT microglia ($***P < 0.0005$), in agreement with our prior work [27,32].

The primary microglia assays we use are enriched such that approximately 95% of the cells are microglia. Therefore, we postulated that an interaction between astrocytes and microglia might account for the *in vivo* findings, which would not be observed when microglia were in isolation from astrocytes. To test this potential mechanism, we measured the production of TNF α in mixed glia cultures (astrocytes and microglia) treated with LPS and found that the levels of TNF α were also significantly lower in the mixed glia from p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice compared to the WT mice (Figure 1B). Thus, primary microglia or mixed glia cultures derived from p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice show significantly less TNF α induction in response to LPS stimulation compared to WT mice at all time points examined from 2 to 24 hr. Overall, these results do not explain the *in vivo* observation of enhanced acute proinflammatory cytokine response following TBI in p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice.

***In vivo*, p38 α is efficiently deleted in acutely isolated adult peritoneal macrophages, but not in acutely isolated microglia from p38 $\alpha^{\Delta\text{LysM-Cre}}$ knockout mice**

Recently, Butovsky *et al.* (2014) reported the gene expression signature of acutely isolated lymphocytes (B and T cells), myeloid cells (that is, dendritic cells, macrophages, and microglia), as well as astrocytes, oligodendrocytes, and neurons [38]. Microglia were found to express 10 times higher levels of p38 α (MAPK14) compared to astrocytes, oligodendrocytes, or neurons. In addition, microglia expressed two- to threefold more p38 α than other tissue macrophage populations [38]. We previously reported [27,32] that p38 α is efficiently depleted in primary microglia from p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice. However, the LysM-Cre mouse line may not have equal efficiency at knocking down p38 α *in vivo* in all myeloid cell populations. For example, the recombination efficacy of the LysM-Cre in microglia in the adult brain has been shown to be between 20 and 45%, depending on the brain region and the method used to assess the recombination [39]. Since both the relative expression of p38 α and the efficiency of the LysM-Cre promoter can vary in different myeloid cell populations, we measured the level of p38 α suppression in acutely isolated peritoneal macrophages and cortical microglia from WT and p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice.


Using CCR2 and CX3CR1 as markers for macrophages and microglia, respectively, we found high levels of CCR2 in the peritoneal macrophages, and high levels of CX3CR1 in the microglia cell fraction (Figure 2), consistent with previous studies [40]. The gene expression

of CCR2 and CX3CR1 indicated that our isolation resulted in enrichment of the target cell populations. In the WT mice, we confirmed that expression of p38 α was higher in the microglia fraction compared to the macrophage population (Figure 2), as previously reported [38]. Interestingly, in the p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice, macrophages but not microglia were found to have efficient deletion of p38 α (Figure 2). It should be noted that CCR2 and CX3CR1 expression alone does not indicate whether cells other than macrophages or microglia are present in our cell fractions; therefore these results may underestimate the percentage of microglia that may be p38 α deficient.

These findings would appear to be in disagreement with our previous reports that p38 α is efficiently depleted in primary microglia from p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice [27,32]. One potential explanation for the discrepancy is the culture conditions, as primary microglia are activated by removal from the CNS tissue environment. The *in vitro* activation associated with the primary culture could be sufficient to increase the expression of LysM, and the 10 to 14 d of culture would be enough time for the Cre/loxP recombination to result in a loss of p38 α . Therefore, microglia in culture appear to be more similar to peritoneal macrophages than to acutely isolated microglia from adult mice in terms of efficiency of p38 α knockdown.

The proinflammatory cytokine response to intracerebroventricular (ICV) injection of LPS is not suppressed in the p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice

We next postulated that the difference in the acute proinflammatory cytokine response between our *in vivo* diffuse TBI model and our *in vitro* LPS model could be the result of different stressor stimuli. To test this idea, we induced a sterile inflammatory response in the CNS by the injection of 25 ng of LPS into the right lateral ventricle of WT mice and p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice, thus mirroring our *in vitro* LPS assays. Mice were euthanized at 6 hr post-injection, which was determined by a preliminary experiment (data not shown) as the peak of the IL-1 β response in the ipsilateral hippocampus following ICV LPS. Figure 3A shows that levels of three proinflammatory cytokines (IL-1 β , IL-6, and TNF α) were increased in the hippocampus at 6 hr after ICV LPS administration. IL-1 β and TNF α showed no difference between the LPS-stimulated WT and p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice. The only difference between LPS-stimulated WT and p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice in CNS responses was a significantly higher up-regulation of IL-6 ($t_{(14)} = 2.246$ $p = 0.014$) in the cortex, as well as in the serum of p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice compared to WT mice (Figure 3B: ($t_{(14)} = 3.635$ $p = 0.0027$)). We replicated this experiment twice with 100 ng and 1 μg of LPS and found no difference in IL-1 β levels between



$p38\alpha^{\Delta LysM-Cre}$ KO mice compared to WT mice at 6 hr post-injection (data not shown), indicating that the LPS dose was not a critical variable. We previously found that IL-1 β , IL-6 and TNF α were increased both at the protein and mRNA level in the cortex following a diffuse TBI in $p38\alpha^{\Delta LysM-Cre}$ KO mice [28]. Therefore, the ICV LPS model more closely resembled our *in vivo* diffuse TBI model, than our *in vitro* LPS model. However, the ICV LPS model did not mirror the overproduction of IL-1 β and TNF α seen in the *in vivo* diffuse TBI model. Nevertheless, the ICV LPS model did not show a reduced proinflammatory cytokine response in the $p38\alpha^{\Delta LysM-Cre}$ KO mice, as seen in the *in vitro* LPS model. A difference in the stressor stimuli is thus not able to explain the acutely elevated levels of proinflammatory cytokines in the CNS following diffuse TBI in $p38\alpha^{\Delta LysM-Cre}$ KO mice [28]. As microglia in the $p38\alpha^{\Delta LysM-Cre}$ KO mice are not p38 α deficient *in vivo*, the overproduction of IL-6 in response to ICV LPS may come from systemic cells, or circulating factors.


The CNS proinflammatory cytokine response to systemic lipopolysaccharide is enhanced, while the systemic proinflammatory response is suppressed in the $p38\alpha^{\Delta LysM-Cre}$ knockout mice

To test the hypothesis that an altered systemic inflammatory response may account for the increased proinflammatory cytokine response in the CNS following diffuse TBI in $p38\alpha^{\Delta LysM-Cre}$ KO mice, we used an intraperitoneal (IP) LPS model. WT and $p38\alpha^{\Delta LysM-Cre}$ KO mice were injected IP with 5 mg/kg LPS (3×10^6 EU/mg). Additionally, to further explore differences in the temporal patterning of the proinflammatory cytokine response in the WT and $p38\alpha^{\Delta LysM-Cre}$ KO mice, serum and cortical tissue were harvested at different time points after

LPS insult. In agreement with previous studies [29], we found a significantly lower serum cytokine response to LPS in the $p38\alpha^{\Delta LysM-Cre}$ KO mice compared to WT mice (Figure 4A). At 1 hr post-injection, the $p38\alpha^{\Delta LysM-Cre}$ KO mice had significantly less IL-6 ($t_{(8)} = 5.062$ $P = 0.001$) and TNF α ($t_{(8)} = 2.479$ $P = 0.0382$) compared to the WT mice. In agreement with our previous study in the diffuse TBI model [28], we found a significantly elevated peak LPS-induced CNS cytokine response in the $p38\alpha^{\Delta LysM-Cre}$ KO mice compared to the WT mice (Figure 4B). At 6 hr post-injection, levels of IL-1 β ($t_{(16)} = 3.823$ $P = 0.0015$), IL-6 ($t_{(7)} = 3.526$ $P = 0.0097$), and TNF α ($t_{(16)} = 2.328$ $P = 0.0333$) were significantly higher in the $p38\alpha^{\Delta LysM-Cre}$ KO compared to the WT mice. In an independent experiment, we used a lower dose of LPS (1 mg/kg) and still found a significant increase in IL-1 β and IL-6 in the CNS of $p38\alpha^{\Delta LysM-Cre}$ KO mice compared to the WT mice at 6 hr post injection (data not shown). Therefore, the results of the IP LPS model support the hypothesis that systemic cells, or circulating factors dependent on p38 α , are important for limiting the proinflammatory cytokine response in the CNS.

The $p38\alpha^{\Delta LysM-Cre}$ knockout mice fail to produce IL-10 in response to systemic lipopolysaccharide

It has been previously reported that p38 α is involved in the production of two anti-inflammatory cytokines, IL-1ra [41] and IL-10 [42,43]. Therefore, we hypothesized that loss of a paracrine anti-inflammatory response might be responsible for the increased CNS proinflammatory cytokines in the $p38\alpha^{\Delta LysM-Cre}$ KO mice. To test this, we measured serum and cortex levels of IL-1ra and IL-10 after systemic LPS administration to WT and $p38\alpha^{\Delta LysM-Cre}$ KO mice. We found no significant difference in the levels of IL-1ra in the serum between WT and $p38\alpha^{\Delta LysM-Cre}$ KO



mice at any time point measured (Figure 5A). In contrast, we found that the levels of IL-10 in the serum were markedly suppressed in the p38α^{ΔLysM-Cre} KO mice compared to the WT mice following LPS stimulation (Figure 5A). At 1 hr, a highly significant ($t_{(8)} = 6.033$ $P = 0.0003$) suppression of the IL-10 response was seen in the p38α KO mice. The levels of IL-10 in the serum of the p38α^{ΔLysM-Cre} KO mice remained suppressed at the 6 hr ($t_{(8)} = 4.996$ $P = 0.0011$) and 24 hr ($t_{(8)} = 3.727$ $P = 0.0058$) time points compared to the WT mice. In the cortex, there was also


no significant difference in the levels of IL-1ra (Figure 5B) between p38α^{ΔLysM-Cre} KO and WT mice. At 6 hr after LPS, there were lower IL-10 levels in the cortex of p38α^{ΔLysM-Cre} KO mice compared to the WT mice, but the decrease did not reach significance (Figure 5B). Thus, a loss of the systemic IL-10 response in the p38α^{ΔLysM-Cre} KO could be the mechanism by which proinflammatory cytokine response in the CNS becomes exaggerated.

Discussion

The goal of this project was to explore the potential mechanism by which p38α could suppress the acute proinflammatory cytokine response in the CNS. Our prior work, using *in vitro* assays, demonstrated that p38α is critical for the production of IL-1β and TNFα from activated microglia [27,32]. We also previously showed, using a small molecule p38α inhibitor, that suppression of the p38α signal transduction pathway limits the IL-1β response in the CNS to systemic LPS [17,27,44]. Moreover, we found that p38β MAPK is not involved in the CNS cytokine production, or in neurotoxicity induced by LPS inflammatory insult or multiple toxic insults [45,46]. Despite this strong precedent for p38α, we recently found that in a diffuse TBI model the acute proinflammatory cytokine response in the CNS was higher in the p38α^{ΔLysM-Cre} KO mice [28]. Therefore, we sought to determine the mechanism by which p38α could limit the acute proinflammatory response, as this could be critical in the use of p38α as a therapeutic target to treat diseases of the CNS that have inflammation as one component of the pathophysiology.

Collectively, our results suggest that p38α has a critical role in limiting the acute cytokine storm, and the spread of the systemic innate immune response to the CNS, by regulating the production of the anti-inflammatory cytokine IL-10. A peripheral injection of LPS into the peritoneal cavity initiates a systemic innate immune response. There remains a debate of how the peripheral inflammation leads to a neuroinflammatory response. However, acute phase proteins (that is, prostaglandin E2), cytokines (that is, IL-1β) and LPS itself have been suggested to be involved in the mechanism by which a peripheral LPS stimulation induces neuroinflammation [47-51]. In agreement with previous studies [29], we found that levels of the proinflammatory cytokines IL-1β, IL-6, and TNFα were all decreased in the serum of mice where macrophages were deficient in p38α. Despite the decrease in the proinflammatory cytokine response in the serum, the peak of the proinflammatory response in the CNS was significantly higher in the p38α^{ΔLysM-Cre} KO mice. Our data are consistent with a model (Figure 6) whereby systemic LPS induces an acute phase response, including upregulation of both pro- and anti-inflammatory cytokines. In WT mice, the balance of the pro- and anti-inflammatory cytokine

A serum response to IP LPS



B CNS response to IP LPS





Figure 4 Levels of pro-inflammatory cytokines in the serum and cortex of mice following systemic lipopolysaccharide (LPS) stimulation. Wild-type (WT) mice (white) or p38α^{ΔLysM-Cre} knockout (KO) mice (black) were injected intraperitoneally (IP) with 5 mg/kg LPS. The cytokine response in the serum was significantly reduced in the p38α^{ΔLysM-Cre} KO mice (A), while the response in the CNS was increased in the p38α^{ΔLysM-Cre} KO mice compared to the WT mice (B) (n = 5 to 8 per group) (*P < 0.05; **P < 0.01; ***P < 0.001).

A serum response to IP LPS



B CNS response to IP LPS




Figure 5 Levels of anti-inflammatory cytokines in the serum and cortex of mice following systemic lipopolysaccharide (LPS) stimulation. Wild-type (WT) mice (white) or p38α^{ΔLysM-Cre} knockout (KO) mice (black) were injected intraperitoneally (IP) with 5 mg/kg LPS. No significant differences in IL-1ra levels in serum (A) or CNS (B) were detected between WT and p38α^{ΔLysM-Cre} KO mice. Levels of IL-10 in the serum were significantly reduced in the p38α^{ΔLysM-Cre} KO mice (A), and there was a trend for reduced levels of IL-10 in the CNS (B) although the data did not reach statistical significance (n = 5 to 8 per group) (**P < 0.01; ***P < 0.001).




Figure 6 Proposed model for p38 α in myeloid cell cytokine response to systemic lipopolysaccharide (LPS). (A) In wild-type (WT) mice, myeloid cells respond to systemic LPS by producing an acute phase response, including the production of proinflammatory cytokines, as well as the anti-inflammatory cytokine IL-10. The systemic response spreads to the brain where microglia produce cytokines locally in the CNS. (B) In the p38 $\alpha^{\Delta\text{LysM-Cre}}$ knockout (KO) mice, the systemic acute phase response is suppressed such that there is less proinflammatory cytokine production in the blood. However, there is also a loss of the negative feedback provided by the anti-inflammatory IL-10, which we propose leads to an exaggerated acute proinflammatory cytokine response in the CNS.

response, along with other acute phase mechanisms, leads to an increase in cytokines in the CNS (Figure 6A). However, if circulating IL-10 levels are not elevated to counterbalance the increased systemic proinflammatory responses, the spread of the inflammatory response from the periphery to the CNS is exaggerated. This model is also consistent with data from IL-10 deficient mice, in a large number of *in vivo* inflammatory models, which collectively suggest that no other anti-inflammatory cytokine can compensate for the loss of IL-10, because without IL-10, there will be aberrant proinflammatory cytokine overproduction (for review see: [52]). Our results reported here demonstrate that p38 α is required by myeloid cells to produce IL-10 in response to systemic LPS, and without an effective IL-10 response - a defect seen in the p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mouse - neuroinflammation can be exaggerated (Figure 6B).

The systemic balance between pro- and anti-inflammatory factors can influence the proinflammatory cytokine response in the CNS. For example, relevant to the current study, IL-10 is able to block proinflammatory cytokine upregulation in the CNS in response to ICV or IP LPS administration [53,54] and to spinal cord injury [55]. Moreover, IP LPS in IL-10 KO mice results in a significantly exaggerated TNF α , IL-1 β and IL-6 response in the plasma and in the CNS compared to WT mice [56], following a similar pattern to our results. Both microglia and astrocytes express the IL-10 receptor; however, *in vitro* IL-10 was found to have a suppressive effect only on astrocytes and not microglia [57]. The effect of IL-10 on microglia was found to be indirect through the astrocyte production of TGF β [57]. We do not know if this mechanism would also be true in our model, but understanding the crosstalk between microglia and astrocytes in regulating the balance between pro- and anti-inflammatory factors is an important future direction.

The broad anti-inflammatory properties of IL-10 are a result of IL-10's ability to inhibit innate function of macrophages, neutrophils, and dendritic cells including their production of proinflammatory cytokines [52,58]. The present study confirmed previous observations [43,59] that macrophages require p38 α for the production of IL-10, which suggests that the induction of this anti-inflammatory cytokine may limit the CNS inflammatory response to LPS. This finding is in agreement with the role of mitogen- and stress-activated kinases 1 and 2 (MSK1/2) in the production of IL-10 [42,43]. MSK1/2 is activated by p38 and Erk1/2 [60]. Microglia and macrophages deficient in p38 α have been shown to have decreased levels of the activated phospho-MSK1/2 [27,42,43]. Critically, MSK1/2 double KO mice are found to have elevated proinflammatory cytokine responses to LPS, which is IL-10 dependent, as illustrated elegantly by the use of an MSK1/2 / IL-10 triple KO mouse [42]. In addition to the MSK1/2 pathway, p38 α can also regulate IL-10 expression in macrophages via the direct downstream target of p38 α , the MAPK-activated protein kinase 2 [61]. Defining why Erk1/2 is not able to compensate for the loss of p38 α will require future studies to understand potential crosstalk between p38 α and Erk1/2 in regulating IL-10 levels following *in vivo* LPS stimulation.

Conclusions

We and others have previously reported using small molecule inhibitors that suppression of p38 α can reduce neuroinflammation *in vivo* [16,17,19,23,27]. In the current study, we found that the peak proinflammatory cytokine response to systemic LPS *in vivo* is significantly greater in the CNS of p38 $\alpha^{\Delta\text{LysM-Cre}}$ KO mice compared to the WT mice, replicating the result found in a diffuse TBI model [28]. We also found that after systemic stimulation of the

innate immune system with LPS, p38 α was necessary for a stimulus-induced increase in serum levels of the anti-inflammatory cytokine IL-10. Therefore, p38 α appears to be a fulcrum in balancing the systemic innate immune response, by regulating levels of pro- and anti-inflammatory cytokines. While p38 α appears to be a promising therapeutic target, our work here highlights the complexity of such efforts. The observations that p38 α may balance the inflammatory response by acutely attenuating the early proinflammatory cytokine surge, while perpetuating the chronic microglia activation response raise the importance of considerations of an optimal therapeutic window for the efficacy of p38 α inhibitors. Our findings that p38 α -dependent systemic inflammatory responses affect the CNS inflammatory responses also suggest that measurement of systemic IL-10 and TNF α levels in serum or plasma might be a useful tool for monitoring the anti- and pro-inflammatory cytokine balance during the course of treatment of CNS disorders.

Abbreviations

AD: Alzheimer's disease; CNS: central nervous system; ICSV: intracerebroventricular; IP: intraperitoneal; IL-1: interleukin-1; IL-1ra: IL-1 receptor antagonist; ISP: isotonic Percoll; KO: knockout; LPS: lipopolysaccharide; MAPK: mitogen-activated protein kinase; MSD: Meso Scale Discovery; MSK1/2: mitogen-and-stress-activated kinases 1 and 2; RT: reverse transcription; TLR: toll-like receptor; TNF α : tumor necrosis factor alpha; veh: vehicle; WT: wild type.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

ADB and LVE designed the research studies. ADB and BX performed the experiments. ADB and LVE drafted the manuscript with the assistance of the other authors. All authors read and approved the final manuscript.

Acknowledgements

We thank Edgardo Dimayuga and Danielle Goulding for their assistance with various aspects of this work. This research was supported in part by an Alzheimer's Association Zenith award ZEN-09-134506 (LVE) and NIH grants R01 NS064247 (LVE), F32 AG037280 (ADB), and K99 AG044445 (ADB). We are grateful to Dr. Huiping Jiang at Boehringer Ingelheim Pharmaceuticals, Inc. and Dr. Jiahuai Han at The Scripps Research Institute for the kind gifts of the knockout mice.

Author details

¹Sanders-Brown Center on Aging, University of Kentucky, 800 S. Limestone Street, Lexington, KY 40536, USA. ²Department of Anatomy and Neurobiology, University of Kentucky, 800 S. Limestone Street, Lexington, KY 40536, USA. ³Current Affiliation: VA Pittsburgh Healthcare System, 3501 Fifth Avenue, Pittsburgh, PA 15260, USA.

Received: 24 August 2014 Accepted: 29 September 2014

Published online: 10 October 2014

References

- Mrak RE, Griffin WST: Glia and their cytokines in progression of neurodegeneration. *Neurobiol Aging* 2005, **26**:349–354.
- Glass CK, Saijo K, Winner B, Marchetto MC, Gage FH: Mechanisms underlying inflammation in neurodegeneration. *Cell* 2010, **140**:918–934.
- Wyss-Coray T, Rogers J: Inflammation in Alzheimer disease—a brief review of the basic science and clinical literature. *Cold Spring Harb Perspect Med* 2012, **2**:a006346.
- Tansey MG, Goldberg MS: Neuroinflammation in Parkinson's disease: its role in neuronal death and implications for therapeutic intervention. *Neurobiol Dis* 2010, **37**:510–518.
- Das M, Mohapatra S, Mohapatra SS: New perspectives on central and peripheral immune responses to acute traumatic brain injury. *J Neuroinflammation* 2012, **9**:236.
- David S, Kroner A: Repertoire of microglial and macrophage responses after spinal cord injury. *Nat Rev Neurosci* 2011, **12**:388–399.
- Iadecola C, Anrather J: The immunology of stroke: from mechanisms to translation. *Nat Med* 2011, **17**:796–808.
- Van Eldik LJ, Thompson WL, Ralay Ranaivo H, Behanna HA, Martin Watterson D: Glia proinflammatory cytokine upregulation as a therapeutic target for neurodegenerative diseases: function-based and target-based discovery approaches. *Int Rev Neurobiol* 2007, **82**:277–296.
- McCoy MK, Tansey MG: TNF signaling inhibition in the CNS: implications for normal brain function and neurodegenerative disease. *J Neuroinflammation* 2008, **5**:45.
- Li Y, Liu L, Barger SW, Griffin WS: Interleukin-1 mediates pathological effects of microglia on tau phosphorylation and on synaptophysin synthesis in cortical neurons through a p38-MAPK pathway. *J Neurosci* 2003, **23**:1605–1611.
- Sheng JG, Jones RA, Zhou XQ, McGinness JM, Van Eldik LJ, Mrak RE, Griffin WS: Interleukin-1 promotion of MAPK-p38 overexpression in experimental animals and in Alzheimer's disease: potential significance for tau protein phosphorylation. *Neurochem Int* 2001, **39**:341–348.
- Bachstetter AD, Van Eldik LJ: The p38 MAP kinase family as regulators of proinflammatory cytokine production in degenerative diseases of the CNS. *Aging Dis* 2010, **1**:199–211.
- Lee JC, Laydon JT, McDonnell PC, Gallagher TF, Kumar S, Green D, McNulty D, Blumenthal MJ, Heys JR, Landvatter SW, Strickler JE, Mclaughlin MM, Siemens IR, Fisher SM, Livi GP, White JR, Adams JL, Young PR: A protein kinase involved in the regulation of inflammatory cytokine biosynthesis. *Nature* 1994, **372**:739–746.
- Margutti S, Laufer SA: Are MAP kinases drug targets? Yes, but difficult ones. *Chem Med Chem* 2007, **2**:1116–1140.
- Hammaker D, Firestein GS: "Go upstream, young man": lessons learned from the p38 saga. *Ann Rheum Dis* 2010, **69**(Suppl 1):i77–i82.
- Munoz L, Ranaivo HR, Roy SM, Hu W, Craft JM, McNamara LK, Chico LW, Van Eldik LJ, Watterson DM: Novel p38 alpha MAPK inhibitor suppresses brain proinflammatory cytokine up-regulation and attenuates synaptic dysfunction and behavioral deficits in an Alzheimer's disease mouse model. *J Neuroinflammation* 2007, **4**:21.
- Watterson DM, Grum-Tokars VL, Roy SM, Schavocky JP, Bradaric BD, Bachstetter AD, Xing B, Dimayuga E, Saeed F, Zhang H, Stanislawski A, Pelletier JC, Minasov G, Anderson WF, Arancio O, Van Eldik LJ: Development of novel chemical probes to address CNS protein kinase involvement in synaptic dysfunction. *PLoS One* 2013, **8**:e66226.
- Sugino T, Nozaki K, Takagi Y, Hattori I, Hashimoto N, Moriguchi T, Nishida E: Activation of mitogen-activated protein kinases after transient forebrain ischemia in gerbil hippocampus. *J Neurosci* 2000, **20**:4506–4514.
- Piao CS, Kim JB, Han PL, Lee JK: Administration of the p38 MAPK inhibitor SB203580 affords brain protection with a wide therapeutic window against focal ischemic insult. *J Neurosci Res* 2003, **73**:537–544.
- Legos JJ, Erhardt JA, White RF, Lenhard SC, Chandra S, Parsons AA, Tuma RF, Barone FC: SB 239063, a novel p38 inhibitor, attenuates early neuronal injury following ischemia. *Brain Res* 2001, **892**:70–77.
- Barone FC, Irving EA, Ray AM, Lee JC, Kassiss S, Kumar S, Badger AM, White RF, McVey MJ, Legos JJ, Erhardt JA, Nelson AH, Ohlstein EH, Hunter AJ, Ward K, Smith BR, Adams JL, Parsons AA: SB 239063, a second-generation p38 mitogen-activated protein kinase inhibitor, reduces brain injury and neurological deficits in cerebral focal ischemia. *J Pharmacol Exp Ther* 2001, **296**:312–321.
- Jin SX, Zhuang ZY, Woolf CJ, Ji RR: p38 mitogen-activated protein kinase is activated after a spinal nerve ligation in spinal cord microglia and dorsal root ganglion neurons and contributes to the generation of neuropathic pain. *J Neurosci* 2003, **23**:4017–4022.
- Kim SW, Yu YM, Piao CS, Kim JB, Lee JK: Inhibition of delayed induction of p38 mitogen-activated protein kinase attenuates kainic acid-induced neuronal loss in the hippocampus. *Brain Res* 2004, **1007**:188–191.

24. Fisk M, Gajendragadkar PR, Maki-Petaja KM, Wilkinson IB, Cheriyan J: **Therapeutic potential of p38 MAP kinase inhibition in the management of cardiovascular disease.** *Am J Cardiovasc Drugs* 2014, **14**:155–165.
25. Huang G, Wang Y, Vogel P, Kanneganti TD, Otsu K, Chi H: **Signaling via the kinase p38alpha programs dendritic cells to drive TH17 differentiation and autoimmune inflammation.** *Nat Immunol* 2012, **13**:152–161.
26. Munoz L, Ammit AJ: **Targeting p38 MAPK pathway for the treatment of Alzheimer's disease.** *Neuropharmacology* 2010, **58**:561–568.
27. Bachstetter AD, Xing B, de Almeida L, Dimayuga ER, Watterson DM, Van Eldik LJ: **Microglial p38alpha MAPK is a key regulator of proinflammatory cytokine up-regulation induced by toll-like receptor (TLR) ligands or beta-amyloid (Abeta).** *J Neuroinflammation* 2011, **8**:79.
28. Bachstetter AD, Rowe RK, Kaneko M, Goulding D, Lifshitz J, Van Eldik LJ: **The p38alpha MAPK regulates microglial responsiveness to diffuse traumatic brain injury.** *J Neurosci* 2013, **33**:6143–6153.
29. Kang YJ, Chen J, Otsuka M, Mols J, Ren S, Wang Y, Han J: **Macrophage deletion of p38alpha partially impairs lipopolysaccharide-induced cellular activation.** *J Immunol* 2008, **180**:5075–5082.
30. Clausen BE, Burkhardt C, Reith W, Renkawitz R, Forster I: **Conditional gene targeting in macrophages and granulocytes using LysMcre mice.** *Transgenic Res* 1999, **8**:265–277.
31. Clarke S, Greaves DR, Chung LP, Tree P, Gordon S: **The human lysozyme promoter directs reporter gene expression to activated myelomonocytic cells in transgenic mice.** *Proc Natl Acad Sci U S A* 1996, **93**:1434–1438.
32. Xing B, Bachstetter AD, Eldik LJ: **Microglial p38alpha MAPK is critical for LPS-induced neuron degeneration, through a mechanism involving TNFalpha.** *Mol Neurodegener* 2011, **6**:84.
33. Petrova TV, Akama KT, Van Eldik LJ: **Cyclopentenone prostaglandins suppress activation of microglia: Down-regulation of inducible nitric-oxide synthase by 15-deoxy-Delta(12,14)-prostaglandin J(2).** *Proc Natl Acad Sci U S A* 1999, **96**:4668–4673.
34. Davies JQ, Gordon S: **Isolation and culture of murine macrophages.** *Methods Mol Biol* 2005, **290**:91–103.
35. Zhang X, Goncalves R, Mosser DM: **The isolation and characterization of murine macrophages.** *Curr Protoc Immunol* 2008, Chapter 14:Unit 14.1. doi:10.1002/0471142735.im1401s83.
36. Pino PA, Cardona AE: **Isolation of brain and spinal cord mononuclear cells using Percoll gradients.** *J Vis Exp* 2011, **48**:2348.
37. Bachstetter AD, Norris CM, Sompol P, Wilcock DM, Goulding D, Neltner JH, St Clair D, Watterson DM, Van Eldik LJ: **Early stage drug treatment that normalizes proinflammatory cytokine production attenuates synaptic dysfunction in a mouse model that exhibits age-dependent progression of Alzheimer's disease-related pathology.** *J Neurosci* 2012, **32**:10201–10210.
38. Butovsky O, Jedrychowski MP, Moore CS, Cialic R, Lanser AJ, Gabriely G, Koeglsperger T, Dake B, Wu PM, Doykan CE, Fanek Z, Liu L, Chen Z, Rothstein JD, Ransohoff RM, Gygi SP, Antel JP, Weiner HL: **Identification of a unique TGF-beta-dependent molecular and functional signature in microglia.** *Nat Neurosci* 2014, **17**:131–143.
39. Goldmann T, Wieghofer P, Müller PF, Wolf Y, Varol D, Yona S, Brendecke SM, Kierdorf K, Staszewski O, Datta M, Luedde T, Heikenwalder M, Jung S, Prinz M: **A new type of microglia gene targeting shows TAK1 to be pivotal in CNS autoimmune inflammation.** *Nat Neurosci* 2013, **16**:1618–1626.
40. Mizutani M, Pino PA, Saederup N, Charo IF, Ransohoff RM, Cardona AE: **The fractalkine receptor but not CCR2 is present on microglia from embryonic development throughout adulthood.** *J Immunol* 2012, **188**:29–36.
41. Darragh J, Ananieva O, Courtney A, Elcombe S, Arthur JS: **MSK1 regulates the transcription of IL-1ra in response to TLR activation in macrophages.** *Biochem J* 2010, **425**:595–602.
42. Ananieva O, Darragh J, Johansen C, Carr JM, McIlrath J, Park JM, Wingate A, Monk CE, Toth R, Santos SG, Iversen L, Arthur JS: **The kinases MSK1 and MSK2 act as negative regulators of Toll-like receptor signaling.** *Nat Immunol* 2008, **9**:1028–1036.
43. Kim C, Sano Y, Todorova K, Carlson BA, Arpa L, Celada A, Lawrence T, Otsu K, Brissette JL, Arthur JS, Park JM: **The kinase p38 alpha serves cell type-specific inflammatory functions in skin injury and coordinates pro- and anti-inflammatory gene expression.** *Nat Immunol* 2008, **9**:1019–1027.
44. Bachstetter AD, Watterson DM, Van Eldik LJ: **Target engagement analysis and link to pharmacodynamic endpoint for a novel class of CNS-penetrant and efficacious p38alpha MAPK inhibitors.** *J Neuroimmune Pharmacol* 2014, **9**:454–460.
45. Xing B, Bachstetter AD, Van Eldik LJ: **Deficiency in p38beta MAPK fails to inhibit cytokine production or protect neurons against inflammatory insult in in vitro and in vivo mouse models.** *PLoS One* 2013, **8**:e56852.
46. Xing B, Bachstetter AD, Van Eldik LJ: **Inhibition of neuronal p38alpha, but not p38beta MAPK, provides neuroprotection against three different neurotoxic insults.** *J Mol Neurosci* 2014, [Epub ahead of print].
47. Chakravarty S, Herkenham M: **Toll-like receptor 4 on nonhematopoietic cells sustains CNS inflammation during endotoxemia, independent of systemic cytokines.** *J Neurosci* 2005, **25**:1788–1796.
48. Engström L, Ruud J, Eskilsson A, Larsson A, Mackerlova L, Kugelberg U, Qian H, Vasilache AM, Larsson P, Engblom D, Sigvardsson M, Jönsson JI, Blomqvist A: **Lipopolysaccharide-induced fever depends on prostaglandin E2 production specifically in brain endothelial cells.** *Endocrinology* 2012, **153**:4849–4861.
49. Chen Z, Jalabi W, Shpargel KB, Farabaugh KT, Dutta R, Yin X, Kidd GJ, Bergmann CC, Stohlman SA, Trapp BD: **Lipopolysaccharide-induced microglial activation and neuroprotection against experimental brain injury is independent of hematogenous TLR4.** *J Neurosci* 2012, **32**:11706–11715.
50. Quan N: **Immune-to-brain signaling: How important are the blood-brain barrier-independent pathways?** *Mol Neurobiol* 2008, **37**:142–152.
51. Laflamme N, Lacroix S, Rivest S: **An essential role of interleukin-1 beta in mediating NF-kappa B activity and COX-2 transcription in cells of the blood-brain barrier in response to a systemic and localized inflammation but not during endotoxemia.** *J Neurosci* 1999, **19**:10923–10930.
52. Saraiva M, O'Garra A: **The regulation of IL-10 production by immune cells.** *Nat Rev Immunol* 2010, **10**:170–181.
53. Bluthé RM, Castanon N, Pousset F, Bristow A, Ball C, Lestage J, Michaud B, Kelley KW, Dantzer R: **Central injection of IL-10 antagonizes the behavioural effects of lipopolysaccharide in rats.** *Psychoneuroendocrinology* 1999, **24**:301–311.
54. Di Santo E, Adami M, Bertorelli R, Ghezzi P: **Systemic interleukin 10 administration inhibits brain tumor necrosis factor production in mice.** *Eur J Pharmacol* 1997, **336**:197–202.
55. Bethea JR, Nagashima H, Acosta MC, Briceno C, Gomez F, Marcillo AE, Looor K, Green J, Dietrich WD: **Systemically administered interleukin-10 reduces tumor necrosis factor-alpha production and significantly improves functional recovery following traumatic spinal cord injury in rats.** *J Neurotrauma* 1999, **16**:851–863.
56. Krzyszton CP, Sparkman NL, Grant RW, Buchanan JB, Broussard SR, Woods J, Johnson RW: **Exacerbated fatigue and motor deficits in interleukin-10-deficient mice after peripheral immune stimulation.** *Am J Physiol Regul Integr Comp Physiol* 2008, **295**:R1109–R1114.
57. Norden DM, Fenn AM, Dugan A, Godbout JP: **TGFbeta produced by IL-10 redirected astrocytes attenuates microglial activation.** *Glia* 2014, **62**:881–895.
58. Pils MC, Pisano F, Fasnacht N, Heinrich JM, Groebe L, Schippers A, Rozell B, Jack RS, Müller W: **Monocytes/macrophages and/or neutrophils are the target of IL-10 in the LPS endotoxemia model.** *Eur J Immunol* 2010, **40**:443–448.
59. Guma M, Hammaker D, Topolewski K, Corr M, Boyle DL, Karin M, Firestein GS: **Antiinflammatory functions of p38 in mouse models of rheumatoid arthritis: advantages of targeting upstream kinases MKK-3 or MKK-6.** *Arthritis Rheum* 2012, **64**:2887–2895.
60. Arthur JS: **MSK activation and physiological roles.** *Front Biosci* 2008, **13**:5866–5879.
61. Ehling C, Ronkina N, Böhmer O, Albrecht U, Bode KA, Lang KS, Kotlyarov A, Radzioch D, Gaestel M, Häussinger D, Bode JG: **Distinct functions of the mitogen-activated protein kinase-activated protein (MAPKAP) kinases MK2 and MK3: MK2 mediates lipopolysaccharide-induced signal transducers and activators of transcription 3 (STAT3) activation by preventing negative regulatory effects of MK3.** *J Biol Chem* 2011, **286**:24113–24124.

doi:10.1186/s12974-014-0175-6

Cite this article as: Bachstetter et al.: The p38alpha mitogen-activated protein kinase limits the CNS proinflammatory cytokine response to systemic lipopolysaccharide, potentially through an IL-10 dependent mechanism. *Journal of Neuroinflammation* 2014 **11**:175.