

Hindawi Publishing Corporation  
Boundary Value Problems  
Volume 2011, Article ID 750769, 18 pages  
doi:10.1155/2011/750769

## Research Article

# A Quasilinear Parabolic System with Nonlocal Boundary Condition

**Botao Chen,<sup>1</sup> Yongsheng Mi,<sup>1,2</sup> and Chunlai Mu<sup>2</sup>**

<sup>1</sup> College of Mathematics and Computer Sciences, Yangtze Normal University, Fuling, Chongqing 408100, China

<sup>2</sup> College of Mathematics and Physics, Chongqing University, Chongqing 401331, China

Correspondence should be addressed to Chunlai Mu, chunlaimu@yahoo.com.cn

Received 8 May 2010; Revised 25 July 2010; Accepted 11 August 2010

Academic Editor: Daniel Franco

Copyright © 2011 Botao Chen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the blow-up properties of the positive solutions to a quasilinear parabolic system with nonlocal boundary condition. We first give the criteria for finite time blowup or global existence, which shows the important influence of nonlocal boundary. And then we establish the precise blow-up rate estimate. These extend the recent results of Wang et al. (2009), which considered the special case  $m_1 = m_2 = 1, p_1 = 0, q_2 = 0$ , and Wang et al. (2007), which studied the single equation.

## 1. Introduction

In this paper, we deal with the following degenerate parabolic system:

$$u_t = \Delta u^{m_1} + u^{p_1} v^{q_1}, \quad v_t = \Delta v^{m_2} + v^{p_2} u^{q_2}, \quad x \in \Omega, \quad t > 0 \quad (1.1)$$

with nonlocal boundary condition

$$u(x, t) = \int_{\Omega} f(x, y) u(y, t) dy, \quad v(x, t) = \int_{\Omega} g(x, y) v(y, t) dy, \quad x \in \partial\Omega, \quad t > 0, \quad (1.2)$$

and initial data

$$u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), \quad x \in \bar{\Omega}, \quad (1.3)$$

where  $m_i, p_i, q_i > 1, i = 1, 2$ , and  $\Omega \subset R^N$  is a bounded connected domain with smooth boundary.  $f(x, y) \neq 0$  and  $g(x, y) \neq 0$  for the sake of the meaning of nonlocal boundary are nonnegative continuous functions defined for  $x \in \partial\Omega$  and  $y \in \overline{\Omega}$ , while the initial data  $v_0, u_0$  are positive continuous functions and satisfy the compatibility conditions  $u_0(x) = \int_{\Omega} f(x, y)u_0(y)dy$  and  $v_0(x) = \int_{\Omega} g(x, y)v_0(y)dy$  for  $x \in \partial\Omega$ , respectively.

Problem (1.1)–(1.3) models a variety of physical phenomena such as the absorption and “downward infiltration” of a fluid (e.g., water) by the porous medium with an internal localized source or in the study of population dynamics (see [1]). The solution  $(u(x, t), v(x, t))$  of the problem (1.1)–(1.3) is said to blow up in finite time if there exists  $T \in (0, \infty)$  called the blow-up time such that

$$\lim_{t \rightarrow T^-} \left( \|u(\cdot, t)\|_{L^\infty(\Omega)} + \|v(\cdot, t)\|_{L^\infty(\Omega)} \right) = +\infty, \quad (1.4)$$

while we say that  $(u(x, t), v(x, t))$  exists globally if

$$\sup_{t \in (0, T)} \left( \|u(\cdot, t)\|_{L^\infty(\Omega)} + \|v(\cdot, t)\|_{L^\infty(\Omega)} \right) < +\infty \quad \text{for any } T \in (0, \infty). \quad (1.5)$$

Over the past few years, a considerable effort has been devoted to the study of the blow-up properties of solutions to parabolic equations with local boundary conditions, say Dirichlet, Neumann, or Robin boundary condition, which can be used to describe heat propagation on the boundary of container (see the survey papers [2, 3] and references therein). The semilinear case ( $m_1 = m_2 = 1, f \equiv 0, g \equiv 0$ ) of (1.1)–(1.3) has been deeply investigated by many authors (see, e.g., [2–11]). The system turns out to be degenerate if  $m_i > 1 (i = 1, 2)$ ; for example, in [12, 13], Galaktionov et al. studied the following degenerate parabolic equations:

$$\begin{aligned} u_t &= \Delta u^{m_1} + v^{q_1}, & v_t &= \Delta v^{m_2} + u^{p_2}, & (x, t) &\in \Omega \times (0, T), \\ u(x, t) &= v(x, t) = 0, & (x, t) &\in \partial\Omega \times (0, T), \\ u(x, 0) &= u_0(x), & v(x, 0) &= v_0(x), & x &\in \overline{\Omega} \end{aligned} \quad (1.6)$$

with  $m_1 > 1, m_2 > 1, p_2 > 1$ , and  $q_1 > 1$ . They obtained that solutions of (1.6) are global if  $p_2 q_1 < m_1 m_2$ , and may blow up in finite time if  $p_2 q_1 > m_1 m_2$ . For the critical case of  $p_2 q_1 = m_1 m_2$ , there should be some additional assumptions on the geometry of  $\Omega$ .

Song et al. [14] considered the following nonlinear diffusion system with  $m_1 \geq 1, m_2 \geq 1$  coupled via more general sources:

$$\begin{aligned} u_t &= \Delta u^{m_1} + u^{p_1} v^{q_1}, & v_t &= \Delta v^{m_2} + u^{p_2} v^{q_2}, & (x, t) &\in \Omega \times (0, T), \\ u(x, t) &= v(x, t) = \varepsilon_0 > 0, & (x, t) &\in \partial\Omega \times (0, T), \\ u(x, 0) &= u_0(x), & v(x, 0) &= v_0(x), & x &\in \overline{\Omega}. \end{aligned} \quad (1.7)$$

Recently, the genuine degenerate situation with zero boundary values for (1.7) has been discussed by Lei and Zheng [15]. Clearly, problem (1.6) is just the special case by taking  $p_1 = q_2 = 0$  in (1.7) with zero boundary condition.

For the more parabolic problems related to the local boundary, we refer to the recent works [16–20] and references therein.

On the other hand, there are a number of important phenomena modeled by parabolic equations coupled with nonlocal boundary condition of form (1.2). In this case, the solution could be used to describe the entropy per volume of the material (see [21–23]). Over the past decades, some basic results such as the global existence and decay property have been obtained for the nonlocal boundary problem (1.1)–(1.3) in the case of scalar equation (see [24–28]). In particular, in [28], Wang et al. studied the following problem:

$$\begin{aligned} u_t &= \Delta u^m + u^p, & (x, t) \in \Omega \times (0, t), \\ u(x, t) &= \int_{\Omega} f(x, y)u(y, t)dy, & (x, t) \in \partial\Omega \times (0, t), \\ u(x, 0) &= u_0(x), & x \in \bar{\Omega}, \end{aligned} \quad (1.8)$$

with  $m > 1, p > 1$ . They obtained the blow-up condition and its blow-up rate estimate. For the special case  $m = 1$  in the system (1.8), under the assumption that  $\int_{\Omega} f(x, y)dy = 1$ , Seo [26] established the following blow-up rate estimate:

$$(p-1)^{-1/(p-1)}(T-t)^{-1/(p-1)} \leq \max_{x \in \bar{\Omega}} u(x, t) \leq C_1(T-t)^{-1/(\gamma-1)}, \quad (1.9)$$

for any  $\gamma \in (1, p)$ . For the more nonlocal boundary problems, we also mention the recent works [29–34]. In particular, Kong and Wang in [29], by using some ideas of Souplet [35], obtained the blow-up conditions and blow-up profile of the following system:

$$u_t = \Delta u + \int_{\Omega} u^m(x, t)v^n(x, t)dx, \quad v_t = \Delta v + \int_{\Omega} u^p(x, t)v^q(x, t)dx, \quad x \in \Omega, t > 0 \quad (1.10)$$

subject to nonlocal boundary (1.2), and Zheng and Kong in [34] gave the condition for global existence or nonexistence of solutions to the following similar system:

$$u_t = \Delta u + u^m \int_{\Omega} v^n(x, t)dx, \quad v_t = \Delta v + v^q \int_{\Omega} u^p(x, t)dx, \quad x \in \Omega, t > 0 \quad (1.11)$$

with nonlocal boundary condition (1.2). The typical characterization of systems (1.10) and (1.11) is the complete couple of the nonlocal sources, which leads to the analysis of simultaneous blowup.

Recently, Wang and Xiang [30] studied the following semilinear parabolic system with nonlocal boundary condition:

$$\begin{aligned} u_t - \Delta u &= v^p, & v_t - \Delta v &= u^q, & x \in \Omega, & t > 0, \\ au(x, t) &= \int_{\Omega} f(x, y)u(y, t)dy, & v(x, t) &= \int_{\Omega} g(x, y)v(y, t)dy, & x \in \partial\Omega, & t > 0, \\ u(x, 0) &= u_0, & v(x, 0) &= v_0, & x \in \Omega, \end{aligned} \quad (1.12)$$

where  $p$  and  $q$  are positive parameters. They gave the criteria for finite time blowup or global existence, and established blow-up rate estimate.

To our knowledge, there is no work dealing with the parabolic system (1.1) with nonlocal boundary condition (1.2) except for the single equation case, although this is a very classical model. Therefore, the main purpose of this paper is to understand how the reaction terms, the weight functions and the nonlinear diffusion affect the blow-up properties for the problem (1.1)–(1.3). We will show that the weight functions  $f(x, y), g(x, y)$  play substantial roles in determining blowup or not of solutions. Firstly, we establish the global existence and finite time blow-up of the solution. Secondly, we establish the precise blowup rate estimates for all solutions which blow up.

Our main results could be stated as follows.

**Theorem 1.1.** *Suppose that  $\int_{\Omega} f(x, y)dy \geq 1, \int_{\Omega} g(x, y)dy \geq 1$  for any  $x \in \partial\Omega$ . If  $q_2 > p_1 - 1$  and  $q_1 > p_2 - 1$  hold, then any solution to (1.1)–(1.3) with positive initial data blows up in finite time.*

**Theorem 1.2.** *Suppose that  $\int_{\Omega} f(x, y)dy < 1, \int_{\Omega} g(x, y)dy < 1$  for any  $x \in \partial\Omega$ .*

- (1) *If  $m_1 > p_1, m_2 > p_2$ , and  $q_1q_2 < (m_1 - p_1)(m_2 - p_2)$ , then every nonnegative solution of (1.1)–(1.3) is global.*
- (2) *If  $m_1 < p_1, m_2 < p_2$  or  $q_1q_2 > (m_1 - p_1)(m_2 - p_2)$ , then the nonnegative solution of (1.1)–(1.3) exists globally for sufficiently small initial values and blows up in finite time for sufficiently large initial values.*

To establish blow-up rate of the blow-up solution, we need the following assumptions on the initial data  $u_0(x), v_0(x)$

(H1)  $u_0(x), v_0(x) \in C^{2+\mu}(\Omega) \cap (\overline{\Omega})$  for some  $0 < \mu < 1$ ;

(H2) There exists a constant  $\delta \geq \delta_0 > 0$ , such tha

$$\Delta u_0^{m_1} + u_0^{p_1} v_0^{q_1} - \delta u_0^{m_1 k_1 + 1}(x) \geq 0, \Delta v_0^{m_2} + v_0^{p_2} u_0^{q_2} - \delta v_0^{m_2 k_2 + 1}(x) \geq 0, \quad (1.13)$$

where  $\delta_0, k_1$ , and  $k_2$  will be given in Section 4.

**Theorem 1.3.** *Suppose that  $\int_{\Omega} f(x, y)dy \leq 1, \int_{\Omega} g(x, y)dy \leq 1$  for any  $x \in \partial\Omega$ ;  $q_1 > m_2, q_2 > m_1$  and satisfy  $q_2 > p_1 - 1$  and  $q_1 > p_2 - 1$ ; assumptions (H1)–(H2) hold. If the solution  $(u, v)$*

of (1.1)–(1.3) with positive initial data  $u_0, v_0$  blows up in finite time  $T_*$ , then there exist constants  $C_i > 0$  ( $i = 1, 2, 3, 4$ ) such that

$$\begin{aligned} & C_1(T_* - t)^{-(q_1 - p_2 + 1)/(q_2 q_1 - (1 - p_1)(1 - p_2))} \\ & \leq \max_{x \in \bar{\Omega}} u(x, t) \leq C_2(T_* - t)^{-(q_1 - p_2 + 1)/(q_2 q_1 - (1 - p_1)(1 - p_2))}, \quad \text{for } 0 < t < T_*, \\ & C_3(T_* - t)^{-(q_2 - p_1 + 1)/(q_2 q_1 - (1 - p_1)(1 - p_2))} \\ & \leq \max_{x \in \bar{\Omega}} v(x, t) \leq C_4(T_* - t)^{-(q_2 - p_1 + 1)/(q_2 q_1 - (1 - p_1)(1 - p_2))}, \quad \text{for } 0 < t < T_*. \end{aligned} \quad (1.14)$$

This paper is organized as follows. In the next section, we give the comparison principle of the solution of problem (1.1)–(1.3) and some important lemmas. In Section 3, we concern the global existence and nonexistence of solution of problem (1.1)–(1.3) and show the proofs of Theorems 1.1 and 1.2. In Section 4, we will give the estimate of the blow-up rate.

## 2. Preliminaries

In this section, we give some basic preliminaries. For convenience, we denote that  $Q_T = Q \times (0, T)$ ,  $S_T = \partial\Omega \times (0, T)$  for  $0 < T < +\infty$ . As it is now well known that degenerate equations need not possess classical solutions, we begin by giving a precise definition of a weak solution for problem (1.1)–(1.3).

*Definition 2.1.* A vector function  $(u(x, t), v(x, t))$  defined on  $\bar{\Omega}_T$ , for some  $T > 0$ , is called a sub (or super) solution of (1.1)–(1.3), if all the following hold:

- (1)  $u(x, t), v(x, t) \in L^\infty(\Omega_T)$ ;
- (2)  $(u(x, t), v(x, t)) \leq (\geq) (\int_\Omega f(x, t)u(y, t)dy, \int_\Omega g(x, y)v(y, t)dy)$  for  $(x, t) \in S_T$ , and  $u(x, 0) \leq (\geq) u_0(x), v(x, 0) \leq (\geq) v_0(x)$  for almost all  $x \in \Omega$ ;
- (3)

$$\begin{aligned} \int_\Omega u(x, t)\phi(x, t)dx & \leq (\geq) \int_\Omega u(x, 0)\phi(x, 0)dx + \int_0^t \int_{\Omega_\tau} (u\phi_\tau + u^{m_1}\Delta\phi + u^{p_1}v^{q_1}\phi)dx d\tau \\ & \quad - \int_0^t \int_{\partial\Omega} \frac{\partial\phi}{\partial n} \left( \int_\Omega f(x, y)u(y, \tau)dy \right)^{m_1} dS d\tau, \\ \int_\Omega v(x, t)\phi(x, t)dx & \leq (\geq) \int_\Omega v(x, 0)\phi(x, 0)dx + \int_0^t \int_{\Omega_\tau} (v\phi_\tau + v^{m_2}\Delta\phi + v^{p_2}u^{q_2}\phi)dx d\tau \\ & \quad - \int_0^t \int_{\partial\Omega} \frac{\partial\phi}{\partial n} \left( \int_\Omega g(x, y)v(y, \tau)dy \right)^{m_2} dS d\tau, \end{aligned} \quad (2.1)$$

where  $n$  is the unit outward normal to the lateral boundary of  $\Omega_T$ . For every  $t \in [0, T]$  and any  $\phi$  belong to the class of test functions,

$$\Phi \equiv \left\{ \phi \in C(\overline{\Omega_T}); \phi_t, \Delta \phi \in C(\Omega_T) \cap L^2(\Omega_T); \phi \geq 0, \phi(x, t)|_{\partial\Omega \times (0, T)} = 0 \right\}. \quad (2.2)$$

A weak solution of (1.1) is a vector function which is both a subsolution and a supersolution of (1.1)-(1.3).

**Lemma 2.2** (Comparison principle). *Let  $(\underline{u}, \underline{v})$  and  $(\overline{u}, \overline{v})$  be a subsolution and supersolution of (1.1)-(1.3) in  $Q_T$ , respectively. Then  $(\underline{u}, \underline{v}) \leq (\overline{u}, \overline{v})$  in  $\overline{\Omega_T}$ , if  $(\underline{u}(x, 0), \underline{v}(x, 0)) \leq (\overline{u}(x, 0), \overline{v}(x, 0))$ .*

*Proof.* Let  $\phi(x, t) \in \Phi$ , the subsolution  $(\underline{u}, \underline{v})$  satisfies

$$\begin{aligned} \int_{\Omega} \underline{u}(x, t) \phi(x, t) dx &\leq \int_{\Omega} \underline{u}(x, 0) \phi(x, 0) dx + \int_0^t \int_{\Omega_T} (\underline{u} \phi_\tau + \underline{u}^{m_1} \Delta \phi + \underline{u}^{p_1} \underline{v}^{q_1} \phi) dx d\tau \\ &\quad - \int_0^t \int_{\partial\Omega} \frac{\partial \phi}{\partial n} \left( \int_{\Omega} f(x, y) \underline{u}(y, \tau) dy \right)^{m_1} dS d\tau. \end{aligned} \quad (2.3)$$

On the other hand, the supersolution  $(\overline{u}, \overline{v})$  satisfies the reversed inequality

$$\begin{aligned} \int_{\Omega} \overline{u}(x, t) \phi(x, t) dx &\geq \int_{\Omega} \overline{u}(x, 0) \phi(x, 0) dx + \int_0^t \int_{\Omega_T} (\overline{u} \phi_\tau + \overline{u}^{m_1} \Delta \phi + \overline{u}^{p_1} \overline{v}^{q_1} \phi) dx d\tau \\ &\quad - \int_0^t \int_{\partial\Omega} \frac{\partial \phi}{\partial n} \left( \int_{\Omega} f(x, y) \overline{u}(y, \tau) dy \right)^{m_1} dS d\tau. \end{aligned} \quad (2.4)$$

Set  $\omega(x, t) = \underline{u}(x, t) - \overline{u}(x, t)$ , we have

$$\begin{aligned} \int_{\Omega} \omega(x, t) \phi(x, t) dx &\leq \int_{\Omega} \omega(x, 0) \phi(x, 0) dx + \int_0^t \int_{Q_T} (\phi_\tau + \Theta_1(x, s) \Delta \phi + \Theta_2(x, s) \phi \overline{v}^{q_1}) \omega dx d\tau \\ &\quad + \int_0^t \int_{\Omega} \phi \underline{u}^{p_1} (\Theta_3(\underline{v} - \overline{v})) dx d\tau \\ &\quad - \int_0^t \int_{\partial\Omega} \frac{\partial \phi}{\partial n} m_1 \xi^{m_1-1} \left( \int_{\Omega} f(x, y) \omega(y, \tau) dy \right) dS d\tau, \quad t \in (0, T), \end{aligned} \quad (2.5)$$

where

$$\begin{aligned} c\Theta_1(x, t) &\equiv \int_0^1 m_1 (\theta \underline{u} + (1 - \theta) \overline{u})^{m_1-1} d\theta, & \Theta_2(x, t) &\equiv \int_0^1 p_1 (\theta \underline{v} + (1 - \theta) \overline{v})^{p_1-1} d\theta, \\ & & \Theta_3(x, t) &\equiv \int_0^1 q_1 (\theta \underline{v} + (1 - \theta) \overline{v})^{q_1-1} d\theta. \end{aligned} \quad (2.6)$$

Since  $(\underline{u}, \underline{v})$  and  $(\bar{u}, \bar{v})$  are bounded in  $\Omega_T$ , it follows from  $m_1 > 1$ ,  $q_1, p_1 \geq 1$  that  $\Theta_i (i = 1, 2, 3)$  are bounded nonnegative functions.  $\xi$  is a function between  $\int_{\Omega} f(x, y) \underline{u}(x, \tau) dy$  and  $\int_{\Omega} f(x, y) \bar{u}(x, \tau) dy$ . Noticing that  $(\bar{u}, \bar{v})$  and  $(\underline{u}, \underline{v})$  are nonnegative bounded function and  $\partial\phi/\partial n \leq 0$  on  $\partial\Omega$ , we choose appropriate function  $\phi$  as in [36] to obtain that

$$\begin{aligned} \int_{\Omega} \omega(x, t)_+ dx &\leq C_1 \int_{\Omega} \omega(x, 0)_+ dx + C_2 \int_0^t \int_{\Omega} \omega(y, \tau)_+ dy d\tau \\ &+ C_3 \int_0^t \int_{\Omega} [\underline{v} - \bar{v}]_+ dx d\tau \quad (\text{using } \omega(x, 0) = \underline{u}(x, 0) - \bar{u}(x, 0) \leq 0). \end{aligned} \quad (2.7)$$

By Gronwall's inequality, we know that  $\omega(x, t) = \underline{u}(x, t) - \bar{u}(x, t) \leq 0$ ,  $\underline{v}(x, t) \leq \bar{v}(x, t)$  can be obtained in similar way, then  $(\bar{u}, \bar{v}) \geq (\underline{u}, \underline{v})$ .

Local in time existence of positive classical solutions of the problem (1.1)–(1.3) can be obtained using fixed point theorem (see [37]), the representation formula and the contraction mapping principle as in [38]. By the above comparison principle, we get the uniqueness of the solution to the problem. The proof is more or less standard, so is omitted here.  $\square$

*Remark 2.3.* From Lemma 2.2, it is easy to see that the solution of (1.1)–(1.3) is unique if  $p_1, p_2, q_1, q_2 > 1$ .

The following comparison lemma plays a crucial role in our proof which can be obtained by similar arguments as in [24, 38–40]

**Lemma 2.4.** *Suppose that  $w_1(x, t), w_2(x, t) \in C^{2,1}(\Omega_T) \cap C(\bar{\Omega}_T)$  and satisfy*

$$\begin{aligned} w_{1t} - d_1(x, t)\Delta w_1 &\geq c_{11}(x, t)w_1 + c_{21}(x, t)w_2(x, t), \quad (x, t) \in \Omega \times (0, T), \\ w_{2t} - d_2(x, t)\Delta w_2 &\geq c_{12}(x, t)w_2 + c_{22}(x, t)w_1(x, t), \quad (x, t) \in \Omega \times (0, T), \\ w_1(x, t) &\geq \int_{\Omega} c_{13}(x, y)w_1(y, t)dy, \quad (x, t) \in \partial\Omega \times (0, T), \\ w_2(x, t) &\geq \int_{\Omega} c_{23}(x, y)w_2(y, t)dy, \quad (x, t) \in \partial\Omega \times (0, T), \\ w_1(x, 0) &\geq 0, \quad w_2(x, 0) \geq 0, \quad x \in \Omega, \end{aligned} \quad (2.8)$$

where  $c_{ij}(x, t) (i = 1, 2; j = 1, 2, 3)$  are bounded functions and  $d_i(x, t) > 0 (i = 1, 2)$ ,  $c_{2j}(x, t) \geq 0, (x, t) \in \Omega \times (0, T)$ , and  $c_{i3}(x, y) \geq 0 (i = 1, 2), (x, y) \in \partial\Omega \times \Omega$  and is not identically zero. Then  $w_i(x, 0) > 0 (i = 1, 2)$  for  $x \in \bar{\Omega}$  imply that  $w_i(x, t) > 0 (i = 1, 2)$  in  $\Omega_T$ . Moreover, if  $c_{i3}(x, y) \equiv 0 (i = 1, 2)$  or if  $\int_{\Omega} c_{i3}(x, y)dy \leq 1, x \in \partial\Omega$ , then  $w_i(x, 0) \geq 0 (i = 1, 2)$  for  $x \in \bar{\Omega}$  imply that  $w_i(x, t) \geq 0$  in  $\Omega_T$ .

Denote that

$$A = \begin{pmatrix} m_1 - p_1 & -q_1 \\ -q_2 & m_2 - p_2 \end{pmatrix}, \quad l = \begin{pmatrix} l_1 \\ l_2 \end{pmatrix}. \quad (2.9)$$

We give some lemmas that will be used in the following section. Please see [41] for their proofs.

**Lemma 2.5.** *If  $m_1 > p_1, m_2 > p_2$ , and  $q_1 q_2 < (m_1 - p_1)(m_2 - p_2)$ , then there exist two positive constants  $l_1, l_2$ , such that  $Al = (1, 1)^T$ . Moreover,  $A(cl) > (0, 0)^T$  for any  $c > 0$ .*

**Lemma 2.6.** *If  $m_1 < p_1, m_2 < p_2$  or  $q_1 q_2 > (m_1 - p_1)(m_2 - p_2)$ , then there exist two positive constants  $l_1, l_2$ , such that  $Al < (0, 0)^T$ . Moreover,  $A(cl) < (0, 0)^T$  for any  $c > 0$ .*

### 3. Global Existence and Blowup in Finite Time

Compared with usual homogeneous Dirichlet boundary data, the weight functions  $f(x, y)$  and  $g(x, y)$  play an important role in the global existence or global nonexistence results for problem (1.1)–(1.3).

*Proof of Theorem 1.1.* We consider the ODE system

$$\begin{aligned} F'(t) &= F^{p_1} H^{q_1}(t), & H'(t) &= H^{p_2} F^{q_2}(t), & t > 0, \\ F(0) &= a > 0, & H(0) &= b > 0, \end{aligned} \quad (3.1)$$

where  $a = (1/2)\min_{\bar{\Omega}} u_0(x)$ ,  $b = (1/2)\min_{\bar{\Omega}} v_0(x)$ , and we use the assumption  $u_0, v_0 > 0$ .

Set

$$\begin{aligned} F_0 &= \left( \frac{(q_2 - p_1 + 1)^{q_1} (q_1 - p_2 + 1)^{1-p_2}}{(q_1 q_2 - (p_1 - 1)(p_2 - 1))^{q_1 - p_2 + 1}} \right)^{1/(q_1 q_2 - (p_1 - 1)(p_2 - 1))} \\ &\quad \times (T_1 - t)^{-(q_1 - p_2 + 1)/(q_1 q_2 - (p_1 - 1)(p_2 - 1))}, \\ H_0 &= \left( \frac{(q_1 - p_2 + 1)^{q_2} (q_2 - p_1 + 1)^{1-p_1}}{(q_1 q_2 - (p_1 - 1)(p_2 - 1))^{q_2 - p_1 + 1}} \right)^{(1/q_1 q_2 - (p_1 - 1)(p_2 - 1))} \\ &\quad \times (T_2 - t)^{-(q_2 - p_1 + 1)/(q_1 q_2 - (p_1 - 1)(p_2 - 1))}, \end{aligned} \quad (3.2)$$

with

$$\begin{aligned} T_1 &= a^{-(q_1 q_2 - (p_1 - 1)(p_2 - 1))/(q_1 - p_2 + 1)} \left( \frac{(q_2 - p_1 + 1)^{q_1} (q_1 - p_2 + 1)^{1-p_2}}{(q_1 q_2 - (p_1 - 1)(p_2 - 1))^{q_1 - p_2 + 1}} \right)^{1/(q_1 - p_2 + 1)}, \\ T_2 &= b^{-(q_1 q_2 - (p_1 - 1)(p_2 - 1))/(q_2 - p_1 + 1)} \left( \frac{(q_1 - p_2 + 1)^{q_2} (q_2 - p_1 + 1)^{1-p_1}}{(q_1 q_2 - (p_1 - 1)(p_2 - 1))^{q_2 - p_1 + 1}} \right)^{1/(q_2 - p_1 + 1)}. \end{aligned} \quad (3.3)$$

It is easy to check that  $(F_0, H_0)$  is the unique solution of the ODE problem (3.1), then  $q_2 > p_1 - 1$  and  $q_1 > p_2 - 1$  imply that  $(F_0, H_0)$  blows up in finite time. Under the assumption that  $\int_{\Omega} f(x, y) dy \geq 1$ ,  $\int_{\Omega} g(x, y) dy \geq 1$  for any  $x \in \partial\Omega$ ,  $(F_0, H_0)$  is a subsolution of problem



(1.1)–(1.3). Therefore, by Lemma 2.2, we see that the solution  $(u, v)$  of problem (1.1)–(1.3) satisfies  $(u, v) \geq (F_0, H_0)$  and then  $(u, v)$  blows up in finite time.  $\square$

*Proof of Theorem 1.2.* (1) Let  $\Psi_1(x)$  be the positive solution of the linear elliptic problem

$$-\Delta \Psi_1(x) = \epsilon_1, \quad x \in \Omega, \quad \Psi_1(x) = \int_{\Omega} f(x, y) dy, \quad x \in \partial\Omega, \quad (3.4)$$

and  $\Psi_2(x)$  be the positive solution of the linear elliptic problem

$$-\Delta \Psi_2(x) = \epsilon_2, \quad x \in \Omega, \quad \Psi_2(x) = \int_{\Omega} g(x, y) dy, \quad x \in \partial\Omega, \quad (3.5)$$

where  $\epsilon_1, \epsilon_2$  are positive constant such that  $0 \leq \Psi_1(x) \leq 1, 0 \leq \Psi_2(x) \leq 1$ . We remark that  $\int_{\Omega} f(x, y) dy < 1$  and  $\int_{\Omega} g(x, y) dy < 1$  ensure the existence of such  $\epsilon_1, \epsilon_2$ .

Denote that

$$\max_{\Omega} \Psi_1 = \bar{K}_1, \quad \min_{\Omega} \Psi_1 = \underline{K}_1; \quad \max_{\Omega} \Psi_2 = \bar{K}_2, \quad \min_{\Omega} \Psi_2 = \underline{K}_2. \quad (3.6)$$

We define the functions  $\bar{u}, \bar{v}$  as following:

$$\bar{u}(x, t) = \bar{u}(x) = M^{l_1} \Psi_1^{1/m_1}, \quad \bar{v}(x, t) = \bar{v}(x) = M^{l_2} \Psi_2^{1/m_2}, \quad (3.7)$$

where  $M$  is a constant to be determined later. Then, we have

$$\begin{aligned} \bar{u}(x, t) |_{x \in \partial\Omega} &= M^{l_1} \Psi_1^{1/m_1} = M^{l_1} \left( \int_{\Omega} f(x, y) dy \right)^{1/m_1} \\ &> M^{l_1} \int_{\Omega} f(x, y) dy \geq M^{l_1} \int_{\Omega} f(x, t) \Psi_1^{1/m_1}(y) dy = \int_{\Omega} f(x, y) \bar{u}(y) dy. \end{aligned} \quad (3.8)$$

In a similar way, we can obtain that

$$|\bar{v}(x, t)|_{x \in \partial\Omega} > \int_{\Omega} g(x, y) \bar{v}(y) dy, \quad (3.9)$$

here, we used  $0 \leq \Psi_1(x) \leq 1, 0 \leq \Psi_2(x) \leq 1, \int_{\Omega} f(x, y) dy < 1$ , and  $\int_{\Omega} g(x, y) dy < 1$ .

On the other hand, we have

$$\begin{aligned} \bar{u}_t - \Delta \bar{u}^{m_1} - \bar{u}^{p_1} \bar{v}^{q_1} &= M^{l_1 m_1} \epsilon_1 - M^{p_1 l_1 + l_2 q_1} \Psi_1^{p_1/m_1} \Psi_2^{q_1/m_2} \\ &\geq M^{l_1 m_1} \epsilon_1 - M^{p_1 l_1 + l_2 q_1} \bar{K}_1^{p_1/m_1} \bar{K}_2^{q_1/m_2}, \end{aligned} \quad (3.10)$$

$$\begin{aligned} \bar{v}_t - \Delta \bar{v}^{m_2} - \bar{v}^{p_2} \bar{u}^{q_2} &= M^{l_2 m_2} \epsilon_2 - M^{p_2 l_2 + l_1 q_2} \Psi_2^{p_2/m_2} \Psi_1^{q_2/m_1} \\ &\geq M^{l_2 m_2} \epsilon_2 - M^{p_2 l_2 + l_1 q_2} \bar{K}_2^{p_2/m_2} \bar{K}_1^{q_2/m_1}. \end{aligned} \quad (3.11)$$

Let

$$\begin{aligned} M_1 &= \left( \frac{\overline{K}_1^{p_1/m_1} \overline{K}_2^{q_1/m_2}}{\varepsilon_1} \right)^{1/(l_1 m_1 - p_1 - l_2 q_1)}, \\ M_2 &= \left( \frac{\overline{K}_2^{p_2/m_2} \overline{K}_1^{q_2/m_1}}{\varepsilon_2} \right)^{1/(l_2 m_2 - p_2 - l_1 q_2)}. \end{aligned} \quad (3.12)$$

If  $m_1 > p_1, m_2 > p_2$ , and  $q_1 p_2 < (m_1 - p_1)(m_2 - p_2)$ , by Lemma 2.5, there exist positive constants  $l_1, l_2$  such that

$$p_1 l_1 + q_1 l_2 < m_1 l_1, \quad q_2 l_2 + p_2 l_1 < m_2 l_2. \quad (3.13)$$

Therefore, we can choose  $M$  sufficiently large, such that

$$M > \max\{M_1, M_2\}, \quad (3.14)$$

$$M^{l_1} \Psi_1^{(1/m_1)} \geq u_0(x), \quad M^{l_2} \Psi_2^{(1/m_2)} \geq v_0(x). \quad (3.15)$$

Now, it follows from (3.8)–(3.15) that  $(\bar{u}, \bar{v})$  defined by (3.7) is a positive supersolution of (1.1)–(1.3).

By comparison principle, we conclude that  $(u, v) \leq (\bar{u}, \bar{v})$ , which implies  $(u, v)$  exists globally.

(2) If  $m_1 < p_1, m_2 < p_2$  or  $(m_1 - p_1)(m_2 - p_2) < q_1 q_2$ , by Lemma 2.6, there exist positive constants  $l_1, l_2$  such that

$$p_1 l_1 + q_1 l_2 > m_1 l_1, \quad q_2 l_2 + p_2 l_1 > m_2 l_2. \quad (3.16)$$

So we can choose  $M = \min\{M_1, M_2\}$ . Furthermore, assume that  $u_0(x), v_0(x)$  are small enough to satisfy (3.15). It follows that  $(\bar{u}, \bar{v})$  defined by (3.7) is a positive supersolution of (1.1)–(1.3). Hence,  $(u, v)$  exists globally.

Due to the requirement of the comparison principle we will construct blow-up subsolutions in some subdomain of  $\Omega$  in which  $u, v > 0$ . We use an idea from Souplet [42] and apply it to degenerate equations. Let  $\varphi(x)$  be a nontrivial nonnegative continuous function and vanished on  $\partial\Omega$ . Without loss of generality, we may assume that  $0 \in \Omega$  and  $\varphi(0) > 0$ . We will construct a blow-up positive subsolution to complete the proof.

Set

$$\underline{u}(x, t) = \frac{1}{(T-t)^{l_1}} \omega^{(1/m_1)} \left( \frac{|x|}{(T-t)^\sigma} \right), \quad \underline{v}(x, t) = \frac{1}{(T-t)^{l_2}} \omega^{(1/m_2)} \left( \frac{|x|}{(T-t)^\sigma} \right), \quad (3.17)$$

with

$$\omega(r) = \frac{R^3}{12} - \frac{R}{4} r^2 + \frac{1}{6} r^3, \quad r = \frac{|x|}{(T-t)}, \quad 0 \leq r \leq R, \quad (3.18)$$

where  $l_1, l_2, \sigma > 0$  and  $0 < T < 1$  are to be determined later. Clearly,  $0 \leq \omega(r) \leq R^3/12$  and  $\omega(r)$  is nonincreasing since  $\omega'(r) = r(r - R)/2 \leq 0$ . Note that

$$\text{supp } \underline{u}(\cdot, t) = \text{supp } \underline{v}(\cdot, t) = \overline{B(0, R(T-t)^\sigma)} \subset \overline{B(0, RT^\sigma)} \subset \Omega, \quad (3.19)$$

for sufficiently small  $T > 0$ . Obviously,  $(\underline{u}, \underline{v})$  becomes unbounded as  $t \rightarrow T^-$ , at the point  $x = 0$ . Calculating directly, we obtain that

$$\underline{u}_t - \Delta \underline{u}^{m_1}(x, t) = \frac{m_1 l_1 \omega^{1/m_1}(r) + \sigma r \omega'(r) \omega^{(1-m_1)/m_1}}{m_1 (T-t)^{l_1+1}} + \frac{R-2r}{2(T-t)^{m_1+2\sigma}} + \frac{(N-1)(R-r)}{2(T-t)^{m_1 l_1 + \sigma}} \quad (3.20)$$

$$\leq \frac{l_1 (R^3/12)^{1/m_1}}{(T-t)^{l_1+1}} + \frac{NR - (N+1)r}{2(T-t)^{m_1 l_1 + 2\sigma}}, \quad (3.21)$$

notice that  $T < 1$  is sufficiently small.

Similarly, we have

$$\underline{v}_t - \Delta \underline{v}^{m_2}(x, t) \leq \frac{l_2 (R^3/12)^{1/m_2}}{(T-t)^{l_2+1}} + \frac{NR - (N+1)r}{2(T-t)^{m_2 l_2 + 2\sigma}}. \quad (3.22)$$

*Case 1.* If  $0 \leq r \leq NR/(N+1)$ , we have  $\omega(r) \geq (3N+1)R^3/12(N+1)^3$ , then

$$\begin{aligned} \underline{u}^{p_1} \underline{v}^{q_1} &= \frac{\omega^{p_1/m_1} \omega^{q_1/m_2}}{(T-t)^{p_1 l_1 + q_1 l_1}} \geq \frac{(R^3/12)^{(q_1/m_2)}}{(T-t)^{p_1 l_1 + q_1 l_2}} \left( \frac{R^3(3N+1)}{12(N+1)^3} \right)^{p_1/m_1}, \\ \underline{v}^{p_2} \underline{u}^{q_2} &= \frac{\omega^{p_2/m_2} \omega^{q_2/m_1}}{(T-t)^{p_2 l_2 + q_2 l_2}} \geq \frac{(R^3/12)^{(p_1/m_1)}}{(T-t)^{p_2 l_2 + q_2 l_1}} \left( \frac{R^3(3N+1)}{12(N+1)^3} \right)^{q_1/m_2}. \end{aligned} \quad (3.23)$$

Hence,

$$\begin{aligned} \underline{u}_t - \Delta \underline{u}^{m_1}(x, t) - \underline{u}^{p_1} \underline{v}^{q_1} &\leq \frac{l_1 (R^3/12)^{1/m_1}}{(T-t)^{l_1+1}} - \frac{(R^3/12)^{q_1/m_2}}{(T-t)^{p_1 l_1 + q_1 l_2}} \left( \frac{R^3(3N+1)}{12(N+1)^3} \right)^{p_1/m_1}, \\ \underline{v}_t - \Delta \underline{v}^{m_2}(x, t) - \underline{v}^{p_2} \underline{u}^{q_2} &\leq \frac{l_2 (R^3/12)^{1/m_2}}{(T-t)^{l_2+1}} - \frac{(R^3/12)^{p_1/m_1}}{(T-t)^{p_2 l_2 + q_2 l_1}} \left( \frac{R^3(3N+1)}{12(N+1)^3} \right)^{q_1/m_2}. \end{aligned} \quad (3.24)$$

*Case 2.* If  $NR/(N+1) < r \leq R$ , then

$$\begin{aligned} \underline{u}_t - \Delta \underline{u}^{m_1}(x, t) - \underline{u}^{p_1} \underline{v}^{q_1} &\leq \frac{l_1 (R^3/12)^{1/m_1}}{(T-t)^{l_1+1}} + \frac{NR - (N+1)r}{2(T-t)^{m_1 l_1 + 2\sigma}}, \\ \underline{v}_t - \Delta \underline{v}^{m_2}(x, t) - \underline{v}^{p_2} \underline{u}^{q_2} &\leq \frac{l_2 (R^3/12)^{1/m_2}}{(T-t)^{l_2+1}} + \frac{NR - (N+1)r}{2(T-t)^{m_2 l_2 + 2\sigma}}. \end{aligned} \quad (3.25)$$

By Lemma 2.6, there exist positive constants  $l_1, l_2$  large enough to satisfy

$$p_1 l_1 + q_1 l_2 > m_1 l_1 + 1, \quad q_2 l_1 + p_2 l_2 > m_2 l_2 + 1, \quad (m_1 - 1)l_1 > 1, \quad (m_2 - 1)l_2 > 1, \quad (3.26)$$

and we can choose  $\sigma > 0$  be sufficiently small that

$$\sigma < \max \left\{ \frac{p_1 l_1 + q_1 l_2 - m_1 l_1}{2}, \frac{p_2 l_2 + q_2 l_1 - m_2 l_2}{2} \right\}. \quad (3.27)$$

Thus, we have

$$p_1 l_1 + q_1 l_2 > m_1 l_1 + 2\sigma > l_1 + 1, \quad p_2 l_2 + q_2 l_1 > m_2 l_2 + 2\sigma > l_2 + 1. \quad (3.28)$$

Hence, for sufficiently small  $T > 0$ , (3.24) and (3.25) imply that

$$\underline{u}_t - \Delta \underline{u}^{m_1}(x, t) - \underline{u}^{p_1} \underline{v}^{q_1} \leq 0, \quad (x, t) \in \Omega \times (0, T), \quad (3.29)$$

$$\underline{v}_t - \Delta \underline{v}^{m_2}(x, t) - \underline{v}^{p_2} \underline{u}^{q_2} \leq 0, \quad (x, t) \in \Omega \times (0, T). \quad (3.30)$$

Since  $\varphi(0) > 0$  and  $\varphi(x)$  is continuous, there exist two positive constants  $\rho$  and  $\varepsilon$  such that  $\varphi(x) \geq \varepsilon$ , for all  $x \in B(0, \rho) \subset \Omega$ . Choose  $T$  small enough to insure  $B(0, RT^\sigma) \subset B(0, \rho)$ , hence  $\underline{u} \leq 0, \underline{v} \leq 0$  on  $\partial\Omega \times (0, T)$ . Under the assumption that  $\int_{\Omega} f(x, y) dy < 1$  and  $\int_{\Omega} g(x, y) dy < 1$  for any  $\partial\Omega$ , we have  $\underline{u}(x, t) \leq \int_{\Omega} f(x, y) \underline{u}(y, t) dy, \underline{v}(x, t) \leq \int_{\Omega} f(x, y) \underline{v}(y, t) dy$  and  $x \in \partial\Omega \times (0, T)$ . Furthermore, choose  $u_0(x), v_0(x)$  so large that  $u_0(x) > \underline{u}(x, 0), v_0(x) > \underline{v}(x, 0)$ . By comparison principle, we have  $(\underline{u}, \underline{v}) \leq (u, v)$ . It shows that solution  $(u, v)$  to (1.1)–(1.3) blows up in finite time.  $\square$

#### 4. Blow-Up Rate Estimates

In this section, we will estimate the blow-up rate of the blow-up solution of (1.1). Throughout this section, we will assume that

$$q_1 > m_2, \quad q_2 > m_1 \quad \text{and satisfy} \quad q_2 > p_1 - 1, \quad q_1 > p_2 - 1. \quad (4.1)$$

To obtain the estimate, we firstly introduce some transformations. Let  $U(x, t) = u^{m_1}(x, t), V(x, t) = (m_2/m_1)^{m_2/(m_2-1)} v^{m_2}(x, t)$ , then problem (1.1)–(1.3) becomes

$$\begin{aligned} U_t &= U^{r_1}(\Delta U + aU^{p_3}V^{q_3}(x, t)), & V_t &= V^{r_2}(\Delta V + bV^{p_4}U^{q_4}(x, t)), & x \in \Omega, t > 0, \\ U(x, t) &= \left( \int_{\Omega} f(x, y) U^{m_3}(y, t) dy \right)^{m_1}, & V(x, t) &= \left( \int_{\Omega} g(x, y) V^{m_4}(y, t) dy \right)^{m_2}, & (4.2) \\ & & & & x \in \partial\Omega, t > 0, \end{aligned}$$

$$U(x, 0) = U_0(x), \quad V(x, 0) = V_0(x), \quad x \in \Omega,$$

where  $U_0(x) = u_0^{m_1}(x)$ ,  $V_0(x) = (m_2/m_1)^{m_2/(m_2-1)} v_0^{m_2}(x)$ ;  $m_3 = 1/m_1 < 1$ ,  $m_4 = 1/m_2 < 1$ ;  $p_3 = p_1/m_1$ ,  $q_3 = q_1/m_2$ ,  $p_4 = p_2/m_2$ ,  $q_4 = q_2/m_1$ ;  $0 < r_1 = (m_1-1)/m_1 < 1$ ,  $0 < r_2 = (m_2-1)/m_2 < 1$ ;  $a = (m_1/m_2)^{q_1/(m_1-1)}$ ,  $b = (m_1/m_2)^{(p_2-m_2)/(m_2-1)}$ . By the conditions (4.1), we have  $q_3 > 1$ ,  $q_4 > 1$  and satisfy that  $q_4 - p_3 - r_1 + 1 > 0$ ,  $q_3 - p_4 - r_2 + 1 > 0$ . Under this transformation, assumptions (H1)-(H3) become

(H1')  $U_0(x), V_0(x) \in C^{2+\mu}(\Omega) \cap \overline{(\Omega)}$ , for some  $0 < \mu < 1$ ;

(H2') there exists a constant  $\delta \geq \delta_0 > 0$ , such that

$$\Delta U_0 + aU_0^{p_3}V_0^{q_3} - \delta U_0^{k_1+1-r_1}(x) \geq 0, \quad \Delta V_0 + bV_0^{p_4}U_0^{q_4} - \delta V_0^{k_2+1-r_2}(x) \geq 0, \quad (4.3)$$

where  $\delta_0, k_1, k_2$  will be given later.

By the standard method [16, 42], we can show that system (4.2) has a smooth nonnegative solution  $(U, V)$ , provided that  $U_0, V_0$  satisfy the hypotheses (H1)'-(H2)'. We thus assume that the solution  $(U, V)$  of problem (4.2) blows up in the finite time  $T_*$ . Denote  $M_1(t) = \max_{\overline{\Omega}} U(x, t)$ ,  $M_2(t) = \max_{\overline{\Omega}} V(x, t)$ . We can obtain the blow-up rate from the following lemmas.

**Lemma 4.1.** *Suppose that  $U_0(x), V_0(x)$  satisfy (H1)'-(H2)', then there exists a positive constant  $K_1$  such that*

$$\begin{aligned} & M_1(t)^{q_4-p_3-r_1+1} + M_2(t)^{q_3-p_4-r_2+1} \\ & \geq C_1(T_* - t)^{-((q_3-p_4-r_2+1)(q_4-p_3-r_1+1))/(q_3 q_4 - (1-r_1-p_3)(1-r_2-p_4))}. \end{aligned} \quad (4.4)$$

*Proof.* By (4.2), we have (see [43])

$$M_1' \leq aM_1^{p_3+r_1}M_2^{q_3}, \quad M_2' \leq bM_1^{q_4}M_2^{p_4+r_2}. \quad (4.5)$$

Noticing that  $q_4 - p_3 - r_1 + 1 > 0$  and  $q_3 - p_4 - r_2 + 1 > 0$ , hence we have

$$\begin{aligned} & \left( M_1^{q_4-p_3-r_1+1}(t) + M_2^{q_3-p_4-r_2+1}(t) \right)' \\ & \leq (a(q_4 - p_3 - r_1 + 1) + b(q_3 - p_4 - r_2 + 1))M_1^{q_4}(t)M_2^{q_3}(t) \\ & \leq C_2 \left( M_1^{q_4-p_3-r_1+1}(t) + M_2^{q_3-p_4-r_2+1}(t) \right)^{\frac{((q_4-p_3-r_1+1)q_3 + (q_3-p_4-r_2+1)q_4)}{((q_4-p_3-r_1+1)(q_3-p_4-r_2+1))}}, \end{aligned} \quad (4.6)$$

by virtue of Young's inequality. Integrating (4.6) from  $t$  to  $T_*$ , we can obtain (4.4).  $\square$

**Lemma 4.2.** *Suppose that  $U_0, V_0$  satisfy (H1)'-(H2)',  $(U, V)$  is a solution of (4.2). Then*

$$U_t - \delta U^{k_1+1} \geq 0, \quad V_t - \delta V^{k_2+1} \geq 0, \quad (x, t) \in \Omega \times (0, T_*), \quad (4.7)$$

where

$$\begin{aligned}
 k_1 &= \frac{q_4 q_3 - (1 - r_1 - p_3)(1 - r_2 - p_4)}{q_3 - r_2 - p_4 + 1}, & k_2 &= \frac{q_4 q_3 - (1 - r_1 - p_3)(1 - r_2 - p_4)}{q_4 - r_1 - p_3 + 1}, \\
 \delta_1 &= \frac{ak_1(1 + k_1 - p_3)}{r_1(2k_1 + 1 - r_1 - p_3)} \left( \frac{1 + k_1 - p_3}{q_3 + k_2} \right)^{(q_3(2k_1+1-r_1-p_3))/(k_1(q_3+k_2))}, \\
 \delta_2 &= \frac{bk_2(1 + k_2 - p_4)}{r_2(2k_2 + 1 - r_2 - p_3)} \left( \frac{1 + k_2 - p_4}{q_4 + k_1} \right)^{(q_4(2k_2+1-r_2-p_4))/(k_2(q_4+k_1))}, \\
 \delta &> \delta_0 = \max\{|\delta_1|, |\delta_2| > 0\}.
 \end{aligned} \tag{4.8}$$

*Proof.* Set  $J_1(x, t) = U_t - \delta U^{k_1+1}$ ,  $J_2(x, t) = V_t - \delta V^{k_2+1}$ ,  $(x, t) \in \Omega \times (0, T_*)$ , a straightforward computation yields

$$\begin{aligned}
 J_{1t} - U^{r_1} \Delta J_1 - (2\delta r_1 U^{k_1} + ap_3 U^{r_1+p_3-1} V^{q_3}) J_1 - aq_3 U^{r_1+p_3} V^{q_3-1} J_2 \\
 = r_1 U^{-1} J_1^2 + \delta k_1(k_1 + 1) U^{k_1+r_1-1} |\nabla U|^2 + r_1 \delta^2 U^{2k_1+1} \\
 + aq_3 \delta U^{r_1+p_3} V^{q_3+k_2} - a\delta(1 + k_1 - p_3) U^{k_1+r_1+p_3} V^{q_3} \\
 \geq r_1 \delta^2 U^{2k_1+1} + aq_3 \delta U^{r_1+p_3} V^{q_3+k_2} - a\delta(1 + k_1 - p_3) U^{k_1+r_1+p_3} V^{q_3}.
 \end{aligned} \tag{4.9}$$

If  $1 + k_1 \leq p_3$ , obviously we have

$$J_{1t} - U^{r_1} \Delta J_1 - (2\delta r_1 U^{k_1} + ap_3 U^{r_1+p_3-1} V^{q_3}) J_1 - aq_3 U^{r_1+p_3} V^{q_3-1} J_2 \geq 0. \tag{4.10}$$

Otherwise, noticing that  $k_1/(2k_1 + 1 - r_1 - p_3) + q_3/(q_3 + k_2) = 1$ , by virtue of Young's inequality,

$$U^{k_1} V^{q_3} \leq \frac{k_1}{2k_1 + 1 - r_1 - p_3} (\theta U^{k_1})^{(2k_1+1-r_1-p_3)/k_1} + \frac{q_3}{q_3 + k_2} \left( \frac{V^{q_3}}{\theta} \right)^{(q_3+k_2)/q_3}, \tag{4.11}$$

where  $\theta = ((k_1 + 1 - p_3)/(q_3 + k_2))^{q_3/(q_3+k_2)}$ , we have

$$\begin{aligned}
 J_{1t} - U^{r_1} \Delta J_1 - (2\delta r_1 U^{k_1} + ap_3 U^{r_1+p_3-1} V^{q_3}) J_1 - aq_3 U^{r_1+p_3} V^{q_3-1} J_2 \\
 \geq r_1 \delta^2 U^{2k_1+1} + aq_3 \delta U^{r_1+p_3} V^{q_3+k_2} - a\delta(1 + k - p_3) U^{k_1+r_1+p_3} V^{q_3} \\
 \geq r_1 \delta(\delta - \delta_1) U^{2k_1} \geq 0.
 \end{aligned} \tag{4.12}$$

Similarly, we also have

$$J_{2t} - V^{r_2} \Delta J_2 - (2\delta r_2 V^{k_2} + bp_4 V^{r_2+p_4}) J_2 - bq_4 V^{r_2+p_4} U^{q_4-1} J_1 \geq 0. \tag{4.13}$$

Fix  $(x, t) \in \partial\Omega \times (0, T_*)$ , we have

$$\begin{aligned} J_1(x, t) &= U_t - \delta U^{k_1+1} \\ &= \left( \int_{\Omega} f(x, y) u(y, t) \right)^{m_1-1} \left( \int_{\Omega} m_1 f(x, y) u_t(y, t) dy - \delta \left( \int_{\Omega} f(x, y) u(y, t) dy \right)^{\lambda} \right), \end{aligned} \quad (4.14)$$

where  $\lambda = m_1 k_1 + 1 > 1$ . Since  $U_t(x, t) = J_1(x, t) + \delta U^{k_1+1}$ , we have

$$\begin{aligned} & \int_{\Omega} m_1 f(x, y) u_t(y, t) dy - \delta \left( \int_{\Omega} f(x, t) u(y, t) dy \right)^{\lambda} \\ &= \int_{\Omega} f(x, y) U^{(1-m_1)/m_1} J_1(y, t) dy \\ &+ \delta \left( \int_{\Omega} f(x, y) U^{\lambda/m_1}(y, t) - \left( \int_{\Omega} f(x, y) U^{1/m_1}(y, t) dy \right)^{\lambda} \right). \end{aligned} \quad (4.15)$$

Noticing that  $0 < \Phi(x) = \int_{\Omega} f(x, y) dy \leq 1, x \in \partial\Omega$ , by virtue of Jensen's inequality, we have

$$\begin{aligned} & \int_{\Omega} f(x, y) U^{\lambda/m_1}(y, t) dy - \left( \int_{\Omega} f(x, y) U^{1/m_1}(y, t) dy \right)^{\lambda} \\ & \geq \int_{\Omega} f(x, y) dy \left( \frac{\int_{\Omega} f(x, y) U^{1/m_1}(y, t) dy}{\int_{\Omega} f(x, y) dy} \right)^{\lambda} - \left( \int_{\Omega} f(x, t) U^{1/m_1}(y, t) \right)^{\lambda} \\ & \geq \Phi(x) \left( \int_{\Omega} f(x, t) U^{1/m_1}(y, t) \frac{dy}{\Phi(x)} \right)^{\lambda} - \left( \int_{\Omega} f(x, y) U^{1/m_1}(y, t) dy \right)^{\lambda} \\ & = \left( \frac{1}{\Phi(x)^{\lambda-1}} \right) \left( \int_{\Omega} f(x, y) U^{1/m_1}(y, t) dy \right)^{\lambda} \geq 0, \end{aligned} \quad (4.16)$$

here, we used  $\lambda > 1$  and  $0 < \Phi(x) \leq 1$  in the last inequality. Hence  $(x, t) \in \partial\Omega \times (0, T_*)$ ,

$$J_1(x, t) \geq \left( \int_{\Omega} f(x, t) U^{1/m_1}(y, t) dy \right)^{m_1-1} \int_{\Omega} f(x, y) U^{(1-m_1)/m_1}(y, t) J_1 dy. \quad (4.17)$$

Similarly, we also have

$$J_2(x, t) \geq \left( \int_{\Omega} g(x, t) V^{1/m_2}(y, t) dy \right)^{m_2-1} \int_{\Omega} g(x, y) V^{(1-m_2)/m_2}(y, t) J_2 dy. \quad (4.18)$$

On the other hand,  $(H1)'$ - $(H2)'$  imply that  $J_1(x, 0) \geq 0, J_2(x, 0) \geq 0, x \in \Omega$ . Combined inequalities (4.12)-(4.18) and Lemma 2.4, we obtain  $J_1 \geq 0, J_2 \geq 0$ , that is, (4.7) holds. Integrating (4.7) from  $t$  to  $T_*$ , we conclude that

$$\begin{aligned} M_1(t) &\leq C_3(T_* - t)^{-(q_3 - p_4 - r_2 + 1)/(q_4 q_3 - (1 - r_1 - p_3)(1 - r_2 - p_4))}, \\ M_2(t) &\leq C_4(T_* - t)^{-(q_4 - p_3 - r_1 + 1)/(q_4 q_3 - (1 - r_1 - p_3)(1 - r_2 - p_4))}. \end{aligned} \quad (4.19)$$

where  $C_3, C_4$  are positive constants independent of  $t$ . It follows from Lemma 4.1 and (4.19), we have the following lemma.

**Lemma 4.3.** *Suppose that  $U_0(x), V_0(x)$  satisfy  $(H1)'$ - $(H3)'$ . If  $(U, V)$  is the solution of system (4.2) and blows up in finite time  $T_*$ , then there exist positive constants  $C_i (i = 3, 4, 5, 6)$  such that*

$$\begin{aligned} C_5 &\leq \max_{x \in \Omega} U(x, t)(T_* - t)^{(q_3 - p_4 - r_2 + 1)/(q_4 q_3 - (1 - r_1 - p_3)(1 - r_2 - p_4))} \leq C_3, \quad \text{for } 0 < t < T_*, \\ C_6 &\leq \max_{x \in \Omega} V(x, t)(T_* - t)^{(q_4 - p_3 - r_1 + 1)/(q_4 q_3 - (1 - r_1 - p_3)(1 - r_2 - p_4))} \leq C_4, \quad \text{for } 0 < t < T_*. \end{aligned} \quad (4.20)$$

According the transform and Lemma 4.3, we can obtain Theorem 1.3. □

## Acknowledgments

The authors would like to thank the anonymous referees for their suggestions and comments on the original manuscript. This work was partially supported by NSF of China (10771226) and partially supported by the Educational Science Foundation of Chongqing (KJ101303), China.

## References

- [1] J. I. Diaz and R. Kersner, "On a nonlinear degenerate parabolic equation in infiltration or evaporation through a porous medium," *Journal of Differential Equations*, vol. 69, no. 3, pp. 368–403, 1987.
- [2] K. Deng and H. A. Levine, "The role of critical exponents in blow-up theorems: the sequel," *Journal of Mathematical Analysis and Applications*, vol. 243, no. 1, pp. 85–126, 2000.
- [3] H. A. Levine, "The role of critical exponents in blowup theorems," *SIAM Review*, vol. 32, no. 2, pp. 262–288, 1990.
- [4] H. Chen, "Global existence and blow-up for a nonlinear reaction-diffusion system," *Journal of Mathematical Analysis and Applications*, vol. 212, no. 2, pp. 481–492, 1997.
- [5] M. Escobedo and M. A. Herrero, "Boundedness and blow up for a semilinear reaction-diffusion system," *Journal of Differential Equations*, vol. 89, no. 1, pp. 176–202, 1991.
- [6] M. Escobedo and H. A. Levine, "Critical blowup and global existence numbers for a weakly coupled system of reaction-diffusion equations," *Archive for Rational Mechanics and Analysis*, vol. 129, no. 1, pp. 47–100, 1995.
- [7] H. A. Levine, "A Fujita type global existence–global nonexistence theorem for a weakly coupled system of reaction-diffusion equations," *Zeitschrift für Angewandte Mathematik und Physik*, vol. 42, no. 3, pp. 408–430, 1991.
- [8] F. Quirós and J. D. Rossi, "Non-simultaneous blow-up in a semilinear parabolic system," *Zeitschrift für Angewandte Mathematik und Physik*, vol. 52, no. 2, pp. 342–346, 2001.
- [9] M. Wang, "Global existence and finite time blow up for a reaction-diffusion system," *Zeitschrift für Angewandte Mathematik und Physik*, vol. 51, no. 1, pp. 160–167, 2000.



- [10] S. Zheng, "Global boundedness of solutions to a reaction-diffusion system," *Mathematical Methods in the Applied Sciences*, vol. 22, no. 1, pp. 43–54, 1999.
- [11] S. Zheng, "Global existence and global non-existence of solutions to a reaction-diffusion system," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 39, no. 3, pp. 327–340, 2000.
- [12] V. A. Galaktionov, S. P. Kurdyumov, and A. A. Samarskii, "A parabolic system of quasilinear equations. I," *Differentsial'nye Uravneniya*, vol. 19, no. 12, pp. 2123–2140, 1983.
- [13] V. A. Galaktionov, S. P. Kurdyumov, and A. A. Samarskii, "A parabolic system of quasilinear equations. II," *Differentsial'nye Uravneniya*, vol. 21, no. 9, pp. 1049–1062, 1985.
- [14] X. Song, S. Zheng, and Z. Jiang, "Blow-up analysis for a nonlinear diffusion system," *Zeitschrift für Angewandte Mathematik und Physik*, vol. 56, no. 1, pp. 1–10, 2005.
- [15] P. Lei and S. Zheng, "Global and nonglobal weak solutions to a degenerate parabolic system," *Journal of Mathematical Analysis and Applications*, vol. 324, no. 1, pp. 177–198, 2006.
- [16] Z. Duan, W. Deng, and C. Xie, "Uniform blow-up profile for a degenerate parabolic system with nonlocal source," *Computers & Mathematics with Applications*, vol. 47, no. 6-7, pp. 977–995, 2004.
- [17] Z. Li, C. Mu, and Z. Cui, "Critical curves for a fast diffusive polytropic filtration system coupled via nonlinear boundary flux," *Zeitschrift für Angewandte Mathematik und Physik*, vol. 60, no. 2, pp. 284–296, 2009.
- [18] Z. Li, Z. Cui, and C. Mu, "Critical curves for fast diffusive polytropic filtration equations coupled through boundary," *Applicable Analysis*, vol. 87, no. 9, pp. 1041–1052, 2008.
- [19] J. Zhou and C. Mu, "On the critical Fujita exponent for a degenerate parabolic system coupled via nonlinear boundary flux," *Proceedings of the Edinburgh Mathematical Society. Series II*, vol. 51, no. 3, pp. 785–805, 2008.
- [20] J. Zhou and C. Mu, "The critical curve for a non-Newtonian polytropic filtration system coupled via nonlinear boundary flux," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 68, no. 1, pp. 1–11, 2008.
- [21] W. A. Day, "A decreasing property of solutions of parabolic equations with applications to thermoelasticity," *Quarterly of Applied Mathematics*, vol. 40, no. 4, pp. 468–475, 1983.
- [22] W. A. Day, *Heat Conduction within Linear Thermoelasticity*, vol. 30 of *Springer Tracts in Natural Philosophy*, Springer, New York, NY, USA, 1985.
- [23] A. Friedman, "Monotonic decay of solutions of parabolic equations with nonlocal boundary conditions," *Quarterly of Applied Mathematics*, vol. 44, no. 3, pp. 401–407, 1986.
- [24] K. Deng, "Comparison principle for some nonlocal problems," *Quarterly of Applied Mathematics*, vol. 50, no. 3, pp. 517–522, 1992.
- [25] C. V. Pao, "Asymptotic behavior of solutions of reaction-diffusion equations with nonlocal boundary conditions," *Journal of Computational and Applied Mathematics*, vol. 88, no. 1, pp. 225–238, 1998.
- [26] S. Seo, "Blowup of solutions to heat equations with nonlocal boundary conditions," *Kobe Journal of Mathematics*, vol. 13, no. 2, pp. 123–132, 1996.
- [27] S. Seo, "Global existence and decreasing property of boundary values of solutions to parabolic equations with nonlocal boundary conditions," *Pacific Journal of Mathematics*, vol. 193, no. 1, pp. 219–226, 2000.
- [28] Y. Wang, C. Mu, and Z. Xiang, "Blowup of solutions to a porous medium equation with nonlocal boundary condition," *Applied Mathematics and Computation*, vol. 192, no. 2, pp. 579–585, 2007.
- [29] L. Kong and M. Wang, "Global existence and blow-up of solutions to a parabolic system with nonlocal sources and boundaries," *Science in China. Series A*, vol. 50, no. 9, pp. 1251–1266, 2007.
- [30] Y. Wang and Z. Xiang, "Blowup analysis for a semilinear parabolic system with nonlocal boundary condition," *Boundary Value Problems*, vol. 2009, Article ID 516390, 14 pages, 2009.
- [31] Y. Wang, C. Mu, and Z. Xiang, "Properties of positive solution for nonlocal reaction-diffusion equation with nonlocal boundary," *Boundary Value Problems*, vol. 2007, Article ID 64579, 12 pages, 2007.
- [32] H.-M. Yin, "On a class of parabolic equations with nonlocal boundary conditions," *Journal of Mathematical Analysis and Applications*, vol. 294, no. 2, pp. 712–728, 2004.
- [33] Y. Yin, "On nonlinear parabolic equations with nonlocal boundary condition," *Journal of Mathematical Analysis and Applications*, vol. 185, no. 1, pp. 161–174, 1994.
- [34] S. Zheng and L. Kong, "Roles of weight functions in a nonlinear nonlocal parabolic system," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 68, no. 8, pp. 2406–2416, 2008.
- [35] P. Souplet, "Uniform blow-up profiles and boundary behavior for diffusion equations with nonlocal nonlinear source," *Journal of Differential Equations*, vol. 153, no. 2, pp. 374–406, 1999.

- [36] J. R. Anderson, "Local existence and uniqueness of solutions of degenerate parabolic equations," *Communications in Partial Differential Equations*, vol. 16, no. 1, pp. 105–143, 1991.
- [37] W. A. Day, "Extensions of a property of the heat equation to linear thermoelasticity and other theories," *Quarterly of Applied Mathematics*, vol. 40, no. 3, pp. 319–330, 1982.
- [38] Z. Lin and Y. Liu, "Uniform blowup profiles for diffusion equations with nonlocal source and nonlocal boundary," *Acta Mathematica Scientia. Series B*, vol. 24, no. 3, pp. 443–450, 2004.
- [39] C. V. Pao, *Nonlinear Parabolic and Elliptic Equations*, Plenum Press, New York, NY, USA, 1992.
- [40] C. V. Pao, "Blowing-up of solution for a nonlocal reaction-diffusion problem in combustion theory," *Journal of Mathematical Analysis and Applications*, vol. 166, no. 2, pp. 591–600, 1992.
- [41] W. Deng, "Global existence and finite time blow up for a degenerate reaction-diffusion system," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 60, no. 5, pp. 977–991, 2005.
- [42] P. Souplet, "Blow-up in nonlocal reaction-diffusion equations," *SIAM Journal on Mathematical Analysis*, vol. 29, no. 6, pp. 1301–1334, 1998.
- [43] A. Friedman and B. McLeod, "Blow-up of positive solutions of semilinear heat equations," *Indiana University Mathematics Journal*, vol. 34, no. 2, pp. 425–447, 1985.