

Acta vet. scand. 2001, 42, 31-44.

Plants as De-Worming Agents of Livestock in the Nordic Countries: Historical Perspective, Popular Beliefs and Prospects for the Future

By P.J. Waller¹, G. Bernes², S.M. Thamsborg³, A. Sukura⁴, S.H. Richter⁵, K. Ingebrigtsen⁶ and J. Höglund⁷

¹Dept. Parasitology (SWEPAR), National Veterinary Institute, Uppsala, Sweden, ²Dept. Agric. Research Nth. Sweden, SLU, Umeå, Sweden, ³Dept. Anim. Science and Animal Health, The Royal Veterinary and Agricultural University, Frederiksberg, Denmark, ⁴Dept. Basic Vet. Science, Univ. Helsinki, Finland, ⁵Inst. Exp. Pathology, Univ. Iceland, Keldur, Reykjavik, Iceland, ⁶Dept. Pharmacology, Microbiology and Food Hygiene, Norwegian School Vet. Medicine, Oslo, Norway, and ⁷SWEPAR, Swedish University of Agricultural Sciences, Uppsala, Sweden.

Waller PJ, Bernes G, Thamsborg SM, Sukura A, Richter SH, Ingebrigtsen K, Höglund J: Plants as de-worming agents of livestock in the Nordic Countries: historical perspective, popular beliefs and prospects for the future. Acta vet. Scand. 2001, 42, 31-44. – Preparations derived from plants were the original therapeutic interventions used by man to control diseases (including parasites), both within humans and livestock. Development of herbal products depended upon local botanical flora with the result that different remedies tended to develop in different parts of the world. Nevertheless, in some instances, the same or related plants were used over wide geographic regions, which also was the result of communication and/or the importation of plant material of high repute. Thus, the Nordic countries have an ancient, rich and diverse history of plant derived anthelmintic medications for human and animal use. Although some of the more commonly used herbal de-wormers were derived from imported plants, or their products, many are from endemic plants or those that thrive in the Scandinavian environment. With the advent of the modern chemotherapeutic era, and the discovery, development and marketing of a seemingly unlimited variety of highly efficacious, safe synthetic chemicals with very wide spectra of activities, herbal remedies virtually disappeared from the consciousness – at least in the Western world. This attitude is now rapidly changing. There is a widespread resurgence in natural product medication, driven by major threats posed by multi-resistant pest, or disease, organisms and the diminishing public perceptions that synthetic chemicals are the panacea to health and disease control. This review attempts to provide a comprehensive account of the depth of historical Nordic information available on herbal de-wormers, with emphasis on livestock and to provide some insights on potentially rewarding areas of "re-discovery" and scientific evaluation in this field.

plant anthelmintics; herbal remedies; helminth parasites; livestock; man.

Introduction

The use of plants, or their extracts, for treatment of gastro-intestinal parasites in humans and livestock is steeped in antiquity. It was Claudius Galēnus (AD 130-200), a Greek physician of Pergamon, who received notoriety for applying medicines prepared from vegetable substances by infusion, or decoction. These became known generically as "galenical" drugs, or preparations, and established the foundation

for modern veterinary pharmacology. It is with some interest to note that the approach taken by Galēnus was in contrast to the Hippocrates / Paracelsus school of thinking which espoused treating "like with like" (*similia similibus curantur*), that later formed the basis for Hahneman's concepts of homeopathy (Vaarst 1996).

In medieval times, plants with reputed anthel-

mintic properties were often mixed with mineral salts (arsenic, copper etc.), or more esoteric materials (blood, faeces, fluids from reptiles, wild animals etc.) to form quite bizarre and often hazardous concoctions – for both parasites and hosts alike. With time, trial and error, such preparations were refined in an attempt to at least moderate the undesirable consequences to the host, but with the advent of safer and more effective synthetic anthelmintic compounds, they rapidly disappeared from the veterinary anthelmintic market (Gibson 1980). Nevertheless, it is of interest to note that the World Health Organisation has recently estimated that 80% of the population of developing countries rely on traditional medicine, mostly plant drugs, for their primary health care needs (Danøe & Bøgh 2000). Also in a global context, modern pharmacopoeia still contain in the order of 25% drugs derived from plants and many others which are synthetic analogues built on prototype compounds isolated from plants.

However, there has been a resurgence of interest in traditional health practices throughout the world. In veterinary medicine, this interest encompasses ethnobotany and the use of herbal remedies. Workshops, forums and conferences are occurring with increasing regularity, which is obviously being driven by a concomitant increasing level of research activity. The forces responsible for this momentum are manifold. These include the perception that "natural is nice", concerns of synthetic drug residues in the environment and the food chain, and particularly the spectre of rapid emergence of multiple resistant pest organisms through misuse and overuse of these modern drugs.

Within the Nordic countries, the same holds true. This region of the world is at the vanguard of the organic farming movement, with major emphasis on livestock production. Many progressive and educated farmers who have chosen to farm livestock organically are well aware of

the importance of nematode parasites affecting the productivity of their animals and adopt grazing strategies aimed at evading, or mitigating the effects of parasites in their animals (Svensson et al. 2000). However, others show less awareness and management of grazing livestock is largely determined by herbage supply and not by parasitological considerations (Vaarst et al. 1996). One specific strategy is to graze livestock on plants with purported anthelmintic properties. For example, a survey of Danish organic dairy farmers with 255 respondents, revealed the use of herbs in the leys on 26% of the farms. Of these, caraway (*Carum carvi*) [56%], parsley (*Petroselinum crispum*) [20%], chicory (*Cichorium intybis*) [10%], chervil (*Anthriscus cerefolium*) and dill (*Aneethum graveolens*) [14%], used singly or in combination, were the most commonly used plants (Smidt 1997). For the purposes of deworming, the feeding of wormwood (*Artemisia absinthium*), mugwort (*A. vulgaris*), chicory (*Cichorium intybis*) and common tansy (*Tanacetum vulgare*) were expected by farmers to have anthelmintic properties (Smidt 1997).

In addition, there has been a major commitment to the development of sustainable disease control systems, which by definition means the integration of a range of non - chemical methods with the minimal use of drugs, to insure continued effectiveness for the foreseeable future. Control of internal parasites of livestock by these means has been a major focus of research activities in the Nordic countries for several years (Thamsborg et al. 1999).

Research in the Nordic countries on anthelmintic properties, or protective effects, of local plants against helminth parasites of livestock is very recent and limited at this stage to studies on tanniferous forages (Kahiya et al. 1999, Bernes et al. 2000). However, many of the earliest written reports of anthelmintic properties of plants originate from this region of the world

(eg., the writings of Henrik Harpestræng in the early 13th century). The purpose of this review is to provide a historical perspective and to present our view of the potential and possibilities for the use of plants that are endemic, or thrive in the Nordic environment, as de-worming agents. Due to the relatively wide variation in climate, soil, altitude etc. within this limited area, the total number of plant species is quite large.

Also, a brief consideration is made of the recent developments in the use of specialized crops, the so-called "nutraceuticals", which are bioactive crops that are either grazed, or fed after preservation, with the main purpose of preventing or curing disease. The ultimate objective would be to use these plants as additional means for the further development and refinement of sustainable parasite control systems of livestock.

Plants and parasites

There is an inextricable association between plants and parasites of livestock. Pastures provide the link between the free-living and the parasitic phases of helminth parasites for all grazing animals. At different stages of growth, pasture species may facilitate or impede the survival of free-living populations, the establishment of parasite burdens and lessen or intensify the effects of parasitism on the host. Competent management of pastures is needed for the twin goals of efficient conversion of herbage to animal products and effective control of gastrointestinal parasites.

Indirect effects of plants on parasites

It is reasonable to assume that the height, density and form of plant growth could affect the micro-environment of the free-living stages of parasites and thus play a role in transmission of infection. Because of the necessity of moisture

films for movement of nematode larvae from faecal deposits to herbage, it might be expected that pastures consisting largely of erect, tall growing species would provide less protection from desiccation, and from removal by heavy rain, than those consisting primarily of prostrate species. Indeed, a factor considered to mitigate the high rates of contamination following increased stocking rates is the decreased availability of herbage, which provide conditions less favourable for the development and survival of the free-living stages of nematode parasites (Thamsborg *et al.* 1996). Although there has been a number of such ecological studies, little has been done in the Nordic countries and this aspect is outside the scope of this review. However, it is important to recognise these indirect effects of herbage when any grazing studies are conducted on plants with purported anthelmintic properties. It must be established that observed effects are directly plant related and not due to effects on development and survival of the free living stages of parasites, or the provision of an enhanced level of nutrition which helps the animal to rapidly mount an immune response against incoming or resident parasite infections.

Direct effects of plants on parasites

This relates specifically to anthelmintic properties of plants. Although there is a large and diverse range of herbal de-wormers that are used throughout the world, particularly in the Asian and African countries, generally there is a lack of scientific validation of the purported anthelmintic effects of these products. Evidence to date is almost entirely anecdotal. This also applies to the reports of de-worming properties of herbal preparations that have either Nordic origins, or a long history of use in this region (for some historical background, see Grove 1990). These can be broadly classified into the following classes (see Table 1):

Table 1. Plants said to have been used against internal parasites in the Nordic countries; their scientific names and names in English and in the Nordic languages.

Plant family	Scientific name	English	Swedish	Norwegian	Danish	Finnish	Icelandic	Host	Target	Part used	Reference
Lichens	<i>Cetraria islandica</i>	Ice-land moss	islands-lav	islands-lav	lav *	jäkälä	fiatlaggräs *	M	H	W	8, 9, 33, 36, 37, 41
Ferns	<i>Pteridium aquilinum</i>	bracken fern	örnbräken *	einstepe *	ørnebregne *	sanamajaka *	amarburkni	M	H,C	R	9, 21, 35, 45
<i>Dryopteris carthusiana</i>	toothed wood fern	skogsbräken	broddteig	mangeløv	meislaelader	meislaelader *					31
<i>D. expansa</i>	spring wood fern	nordbräken	sauteig	finblad	isovaljejuuri *	flekkaþurkni					31
<i>D. filix-mas</i>	male fern	träjon *	orneteig *	almindelig mangelsøv *	kvikkvaljevjuuri *	stóri þurkni *	M,A,S,H	H,N,C,T	R		4, 5, 8, 9, 10, 30, 31, 33, 35, 44, 45, 53, 54, 70
Polypodiaceae	<i>Polypodium vulgare</i>	polypody	stensöla *	sisselrot	engelsød	kallioimarre *	köldugras	M	H,C	R	45, 49
Coniferous trees	<i>Picea abies</i>	spruce	gran *	rodgran	kuusi *	rauðgröni	M,A	H	H	J	29, 55
<i>Pinus sylvestris</i>	pine	tall *	furu	skovfyr	mänty *	skógarfura	M	C	H,P,T	45, 49	
<i>Juniperus communis</i>	juniper	en *	emer *	enebaer *	kataja *	emer *	M,A,C,H	H,N,C	B,R,P	9, 14, 25, 34, 44, 45, 53	
<i>J. sabina</i>	savin	sävenboom		sevenboom *	rohtokataja	sabimueinir	C,H,P		W		9
Trees, shrubs	<i>Populus tremula</i>	aspen	asp *	bævreasp	haapa *	ösp	M,H	H,N,C	L,P		34, 45
<i>Salix spp</i>	willow	säl	vier	pil *	paju *	viðir *	M,S,H	H,C,T	L,P		9, 33, 45
<i>Myrica gale</i>	sweet gale	pors	pors *	ors	suomyrtti	mjaðartving	M,H	H	W,L,F		9, 35
<i>Alnus glutinosa</i>	alder	al	or	el *	tervaloppi	rauðelri	S	T	H		9
<i>Betula pendula</i>	silver birch	värbjörk	hengebjörk	vorte-birk	raudskoivu *	M	C	L,J,P	45		45
<i>B. pubescens</i>	downy birch	glasbjörk	björk	dun-brk *	hieskoivu *	birki *	M,S	H,C	L,J,P,C		9, 33, 37, 45, 50
<i>Corylus avellana</i>	hazel bush	hassel	solbaer	hassel	pähkinäpensas	heslivður	H	L			9
<i>Ribes nigrum</i>	blackcurrant	svaria vinbär	solbaer	solbaer	musta viinimarja *	sölberjanunni	M	C	B		45
<i>R. rubrum</i>	redcurrant	röda vinbär	hagerrips	ribs *	punainen viinimarja	rifs	M				9
<i>Prunus padus</i>	bird cherry	hægg	hegg	hiere	tuomi *	heggur	M	C	B,P		45
<i>Pyrus communis</i>	pear	päron	pære	pære	päärynä	peruré	S	N	O		9
<i>Sorbus aucuparia</i>	rowan	rönn	rogn	ron	phlajaja	reymir	M	C	P		45
<i>Betula sempervirens</i>	boxwood	buxbom	buxbom *	buxbom	puksupu	fagurim	H	C	L		9
<i>Fraxinus alnus</i>	blackweed	brakved	trollhegg	torst	paansama *	porstaré	M	C	P		45, 53
<i>Daphne mezereum</i>	mezeron	tibast	tybsast *	pebertree	näsä *	löfraré	M	H,C	B,P		34, 35, 45, 55
<i>Fraxinus excelsior</i>	ash tree	ask *	ask	ask	saarni *	askur	M	H,C	L,P		45, 49
<i>Sambucus nigra</i>	black elder	fläder	svarthyll	hyld *	selja *	svartyllir	M,S	C,T	B		9, 45

Host (if specified): M = human, A = animal, C = cattle, S = sheep, H = horse, P = pig.

Target (if specified): H = helminths, N = nematodes, C = cestodes, T = trematodes, P = protozoa.

Part used (if specified): W = whole plant, L = leaves, F = flowers, S = seeds, B = berries, H = shoots or buds,

R = root or tuber, J = sap, P = phloem or bark, O = wood, C = charcoal or ashes, T = tar.

* Indicates that the plant is mentioned in literature from that Nordic country.

Note: Plants included in this table are based on generally historical reports. This does not imply endorsement by the authors in either their effectiveness against parasites, or safety for human and/or animal use.

Table 1. Continued

Plant family	Scientific name	English	Swedish	Norwegian	Danish	Finnish	Icelandic	Host	Target	Part used	Reference
Herbaceous plants											
<i>Cannabaceae</i>	<i>Humulus lupulus</i>	common hops	humle *	humle	humle *	humala *	humall	M,S	H,C,T	W,R,C	9, 45, 49
<i>Urticaceae</i>	<i>Urtica dioica</i>	stinging nettle	brämässla *	stornesle	stor nælde *	nokkonen	strömletta	M,S	H,T	W,S	9, 23, 44, 49
	<i>U. urens</i>	small nettle	ettelmässla	smånesle *	liden nælde	rautakononen	brenninetla *	M	H	W,S	41, 55
<i>Polygonaceae</i>	<i>Rumex acetosa</i>	green sorrel	ängssyra	engärsyre	rodne *	suolähinä *	tunsúra	M,P	H,C	L,F,R	9, 45
<i>Chenopodiaceae</i>	<i>Chenopodium ambrosioides</i>	goosefoot	citronmälla	melde	gäsefod	savikka *					31
	<i>C. bonus-henricus</i>	good king Henry	lungrot *	stolt henrik	stolthemiks-gäsefod	hyvänheikinsavikka	hinriksnjöfi	M	H,C	W	49
	<i>C. rubrum</i>	red goosefoot	rödmälla *	rodmelde	rod gäsefod	punasavikka		M	H	W	49
<i>Portulacaceae</i>	<i>Portulaca oleracea</i>	common purslane	portlak *	portullakk	portullak	vihannesportullakka	portullakka	M	H	W	44
<i>Caryophyllaceae</i>	<i>Agrostemma githago</i>	corncockle	klätt	klinte	klinte *	aurankukka	akursjama	M			9
<i>Ranunculaceae</i>	<i>Hepatica nobilis</i>	hepatica	bläsippa	blåveis	anemone *	sintuokko	blå-anemöna	M			9
<i>Fumariaceae</i>	<i>Corydalis intermedia</i>	corydalis	smånnäveört *	lerkespore	lerkespore	hentokiturunkammi *		M	H,C	R	45, 49
	<i>Fumaria officinalis</i>	common fumitory	jordrok *	jordroyk	lægge- jordrog	peltoemaikki	reykjurt	M	C	W	20
<i>Brassicaceae</i>	<i>Armoracia rusticana</i>	horseradish	pepparrot	pepparrot	peberrod *	piparijuuri	piparrot *	H	H	R	9, 41
	<i>Brassica spp.</i>	cabbage	kål *	kål	kål *	kaali, nauris *		M	H,C	S,R	9, 44, 45
	<i>Cardamine hirsuta</i>	hairy bitter-cress	ängsbräsma	rosetikarse	ros et	mäkitiukka	lambaklukkka *		H	L	8, 36, 37
	<i>C. pratensis</i>	cuckooflower	polarbräsma	polarikarse	polarkarse	poljanlitukka	hrafnaklukkka *	M	H	L	33, 41
	<i>polemonioides</i>										
<i>Descurainia sophia</i>		flixweed	stillfrö *	hundesennep	finbladet	liitullii *	þefjurt	M	H,N,C	S	22, 45, 49
<i>Lepidium sativum</i>		garden cress	kryddkrasse *	mårkarse	karse *	vihanneskrassi	karsi	M,S	H,T	W,S	9, 44
<i>Raphanus sativus niger</i>		garden radish	rätikka *	reddik	reddikke *	retikka	radisa	C,H	H	R	9, 44
<i>Sisymbrium officinale</i>		hedge mustard	vägsenap	vegsennep	vejsennep *	rohtopenaruoho	göudeurt	M			9
<i>Cruciferae</i>	<i>Rhodiola rosea</i>	roseroot	rosenrot	rosennep	rosennep	ruusujuuri	burnirot	C	H	R	55
	<i>Sampervivum tectorum</i>	house leek	taklök *	teklök	huslög *	mehtähti	húslautaur	M	C	W	9, 44
<i>Rosaceae</i>	<i>Agrimonia eupatoria</i>	agrimony	snåbörre *	åkermåne	agermåne *	maarianverjuuri		M	H	W	9, 44, 49
	<i>Fragaria vesca</i>	wood strawberry	smultron	markjordber	jordber *	mansikka *	villijar/æber	M	C	B	9, 45
	<i>Potentilla anserina</i>	silverweed	gåsört *	gåsennep	gäse-potentil	kerohahikki	tágamura	M	H	W	44
	<i>P. erecta</i>	tormentil	blodört *	tepperot	tormentil *	rätvävä	blómura	M,S	H,T	W,R	9, 44
	<i>Rubus chamaemorus</i>	cloudberry	ljorrton	molte	multibeber	lakka *	multuber	M	C	B	45
	<i>R. idaeus</i>	red raspberry	hallon	bringebær	hindeber *	vadelma *	hindber	M	C	B	9, 45
	<i>Sanguisorba officinalis</i>	great burnet	blodtopp *	bringebær	lægge kvessurt	punaluppio	blóðkollur	H	H	R	30
<i>Fabaceae</i>	<i>Vicia sativa</i>	common vetch	forderveiker	förvikke	vikke *	vikkeri	fóurflækja	H		W	9
<i>Rutaceae</i>	<i>Ruta graveolens</i>	common rue	virruta	virrute	rude *	ruutakasvi	rúðjurt	M			9
<i>Chusiaceae</i>	<i>Hypericum perforatum</i>	perforate St-John's-wort	ákta johannesört	prikkperikum	prikkbladet perikon *	måkkuisma	jónsmessurummi	M	H	F,H	9, 49, 51
	<i>H. maculatum</i>	imperfornate St-John's-wort	fyrkantig johannesört	frkant-perikum	kanet perikon	såimåkuisma *	flækjågullrunni	M	C		45
<i>Violaceae</i>	<i>Viola odorata</i>	sweet violet	luktvíol	marstíol	marst-víol *	nuoksuorvokki	ilmfjola	M			9
<i>Cucurbitaceae</i>	<i>Cucumis sativus</i>	cucumber	gurka	agurk	agurk	kurkku *	agúrka	M	C	S	45

Table 1. Continued

Plant family	Scientific name	English	Swedish	Norwegian	Danish	Finnish	Icelandic	Host	Target	Part used	Reference
	<i>Cucurbita pepo</i>	pumpkin	pumpa	gresskar	mandel-gresskar *	kupitsa *	grasker	M	N,C	S	9, 45
<i>Apiaceae</i>	<i>Anethum graveolens</i>	dill	dill	dill	dill *	tilli	dill	C	H	S,R	9
	<i>Angelica archangelica</i>	angelica	kvanne	kvaan	kvan	vänönputki	hvönn *	M,H	H	S,R	8, 32, 33, 36, 37, 41
	<i>Anthriscus cerefolium</i>	chervil	dansk körvel *	hagesjørvel	korvel *	maustekriveli	garðakerfyll	M	H	W	9, 44
	<i>Apium graveolens</i>	celeriace	selleri	hageselleri	selleri *	selleri	bláaþelja	M	H	W	9
	<i>Carum carvi</i>	caraway	kummin	karve *	kommen *	kumina *	kümen	M	H,C	W,S	9, 34, 45
	<i>Coriandrum sativum</i>	coriander	koriantar	koriantar	koriantar *	koriantari	koriantara	M	H,C	R	9
	<i>Daucus carota</i>	carrot	morot *	gulerot *	gulerot *	potkama *	gulrot	M,P	H,C	R	9, 35, 44, 45, 49, 55
	<i>Foeniculum vulgare</i>	fennel	fänkål	fennikel	fennikel *	fenkoli	fennikka	S	T		9
	<i>Levisticum officinale</i>	lovage	libbstikka *	løpstikke	løvstikke *	liperi	trolliarygg/	M	H	S	9, 44
	<i>Persicoffinum crispum</i>	parsley	persilja	persille	persille *	persilja *	steinselja	C	P	H	29
<i>Ericaceae</i>	<i>Calluna vulgaris</i>	heather	ljung *	rosslyng	lyng *	kanerva	beitilyng	S	T	B	9, 49
	<i>Vaccinium myrtillus</i>	blueberry	blåbær	blåbær	blåbær	mustikka *	aalbláberja-lyng	M	C	B	45
	<i>V. oxycoccos</i>	cranberry	tranebær	tranebær	tranebær	kapalo *		M	C	B	45
	<i>V. uliginosum</i>	bog bilberry	odon	blokkebær	mosebølle	juolukka *	bláberjalyng	M	C	B	45
	<i>V. vitis-idaea</i>	lingonberry	lingon	tyrtebær	tyrtebær	puolukka *	rauðberjalyng	M	C	B	45
<i>Gentianaceae</i>	<i>Centaurium litorale</i>	seaside centaury	kustarun	tusengyliden	tusingsyliden *	isoramiasappi	koldublióm	H			9
	<i>Gentianaella campesstris</i>	field gentian	fältgentiana	bakkesøte	breðhegret	keokatkerø	mariauvöundur *	H	H	R	8, 32, 33, 36, 37
<i>Menyanthaceae</i>	<i>Menyanthes trifoliata</i>	marsh trefoil	vattenklöver *	bukkeblad *	bukkeblad *	raate *	horblaöka *	M,S,H,P	H,C,T	W,L,R	4, 8, 9, 32, 33, 36, 37, 41, 45, 52, 55
<i>Verbenaceae</i>	<i>Verbena officinalis</i>	vervain	läkeverbena *	jernurt	jernurt	tarha verbana	jármurt	M	H	W,F	44
<i>Lamiaceae</i>	<i>Glachona hederacea</i>	ground-ivy	jordreva	krossknapp	konsknapp *	maahumala	krosshuappur	H		W	9
	<i>Hyssopus officinalis</i>	hyssop	isop *	isop	isop *	isoppi	isöpur	M	H,N	W	9, 44
	<i>Leonurus cardiaca</i>	motherwort	hjärtstilla	lovemale	almindelig	nukula		M			9
	<i>Melissa officinalis</i>	lemon balm	citronmeliss	sitronmelisse	hertespann *	hertensfyv *	bjartafró	C			9
	<i>Mentha crispata</i>	mint	krasmynta *	kattemynte	mynte *	miintu		M	N	W	9, 44
	<i>Nepeta cataria</i>	catnip	kattmynta *	liekjakatte-mynte	katteurt *	aitokkissanmiintu	kattarmynta	M	H	W	9, 44
	<i>Origanum vulgare</i>	wild majoram	kungsmynta	bergmynte	almindelig	mäkmeitami	kjarmynta	M			9
	<i>Salvia spp</i>	sage	salvia	salvie	salvie *	salvia	salvia	S	T	W	9
	<i>Thymus serpyllum</i>	wild thyme	backtimjan *	kryptimian	smalbladet timian	kangasajuruoho	blöðberg	M	H	W	38
	<i>T. vulgaris</i>	garden thyme	trädgårds-timjan *	kryddertimian	havetimian	timjami *	garðablöðberg	M	H,C	W	38, 45
<i>Solanaceae</i>	<i>Hyoscyomus niger</i>	henbane	bolnört	bulneurt	bulneurt *	hullukaali	skollarót	H		S	9
	<i>Nicotiana rustica</i>	small tobacco	tobak	tobakk	tobak *	tupakka	bönda-töbaksjurt	S	T	R	9
	<i>Solanum dulcamara</i>	woody nightshade	besksöa	slyngsotvier	bittersød	punakoiso	eiuritaekja	M			9
	<i>S. nigrum</i>	black nightshade	natskatta	svart sotvier	sort	mustakoiso	hümjurt	H		L,B	9
	<i>S. tuberosum</i>	potato	potatis	potet	kartoffel	peruna *	kartaffa	M	C	R	45

Table 1. Continued

Plant family	Scientific name	English	Swedish	Norwegian	Danish	Finnish	Icelandic	Host	Target	Part used	Reference
Scrophulariaceae	<i>Rhinanthus spp</i>	yellow rattle	skallra	kall	skjaller *	laukka	fokasjóður	M			9
	<i>Veronica anagallis-aquatica</i>	blue water speedwell	vattenveronika	vassveronika	areprits	komantädäcke	laugdepla *	M	H	W	8, 36, 37
	<i>V. chamaedrys</i>	germander speedwell	teveronika *	teskjegge-veronika	areprits	numtiädäcke	völudepla	M	H	W	44
<i>Plantaginaceae</i>	<i>Plantago lanceolata</i>	English plantain	svartkämpar	smalkjemppe *	lancet-vejbred	hemarátamo	selgrest *	M	H	L, R	8, 33, 35, 36, 41
	<i>P. major</i>	great plantain	groblad	groblad *	vejbred *	pihaaratamo	græjstira *	M	H	W, L, R	9, 35, 37, 41, 55
	<i>P. maritima</i>	sea plantain	gulkämpar	strandkjemppe	strand-vejbred	meriaratamo	kattartunga *	M	H	L, R	8, 36
<i>Valerianaceae</i>	<i>Valeriana officinalis</i>	valerian	vänderot *	lege-vendelrot	legebaldrian *	rohovirmajauri	garðabrúða	M, H	H	W, R	4, 9, 44, 49
<i>Asteraceae</i>	<i>Achillea millefolium</i>	yarrow	rölleka *	ryllike *	siakäräsämö	vallhumall		M	H	W, L, F	9, 34, 35, 38
	<i>Arenaria abrotanum</i>	southernwood	abrodd *	abrodd	ambra	aaprottimaruna		M	N	W	44
	<i>A. absinthium</i>	wormwood	malört *	ekte malurt *	malurt *	mali *		M, C, S, H	H, C	W, L, S	5, 9, 25, 34, 44, 45, 49, 53, 55, 58
	<i>A. vulgaris</i>	mugwort	gräbo *	burot	gräbynke *	pujo		M	H	W	9, 38, 49, 58
	<i>Ononis benedictus</i>	blessed thistle	kardbenedik * *	kardbenedik *	korbenedik *	kanvasohdake		M	H	W, L	9, 35, 44, 55
	<i>Helianthus annuus</i>	sunflower	solros	solsikke	solsikke	auringonkukka *	sólfífill	M	C	S	45
	<i>Inula helenium</i>	elecampane	átandsrot	átantrot *	legeatiant *	isohirveenjauri	hálsurt	M	H	R	9, 55
	<i>Matricaria maritima</i>	sea mayweed	kustbaldersbrá	strand-balderbrá	kamille *	merisaunio	baldursbrá *	M	H	L, F	8, 9, 33, 36, 37, 41
	<i>Senecio vulgaris</i>	common groundsel	korsört *	ákersvineblom	almindelig brandbeeger *	peltovilakko *	krossfífill	M, C	C	L	9, 22, 45
	<i>Silphium maritimum</i>	milk thistle	marististel	marististel	maricidsel *	maarianohdake *	maripustill	H	H	S	9, 53
	<i>Tanacetum balsamita</i>	alecost	balsamblad	balsam *	okseoje	palsamipäivän-kakkara		M	H	W	35
<i>Liliaceae</i>	<i>T. parthenium</i>	feverfew	mattam *	mattam	mattam	reunuspäivinkakkara		M	H	S	44
	<i>T. vulgare</i>	common tansy	reinfana *	reinfam *	reinfam *	pietaryrtti *	regnfang	M, A, S, H	H, N, C, T	W, L, F, S	1, 4, 9, 10, 19, 30, 34, 35, 44, 45, 49, 53, 55, 58, 70
<i>Liliaceae</i>	<i>Allium cepa</i>	onion	rödlök *	rodlok *	rödlög *	puunasipuli	rauðlaukur	M	H	W	9, 35, 44, 55
	<i>A. porrum</i>	leek	purjölök	purjölök *	purlög	purjo	bláðlaukur	M	H	W	34
	<i>A. sativum</i>	garlic	vitlök *	hvítlok *	hvídlög *	valkosipuli *	hvítlaukur	M, A	W, N, C	W	4, 9, 35, 44, 45, 55, 70
	<i>A. ursinum</i>	ransom	ranslök	ranslök *	rams-lög	karhumakka	bjarrautlaukur	M	H	W	34
	<i>Asparagus officinalis</i>	asparagus	sparris	aspargos	aspargos	ruokoparsa		M			9
<i>Araceae</i>	<i>Iris pseudacorus</i>	iris	svarðslíja *	sverdlíje	gul sverdlíje *	keltakurjenniikka	gula sverðlíja	C, P	H	R	9, 30
	<i>Acorus calamus</i>	sweetflag	kalmus	kalmustrot	kalmus	kalmjuurt *	kalmusrót	M	C	R	45
Grasses											
<i>Poaceae</i>	<i>Elytrigia repens</i>	couch grass	kvíkrot *	kveke	almindelig kvikgress *	juola vehmä	húsapuntur	M	H	W, R	9, 18
	<i>Hordeum vulgare</i>	barley	korn	bygg	bygg	ohra	bygg	M			9
	<i>Secale cereale</i>	rye	råg	rug	rug *	ruis	rúgur	M			9

Lichens and Ferns: One of the plants most commonly mentioned in the Nordic literature is male fern (*Dryopteris filix-mas*), a common fern that is widespread throughout the Northern hemisphere. Extracts from powdered rhizomes were first used by the Greeks (circa 350-250 BC) to treat tapeworm infections. This product (oil of aspidium) became an established product in many Pharmacopoeia of the Western World and was sold until the end of the 1940s. A number of active compounds have been isolated from this product, but it appears that the anthelmintic constituent is filicic acid.

Trees and Shrubs: Reports are few on the extensive use of trees and shrubs specifically to feed to livestock as treatment against parasites. However, products of willow (*Salix* spp.) have been widely used as analgesics or antipyretics in humans, probably attributable to the content of salicin and derivatives. *Salix* spp. also has a reputation as an anthelmintic for humans and livestock. Horses fed leaves are not supposed to get worms and a decoction of the bark is efficacious against flukes (trematode parasites) and diarrhoea in sheep (Brøndegård 1980).

Herbaceous Plants: There is a great variety of these plant types that has been used as deworming preparations. Whilst most of those mentioned in Table 1 thrive in the Nordic environment, many originated from other countries. Possibly one of the most widespread and commonly used herbal anthelmintic is oil of chenopodium, derived from *Chenopodium ambrosioides*, popularly known as American wormseed, or goosefoot. Archeological and ethnological studies suggest that this material has been used for many centuries. It is of passing interest that in the early eighteenth century, Peter Kalm (1715-1779), a Swedish botanist and traveller, reported that it was used by both the indigenous inhabitants and European settlers in the American colonies for the treatment of *Ascaris* infections. Plants were taken to Europe,

cultivated widely, and were soon in common usage. The active principle, ascaridol, a volatile terpene, was isolated and eventually synthesized. However, in the Nordic countries, *Chenopodium* is not one of the most commonly mentioned plant families.

Some of the plants mentioned are now commonly used as spices eg. caraway (*Carum carvi*), thyme (*Thymus* spp) and mint (*Mentha* spp). These have been found in Russian studies to have effect against *Trichostrongylus* larvae *in vitro* and also in sheep (Gadzhiev & Eminov 1986, Eminov 1982).

Members of the family *Asteraceae* have also a prominent position in the herbal deworming literature. The Romans used dried, unexpanded flower heads obtained from several species of the genus *Artemisia* in the first century, for the treatment of *Ascaris*, *Enterobius* and tapeworm infections. The name given for this herbal preparation was *semen-contra vermes* (semen against worms), apparently because of its superficial resemblance to semen. It became an important member of the European pharmacopoeia until the early 20th century. The active principle was found to be the sesquiterpene lactone, santonin. More recent pharmacological studies have demonstrated the pharmacological basis of this chemical. Low concentrations of santonin are reported to have a selective toxic action on the ganglion located in the nerve ring of *Ascaris* spp. (Sollmann 1957). Against other nematodes, such as *Oxyuris* spp and the cestodes, santonin is not effective (Steiniegger & Hänsel 1972). Pharmacological studies investigating the specific effects of santonin-containing herbal preparations are not known. This is probably because santonin had been isolated and used as a vermifuge as early as 1830. Due to its narrow therapeutic window (safety index) and toxicity, the crude drug santonin is no longer used (Reynolds & Prasad 1982, Tyler et al. 1988, De Smet 1997).

Another member of the *Asteraceae* family, more widely used in the Nordic countries, is common tansy (*Tanacetum vulgare*). The active component is claimed to be thujon. *In vitro* studies have shown an effect of this plant on *Trichostrongylus* and *Ostertagia circumcincta* spp. (Gadzhiev & Eminov 1986, Eminov 1982). Vegetables, such as carrot (*Daucus carota*), brassicas (*Brassica* spp), the onion group (*Allium* spp.), as well as all kinds of berries have had widespread use against parasites in the Nordic as well as most other countries. Seeds of pumpkin and cucumber (Cucurbitaceae) have been used in tropical America for centuries as a treatment of tapeworm infections. From there the popularity of this remedy spread to Europe. The active component, cucurbitine, was identified as an amino acid (3-amino 3 carboxy pyrrolidin). Leaves from another tropical plant, tobacco (*Nicotiana rustica*), have enjoyed universal popularity and latterly notoriety for use in smoking. However, infusions of this plant, or synthetic analogues (e.g. nicotine sulphate) were commonly used as anti-nematode preparations in ruminant livestock up until the advent of the modern broad spectrum anthelmintics in the mid 1950s. Both these plants have also been grown and used as anthelmintics in the Nordic countries.

Pasture plants: The possible use of specialised crops to control nematode infections in grazing ruminants has attracted considerable research interest in recent years. Bioactive plants or forages with secondary metabolites, particularly legumes with a high content of proanthocyanidins (condensed tannins) e.g. sulla (*Hedysarum coronarium*) or lotus major (*Lotus pedunculatus*) have been reported to reduce worm burdens in grazing lambs by up to 50% (Niezen *et al.* 1995). An *in vivo* anthelmintic effect has also been observed using quebracho, a condensed tannins extract, as a single high dose against sheep nematodes (Athanasidou *et al.*

1999) and the capacity of purified condensed tannins from Danish legumes to kill nematode larvae *in vitro* has been demonstrated (Kahiya *et al.* 1999). However, in several field studies it has been difficult to relate anti-parasitic effects to the actual amounts of condensed tannins (e.g. Niezen *et al.* 1998). A complicating factor is that condensed tannins are a poorly defined group of compounds (basically polymers capable of covalently binding protein) making standardised determinations in plant material difficult.

It has been postulated that the beneficial effects of tanniferous plants against internal parasites could be due to one, or a combination, of the following factors:

- Tanniferous plants increase the supply and absorption of digestible protein by animals. This is achieved by tannins forming non-biodegradable complexes with protein in the rumen, which dissociate at low pH in the abomasum to release more protein for metabolism in the small intestine of ruminants – in other words, "nature's protected protein." This indirectly improves host resistance and resilience to nematode parasite infections.
- Tannins have a direct anthelmintic effect on resident worm populations in animals.
- Tannins and/or metabolites in dung have a direct effect on the viability of the free-living stages (development of eggs to infective larval stages).

Although there is some evidence to support each of these above claims (for review, see Kahn & Diaz-Hernandez 1999), we believe that the data are by no means clear-cut (Bernes *et al.* 2000).

Limitations with using plants as natural anthelmintics

It is not a simple matter of just growing these plants and expecting them to be used in a natural parasite control system. In a longer perspec-

tive, many issues need to be considered. These include, whether the chosen plants are amenable to cultivation and if so by what means (pure stands or as mixed leys), ease of harvesting seeds and thus their commercial availability, and means of use or administration (grazing, or individual stable feeding – short, or long term). In addition, factors such as palatability, stability, biodegradability of active compounds in preserved products, whether these are to be used curatively, or preventively, need to be considered. Finally, dosage may be difficult to control and the possibility of toxic side effects in animals needs to be considered.

Some of the potential candidate plants cannot withstand trampling by livestock, are poor competitors with other pasture species in mixed grazing swards (e.g. *Lotus* spp., Beuselinck & Grant 1995), or they are preferentially sought out by grazing animals and thus easily succumb to even light grazing pressure (e.g. *H. coronarium*, Niezen et al. 1995).

Plants which have a high content of known direct-acting parasiticides (e.g. santonin in wormseed) may be effective for short-term "curative" use eg. a short grazing interval on a "deworming" paddock before a pasture change. In other cases, plants may have to constitute a substantial proportion of the feed and may therefore be used in a preventive fashion mixed with grass and clover in larger grazing areas, or in pure stands for rotational grazing.

A word of warning – plant toxicity

The whole animal kingdom is dependent on the use of plant material. Plants have probably covered much of our planet throughout the history of multicellular life. This implies that herbivores have been too few in number to consume all the food available (Hairston et al. 1960), and probably more importantly, that some plants have evolved defence mechanisms against being eaten by herbivorous animals (Murdoch

1966). One of their defence principles is the production of chemical compounds, which may be harmful or distasteful to potential herbivores. The fact that certain plants could have adverse effects on man and livestock has been known since ancient times. Likewise, it has been recognised that some plants could be of benefit in disease conditions. These two aspects of the plant kingdom, the beneficial and the harmful properties of plants, strongly related to dosage, are described in the early medical literature of classical Greece and Rome (Hippocrates, Theophrastus, Dioscorides).

Modern scientific literature on plant effects on livestock deals mainly with adverse effects, and less attention has been paid to the curative potential of plant material. The complex nature of this discipline is reflected in the difficulties in classification of poisonous plants. Attempts to classify them according to the chemical nature of their active constituents are met with the obstacles that these may be either a single substance or a number of substances with wide differences in chemical properties. Accordingly, a chemical classification will lead to considerable overlapping with some plants featuring in several chemical groups. Albeit these difficulties, the majority of recent textbooks group the poisonous plants according to their known toxic constituents (Cooper & Johnson 1998). These include a vast range of compounds that may be classified as alkaloids, glycosides, nitrates, oxalates, photodynamic substances, thiaminases, local irritants and phytoestrogens. The most reputed plants in the Nordic flora with reported responsibility for livestock poisoning include wolf's-bane (*Aconitum lycoctonum*) (alkaloids), cowbane (*Cicuta virosa*) (alkaloids), groundsel (*Senecio* spp.) (alkaloids), yew (*Taxus baccata*) (alkaloids), brassicas (*Brassica* spp.) (S-methyl cystein sulphoxide, progoitrin, nitrates, amongst others), foxglove (*Digitalis purpurea*) (digitalis-glycosides), bog

asphodel (*Narthecium ossifragum*) (saponin-glycosides), sorrel (*Rumex* spp.) (oxalates), St. John's-wort (*Hypericum* spp.) (photodynamic substances), bracken (*Pteridium aquilinum*) (thiaminases, alkaloids), horsetail (*Equisetum* spp.) (thiaminases) and *Ranunculus* spp. (local irritants) (Søli 1981, Laksesvala & Dishington 1983, Nærland 1984, Solberg 1984, Ulvund 1984, Waldeland 1984, Øverås 1984, Andersson *et al.* 1989, Sivertsen *et al.* 1992, Flåøyen *et al.* 1995, Flåøyen & Frøslie 1997). Several of these plants appear in Table 1 and thus are examples of both beneficial (according to popular belief!) and harmful properties of plants. One should be aware that there is some structural overlap between these simple chemical groups and that other classifications are possible. Furthermore, it must be borne in mind that little is known of the toxic constituents of many plants and that the poisonous properties of a plant may be due to more than one substance.

It is also important to understand that natural, often genetically determined, variations exist between different populations of some plant species, which may affect their toxic potential. An example of this is birdsfoot trefoil (*Lotus corniculatus*), in relation to cyanogenic glycosides. In different populations (cultivars), these substances may be absent, or the enzymes that break them down to release hydrocyanic acid may be absent, or both of them may be present. Consequently, if one of these constituents is lacking, the plant is harmless, and only when both are present the plant is poisonous. This knowledge is essential to make a rational selection of non-toxic cultivars of *L. corniculatus* as an important constituent for pasture leys.

Like any 'new' area of treatment against target pest organisms, the use of plant materials to control internal parasites should be tempered with common sense and based on scientific validation of a useful measure of efficacy, which at the same time is host benign.

Conclusion

This review provides ample evidence that a considerable amount of information relating to the use of plant material as de-worming preparations for man and his livestock in the Nordic countries, is available. However, almost all of these reports are historical and/or anecdotal. Evidence for effectiveness of plant de-worming preparations has been rarely obtained and little has been made available in scientific publications. With respect to increasing interest in the therapeutic use of natural products, we believe that it is important that a systematic evaluation is made of the botanical resources of the Nordic countries in relation to the purported de-worming properties of those plants that are endemic, or thrive, in this region of the world.

References

- Ahonen U: Fytoterapian käsikirja (Manual for Phytotherapy). Gummerus kirjapaino Oy, Saarijärvi, 1997.
- Andersson G, Hedhammar Å, Holmgren A, Persson H, Tjälve H: Förgiftningspanoramat hos djur. In: Förgiftningar hos djur (Poisoning panorama in animals. In: Poisoning in animals). Svensk Veterinärtidning 1989, 41, Supplement 19, 5-37.
- Athanasiadou S, Kyriazakis I, Coop RL, Jackson F: Evidence for direct anthelmintic effect of condensed tannins. WAAVP Conf. Copenhagen. 1999, f.5.04.
- Bergmark M: Vallört och vitlök (Comfrey and garlic). Prisma, Stockholm, 1967.
- Bergmark M: Lust och lidande (Lust and suffering). Natur & Kultur, Stockholm, 1968.
- Bernes G, Waller PJ, Christensson D: The effect of birdsfoot trefoil (*Lotus corniculatus*) and white clover (*Trifolium repens*) in mixed pasture swards on incoming and established worm infections in young lambs. Acta. Vet. Scand. 2000, 41, 351-361.
- Beuselinck PR, Grant WF: Birdsfoot trefoil. In: (Barnes RF, Miller DA, Nelson CJ, eds). Forages, vol. 1: An Introduction to Grassland Agriculture. 5th edit. Iowa State University Press, Ames, Iowa, USA, 1995, 237-248.
- Bjarnason A: Um íslenskar drykkurtir, söfnun þeirra, geymslu, nytsemi, verkanir og tilreiðslu (About

- Icelandic plants used in drinks, their collection, storage, usefulness, effects and preparation). Prentsmiðja Norður- og Austuramtsins hjá H. Helgasyni, 1860.
- Brøndegaard VJ*: Folk og Flora – Dansk etnobotanik, vol. 1-4 (People and flora - Danish ethnobotany, vol. 1-4). Rosenkilde og Bagger, 1980.
- Cantell S, Saarnio V*: Suomen myrkylliset ja lääkekasvit, niiden vaikuttavat aineet, vaikutukset elimistöön, lääkkeinä käyttö sekä rohdoksis keräily ja viljely (Toxic and medicinal plants in Finland. Effective ingredients, function and use in medicine. Cultivation and collection). Karisto Oy, Hämeenlinna, 1936.
- Cooper MR, Johnson AW*: Poisonous plants and fungi in Britain. 2nd edit. The Stationary Office, London, 1998.
- Danøe R, Bøgh HB*: Usage of herbal medicine against helminths in livestock. An old tradition gets its renaissance. World Animal Review 1999, 93, 60-67.
- De Smet PAGM (Ed)*: Adverse Effects of Herbal Drugs. Springer-Verlag, Berlin Heidelberg New York, 1997, 3.
- Einarsson, G*: Um sauðfænað (About sheep). Prentsmiðja Ísafoldar, Reykjavík, 1879.
- Eminov RS*: Effect of certain pasture plants on gastrointestinal nematodes of sheep. Sov. Agric. Sci. 1982, 1, 72-74.
- Flåøyen A, Binde M, Bratberg B, Dønne B, Fjølstad M, Grønstøl H, Hassan H, Mantle PG, Landsverk T, Schönheit J, Tønnessen MH*: Nephrotoxicity of *Nartheceum ossifragum* in cattle in Norway. Vet. Rec. 1995, 137, 259-263.
- Flåøyen A, Frøslie A*: Photosensitization disorders. In: Handbook of plant and fungal toxicants. D'Mello JPF ed., CRC Press, Boca Raton, 1997, 191-204.
- Fogelfors H*: Nyttiga ogräs, del 2 (Useful weeds, part 2). Lantmannen, 1984, 3, 19.
- Fogelfors H*: Nyttiga ogräs, del 7 (Useful weeds, part 7). Lantmannen, 1984, 8, 51.
- Fogelfors H*: Nyttiga ogräs, del 10 (Useful weeds, part 10). Lantmannen, 1984, 11, 31.
- Fogelfors H*: Nyttiga ogräs, del 16 (Useful weeds, part 16). Lantmannen, 1984, 18, 38.
- Fogelfors H*: Nyttiga ogräs, del 17 (Useful weeds, part 17). Lantmannen, 1984, 21, 44.
- Fogelfors H*: Nyttiga ogräs, del 22 (Useful weeds, part 22). Lantmannen, 1985, 3, 52.
- Gadzhiev YG, Eminov RS*: Action of medicinal plants on gastrointestinal nematodes of sheep. Byuletén Vsesoyuznogo Instituta Gelmintologii im. K.I. Skryabina, 1986, 44, 12-16.
- Ganander C*: Eläinden Tauti-kirja, kolmas ylösspano (Disease of Animals). 3rd edit. J. D. Frendell ja poika, Helsinki & Turku, 1829.
- Gibson TE*: Factors influencing the application of anthelmintics in practice. Vet. Parasitol. 1980, 6, 241-245.
- Grove DI*: The discovery and development of anthelmintics. In: A History of Human Helminthology DI Grove (ed). C.A.B. International, Oxford, 1990, 75-101.
- Hairston NG, Smith FE, Slobodkin LB*: Community structure, population control, and competition. American Naturalist 1960, 94, 421-425.
- Hako M*: Kansanomainen lääkintätietous (Medicine in Folklore). 2nd edit. Suomalaisen Kirjallisuuden Seuran Toimituksia 229:4, KJ Gummerus OY, Jyväskylä, 1982.
- Hewe N*: Välsignade växter (Blessed plants). Stockholm, 1939.
- Hiltunen R, Holm Y*: Luonnonlääkkeet (Natural Medicine). Helsingin yliopisto oppimateriaaleja 27. Painotalo Miktor, Helsinki, 1994.
- Hjaltalín J*: Lækninga-Bók um þá helztu kvilla á kvikfænaði, samantekin fyrir Íslendinga og lögðu eptir þörfum þeirra, af Jóni Hjaltalín, Studios. Chirurg. et Medic. (Medical book about the main ailments of livestock, compiled for the Icelanders and adapted to their needs, by Jóni Hjaltalín, Studios. Chirurg. et Medic.). Kaupmannahöfn, 1837.
- Hjaltalín J, Oddur J*: Íslensk grasafæði (Icelandic botany). Gefin út að tilhlutun Hins íslenska bókmenntafélags, Kaupmannahöfn, 1830.
- Høeg OA*: Planter og tradisjon (Plants and tradition). Universitetsforlaget, Oslo-Bergen-Tromsø, 1976.
- Høeg OA*: Våre medisinske planter (Our medical plants). Forlaget Det Beste, 1984.
- Jónsson BL*: Íslenskar lækninga- og drykkjarjurtir (Icelandic medical plants and plants used as drinks). 13. Rit Náttúrulekningafélags Íslands, Reykjavík, 1977.
- Jónsson J*: Lítil ritgjörð um nytsemi nokkurra íslenskra jurta eptir ymsa höfunda (A booklet about the usefulness of some Icelandic plants by various authors). Einar Þórðarson, Reykjavík, 1880.
- Juneby HB*: Junebys medicinalväxter (Juneby's medical plants). Reformförlaget, Malmö, 1984.
- Kahiya CJ, Mukaratirwa S, Thamsborg SM, Ndlovo LR*: Anthelmintic effects of proanthocyanidins and related polyphenolics. Proc. WAAVP Conf.

- Copenhagen, 1999, g 6.86.
- Kahn LP, Diaz-Hernandez A*: Tannins with anthelmintic properties. International Workshop on Tannins in Livestock and Human Nutrition. Waite Campus, University of Adelaide. ACIAR Proceedings No 92, 1999, 130-139.
- Kristjánsson G, Sigfússon B (Ed.)*: Rit Björns Halldórssonar í Sauðlauksdal (The writings of Björn Halldórsson of Sauðlauksdalur). Búnaðarfélag Íslands, Reykjavík, 1983.
- Laksesvala B, Dishington IW*: Bog asphodel (*Narthe-cium ossifragum*) as a cause of photosensitisation in lambs in Norway. *Vet. Rec.* 1983, 112, 375-378.
- Murdoch WW*: Community structure, population control, and competition: A critique. *American Naturalist* 1966, 100, 219-226.
- Månsson A*: En mycket nyttigh örta-book (A very useful book of herbs). 1642. Rediviva, Stockholm, 1987.
- Naakka-Korhonen M*: Vaivasta taudiksi. Lapamaton liittyvä kansanparannus erityisesti pohjoiskarjalaisen aineiston valossa (Dissertation: From Illness to Disease. Ethnomedicine associated with broad tapeworm with particular reference to material from Northern Karelia, Finland). Suomalaisen Kirjallisuuden Seuran Toimituksia 666. Kirjakas/Kirjapaino Raamattutalo, Pieksämäki, 1997.
- Niezen JH, Waghorn GC, Charleston WAG*: Establishment and fecundity of *Ostertagia circumcincta* and *Trichostrongylus colubriformis* in lambs fed lotus (*Lotus pedunculatus*) or perennial ryegrass (*Lolium perenne*). *Vet. Parasitol.* 1998, 78, 13-21.
- Niezen JH, Waghorn TS, Charleston WAG, Waghorn GC*: Growth and gastrointestinal nematode parasitism in lambs grazing either lucerne (*Medicago sativa*) or sulla (*Hedysarum coronarium*) which contains condensed tannins. *J. Agric. Sci. Camb.* 1995, 125, 281-289.
- Nærland G*: Planteforgiftninger hos drøvtyggere i Rogaland. Et tilbakeblikk. In: Giftige planter og planteforgiftninger hos dyr i Rogaland (Plant poisoning in ruminants in Rogaland. A retrospect. In: Poisonous plants and plant poisoning in animals in Rogaland). Rogaland Veterinærforenings seminar på Utstein Kloster 1984, 6-9.
- Ohlmarks Å*: Linnés hälsoörter (The medical plants of Linné). Sjöstrands förlag, Stockholm, 1985.
- Ólafsson E*: Ferðabók Eggerts Ólafssonar og Bjarna Pálssonar. Um ferðir þeirra á Íslandi árin 1752-1757. I og II bindi (The travel book of Eggert Ólafsson and Bjarni Pálsson. About their travels in Iceland in 1752-1757. Vol. I and II). Bókauktáfang Órn og Örlygur h.f., Reykjavík, 1981.
- Pettersson B, Tunón H, Svanberg I*: Johannesört (St John's wort). *Fauna och Flora* 1998, 93, 202-206.
- Pétursson J*: Jóns Péturssonar Handlæknis Norðlendinga frá 1775 til 1801 Lækninga-Bók fyrir almúga (The medical book of Jón Pétursson for the public). Yfirlésin, aukin og endurbætt af Landphysikus Jóni Þorsteinssyni og Handlækni Sveini Pálssyni. Útgefin af Þorsteini Jónssyni Stud. Theol., Kaupmannahöfn, 1834.
- Piippo S*: Luonnon lääkekasvit (Medicinal Herbs in Nature). Helsingin yliopiston kasvitieteen monistea 144. Helsinki, 1996.
- Rautavaara T*: Mihin kasvimme kelpaavat, ruokaa, ryytiä ja rohtoa luonnosta (The use of our plants: Food, Drugs and Herbs from Nature). 3rd edit. Werner Söderström OY, Porvoo-Helsinki-Juva, 1976.
- Reichborn-Kjennerud I*: Våre folkemedisinske lægeurter (Our medical herbs). Centraltrykkeriet, Kristiania, 1922.
- Reynolds JEF, Prasad AB (Eds)*: Martindale The Extra Pharmacopoeia. 28th edit. The Pharmaceutical Press, London, 1982.
- Sivertsen T, Øvernes G, Karlsen B, Söli N*: Forgiftninger hjå norske husdyr i 1990 (Poisoning in Norwegian domestic animals in 1990). *Norsk Veterinærtidsskrift* 1992, 104, 173-182.
- Smidt D*: Brugen af urter i græsmarken til husdyr (Use of herbs in pasture). KVL, M.Sc. Thesis, 1997.
- Solberg I*: Aktuelle planteforgiftninger hos storfe og hest. In: Giftige planter og planteforgiftninger hos dyr i Rogaland (Plant poisoning in cattle and horses. In: Poisonous plants and plant poisoning in animals in Rogaland). Rogaland Veterinærforenings seminar på Utstein Kloster 1984, 16-20.
- Sollmann T*: A manual of Pharmacology and its Application to Therapeutics and Toxicology. 8th edit. W.B Saunders Company, Philadelphia, 1957.
- Steinegger E, Hänsel R*: Lehrbuch der Pharmacognosie (Textbook of pharmacognosy). 3rd edit. Springer Verlag, Berlin, 1972.
- Svensson C, Hessle A, Höglund J*: Parasite control methods in organic and conventional dairy herds in Sweden. *J. Livestock. Prod.* 2000, 66, 57-69.
- Söli NE*: Planteforgiftninger hos sau med særlig henblik på norske forhold (Plant poisoning in sheep

- with focus on the situation in Norway). Norsk Veterinærtidsskrift 1981, 93, 87-91.
- Thamsborg SM, Jørgensen RJ, Waller PJ, Nansen P:* The influence of stocking rate on gastrointestinal nematode infections of sheep over a two-year grazing period. Vet. Parasitol. 1996, 67, 207-224.
- Thamsborg SM, Roepstorff A, Larsen M:* Integrated and biological control of parasites in organic and conventional production systems. Vet. Parasitol. 1999, 84, 169-186.
- Tyler VE, Brady LR, Robbers JE:* Pharmacognosy. 9th edit. Lea and Fabiger, Philadelphia, 1988.
- Ulvund MJ:* Alveld og andre sjukdommer med symptomer på fotosensibilitet hos sau. In: Giftige planter og planteforgiftninger hos dyr i Rogaland ("Alveld" (hepatogenous photosensitization) and other diseases with symptoms of photosensitivity in sheep. In: Poisonous plants and plant poisoning in animals in Rogaland). Rogaland Veterinærforenings seminar på Utstein Kloster 1984, 23-29.
- Vaarst M:* Veterinær homøopati: baggrund, principper og anvendelse med speciel fokus på økologiske malkekvægsbesætninger – et litteraturreview (Veterinary homeopathy: background, principles and use, with special focus on organic dairy herds – a literature review). Report from Danish Institute of Animal Science No. 731, 1996.
- Vaarst M, Ploeger H, Thamsborg SM, Sørensen JT:* Organic dairy farming and nematode parasitism. Infection patterns among replacement heifers in dairy herds not preventively treated with anthelmintics. In: Livestock farming systems. (Dent, McGregor, Sibbald eds.) Wageningen Press. EAAP Publication 1996, No. 79, 85-90.
- Vennerholm J, Dahlström H, Stålfors H:* Husdjurens sjukdomar (Diseases of domestic animals). Albert Bonniers förlag, Stockholm, 1920.
- Waldeland H:* Aktuelle planteforgiftninger hos sau i Rogaland. In: Giftige planter og planteforgiftninger hos dyr i Rogaland (Plant poisoning in sheep in Rogaland. In: Poisonous plants and plant poisoning in animals in Rogaland). Rogaland Veterinærforenings seminar på Utstein Kloster 1984, 30-37.
- Øverås J:* Forgiftning med planter som gir hemolytisk anemi og/eller methemoglobinemi. In: Giftige planter og planteforgiftninger hos dyr i Rogaland (Poisoning with plants leading to haemolytical anaemia and/or methemoglobinaemia. In: Poisonous plants and plant poisoning in animals in Rogaland). Rogaland Veterinærforenings seminar på Utstein Kloster 1984, 13-16.

Sammanfattning

Växter för avmaskning av husdjur i de nordiska länderna: historiskt perspektiv, folklig tro och möjlig framtida användning.

Att söka efter substanser inom växtriket som ger bot och lindring mot sjukdomar (däribland parasiter) såväl hos sig själv som sin boskap är något som har varit föremål för mänsklig aktivitet sedan urminnes tider. Framställningen av örtmediciner har varit beroende av den lokala floran, vilket har medfört att olika preparat utvecklats i olika delar av världen. Dock har i vissa fall samma eller närbesläktade arter använts i många länder. Det har även varit vanligt med import av växter med högt anseende. De nordiska länderna har historiskt sett haft ett rikt och varierat utbud av anthelmintika från växtvärlden, för bruk till såväl människor som husdjur. Detta har varit såväl importerade preparat som endemiskt förekommande växter och växter som kan odlas i Norden. Mycket kändedom och erfarenhet av naturligt förekommande läkeväxter har emellertid gått förlorad i västvärlden. Detta som en konsekvens av utvecklingen och lanseringen av en mängd effektiva, säkra, syntetiskt framställda läkemedel med bred verkan. Under senare år har dock intresset till dessa produkter i viss mån förändrats och det finns ett förnyat intresse för olika typer av naturläkemedel. Det beror delvis på att vissa preparat blivit verkningslösa till följd av att resistens utvecklats hos sjukdomsorganismerna. Avsikten med denna litteraturoversikt är att redovisa uppgifter framförallt från äldre nordiska källor om vilka växter som har använts i avmaskande syfte. En diskussion förs också om på vilka områden ny forskning kan ge värdefull och användbar kunskap.

(Received June 5, 2000; accepted September 8, 2000).

Reprints may be obtained from: P.J. Waller, Department of Parasitology (SWEPAR), Statens Veterinärmedicinska Anstalt, S-751 889 Uppsala, Sweden. E-mail: Peter.Waller@sva.se, tel: +46 18 67 41 27, fax: +46 18 30 91 62.