



METHODOLOGY

Open Access

Optimizing plant transporter expression in *Xenopus* oocytes

Huimin Feng^{1,2}, Xiudong Xia¹, Xiaorong Fan^{1,2*}, Guohua Xu¹ and Anthony J Miller^{2*}

Abstract

Background: Rapid improvements in DNA synthesis technology are revolutionizing gene cloning and the characterization of their encoded proteins. *Xenopus laevis* oocytes are a commonly used heterologous system for the expression and functional characterization of membrane proteins. For many plant proteins, particularly transporters, low levels of expression can limit functional activity in these cells making it difficult to characterize the protein. Improvements in synthetic DNA technology now make it quick, easy and relatively cheap to optimize the codon usage of plant cDNAs for *Xenopus*. We have tested if this optimization process can improve the functional activity of a two-component plant nitrate transporter assayed in oocytes.

Results: We used the generally available software (<http://www.kazusa.or.jp/codon/>; <http://genomes.urv.es/OPTIMIZER/>) to predict a DNA sequence for the plant gene that is better suited for *Xenopus laevis*. Rice *OsNAR2.1* and *OsNRT2.3a* DNA optimized sequences were commercially synthesized for *Xenopus* expression. The template DNA was used to synthesize cRNA using a commercially available kit. Oocytes were injected with cRNA mixture of optimized and original *OsNAR2.1* and *OsNRT2.3a*. Oocytes injected with cRNA obtained from using the optimized DNA template could accumulate significantly more NO₃⁻ than the original genes after 16 h incubation in 0.5 mM Na¹⁵NO₃. Two-electrode voltage clamp analysis of the oocytes confirmed that the codon optimized template resulted in significantly larger currents when compared with the original rice cDNA.

Conclusion: The functional activity of a rice high affinity nitrate transporter in oocytes was improved by DNA codon optimization of the genes. This methodology offers the prospect for improved expression and better subsequent functional characterization of plant proteins in the *Xenopus* oocyte system.

Keywords: DNA optimization, *Xenopus* oocyte, Plant, Nitrate transporter, Uptake, Electrophysiology

Background

Heterologous expression systems are often used for the functional characterization of a gene. *Xenopus laevis* oocytes are widely used to express membrane proteins and channels. Over twenty years ago, the first plant membrane proteins were expressed in oocytes and these were a hexose transporter and a K⁺ channel [1,2]. Since then, many plant membrane proteins including carriers [3-5], channels [6-9] and aquaporins [10-13] have been successfully expressed in oocytes. Oocyte expression was used to demonstrate function for the first plant nitrate transporter (Chl1, AtNRT1.1

or AtNPF6.3) that was identified and later for many more family members [3,14-20]. Some of the plant NRT2 nitrate transporter family members require a second gene NAR2 for function and this requirement was demonstrated using oocyte expression [4,5,21-23]. The high affinity rice nitrate transporter, OsNRT2.3a needs a partner protein, OsNAR2.1 for function in oocytes [22,23].

Although all organisms generally share the same genetic code, each genus has evolved a slightly different pattern of codon usage. Heterologous protein expression in a foreign host may be diminished by factors such as biased codon usage, GC content and repeat sequences. To overcome these limitations, codon optimization can be used to enhance gene expression in various host cells. Heterologous synthetic genes with codon optimization showed increased expression levels in various organisms including *E. coli* [24,25], yeast [26] and mammalian cells [27,28]. For many

* Correspondence: xiaorongfan@njau.edu.cn; tony.miller@jic.ac.uk

¹State Key Laboratory of Crop Genetics and Germplasm Enhancement, College of Resources and Environmental Sciences, Nanjing Agricultural University, Nanjing 210095, PR China

²Department of Metabolic Biology, John Innes Centre, Norwich Research Park NR4 7UH, UK

plant transporters expressed in *Xenopus* oocytes, the low levels of expression can often limit the functional assay, making the detailed characterization of the protein difficult. In the past, it was speculated that differing codon bias may explain the very low levels of expression of some plant proteins in oocytes [29]. Improvements in DNA synthesis technology have enabled the technique to be used for cost-effective gene cloning. Commercial suppliers make it possible to obtain the synthetic DNA with codon optimization in just a few weeks. In this study, DNA of the rice genes *OsNAR2.1* and *OsNRT2.3a* were codon optimized and synthesized for oocyte expression. The cRNA of *OsNAR2.1* and *OsNRT2.3a* were then synthesized using a commercially available kit. We compared how this process may improve the functional activity of plant nitrate transporter proteins expressed in oocytes. The nitrate transport activity was assayed using ^{15}N -enriched nitrate uptake and the two-electrode voltage clamp technique.

Results and discussion

Codon optimization of *OsNAR2.1* and *OsNRT2.3a*

There are some general rules that emerge from the analysis of the preferred codons in *Xenopus* and these can be used to optimize a gene sequence for expression in oocytes [30]. Using the codon usage bias software for *Xenopus* (<http://www.kazusa.or.jp/codon/>; <http://genomes.urv.es/OPTIMIZER/>), the DNA gene sequence was optimized for *OsNAR2.1* (LOC_Os02g38230) and *OsNRT2.3a* (LOC_Os01g50820) and the resulting DNA sequences were synthesized by the Genescript Company and named *syn-OsNAR2.1* and *syn-OsNRT2.3a*. After software optimization, the predicted GC content of *syn-OsNAR2.1* and *syn-OsNRT2.3a* was adjusted from 72.0 to 52.6% and 67.2 to 49.0% respectively, when compared with the original


genes (Table 1). This change now makes the plant genes synthetic DNA much closer to the typical 50% GC content found in *Xenopus* [31]. For both synthetic DNAs the melting temperature (T_m) was decreased and the number of repeat sequences was decreased in *syn-OsNAR2.1* (see Table 1). Sequence alignment of the open reading frames showed that *syn-OsNAR2.1* and *syn-OsNRT2.3a* shared 73% and 74% identity with the original genes (Figure 1), but the amino acid sequences did not change after optimization (see Additional file 1).

Nitrate uptake of oocytes

The original and synthetic (optimized) *OsNAR2.1* and *OsNRT2.3a* were subcloned in to expression vector pT7Ts [30] and then used as template to synthesize mRNA. Mixed mRNA of either synthetic genes (*syn-OsNAR2.1*: *syn-OsNRT2.3a*) or the original genes (*ori-OsNAR2.1*: *ori-OsNRT2.3a*) were injected into oocytes. Both RNA mixes were injected at the same ratio (*OsNAR2.1*:*OsNRT2.3a*, 25:50 ng). We used a colorimetric method to assay the amount of nitrate accumulated inside the oocyte. After 16 h incubation in MBS containing 0.5 mM NaNO_3 , oocytes injected with mRNA of synthetic genes showed increased NO_3^- uptake when compared with the original genes (Figure 2). Similar results were obtained in 5 mM NaNO_3 (Additional file 2). These data did not show a significant difference between the original genes and water injected oocytes. In another set of experiments, injected oocytes were incubated in MBS containing 0.5 mM $\text{Na}^{15}\text{NO}_3^-$ for 8 and 16 h. Compared to original genes, $^{15}\text{NO}_3^-$ uptake of single oocyte injected with synthetic genes were greatly enhanced after 8 h and 16 h incubation (Figure 3). Individual oocytes injected with synthetic gene mRNAs generally showed much greater $^{15}\text{NO}_3^-$ uptake

Table 1 DNA sequence parameters of optimized plant transporter genes *OsNAR2.1* and *OsNRT2.3a*



	<i>OsNAR2.1</i>		<i>OsNRT2.3a</i>	
	Synthetic	Original	Synthetic	Original
GC content (%)	52.6	71.0	49.0	67.2
Max direct repeat	Size:	8	10	13
	Distance:	7	984	993
	Frequency:	2	2	2
Max inverted repeat	Size:	10	12	11
	Tm	27.3	48.8	53.6
Repeat sequence	Start positions	3, 475	230, 1497	764, 615
	Max dyad repeat	size	9	10
	Tm	27.5	31.9	50.2
	Start positions	151, 450	198, 1156	309, 1442
Cis-acting element	PolyT	0	0	0
	PolyA	0	0	0




the optimized DNA. These data demonstrate the significant methodological advantage of using a template DNA that has been optimized for *Xenopus* expression.

Enhanced transporter activity in oocytes

The expression of foreign proteins in oocytes has long been known to be improved by the inclusion of a polyA tail and the use of expression vectors that include *Xenopus*



globin flanking UTR (untranslated region) sequence alongside the foreign DNA [1,29,30]. The polyA tail is recognized to improve mRNA stability and lifetime in the oocyte thereby improving the production of a foreign protein. The frog globin flanking UTR sequence is thought to make the heterologous cDNA more *Xenopus*-like and therefore thought to improve translation [29,30]. Similarly the mRNA produced from the synthetic DNA is more like *Xenopus* message and this has resulted in improved translational efficiency in the oocyte. For *OsNAR2.1* the removal of some repeat sequence (Table 1) may also have given better translation of the foreign protein. Making the GC content more like the 50% found in *Xenopus* [31] is likely to improve expression of plant genes. In *Arabidopsis* the GC content was reported as 44%, on the other hand in maize, rice and barley the figure was higher at >60% [32]. Mammals usually have around 44% GC in their coding sequence and experimental work directly comparing low-GC genes with their high-GC counterparts showed 100-fold greater expression in the GC-rich genes [33]. This study also



showed that the mRNA degradation rate was independent of the GC content.

Together these data clearly show the methodological advantage for plant genes of using a synthetic template that has codon usage more like that found in *Xenopus*. It is widely accepted that optimized codons help to achieve faster translation rates and high accuracy in bacteria, yeast and mammalian cells [24-28,34] and we now show this has advantages for plant genes in *Xenopus* oocytes too.

Conclusion

In this study, rice nitrate transporter *OsNAR2.1* and *OsNRT2.3a* were codon optimized for *Xenopus laevis*. Nitrate transport activity was analyzed and compared between oocytes injected with different sources of template DNA. The optimization changes the DNA, but not the protein sequence. Compared with the original plant genes, oocytes injected with optimized genes had increased nitrate uptake and larger currents in electrophysiological analyses suggesting that there was an increased level of protein expression. Taken together, these data show that the codon optimized template can give much improved expression and therefore provides a big advantage when aiming to functionally characterize a plant transporter protein in the *Xenopus* oocyte system. Although this may not be the case for all plant transporter genes the relatively cheap cost of DNA synthesis now makes this worthwhile when using oocyte expression.

Methods

Cloning and mRNA synthesis of *OsNAR2.1* and *OsNRT2.3a*

OsNAR2.1 and *OsNRT2.3a* were codon optimized and synthesized by the Genescript Company. cDNAs were then

subcloned into the *Bgl*III and *Spe*I sites of the oocyte expression vector pT7TS [35] using a directional cloning method. Original *OsNAR2.1* and *OsNRT2.3a* construct are as described previously [22]. Plasmid was linearized by *Bam*H I (Roche) and purified by PCR purification kit (QIAGEN). The mRNA synthesis kit (mMESSAGE mMACHINE[®] T7 Kit, Ambion) was used to synthesize the mRNA of all genes. A compressed air system (Harvard) was used for injection [1]. Glass tips were calibrated for injection using known volumes and 1 μ l mRNA mixtures (0.5 μ g *OsNAR2.1* and 1 μ g *OsNRT2.3a* in total) were used to inject around 22–25 oocytes. Thus mRNA mixtures were injected as 25 ng: 50 ng, *OsNAR2.1*: *OsNRT2.3a*, and this weight ratio was chosen to reflect the differing molecular sizes and give similar molecular ratios [22,23]. Oocytes from three different frogs were used for the data shown.

NO_3^- accumulation and $^{15}\text{N-NO}_3^-$ uptake in oocytes

After injection, the oocytes were incubated for 2 days in NO_3^- free MBS solution (88 mM NaCl, 1 mM KCl, 2.4 mM NaHCO_3 , 0.71 mM CaCl_2 , 0.82 mM MgSO_4 and 15 mM HEPES, pH 7.4). The solution contains 10 g/ml sodium penicillin and 10 g/ml streptomycin sulphate. For NO_3^- measurement, oocytes were incubated in MBS solution containing 0.5 mM NaNO_3 at 18°C for 16 h. After incubation, oocytes were washed with NO_3^- free MBS for four times. Four oocytes were collected as one sample in 1.5 mL tube, and 100 μ l H_2O was added into the tube. Lyses the oocytes and centrifuge the tube at 13000 rpm. The supernatants (40 μ l) were collected for NO_3^- assay using the kit (Nitrate/Nitrite Colorimetric Assay kit, Cayman).

For $^{15}\text{N-NO}_3^-$ measurement, oocytes were incubated in MBS solution containing 0.5 mM $\text{Na}^{15}\text{NO}_3$ with a 99% atom excess of ^{15}N for 8 h and 16 h. oocytes were washed four times with ice-cold 0.5 mM NaNO_3 MBS. Single oocyte was transferred to an empty tin capsule and then dried at 60°C for one week. Analysis for total ^{15}N content using a continuous-flow isotope ratio mass spectrometer coupled with a carbon nitrogen elemental analyzer (ANCA-GSLMS; PDZ Europa). The delta- ^{15}N was calculated as described previously [4].

Two-electrode voltage clamp analysis

The nitrate-elicited currents were recorded in oocyte using two-electrode voltage clamp method (pClamp 10.2, Axon). The oocytes were incubated in nitrate-free MBS and then treated with MBS containing 0.5 mM sodium nitrate. Membrane potential of oocytes was pulsed from 0 to -160 mV with 20 mV incremental steps. The currents were recorded to obtain current-voltage curves [2-4].

Additional files

Additional file 1: Figure S1. Alignment of synthetic and original OsNAR2.1 and OsNRT2.3a amino acid sequences showing they encoded identical proteins.

Additional file 2: Nitrate accumulation in *Xenopus* oocytes in different concentration of nitrate. Oocytes were incubated in MBS with 0.5 mM and 5 mM NaNO₃ for 16 h and washed four times with NO₃⁻ free MBS solution. Four oocytes were pooled for each sample. The values are means SE of four replicates with a and b indicating the statistical significance at $p \leq 0.05$.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AJM and XF conceived the study. HF, XX and XF carried out the experiments. HF drafted the manuscript and all authors approved the final version.

Acknowledgements

This work was funded by the China 973 Program (grant number 2011CB100300), the National Natural Science Foundation, the PAPD and 111 Project (grant number 12009). AJM is supported by grant funding BB/JJ004553/1 from the BBSRC and the John Innes Foundation. The oocytes used in this work were kindly donated by the Gurdon Institute, Cambridge, UK.

Received: 2 October 2013 Accepted: 13 December 2013

Published: 20 December 2013

References

- Boorer KJ, Forde BG, Leigh RA, Miller AJ: **Functional expression of a plant plasma membrane transporter in *Xenopus* oocytes.** *FEBS Lett* 1992, **302**:166–168.
- Cao YW, Anderova M, Crawford NM, Schroeder JI: **Expression of an outward-rectifying potassium channel from maize mRNA and complementary RNA in *Xenopus* oocytes.** *Plant Cell* 1992, **4**:961–969.
- Zhou JJ, Theodoulou FL, Muldin I, Ingemarsson B, Miller AJ: **Cloning and functional characterization of a *Brassica napus* transporter which is able to transport nitrate and histidine.** *J Biol Chem* 1998, **273**:12017–12023.
- Tong YP, Zhou JJ, Li ZS, Miller AJ: **A two-component high-affinity nitrate uptake system in barley.** *Plant J* 2005, **10**:1365–1374.
- Kotur Z, Mackenzie N, Ramesh S, Tyerman SD, Kaiser BN, Glass AD: **Nitrate transport capacity of the *Arabidopsis thaliana* NRT2 family members and their interactions with AtNAR2.1.** *New Phytol* 2012, **194**:724–731.
- Schachtman DP, Schroeder JI, Lucas WJ, Anderson JA, Gaber RF: **Expression of an inward-rectifying potassium channel by the *Arabidopsis* KAT1 Cdna.** *Science* 1992, **258**:1654–1658.
- Chilcott TC, Shartzner SF, Iverson MW, Garvin DF, Kochian LV, Lucas WJ: **Potassium transport kinetics of KAT1 expressed in *Xenopus* oocytes: a proposed molecular structure and field effect mechanism for membrane transport.** *CR Acad Sci Paris* 1995, **318**:761–771.
- Véry A, Gaymard F, Bosseux C, Sentenac H, Thibaud J: **Expression analysis of a cloned plant K⁺ channel in *Xenopus* oocytes analysis of macroscopic currents.** *Plant J* 1995, **7**:321–332.
- Wang Y, Wu WH: **Potassium transport and signaling in higher plants.** *Annu Rev of Plant Biol* 2013, **64**:451–476.
- Daniels MJ, Mirkov TE, Chrispeels MJ: **The plasma membrane of *Arabidopsis thaliana* contains a mercury insensitive aquaporin that is a homolog of the tonoplast water channel protein TIP.** *Plant Physiol* 1994, **106**:1325–1333.
- Johansson I, Karlsson M, Shukla VK, Chrispeels MJ, Larsson C, Kjellbom P: **Water transport activity of the plasma membrane aquaporin PM28A is regulated by phosphorylation.** *Plant Cell* 1998, **10**:451–459.
- Higuchi T, Suga S, Tsuchiya T, Hisada H, Morishima S, Okada Y, Maeshim M: **a, Molecular cloning, water channel activity and tissue specific expression of two isoforms of the radish vacuolar aquaporin.** *Plant Cell Physiol* 1998, **39**:905–913.
- Hu W, Yuan QQ, Wang Y, Cai R, Deng XM, Wang J, Zhou SY, Chen LH, Huang C, Ma ZB, Yang GX, He GY: **Overexpression of a wheat aquaporin gene, TaAQP8, enhances salt stress tolerance in transgenic tobacco.** *Plant and Cell Physiol* 2012, **53**:2127–2141.
- Liu KH, Huang CY, Tsay YF: **CHL1 is a dual-affinity nitrate transporter of *Arabidopsis* involved in multiple phases of nitrate uptake.** *Plant Cell* 1999, **11**:865–874.
- Lin CM, Koh S, Stacey G, Yu SM, Lin TY, Tsay YF: **Cloning and functional characterization of a constitutively expressed nitrate transporter gene, OsNRT1, from rice.** *Plant Physiol* 2000, **122**:379–388.
- Chiu CC, Lin CS, Hsia AP, Su RC, Lin HL, Tsay YF: **Mutation of a nitrate transporter, AtNRT1.4, results in a reduced petiole nitrate content and altered leaf development.** *Plant Cell Physiol* 2004, **45**:1139–1148.
- Almagro A, Lin SH, Tsay YF: **Characterization of the *Arabidopsis* nitrate transporter NRT1.6 reveals a role of nitrate in early embryo development.** *Plant Cell* 2008, **20**:3289–3299.
- Fan SC, Lin CS, Hsu PK, Lin SH, Tsay YF: **The *Arabidopsis* nitrate transporter NRT1.7, expressed in phloem, is responsible for source-to-sink remobilization of nitrate.** *Plant Cell* 2009, **21**:2750–2761.
- Li JY, Fu YL, Pike SM, Bao J, Tian W, Zhang Y, Li HM, Huang J, Li LG, Schroeder JI, Gassmann W, Gong JM: **The *Arabidopsis* nitrate transporter NRT1.8 functions in nitrate removal from the xylem sap and mediates cadmium tolerance.** *Plant Cell* 2010, **22**:1633–1646.
- Wang YY, Tsay YF: ***Arabidopsis* nitrate transporter NRT1.9 is important in phloem nitrate transport.** *Plant Cell* 2011, **23**:1945–1957.
- Zhou JJ, Fernandez E, Galvan A, Miller AJ: **A high affinity nitrate transport system from *Chlamydomonas* requires two gene products.** *FEBS Lett* 2000, **466**:225–227.
- Feng HM, Yan M, Fan XR, Li BZ, Shen QR, Miller AJ, Xu GH: **Spatial expression and regulation of rice high-affinity nitrate transporters by nitrogen and carbon status.** *J Exp Bot* 2011, **62**:2319–2332.
- Yan M, Fan XR, Feng HM, Miller AJ, Sheng QR, Xu GH: **Rice OsNAR2.1 interacts with OsNRT2.1, OsNRT2.2 and OsNRT2.3a nitrate transporters to provide uptake over high and low concentration ranges.** *Plant Cell Environ* 2011, **34**:1360–1372.
- Burgess-Brown NA, Sharma S, Sobott F, Loenarz C, Oppermann U, Gileadi O: **Codon optimization can improve expression of human genes in *Escherichia coli*: A multi-gene study.** *Protein Expr Purif* 2008, **59**:94–102.
- Maertens B, Spriestersbach A, von Groll U, Roth U, Kubicek J, Gerrits M, Graf M, Liss M, Daubert D, Wagner R, Schäfer F: **Gene optimization mechanisms: a multi-gene study reveals a high success rate of full-length human proteins expressed in *Escherichia coli*.** *Protein Sci* 2010, **19**:1312–1326.
- Tu YB, Wang YQ, Wang G, Wu J, Liu YG, Wang SJ, Jiang CG, Cai XH: **High-level expression and immunogenicity of a porcine circovirus type 2 capsid protein through codon optimization in *Pichia pastoris*.** *Appl Microbiol and Biotechnol* 2013, **97**:2867–2875.
- Wurm FM: **Production of recombinant protein therapeutics in cultivated mammalian cells.** *Nat Biotechnol* 2004, **22**:1393–1398.
- Fath S, Bauer A, Liss M, Spriestersbach A, Maertens B, Hahn P, Ludwig C, Schäfer F, Graf M, Wagner R: **Multiparameter RNA and codon optimization: a standardized tool to assess and enhance autologous mammalian gene expression.** *PLoS One* 2011, **6**:e17596.
- Miller AJ, Zhou J-J: ***Xenopus* oocytes as an expression system for plant transporters.** *Biochim Biophys Acta* 2000, **1465**:343–358.
- Musto H, Cruveiller S, D'Onofrio G, Romero H, Bernardi G: **Translational selection on codon usage in *Xenopus laevis*.** *Mol Biol Evol* 2001, **18**:1703–1707.
- Amin NM, Tandon P, Osborne Nishimura E, Conlon FL: **RNA-seq in the tetraploid *Xenopus laevis* enables genome-wide insight in a classic developmental biology model organism.** *Methods* 2013. (in press) <http://dx.doi.org/10.1016/j.jymeth.2013.06.009>.
- Carels N, Bernardi G: **Two classes of genes in plants.** *Genetics* 2000, **154**:1819–1825.
- Kudla G, Lipinski L, Caffin F, Helwak A, Zyllic M: **High guanine and cytosine content increases mRNA levels in mammalian cells.** *PLOS Biology* 2006, **4**(e180):0933–0942.
- Gustafsson C, Govindarajan S, Minshull J: **Codon bias and heterologous protein expression.** *Trends Biotechnol* 2004, **22**:346–353.
- Cleaver O, Patterson KD, Krieg PA: **Overexpression of the tinman-related genes XNkx-2.5 and XNkx-2.3 in *Xenopus* embryos results in myocardial hyperplasia.** *Development* 1996, **122**:3549–3556.

doi:10.1186/1746-4811-9-48

Cite this article as: Feng et al.: Optimizing plant transporter expression in *Xenopus* oocytes. *Plant Methods* 2013 **9**:48.